

ELLIPTIC CURVES WITH TORSION GROUP $\mathbb{Z}/8\mathbb{Z}$ OR $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$

ANDREJ DUJELLA AND JUAN CARLOS PERAL

ABSTRACT. We show the existence of families of elliptic curves over \mathbb{Q} whose generic rank is at least 2 for the torsion groups $\mathbb{Z}/8\mathbb{Z}$ and $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$. Also in both cases we prove the existence of infinitely many elliptic curves, which are parameterized by the points of an elliptic curve with positive rank, with such torsion group and rank at least 3.

These results represent an improvement of previous results by Campbell, Kulesz, Lecacheux, Dujella and Rabarison where families with rank at least 1 were constructed in both cases.

1. INTRODUCTION

The construction of families of elliptic curves having high rank often is based in two basic strategies as mentioned by Elkies in [El].

a) The Neron method studies the pencil of cubics passing through a set of nine rational random points and then looks for independence. See [Sh] for a good description of the method. Families of rank up to 10 were constructed in this way.

b) The Mestre method uses polynomial identities forcing the existence of rational points in the curve and then searches for independence conditions. In this way Mestre was able to construct a rank 11 curve over $\mathbb{Q}(u)$, see [Me].

In our case we want that the curve has a predetermined torsion group, $\mathbb{Z}/8\mathbb{Z}$ or $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$, so we should start from the general model for such torsion and then try to impose the existence of new points. One way to do this is looking for good quadratic sections. Also the method developed by Lecacheux [Le1], [Le2], can be used in these cases. In her method Lecacheux uses fibrations of the corresponding surfaces, (as the ones given explicitly in Beauville [Be], in Bertin and Lecacheux [BL] or in Livne and Yui [LY]) in order to find elliptic curves with positive rank over $\mathbb{Q}(u)$. Another useful tool in order to get high rank curves over $\mathbb{Q}(u)$ are the diophantine triples, in fact for the torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ we construct a rank 4 family, the current record for this torsion group, see [DP].

In the case of torsion group $\mathbb{Z}/8\mathbb{Z}$ families having rank at least 1 over $\mathbb{Q}(u)$ has been found by several authors, see [Ku], [Le1], [Le2] and [Ra]. In this paper we prove the existence of two elliptic curves having this torsion group and rank at least 2 over $\mathbb{Q}(u)$, also the existence of infinitely many elliptic curves over \mathbb{Q} with this torsion group and rank at least 3, parametrized by the points of an elliptic curve with positive rank.

Elliptic curves with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ and rank at least 1 over $\mathbb{Q}(u)$ have been constructed by several authors, see [Ca], [Ku], [Le1], [Du1] and [Ra]. Here we prove the existence of three elliptic curves with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ and whose rank over $\mathbb{Q}(u)$ is at least 2. We prove also the existence of infinitely many elliptic curves with this torsion group and rank at least 3 over \mathbb{Q} , parametrized by the points of an elliptic curve of positive rank.

2010 *Mathematics Subject Classification.* 11G05.

A. D. was supported by the Ministry of Science, Education and Sports, Republic of Croatia, grant 037-0372781-2821. J. C. P. was supported by the UPV/EHU grant EHU 10/05.

We describe first the model for the elliptic curves having each of these torsion groups. Then, in both cases, we show the existence of several families having rank at least 1 over $\mathbb{Q}(u)$. We finally show the existence of two families in the $\mathbb{Z}/8\mathbb{Z}$ case and three families in the case of torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ having rank at least 2 over $\mathbb{Q}(u)$. In all the cases we describe the coefficients of the families and the coordinates of independent points.

We also exhibit some examples of curves with high rank. The current records both for families and for curves can be found in [Du2]. One way to find high rank elliptic curves over \mathbb{Q} is the construction of elliptic curves with positive rank over $\mathbb{Q}(u)$, as high as possible, and then searching for good specializations with adequate sieving tools such as the Mestre-Nagao sums for example.

2. TORSION GROUP

2.1. Curves with torsion group $\mathbb{Z}/8\mathbb{Z}$. The Tate's normal form for an elliptic curve is given by

$$E(b, c) : y^2 + (1 - c)xy - by = x^3 - bx^2$$

(see [Kn]). It is nonsingular if and only if $b \neq 0$. Using the addition law for $P = (0, 0)$ and taking $d = b/c$ we have

$$\begin{aligned} 4P &= (d(d-1), d^2(c-d+1)), \\ -4P &= (d(d-1), d(d-1)^2) \end{aligned}$$

so P is a torsion point of order 8 for b and c as follows

$$\begin{aligned} b &= (2v-1)(v-1), \\ c &= \frac{(2v-1)(v-1)}{v} \end{aligned}$$

with v a rational, see [Kn]. For these values of b and c we can write the curve in the form $y^2 = x^3 + A_8(v)x^2 + B_8(v)x$ where

$$\begin{aligned} A_8(v) &= 1 - 8v + 16v^2 - 16v^3 + 8v^4, \\ B_8(v) &= 16(-1+v)^4v^4. \end{aligned}$$

Writing the curve in this form is a convenient way to search for candidates for new rational points. In fact their x -coordinates should be either divisors of B or rational squares times divisors of B .

2.2. Curves with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$. Using again the Tate's normal form and the addition law for the point $P = (0, 0)$ we find

$$\begin{aligned} 3P &= (c, b-c), \\ -3P &= (c, c^2). \end{aligned}$$

It follows that P will be a torsion point of order 6 for $b = c + c^2$. For this value of b we write the curve in the form $y^2 = x^3 + A_6(c)x^2 + B_6(c)x$. We get

$$\begin{aligned} A_6(c) &= -1 + 6c - 3c^2, \\ B_6(c) &= -16c^3. \end{aligned}$$

In the new coordinates the torsion point of order 6 becomes $(-4c, 4c(1+c))$. Now we use the fact that the curves with torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ have a model $y^2 = x(x-m)(x-n)$. So in order to get a curve with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ it is enough that in the family above the polynomial

$$x^3 + A_6(c)x^2 + B_6(c)x = x(-16c^3 + x + 6cx - 3c^2x + x^2)$$

factorizes into linear factors. So the discriminant $\Delta = (1+c)^3(1+9c)$ of the second order polynomial must be a square, hence

$$c = \frac{-v^2 + 1}{2(3v - 5)}.$$

For these values of c the corresponding curves have torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ and can be written as $y^2 = x^3 + A_{26}(v)x^2 + B_{26}(v)x$ where

$$\begin{aligned} A_{26}(v) &= 37 - 84v + 102v^2 - 36v^3 - 3v^4, \\ B_{26}(v) &= 32(-1+v)^3(1+v)^3(-5+3v). \end{aligned}$$

The torsion point of order 6 transforms into

$$(8(-1+v)(1+v)(-5+3v), 8(-3+v)^2(-1+v)(1+v)(-5+3v)).$$

Remark. The cases treated here, jointly with the general curve having torsion $\mathbb{Z}/7\mathbb{Z}$ and whose model is

$$y^2 = x^3 + x^2(1 - 2t + 3t^2 + 6t^3 + t^4) + x(-8t^2(1+t)(-1+t+t^2)) + 16t^4(1+t)^2,$$

are the three cases in which the general model for such torsion group is a $K3$ surface.

3. RANK 1 FAMILIES

3.1. The case of torsion group $\mathbb{Z}/8\mathbb{Z}$. For this torsion group we show ten conditions upon v leading to rank 1 families. Some of them were already known due to the authors quoted before. We first list eighth values of x_i which becomes the x -coordinate of a new point once we specialize to the corresponding values of v_i , $i = 1 \dots, 8$. We include another two values, v_9 and v_{10} , found by Lecacheux by using adequate fibration of the general model with torsion $\mathbb{Z}/8\mathbb{Z}$.

$$\begin{aligned} x_1 &= \frac{-16v^4(1-4v+2v^2)}{(-1+4w)^2}, & v_1 &= \frac{1+w^2}{3-2w+w^2}, \\ x_2 &= \frac{-(-1+v)^4(-5+8v)(-5+18v)}{4(-2+3v)^2}, & v_2 &= \frac{5(1+w^2)}{2(9+4w^2)}, \\ x_3 &= \frac{-4(-3+v)(-1+v)^2v^4(-1+3v)}{(1-4v+2v^2)^2}, & v_3 &= \frac{1+3w^2}{3+w^2}, \\ x_4 &= 16(-1+v)^2v^2(1-2v+2v^2), & v_4 &= \frac{(-2+w)w}{-2+w^2}, \\ x_5 &= \frac{-64(-1+v)^2v^2(-1-v+v^2)}{(-1-4v+4v^2)^2}, & v_5 &= \frac{(-2+w)w}{1+w^2}, \\ x_6 &= -(-1+v)^2(1-6v+4v^2), & v_6 &= \frac{2-2w+w^2}{4+w^2}, \\ x_7 &= 4v^4, & v_7 &= \frac{-5+w^2}{4(1+w)}, \\ x_8 &= \frac{-(-1+v)^2(-5+2v)^2(25-70v+36v^2)}{(-7+6v)^2}, & v_8 &= \frac{34-6w+w^2}{36+w^2}. \end{aligned}$$

$$v_9 = \frac{w^2 + 12}{2(w^2 + 4)}, \quad v_{10} = \frac{-2w}{1 - w + w^2}.$$

Details of one of this cases is given in the next section. In every case the new point is of infinite order so the rank of the corresponding curve is at least 1 over $\mathbb{Q}(w)$.

3.2. The case of torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$. We have found several conditions upon v leading to rank 1 families. In the next table we list nine values of x_i which becomes the x -coordinate of a new point once we specialize to the value of $v_i, i = 1 \dots, 9$. As before some of them were already known due to the authors quoted above.

$$\begin{array}{ll}
x_1 = 8(-1+v)^3(1+v), & v_1 = \frac{3(-1+w)(1+w)}{-29-8w+w^2}, \\
x_2 = 4(1+v)^3, & v_2 = \frac{3(-3+w)(3+w)}{-45-24w+w^2}, \\
x_3 = 2(-1+v)(1+v)^2(-5+3v), & v_3 = \frac{-7+w^2}{1-4w+w^2}, \\
x_4 = -16(-1+v)^2(1+v), & v_4 = \frac{-11+w^2}{5-4w+w^2}, \\
x_5 = 16(-5+3v)(3v-7)^2, & v_5 = \frac{3(261+w^2)}{153-24w+w^2}, \\
x_6 = 16(1+v)(v-5)^2, & v_6 = \frac{135-w^2}{141+24w+w^2}, \\
x_7 = \frac{4(-1+v)^2(1+v)^2(41-54v+49v^2)}{(-1+3v)^2}, & v_7 = \frac{41-w^2}{2(27+7w)}, \\
x_8 = (-5+3v)(3v-1)^2, & v_8 = \frac{3}{5-w^2}, \\
x_9 = \frac{2(v-1)(v+1)^3(3v-1)^2}{(2v+2)^2}, & v_9 = \frac{-7-2w^2}{3(-3+2w^2)}.
\end{array}$$

More details are given in another section. In every case the new point is of infinite order so the rank of the curve is at least 1 over $\mathbb{Q}(w)$.

4. RANK 2 FAMILIES FOR THE TORSION GROUP $\mathbb{Z}/8\mathbb{Z}$

4.1. A family with rank 1. We present here some details for the family in which we have found two subfamilies with torsion $\mathbb{Z}/8\mathbb{Z}$ and generic rank at least 2. It corresponds to the third entry in the table of rank 1 families above and it is a reparametrization of one of the families in [Le1]. By inserting in the general family $y^2 = x^3 + A_8(v)x^2 + B_8(v)x$ the value $v = v_3(w)$ we get the rank 1 family given by $y^2 = x^3 + AA_8(w)x^2 + BB_8(w)x$ where

$$\begin{aligned}
AA_8(w) &= -31 - 148w^2 + 214w^4 - 116w^6 + 337w^8, \\
BB_8(w) &= 256(-1+w)^4(1+w)^4(1+3w^2)^4.
\end{aligned}$$

By searching on several homogeneous spaces of the associate curve we have found the possibility of imposing two new conditions which lead to new points. The values x_1 and x_2 jointly with the specialization of the parameter are

$$\begin{aligned}
x_1 &= \frac{(-1+w)^2(1+w)^2(5+7w^2)^2(11+25w^2)}{16}, & w_1 &= \frac{11-u^2}{10u}, \\
x_2 &= \frac{(-1+w)^2(1+w)^2(1+11w^2)^2(7+29w^2)}{16w^2}, & w_2 &= \frac{29-12u+u^2}{-29+u^2}.
\end{aligned}$$

With these specializations we get two different families of rank at least 2 over $\mathbb{Q}(u)$.

4.2. First family with rank 2 and torsion group $\mathbb{Z}/8\mathbb{Z}$. Once we insert w_1 into the coefficients AA_8, BB_8 we get as new coefficients AAA_8, BBB_8 given by

$$\begin{aligned} AAA_8 &= 337u^{16} - 41256u^{14} + 4047356u^{12} - 288332632u^{10} + 2363813190u^8 \\ &\quad - 34888248472u^6 + 59257339196u^4 - 73087520616u^2 + 72238942897, \\ BBB_8 &= 256(363 + 34u^2 + 3u^4)^4(11 + u)^4(-11 + u)^4(-1 + u)^4(1 + u)^4. \end{aligned}$$

The x -coordinates of two independent infinite order points are

$$\begin{aligned} X_1 &= \frac{2^{12}5^2(-11 + u)^2(-1 + u)^2u^2(1 + u)^2(11 + u)^2(-11 + u^2)^2(363 + 34u^2 + 3u^4)^4}{(102487 - 303468u^2 + 43482u^4 - 2508u^6 + 7u^8)^2}, \\ X_2 &= \frac{(-11 + u)^2(-1 + u)^2(1 + u)^2(11 + u)^2(11 + u^2)^2(847 + 346u^2 + 7u^4)^2}{64u^2}. \end{aligned}$$

The x -coordinates of the torsion point of order 8 is:

$$T_1 = -8(-11 + u)(-1 + u)(1 + u)(11 + u)(363 + 34u^2 + 3u^4)^3.$$

That the rank of this curve is at least 2 over $\mathbb{Q}(u)$ can be proved using a specialization argument, since the specialization map is a homomorphism.

4.3. Second family with rank 2 and torsion group $\mathbb{Z}/8\mathbb{Z}$. Once we insert w_2 into the coefficients AA_8, BB_8 we get as new coefficients aaa_8, bbb_8 given by

$$\begin{aligned} aaa_8 &= 500246412961 - 2069985157080u + 3162080774436u^2 - 2895517882032u^3 + \\ &\quad 1873181389706u^4 - 906769167048u^5 + 333391978480u^6 - 93284915496u^7 + \\ &\quad 19860033555u^8 - 3216721224u^9 + 396423280u^{10} - 37179432u^{11} + \\ &\quad 2648426u^{12} - 141168u^{13} + 5316u^{14} - 120u^{15} + u^{16}, \\ bbb_8 &= 256(-6 + u)^4u^4(-29 + 6u)^4(841 - 522u + 137u^2 - 18u^3 + u^4)^4. \end{aligned}$$

The x -coordinates of two independent infinite order points are

$$\begin{aligned} &\frac{64(-6 + u)^2u^2(-29 + 6u)^2(-29 + u^2)^2(29 - 12u + u^2)^2(841 - 522u + 137u^2 - 18u^3 + u^4)^4}{(707281 - 292668u - 200158u^2 + 168432u^3 - 46685u^4 + 5808u^5 - 238u^6 - 12u^7 + u^8)^2}, \\ &\frac{(-6 + u)^2u^2(-29 + 6u)^2(87 - 29u + 3u^2)^2(2523 - 1914u + 541u^2 - 66u^3 + 3u^4)^2}{4(29 - 12u + u^2)^2}. \end{aligned}$$

The x -coordinate of the torsion point of order 8 is:

$$T_2 = 8(-6 + u)u(-29 + 6u)(841 - 522u + 137u^2 - 18u^3 + u^4)^3.$$

Remark. When we use $w_3 = \frac{(-3+u)(3+u)}{7-6u}$ and $w_4 = \frac{(-3+u)(3+u)}{11-6u}$ in the family of rank at least 1 corresponding to v_{10} we get two families of rank at least 2. They are a reparametrization of the two families above. So we have the following fact, when we specialize in the general family with torsion group $\mathbb{Z}/8\mathbb{Z}$ to $v_3 = \frac{1+3w^2}{3+w^2}$ and to $v_{10} = \frac{-2w}{1-w+w^2}$ we get two different families having rank at least 1. When we use in the first the values $w_1 = \frac{11-u^2}{10u}$ and $w_2 = \frac{29-12u+u^2}{-29+u^2}$ we get two families of

rank at least 2 that are a reparametrization of the rank 2 families that we get by using v_{10} followed by $w_3 = \frac{(-3+u)(3+u)}{7-6u}$ and $w_4 = \frac{(-3+u)(3+u)}{11-6u}$.

So at the end with the changes v_3 follow by w_1 and w_2 and v_{10} follow by w_3 and w_4 we reach the same families of rank at least 2 even though the families of rank at least 1 of the intermediate step are different.

4.4. Rank 3 for the torsion group $\mathbb{Z}/8\mathbb{Z}$. It can be proved that there exist infinitely many elliptic curves with torsion group $\mathbb{Z}/8\mathbb{Z}$ parametrized by the points of a positive rank elliptic curve. In fact it is enough to see that the equation $w_1(r) = w_2(s)$, i.e.:

$$\frac{11 - r^2}{10r} = \frac{29 - 12s + s^2}{-29 + s^2}$$

has infinitely many solutions. This is the same as to solve

$$319 + 290r - 29r^2 - 120rs - 11s^2 + 10rs^2 + r^2s^2 = 0$$

in rational terms, so the discriminant $\Delta = 3509 + 62r^2 + 29r^4$ has to be a square. But $t^2 = 3509 + 62r^2 + 29r^4$ has a solution, $(r, t) = (1, 60)$ for example, hence it is equivalent to the cubic $y^2 = x^3 - 463x^2 + 45936x$ whose rank is 2 as proved with `mwrank` [Cr]. This, jointly with the independence of the corresponding points, implies the existence of infinitely many solutions parametrized by the points of the elliptic curve, see [Le1] or [Ra] for this kind of argument.

5. RANK 2 FAMILIES FOR THE TORSION GROUP $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$

5.1. A family with rank 1. We present here some details for the family of rank 1 in which we have found two subfamilies of generic rank at least 2. It corresponds to the eighth entry in the table above. We have found the possibility of converting $x_8 = (-5 + 3v)(3v - 1)^2$ in the x -coordinate of a new point by considering the homogenous space $(U, V) = (3v - 1, 1)$ of the initial family

$$y^2 = x^3 + (37 - 84v + 102v^2 - 36v^3 - 3v^4)x^2 + (32(-1 + v)^3(1 + v)^3(-5 + 3v))x.$$

The condition that has to be fulfilled is that $v(-3 + 5v)$ converts into a square, hence we get $v_8 = \frac{3}{5-w^2}$. Once we insert v_8 in the preceding family and take off denominators we have the family $y^2 = x^3 + AA_{26}(w)x^2 + BB_{26}(w)x$ where

$$\begin{aligned} AA_{26}(w) &= 9472 - 7808w^2 + 2688w^4 - 488w^6 + 37w^8, \\ BB_{26}(w) &= 32(-8 + w^2)^3(-5 + w^2)(-2 + w^2)^3(-16 + 5w^2). \end{aligned}$$

The point of infinite order is

$$\begin{aligned} P = & \left(-(-5 + w^2)(4 + w^2)^2(-16 + 5w^2), \right. \\ & \left. 27(-2 + w)^2w(2 + w)^2(-5 + w^2)(4 + w^2)(-16 + 5w^2) \right) \end{aligned}$$

and the torsion point of order 6 is

$$\begin{aligned} T = & \left(8(-8 + w^2)(-5 + w^2)(-2 + w^2)(-16 + 5w^2), \right. \\ & \left. 72(-2 + w)^2(2 + w)^2(-8 + w^2)(-5 + w^2)(-2 + w^2)(-16 + 5w^2) \right) \end{aligned}$$

By searching on several homogeneous spaces we have found the possibility of imposing two new conditions which lead to new points, hence, in case of independence, to a couple of rank 2 families. The values x_1 and x_2 jointly with the specialization of the parameter are

$$\begin{aligned} x_1 &= 2(-8 + w^2)^2(-2 + w^2)^3, & w_1 &= \frac{2(7 + u^2)}{-7 - 2u + u^2}, \\ x_2 &= (-8 + w^2)(-5 + w^2)(-16 + 5w^2)w^4, & w_2 &= \frac{5 - 2u + u^2}{-5 + u^2}. \end{aligned}$$

5.2. First family with rank 2. In order to force $2(-8 + w^2)^2(-2 + w^2)^3$ to be the x -coordinate of a new point it is enough to solve $2w^2 - 7 = M^2$. This is achieved with $w_1 = \frac{2(7+u^2)}{-7-2u+u^2}$, and the corresponding family is $y^2 = x^3 + AAA_{26}(u)x^2 + BBB_{26}(u)x$ where

$$\begin{aligned} AAA_{26}(u) &= -2(5764801 + 6588344u - 21647416u^2 + 29445864u^3 - 9604u^4 + \\ &\quad 27969592u^5 - 44631944u^6 + 9779112u^7 + 5909830u^8 - 1397016u^9 \\ &\quad - 910856u^{10} - 81544u^{11} - 4u^{12} - 1752u^{13} - 184u^{14} - 8u^{15} + u^{16}), \\ BBB_{26}(u) &= (-7 - 10u + u^2)^3(-7 + 2u + u^2)^3(49 + 140u - 106u^2 - 20u^3 + u^4) \\ &\quad (49 - 28u + 38u^2 + 4u^3 + u^4)^3(49 - 112u + 110u^2 + 16u^3 + u^4). \end{aligned}$$

The x -coordinates of the two infinite order points are

$$\begin{aligned} X_1 &= (49 + 140u - 106u^2 - 20u^3 + u^4)(49 + 14u + 2u^2 - 2u^3 + u^4)^2 \times \\ &\quad (49 - 112u + 110u^2 + 16u^3 + u^4), \\ X_2 &= \frac{(-7 - 10u + u^2)^2(-7 + 2u + u^2)^2(49 - 28u + 38u^2 + 4u^3 + u^4)^3}{(-7 - 2u + u^2)^2}, \end{aligned}$$

and the x -coordinate of the torsion point of order 6 is

$$\begin{aligned} T &= (-7 - 10u + u^2)(-7 + 2u + u^2)(49 + 140u - 106u^2 - 20u^3 + u^4) \times \\ &\quad (49 - 28u + 38u^2 + 4u^3 + u^4)(49 - 112u + 110u^2 + 16u^3 + u^4). \end{aligned}$$

The specialization for $u = 2$ gives an elliptic curve with rank 2 and the specialized points are independent so the fact that specialization is a homomorphism implies that the rank of the curve $y^2 = x^3 + AAA_{26}(u)x^2 + BBB_{26}(u)x$ is at least 2 over $\mathbb{Q}(u)$.

5.3. Second family with rank 2. Imposing $(-8 + w^2)(-5 + w^2)(-16 + 5w^2)w^4$ as the x -coordinate of a new point it is equivalent to solve $5w^2 - 4 = M^2$. This is achieved with $w_2 = \frac{5-2u+u^2}{-5+u^2}$.

The corresponding family is $y^2 = x^3 + aaa_{26}(u)x^2 + bbb_{26}(u)x$ where

$$\begin{aligned} aaa_{26}(u) &= 1523828125 + 1171250000u - 3482125000u^2 - 1970850000u^3 + \\ &\quad 3530367500u^4 + 1221154000u^5 - 2018502200u^6 - 238418640u^7 + \\ &\quad 632792782u^8 - 47683728u^9 - 80740088u^{10} + 9769232u^{11} + \\ &\quad 5648588u^{12} - 630672u^{13} - 222856u^{14} + 14992u^{15} + 3901u^{16}, \\ bbb_{26}(u) &= 128(-7 + 2u + u^2)^3(-25 - 10u + 7u^2)^3(25 + 5u - 16u^2 + u^3 + u^4) \times \\ &\quad (25 + 20u - 34u^2 + 4u^3 + u^4)^3(275 + 100u - 230u^2 + 20u^3 + 11u^4) \end{aligned}$$

The x -coordinates of the two infinite order points are

$$\begin{aligned} X_1 &= -4(25 + 5u - 16u^2 + u^3 + u^4)(125 - 20u - 26u^2 - 4u^3 + 5u^4)^2 \times \\ &\quad (275 + 100u - 230u^2 + 20u^3 + 11u^4), \\ X_2 &= \frac{-4(5 - 2u + u^2)^4(-7 + 2u + u^2)(-25 - 10u + 7u^2)}{(-5 + u^2)^2} \times \\ &\quad (25 + 5u - 16u^2 + u^3 + u^4)(275 + 100u - 230u^2 + 20u^3 + 11u^4), \end{aligned}$$

and the x -coordinate of the torsion point of order 6 is

$$\begin{aligned} T &= 32(-7 + 2u + u^2)(-25 - 10u + 7u^2)(25 + 5u - 16u^2 + u^3 + u^4) \times \\ &\quad (25 + 20u - 34u^2 + 4u^3 + u^4)(275 + 100u - 230u^2 + 20u^3 + 11u^4). \end{aligned}$$

5.4. Third family with rank 2. A variant of the model for torsion group $\mathbb{Z}/6\mathbb{Z}$ by Hadano (see [H]) will be used for the construction of another family of curves with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ and rank at least 2. The curves with torsion group $\mathbb{Z}/6\mathbb{Z}$ in [H] have the equation

$$Y^2 = X^3 + X^2(a^2 + 2ab - 2b^2) - X(2a - b)b^3.$$

For this model we first force the existence of a new point in order to have a family with this torsion and rank at least 1, then we choose the parameters that give complete factorization, and so the torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$, and in the resulting family for this torsion and rank 1 we produce a second independent point with a quadratic section.

We observe that with $b = -\frac{(-1+a^2-v)(-1+a^2+v)}{4(-a+a^3-v)}$ we have a point in the curve with x coordinate given by

$$X = -\frac{(1 - 4a + 3a^2 - v)(-1 + a^2 + v)^3}{16(-a + a^3 - v)^2}$$

Once we clear denominators this family can be written as $Y^2 = X^3 + A_{61}(a, v)X^2 + B_{61}(a, v)X$ where

$$\begin{aligned} A_{61} &= 2(-1 + 8a^2 - 10a^4 + 3a^8 + 4av + 8a^3v - 12a^5v + 2v^2 + 6a^4v^2 - 4av^3 - v^4), \\ B_{61} &= (-1 + a^2 - v)^3(1 - 4a + 3a^2 - v)(-1 + a^2 + v)^3(1 + 4a + 3a^2 + v). \end{aligned}$$

The new point becomes

$$P = (-(1 - 4a + 3a^2 - v)(-1 + a^2 + v)^3, 4(1 - 4a + 3a^2 - v)(-a + a^3 - v)(-1 + a^2 + v)^3).$$

For $a \neq 1$, in general, P is of infinite order so the curve has rank at least 1 over $\mathbb{Q}(a, v)$. Now we observe that the complete factorization of the cubic is equivalent to forcing the discriminant of a second degree polynomial to be a square.

The condition is $a(-a + a^3 - v)(-1 + a^2 - av + v^2) = \text{square}$. This can be achieved with $a = v + 1$ followed by $v = \frac{1-w^2}{-3+2w}$. Once we perform these changes we get $Y^2 = X^3 + a_{26}(w)X^2 + b_{26}(w)X$ where

$$\begin{aligned} a_{26} &= 2(-24 - 216w + 1008w^2 - 1596w^3 + 1319w^4 - 648w^5 + 198w^6 - 36w^7 + 3w^8), \\ b_{26} &= (-4 + w)^3(-3 + w)(-2 + w)^3(-1 + w)^3w(1 + w)^3(-7 + 3w)(-2 + 3w). \end{aligned}$$

With these changes the x -coordinate of the infinite order point is

$$X = -\frac{(-4 + w)^3(-2 + w)^3(-1 + w)^2w(1 + w)^2(-2 + 3w)}{(2 - 2w + w^2)^2}$$

So we have a curve with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ and rank at least 1 over $\mathbb{Q}(w)$.

Now we see that $(-4 + w)(-3 + w)(-1 + w)^3(1 + w)^2(-7 + 3w)$ will be a new point on the curve if we force $4 - 9w + 3w^2$ to be a square. This is the same as

choosing $w = -\frac{9+4u}{-3+u^2}$. Once we perform this change and clear denominators we get the following coefficients for the new family

$$\begin{aligned} A_{263} &= -2(157464 - 1889568u - 13594392u^2 - 38047968u^3 - \\ &\quad 62500248u^4 - 69622416u^5 - 57719412u^6 - 38941344u^7 - \\ &\quad 23353995u^8 - 12980448u^9 - 6413268u^{10} - 2578608u^{11} - \\ &\quad 771608u^{12} - 156576u^{13} - 18648u^{14} - 864u^{15} + 24u^{16}), \\ B_{263} &= -(-6+u)^3u(2+u)^3(-1+2u)^3(3+2u)^3(4+3u)(9+4u) \\ &\quad (6+4u+u^2)^3(3+4u+2u^2)^3(21+12u+2u^2)(6+12u+7u^2). \end{aligned}$$

The x -coordinates of the non-torsion points are

$$\begin{aligned} X_1 &= (-6+u)^2u(2+u)^2(-1+2u)(3+2u)(4+3u) \times \\ &\quad (6+4u+u^2)^3(6+12u+7u^2), \\ X_2 &= -\frac{(-6+u)^2(2+u)^2(-1+2u)^3(3+2u)^3(9+4u)}{(45+48u+22u^2+8u^3+2u^4)^2} \times \\ &\quad (6+4u+u^2)^2(3+4u+2u^2)^3(21+12u+2u^2). \end{aligned}$$

A specialization argument, as in Section 5.2, shows that these two points are independent so this curve has rank at least 2 over $\mathbb{Q}(u)$ and torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$.

5.5. Rank 3 for the torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$. Now we prove the existence of infinitely many elliptic curves with rank at least 3 and torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ parametrized by the points of a positive rank elliptic curve. In fact it is enough to see that the equation $w_1(r) = w_2(s)$, i.e.:

$$\frac{2(7+r^2)}{-7-2r+r^2} = \frac{5-2s+s^2}{-5+s^2}$$

has infinitely many solutions. This is the same as solving

$$-35 + 10r - 15r^2 - 14s - 4rs + 2r^2s + 21s^2 + 2rs^2 + r^2s^2 = 0$$

in rational terms, so the discriminant $\Delta = 49 - 7r + 20r^2 + r^3 + r^4$ has to be a rational square. But $t^2 = 49 - 7r + 20r^2 + r^3 + r^4$ has a solution, $(r, t) = (1, 8)$ for example, hence it is birationally equivalent to the cubic $y^2 = x^3 - 43x^2 + 280x$ whose rank is 1 as proved with `mwrnk` [Cr]. This and the independence of the corresponding points, implies the existence of infinitely many solutions parametrized by the points of the elliptic curve.

6. EXAMPLES OF CURVES WITH HIGH RANK

6.1. The case of torsion group $\mathbb{Z}/8\mathbb{Z}$. The highest known rank of an elliptic curve over \mathbb{Q} with torsion group $\mathbb{Z}/8\mathbb{Z}$ is rank 6 curve found by Elkies in 2006. See [Du2] for the details of this curve.

The following list includes examples of rank 5 curves found in the rank 1 families of subsection 3.1. First column indicates the number of the family and the second the value(s) of the parameter producing a rank 5 curve. The indication (L) means

that this curve has been previously found by Lecacheux.

Family number	w values
2	$\frac{287}{109}$,
3	$\frac{73}{83}$, $\frac{37}{157}$,
4	$-\frac{87}{28}$,
5	$\frac{317}{10}$,
6	$-\frac{28}{79}$, $\frac{100}{29}$ (L), $\frac{304}{55}$.

6.2. The case of torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$. The highest known rank of an elliptic curve over \mathbb{Q} with torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ was found by Elkies in 2006. See [Du2] for the details of this curve. It has rank 6 and it can be seen to correspond to the value $u = -\frac{16}{3}$ in the family with rank 2 included in subsection 5.2. For $u = \frac{5}{13}$ we get a curve with rank 5.

The following list includes examples of rank 5 curves found in the rank 1 families of subsection 3.2. First column indicates the number of the family and the second the value(s) of the parameter producing a rank 5 curve. The indication (L) and (D) means that in these cases the curves were found previously by Lecacheux and Dujella respectively.

Family number	w values
1	$\frac{306}{11}$
3	$\frac{53}{90}$ (D), $-\frac{127}{74}$ (L)
5	$\frac{31}{42}$
6	$\frac{13}{43}$, $-\frac{431}{33}$
7	$\frac{115}{6}$, $-\frac{142}{33}$ (D), $-\frac{391}{387}$, $-\frac{1011}{551}$
8	$\frac{302}{161}$
9	$\frac{44}{61}$, $\frac{40}{57}$, $\frac{172}{191}$, $\frac{214}{163}$, $\frac{284}{197}$

REFERENCES

- [Be] A. Beauville, Les familles stables des courbes elliptiques sur P^1 admettent quatre fibres singulieres, C.R. Acad. Sc. Paris. Serie I 294 (1982), 657-660.
- [BL] M.J. Bertin, O. Lecacheux, Elliptic fibrations on the modular surface associated to $\Gamma_1(8)$, Arxiv: 1105.6312v1 (2011)
- [Ca] G. Campbell, Finding elliptic curves defined over \mathbb{Q} of high rank, African americans in Mathematics (Piscataway, NJ, 1996) DIMACS Ser. Discrete Math. Theoret. Compt. Sci., vol 34, AMS, Providence, RI, 1997, p. 107-109.
- [Cr] J. Cremona, Algorithms for Modular Elliptic Curves. Cambridge University Press, Cambridge, 1997.
- [Du1] A. Dujella, On Mordell-Weil groups of elliptic curves induced by Diophantine triples, Glas. Mat. Ser. III 42 (2007), 3-18.

- [Du2] A. Dujella, High rank elliptic curves with prescribed torsion,
<http://web.math.hr/~duje/tors/tors.html>.
- [DP] A. Dujella, J.C. Peral, High rank elliptic curves with torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ induced by
 Diophantine triples, preprint in
<http://web.math.hr/~duje/papers1.html>.
- [El] N. Elkies, Three lectures on elliptic surfaces and curves of high rank, Arxiv: 0709.2908v1
 (2007)
- [H] T. Hadano, Elliptic curves with a torsion point, Nagoya Math. J., 66 (1977) p. 99-108.
- [Kn] A. Knapp, Elliptic curves, Mathematical Notes, Princeton University Press, (1992)
- [Ku] L. Kulesz, Families of elliptic curves of high rank with nontrivial torsion group over \mathbb{Q} , Acta
 Arith. 108 (2003), no. 4, p. 339-356.
- [Le1] O. Lecacheux, Rang de courbes elliptiques, Acta Arith. 109 (2003), no. 2, p. 131-142.
- [Le2] O. Lecacheux, Rang de courbes elliptiques dont le groupe de torsion est non trivial, Ann.
 Sci. Math. Quebec 28 (2004), no 1-2, p. 145-151.
- [LY] R. Livne, N. Yui, The modularity of certain non-rigid Calabi-Yau threefolds, Arxiv:
 0304497v2 (2005)
- [Me] J.F. Mestre, Courbes elliptiques de rang ≥ 11 sur $\mathbb{Q}(t)$, C.R. Acad. Sc. Paris. Serie I 313
 (1991), 139-142.
- [Ra] P. Rabarison, Construction of elliptic curves with high rank and large torsion group,
 (preprint) (2009)
- [Sh] T. Shioda, An infinite family of elliptic curves over \mathbb{Q} with large rank via Neron's method,
 Invent. Math. 106 (1991), 109-119.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, BIJENIČKA CESTA 30, 10000 ZAGREB,
 CROATIA

E-mail address, A. Dujella: duje@math.hr

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DEL PAÍS VASCO, APTDO. 644, 48080 BILBAO,
 SPAIN

E-mail address, J. C. Peral: juancarlos.peral@ehu.es