

## ON DOUBLE COSETS IN FREE GROUPS

RITA GITIK AND ELIYAHU RIPS

**ABSTRACT.** It is shown that for any finitely generated subgroups  $H$  and  $K$  of a free group  $F$ , and for any  $g \in F$  the double coset  $HgK$  is closed in the profinite topology of  $F$ .

A well-known theorem of M. Hall [2] states (in different language) that any finitely generated subgroup of a free group is closed in the profinite topology. We show that a slight modification of its proof [6] yields a stronger result:

**Theorem.** *For any finitely generated subgroups  $H$  and  $K$  of a free group  $F$ , and for any  $g \in F$  the double coset  $HgK$  is closed in the profinite topology of  $F$ .*

A free group  $F = \langle X \rangle$  can be viewed as the fundamental group of a wedge  $W$  of  $|X|$  oriented circles labeled by elements of  $X$ . Subgroups of  $F$  correspond bijectively to based covering spaces of  $W$ , and any covering space of  $W$  is a graph which inherits orientation and labeling of its edges from  $W$ .

Let  $X_0 \subset X$  and let  $\Gamma$  be a subgraph of a covering of  $W$ . An  $X_0$ -component of  $\Gamma$  is a maximal connected subgraph of  $\Gamma$  with all its edges labeled by elements of  $X_0$ .

**Lemma.** *Let  $\Gamma$  be a subgraph of a covering of  $W$  such that  $\Gamma$  has finitely many vertices. There exists an embedding of  $\Gamma$  in a covering  $\Gamma'$  of  $W$  such that  $\Gamma$  and  $\Gamma'$  have the same vertices, and for any  $X_0 \subset X$  distinct  $X_0$ -components of  $\Gamma$  remain distinct in  $\Gamma'$ .*

*Proof.* We give an algorithm for constructing  $\Gamma'$  by adding edges to  $\Gamma$  in a unique way. For any vertex  $v$  of  $\Gamma$  and for any  $x \in X$  the number of edges labeled with  $x$  having an endpoint at  $v$  is either 0, 1 or 2. If the number is 0, we add an edge labeled with  $x$  with both endpoints at  $v$ . If the number is 1 or 2, let  $p_x$  be the maximal path consisting of edges labeled only with  $x$  and with an endpoint at  $v$ . If  $p_x$  has both endpoints at  $v$  we do nothing. Otherwise we add to  $\Gamma$  an edge labeled with  $x$  connecting the endpoints of  $p_x$ . It is clear that the projection from  $\Gamma$  to  $W$  extends uniquely to a covering map from  $\Gamma'$  to  $W$ .

*Remark.* Let  $H$  be a finitely generated subgroup of  $F$ , and let  $f \in F \setminus H$ . Let  $\Gamma$  be the minimal connected subgraph of the covering  $C$  of  $W$  corresponding to  $H$  which contains the core of  $C$  (cf. [6]) and the path  $p$  beginning at the basepoint  $v_0$  of  $C$  whose projection in  $W$  represents  $f$ . Embed  $\Gamma$  in a covering  $\Gamma'$  as in the lemma. Then as  $\Gamma$  has finitely many vertices, so does  $\Gamma'$ , therefore the subgroup  $M$  of  $F$

---

1991 *Mathematics Subject Classification.* 20F32, 20E05, 20E26.

corresponding to  $\Gamma'$  has finite index in  $F$ . As  $p$  is not a closed path in  $\Gamma$ , it remains not closed in  $\Gamma'$ , hence  $f \notin M$ . As  $\Gamma$  is a subgraph of  $\Gamma'$ ,  $H$  is a subgroup of  $M$ , proving M. Hall's theorem (cf. [6]).

*Proof of the theorem.* As  $HgK = H(gKg^{-1})g$ , it is enough to consider the case  $g = 1$ .

By an observation due to P.Kropholler, we can replace  $F$  by a subgroup of finite index (cf. [3]), so we can assume that  $K$  is a free factor of  $F$ ,  $F = K * L$ . Let  $X_1$  and  $X_2$  be sets of free generators of  $K$  and  $L$  respectively, then  $X = X_1 \cup X_2$  is a set of free generators of  $F$ . Let  $f \in F, f \notin HK$ . Our goal is to construct a subgroup  $M$  of finite index in  $F$  such that  $Mf \cap HK = \emptyset$ .

Let  $\Gamma, \Gamma'$  and  $M$  be as in the remark. As  $f \notin HK$ , the  $X_1$ -component of  $v_0$  in  $\Gamma$  does not contain the endpoint of  $p$ , therefore the lemma implies that  $\Gamma'$  has the same property. But the condition  $Mf \cap HK = \emptyset$  is equivalent to the condition that the endpoint of  $p$  does not belong to the  $X_1$ -component of  $v_0$ , proving the theorem.

*Remarks.*

- 1) The first published proof of the theorem is due to G.A. Niblo [3], who found a much more simple and elegant argument than the original proof of the authors [1].
- 2) A more general result saying that for any finitely generated subgroups  $H_1, \dots, H_n$  of a free group  $F$  the set  $H_1 \cdots H_n$  is closed in the profinite topology on  $F$  was obtained by L. Ribes and P.A. Zalesskii [4], and by K. Henckell, S.T. Margolis, J.E. Pin and J. Rhodes [5].
- 3) It is easy to construct a closed subset  $B$  of  $F$  and a finitely generated subgroup  $K$  such that the product  $BK$  is not closed in the profinite topology of  $F$ .

## REFERENCES

- [1] R. Gitik and E. Rips, *On Separability Properties of Groups*, Int J. of Algebra and Computation **5** (1995), 703-717.
- [2] M. Hall, Jr., *Coset Representations in Free Groups*, Trans. AMS **67** (1949), 431-451.
- [3] G.A. Niblo, *Separability Properties of Free Groups and Surface Groups*, J. of Pure and Applied Algebra **78** (1992), 77-84.
- [4] L. Ribes and P.A. Zalesskii, *On the Profinite Topology on a Free Group I*, Bull. LMS **25** (1993), 37-43.
- [5] K. Henckell, S.T. Margolis, J.E. Pin and J. Rhodes, *Ash's Type II Theorem, Profinite Topology and Malcev Products I*, Int J. of Algebra and Computation **1** (1991), 411-436.
- [6] J.R. Stallings, *Topology of Finite Graphs*, Invent. Math. **71** (1983), 551-565.

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM, 91904, ISRAEL

*Current address:* A and H consultants, Ann Arbor, MI 48104

*E-mail address:* ritagtk @ umich.edu

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM, 91904, ISRAEL