

# NONSTATIONARY ANALOGUE BLACK HOLES

GREGORY ESKIN, DEPARTMENT OF MATHEMATICS, UCLA,  
LOS ANGELES, CA 90095-1555, USA

**ABSTRACT.** We study the existence of analogue nonstationary spherically symmetric black holes. The prime example is the acoustic model (cf. [V], [U]). We consider also a more general class of metrics that could be useful in other physical models of analogue black and white holes. We give examples of the appearance of black holes and of disappearance of white holes. We also discuss the relation between the apparent and the event horizons for the case of analogue black holes. In the end we study the inverse problem of determination of black or white holes by boundary measurements for the spherically symmetric nonstationary metrics.

## 1. INTRODUCTION

Let

$$(1.1) \quad \sum_{j,k=0}^n g_{j,k}(x_0, x) dx_j dx_k$$

be a Lorentzian metric with the signature  $(1, -1, \dots, -1)$ , where  $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ ,  $x_0 \in \mathbb{R}$  is the time variable. We assume that  $g_{jk}(x_0, x) \in C^\infty(\mathbb{R}^n \times \mathbb{R})$ ,  $g_{jk}(x_0, x) = \eta_{jk} + O(\frac{1}{|x|})$  uniformly in  $x_0$ ,  $[\eta_{jk}]_{j,k=0}^n$  is the Minkowski tensor. We also assume that  $g_{jk}(x_0, x) - g_{j,k}(\pm\infty, x)$  decay fast uniformly in  $x$  when  $x_0 \rightarrow \pm\infty$ .

Consider the wave equation

$$(1.2) \quad \square_g u(x_0, x) \stackrel{\text{def}}{=} \sum_{j,k=0}^n \frac{1}{\sqrt{(-1)^n g(x_0, x)}} \frac{\partial}{\partial x_j} \left( \sqrt{(-1)^n g} g^{jk}(x_0, x) \frac{\partial u}{\partial x_k} \right) = 0,$$

where  $[g^{jk}]_{j,k=0}^n = ([g_{jk}]_{j,k=0}^n)^{-1}$ ,  $g(x_0, x) = \det[g_{jk}]_{j,k=0}^n$ .

In the case when the metric  $[g_{jk}(x_0, x)]_{j,k=0}^n$  is a solution of the Einstein's equations of general relativity, the solutions  $u(x_0, x)$  of the equation (1.2) describe linear gravitational waves in the background of the metric (1.1).

One of the striking phenomenons of general relativity is the appearance of black holes and white holes.

In general relativity the domain  $D \subset \mathbb{R}^n \times \mathbb{R}$  is called a black hole if no signal (disturbance) inside  $D$  can escape to the spatial infinity.

In this paper we shall consider some classes of nonstationary black and white holes that arise in the study of analogue space-times. Our definitions are similar but different from the definition of black and white holes in general relativity (cf. [W], [FN]).

**Definition 1.1** Let  $D$  be a domain in  $\mathbb{R}^n \times \mathbb{R}$  such that the domains  $D_t = D \cap \{x_0 = t\} \subset \mathbb{R}^n$  are uniformly bounded on  $\mathbb{R}$ . Suppose  $\partial D$  is a characteristic surface for the equation (1.2).

We say that  $D$  is a black hole if no signal (disturbance) inside  $D$  can reach the exterior of  $D$ .

**Definition 1.2** The domain  $D \subset \mathbb{R}^n \times \mathbb{R}$  is called an outer black hole if it is a black hole and any signal (disturbance) outside  $D$  reaches spatial infinity when  $x_0 \rightarrow +\infty$ .

The boundary of a black hole (outer black hole) is called a black hole horizon (outer black hole horizon). The Definition 1.2 is similar to the definition of a black hole in general relativity.

Analogously,  $D$  is a white hole if no signal (disturbance) outside  $D$  can penetrate into the interior of  $D$ , and  $D$  is an outer white hole if it is a white hole and any signal (disturbance) outside  $D$  can reach the spatial infinity when  $x_0 \rightarrow -\infty$ . We introduced the notion of an outer black hole because sometimes there exists more than one black hole. Then only the largest black hole that contains all other will be the outer black hole. Moreover, the outer black holes can be determined by the boundary measurements (cf. §5).

The equation (1.2) describes also linear waves in a moving medium such as the propagation of light in moving dielectric (cf. [G], [LP], [NVV]) or acoustic waves in a moving fluid (cf. [V1], [U], [BLV]). The black holes for such equations are called optical or acoustic black holes. They are also called analogue or artificial black holes. There are other interesting physical models of analogue spacetimes as surface waves models, Bose-Einstein condensates, gravity waves and others (see excellent papers [V1], [V2], where many physical examples are considered). The study of analogue black holes is much simpler than the study of the black holes in the general relativity and physicists expect that it will help them to understand better the black holes of the general relativity. They also expect to study the analogue black holes experimentally. Another road leading to the study of analogue black holes is the attempt to answer the following question (cf. [E2], [E4]): Can one determine the coefficients of the wave equation by the boundary measurements? Such problems are called the inverse hyperbolic problems. The presence of black hole will make the determination of the coefficients inside the black hole impossible.

Now we shall describe the content of the present paper. In §2 we study the event and apparent horizons and the relations between them. In §3 we shall refine the results of [E1] on the existence of black and white holes for stationary metrics in two space dimensions. The novelty here is the consideration of metric with singularity at some point. Although the emphasis in the paper is on the spherically symmetric case we consider in §3 the nonspherically symmetric metrics too, since the proofs in the stationary case are almost the same. In §4 we study the existence of nonstationary black and white holes depending on  $x_0$  and  $|x|$  only. Note that there are only few studies of nonstationary analogue black holes, mostly in the case of one space dimension (cf. [BLSV]). We consider separately the case of acoustic black or white holes and the case of general nonstationary spherically symmetric metrics. We give also an example of black holes appearing at some time  $x_0 = t$  and white holes disappearing at some  $x_0 = t$ . Finally, in §5 we consider the inverse problem of the determination of a nonstationary spherically symmetric black or a white hole knowing the time-dependent Dirichlet-to-Neumann operator on the boundary. Inverse problems for stationary metrics in any dimensions were considered in [E2].

## 2. EVENT HORIZON AND APPARENT HORIZON

In this section we describe black and white holes analytically, extending the proof of Theorem 3.1 of [E1] to the case of nonstationary metrics. We assume that

$$(2.1) \quad g^{00}(x_0, x) > 0, \quad \forall (x_0, x).$$

Let  $S(x_0, x) = 0$  be a smooth surface in  $\mathbb{R}^n \times \mathbb{R}$ , closed in  $\mathbb{R}^n$ ,  $\forall x_0 \in \mathbb{R}$  and the gradient  $(S_{x_1}, \dots, S_{x_n})$  be not zero when  $S(x_0, x) = 0$ . We shall choose outward direction of the gradient. Suppose  $S(x_0, x) = 0$  is a characteristic surface for the equation (1.2), i.e.

$$(2.2) \quad \sum_{j,k=0}^n g^{jk}(x_0, x) S_{x_j}(x_0, x) S_{x_k}(x_0, x) = 0 \quad \text{when} \quad S(x_0, x) = 0.$$

Equation (2.2) has two roots with respect to  $S_{x_0}$ :

$$(2.3)$$

$$S_{x_0}^{\pm} = \left( - \sum_{j=1}^n g^{j0}(x_0, x) S_{x_j} \pm \sqrt{Q(x_0, x)} \right) / g^{00}(x_0, x), \quad \text{where}$$

$$(2.4)$$

$$Q(x_0, x) = \left( \sum_{j=1}^n g^{j0}(x_0, x) S_{x_j} \right)^2 - g^{00}(x_0, x) \sum_{j,k=1}^n g^{jk}(x_0, x) S_{x_j} S_{x_k}.$$

Since (1.2) is strictly hyperbolic we have that  $Q(x_0, x) > 0 \quad \forall (x_0, x)$ . Therefore  $S_{x_0}^-(x_0, x) < S_{x_0}^+(x_0, x)$ .

**Theorem 2.1.** *Let  $d_{x_0}$  be the diameter of  $S(x_0, x) = 0$  in  $\mathbb{R}^n$  for fixed  $x_0$ . Suppose that  $d_{x_0}$  is bounded on  $(-\infty, +\infty)$ . Suppose  $S(x_0, x)$  is a smooth characteristic surface,  $-\infty < x_0 < +\infty$ . Then  $S(x_0, x) = 0$  is a boundary of a black hole if  $S_{x_0}(x_0, x)$  is the smaller root in (2.3), i.e.  $S_{x_0}(x_0, x) = S_{x_0}^-(x_0, x)$ , and  $S(x_0, x) = 0$  is a boundary of a white hole if  $S_{x_0}(x_0, x)$  is the larger root in (2.3), i.e.  $S_{x_0}(x_0, x) = S_{x_0}^+(x_0, x)$ .*

**Proof:** Let  $S(y_0, y) = 0$ , and let  $S_{x_0} = S_{x_0}^+$  be the larger root in (2.3). Denote by  $K_+(y_0, y)$  the forward time-like half-cone in  $\mathbb{R}^n \times \mathbb{R}$  consisting of  $(\dot{x}_0, \dot{x}_1, \dots, \dot{x}_n) \in \mathbb{R}^n \times \mathbb{R}$  such that

$$(2.5) \quad \sum_{j,k=0}^n g_{jk}(y_0, y) \dot{x}_j \dot{x}_k > 0, \quad \dot{x}_0 > 0.$$

Let  $K^+(y_0, y)$  be the half-cone of the dual cone

$$(2.6) \quad \sum_{j,k=0}^n g^{jk}(y_0, y) \xi_j \xi_k > 0, \quad (\xi_0, \dots, \xi_n) \in \mathbb{R}^n \times \mathbb{R},$$

that contains  $(1, 0, \dots, 0)$ . Note that

$$(2.7) \quad \sum_{k=0}^n \dot{x}_k \xi_k > 0$$

for all  $(\dot{x}_0, \dots, \dot{x}_n) \in K_+(y_0, y)$  and all  $(\xi_0, \xi_1, \dots, \xi_n) \in K^+(y_0, y)$ . Vector  $(S_{x_0}^+ + \varepsilon, S_{x_1}^+, \dots, S_{x_n}^+) \in K^+(y_0, y)$  for any  $\varepsilon > 0$  because

$$\begin{aligned} g^{00}(y_0, y)(S_{x_0}^+ + \varepsilon)^2 + 2 \sum_{j=1}^n g^{j0}(y_0, y)(S_{x_0}^+ + \varepsilon)S_{x_j}^+ + \sum_{j,k=1}^n g^{jk}(y_0, y)S_{x_j}^+ S_{x_k}^+ \\ = g^{00}(y_0, y)\varepsilon^2 + 2\varepsilon \left( \sum_{j=0}^n g^{j0} S_{x_j}^+ \right). \end{aligned}$$

We use that  $S^+$  satisfies (2.2). It follows from (2.3) that  $\sum_{j=0}^n g^{j0} S_{x_j}^+ = +\sqrt{Q} > 0$ . Thus  $g^{00}\varepsilon^2 + 2\varepsilon\sqrt{Q} > 0$ . Therefore

$$(2.8) \quad g^{00}(S_{x_0}^+ + \varepsilon)^2 + 2 \sum_{j=1}^n g^{j0}(S_{x_0}^+ + \varepsilon)S_{x_j}^+ + \sum_{j,k=1}^n g^{jk}(y_0, y)S_{x_j}^+ S_{x_k}^+ > 0, \quad \forall \varepsilon > 0.$$

Taking  $\varepsilon > 0$  large we get that the vector

$$(S_{x_0}^+ + \varepsilon, S_{x_1}^+, \dots, S_{x_n}^+) = \varepsilon \left( 1 + \frac{S_{x_0}^+}{\varepsilon}, \frac{S_{x_1}^+}{\varepsilon}, \dots, \frac{S_{x_n}^+}{\varepsilon} \right)$$

can be deformed continuously to the vector  $\varepsilon(1, 0, \dots, 0) \in K^+(y_0, y)$ . Therefore  $(S_{x_0}^+ + \varepsilon, S_{x_1}^+, \dots, S_{x_n}^+)$  belongs to the half-cone  $K^+(y_0, y)$ . It follows from (2.7) that

$$(2.9) \quad (S_{x_0}^+ + \varepsilon)\dot{x}_0 + \sum_{k=1}^n S_{x_k}^+ \dot{x}_k > 0$$

for all  $(\dot{x}_n, \dots, \dot{x}_1) \in K_+(y_0, y)$ ,  $\forall \varepsilon > 0$ . Passing to the limit when  $\varepsilon \rightarrow 0$  we get

$$(2.10) \quad \sum_{j=0}^n S_{x_j}^+ \dot{x}_j \geq 0$$

for all  $(\dot{x}_1, \dots, \dot{x}_n) \in \overline{K}_+(y_0, y)$ . Since  $(S_{x_0}^+, \dots, S_{x_n}^+)$  is an outward normal the inequality (2.10) shows that all forward time-like rays starting at  $(y_0, y)$  are pointed inside the exterior of  $S^+(x_0, x) = 0$ . This proves that  $S^+(x_0, x) = 0$  is a boundary of a white hole.

Consider now the case when  $S(x_0, x) = S^-(x_0, x_0)$ , i.e.  $S_{x_0}^-$  is the smaller root in (2.3). For  $\varepsilon > 0$  we have using (2.3) that  $(-S_{x_0}^- + \varepsilon, -S_{x_1}^-, \dots, -S_{x_n}^-) \in K^+(y)$  and as in (2.9), (2.10) we get that

$$(2.11) \quad \sum_{j=0}^n S_{x_j}^- \dot{x}_j \leq 0$$

for all  $(\dot{x}_0, \dots, \dot{x}_n) \in K_+(y)$ . Since  $(S_{x_0}^-, \dots, S_{x_n}^-)$  is an outward normal to  $S^-(y_0, y) = 0$ , we have that any forward time-like ray starting at  $(y_0, y)$  is pointed inside  $S^-(x_0, x) = 0$ . Therefore  $S^-(x_0, x) = 0$  is a boundary of the black hole.  $\square$

Black or white hole horizons are the notions global in time  $x_0$ . The introduction of the notion of an apparent horizon is a way to get an information about the event horizon (black or white hole) at a given time.

Consider a nonstationary metric in  $\mathbb{R}^n \times \mathbb{R}$ . Fix  $x_0 = t$ . Let  $S(x) = 0$  be a closed smooth surface in  $\mathbb{R}^n \times \{x_0 = t\}$ . We assume that  $S_x(x) \neq 0$  when  $S(x) = 0$  and that  $S_x(x)$  is an outward normal to  $S(x) = 0$ .

Consider the system of null-bicharacteristics (cf. [E1]) with initial data  $x_0(0) = t, x_j(0) = y_j, 1 \leq j \leq n, S(y) = 0, \xi_j(0) = S_{x_j}(y), 1 \leq j \leq$

$n$ . There are two such null-bicharacteristics with

$$(2.12) \quad \xi_0^\pm(0) = \frac{1}{g^{00}(t, y)} \left( - \sum_{j=1}^n g^{j0}(t, y) S_{x_j}(y) \pm \sqrt{Q(t, y)} \right),$$

where  $Q(t, y)$  is the same as in (??). Note that the strict hyperbolicity implies that  $Q(t, y) > 0$ . Thus if

$$(2.13) \quad \sum_{j,k=1}^n g^{jk}(t, y) S_{x_j}(y) S_{x_k}(y) \geq 0$$

then  $Q > 0$  implies  $\sum_{j=1}^n g^{j0}(t, y) S_{x_j}(y) \neq 0$ . We assume in the case of black holes (cf. [E1]) that

$$(2.14) \quad \sum_{j=1}^n g^{j0}(t, y) S_{x_j}(y) < 0 \quad \text{when} \quad S(y) = 0.$$

In the case of white hole we assume that (2.13) holds and

$$(2.15) \quad \sum_{j=1}^n g^{j0}(t, y) S_{x_j}(y) > 0 \quad \text{when} \quad S(y) = 0.$$

Let  $K_+(y)$  and  $K^+(y)$  be the same as in (2.5), (2.6). If (2.13) and (2.14) hold then

$$(2.16) \quad \xi_0^+(0) > 0, \quad \xi_0^-(0) \geq 0.$$

Note that  $\xi_0^-(0) = 0$  if

$$(2.17) \quad \sum_{j,k=1}^n g^{jk}(t, y) S_{x_j}(y) S_{x_k}(y) = 0.$$

We generalize the proof of Theorem 2.1 for the case when (2.17) may not hold.

As in (2.8), (2.9), (2.10) we get that  $\overline{K_+}(y) \subset \Pi^+$ , where  $\Pi^+$  is the half-space

$$(2.18) \quad \xi_0^+ \dot{x}_0 + \sum \dot{x}_k S_{x_k}(y) \geq 0.$$

Analogously to (2.11) we get that  $\overline{K_+}(y) \subset \Pi^-$  where  $\Pi^-$  is the half-space

$$(2.19) \quad \xi_0^- \dot{x}_0 + \sum_{k=1}^n \dot{x}_k S_{x_k}(y) \leq 0.$$

Thus  $\overline{K_+}(y) \subset \Pi^+ \cap \Pi^-$ . Denote by  $D \subset \mathbb{R}^n$  the domain with boundary  $\{S(x) = 0\}$ . If  $\xi_0^+(0) > 0$ ,  $\xi_0^-(0) \geq 0$  for all  $y$  such that  $S(y) = 0$  the domain  $D \times \mathbb{R}$  is a trapped (no escape) region for the stationary metric

$$(2.20) \quad \sum_{j,k=0}^n g_{jk}(t, x) dx_j dx_k,$$

where  $t$  is fixed (“frozen”). When  $\xi_0^-(0) = 0$   $D \times \mathbb{R}$  is the outmost trapped region, i.e. a black hole for the metric (2.20), where  $t$  is fixed.

Analogously, when (2.13) and (2.15) hold, we have  $\xi_0^-(y) < 0$ ,  $\xi_0^+(y) \leq 0$ ,  $S(y) = 0$ . As in (2.18), (2.19) we get that

$$(2.21) \quad \overline{K_+}(y) \subset \Pi_1^- \cap \Pi_1^+,$$

where  $\Pi_1^-$  consists of  $(\dot{x}_0, \dots, \dot{x}_n), \dot{x}_0 > 0$ , satisfying

$$\xi_0^-(y)\dot{x}_0 + \sum_{k=1}^n \dot{x}_k S_{x_k}(y) \leq 0 \text{ and}$$

$$\Pi_1^+ = \{(\dot{x}_0, \dots, \dot{x}_n) : \xi_0^+(y)\dot{x}_0 + \sum_{k=1}^n \dot{x}_k S_{x_k}(y) \geq 0, \dot{x}_0 > 0\}.$$

Therefore for the metric (2.20),  $t$  is fixed, we have that any disturbance in the exterior of  $D \times \mathbb{R}$  can not reach the interior of  $D \times \mathbb{R}$ , i.e.  $D \times \mathbb{R}$  is an antitrapped region. In the case when  $\xi_0^+(y) = 0$   $D \times \mathbb{R}$  is the outmost antitrapped region, i.e.  $S \times \mathbb{R}$  is a white hole horizon for the metric (2.20). Therefore the apparent horizon is the black or white hole horizon for the metric (2.20) when  $x_0 = t$  is fixed. Note that  $D \times \mathbb{R}$  is not necessary a trapped region for the non-stationary metric. Examine the relation between the apparent horizon and the event horizon. Let  $S(x_0, x) = 0$  be a smooth surface in  $\mathbb{R}^{n+1}$  such that  $S(x_0, x) = 0$  is a smooth closed surface in  $\mathbb{R}^n$  for each  $x_0 \in \mathbb{R}$ . Assume that  $\xi_0^+(0) > 0$ ,  $\xi_0^-(0) = 0$  (cf. (2.12)) hold for each  $x_0$  when  $S(x_0, y) = 0$ . Then  $\{S(t, y) = 0\} \times \mathbb{R}$  is the apparent horizon for the metric (2.20) when  $x_0 = t$  is fixed. The surface  $S(x_0, x) = 0$  in  $\mathbb{R}^{n+1}$  is called the dynamic horizon. In §3 we shall study the stationary acoustic metrics and in §4 the nonstationary acoustic metrics with  $A(x_0) \leq A_0 < 0$  (cf. (3.1) and Theorem 4.1). Let  $r = r^+(x_0)$  be the black hole horizon (cf. Theorem 4.1). The dynamic horizon in this case is  $r = |A(x_0)|$ , i.e.  $\{r = |A(t)|\} \times \mathbb{R}$  is apparent horizon for each  $t$ . If  $|A(x_0)|$  is increasing when  $x_0$  is increasing we have that the dynamic horizon is inside the black hole with the boundary  $r = r^+(x_0)$ . If  $|A(x_0)|$  is decreasing when  $x_0$  is increasing then the black hole with the boundary  $r = r^+(x_0)$  is inside the domain bounded by the dynamic horizon.

### 3. STATIONARY BLACK HOLES IN TWO SPACE DIMENSIONS

**3.1. Metric singularity.** In this section we consider the case of  $n = 2$  and the Lorentz metric tensor  $[g_{jk}^7(x)]_{j,k=0}^2$  independent of the time variable  $x_0$ .

Let  $S$  be the ergosphere (cf [E1]), i.e the smooth closed curve  $\Delta(x) = 0$  where  $\Delta(x) = g^{11}(x)g^{22}(x) - (g^{12}(x))^2$ .

Suppose that  $S$  is the boundary of simply connected domain  $\Omega$  and suppose  $O = (0, 0) \in \Omega$ . We assume that  $[g_{jk}(x)]_{j,k=0}^n$  are  $C^\infty$  in  $\Omega \setminus O$ ,  $\Delta(x) < 0$  in  $\Omega \setminus O$  and  $[g_{jk}(x)]_{j,k=0}^2$  has a singularity at  $O$ . Note that the Schwarzschild metric (see, for example, [W]) has a singularity at the origin.

Another example of metric having a singularity is the acoustic metric (cf. [V1]) when

$$(3.1) \quad g^{00} = 1, \quad g^{j0} = g^{0j} = v^j, \quad 1 \leq j \leq 2, \quad g^{jk} = -\delta_{ij} + v^i v^j, \quad 1 \leq j, k \leq 2,$$

where  $v = (v^1, v^2)$  is the velocity of the flow,

$$v = \frac{A}{r}\hat{r} + \frac{B}{r}\hat{\theta}, \quad r = |x|, \quad \hat{r} = \frac{x}{|x|}, \quad \hat{\theta} = \frac{(-x_2, x_1)}{|x|},$$

$A$  and  $B$  are constants. Note that many metrics such as Kerr metric (cf. [W]), Gordon metric (cf. [G], [LP]) have the form (3.1).

In case of general stationary metric in  $\mathbb{R}^2 \times \mathbb{R}$  we assume that when  $|x| < \varepsilon$ ,  $\varepsilon$  is small, the metric tensor  $[g^{jk}]_{j,k=0}^2$  has the following form

$$(3.2) \quad g^{jk}(x) = g_1^{jk}(x) + g_2^{jk}(x), \quad |x| < \varepsilon,$$

where  $g_1^{jk}$  is similar to an acoustic metric,

$$(3.3) \quad g_1^{jk} = v^j v^k, \quad 1 \leq j, k \leq 2, \quad g_1^{j0} = g_1^{0j} = v^j, \quad 1 \leq j \leq 2, \quad g_1^{00} = 0,$$

$$v = (v^1, v^2) = \frac{b_1(\theta)}{r}\hat{r} + \frac{b_2(\theta)}{r}\hat{\theta}, \quad b_j, \quad j = 1, 2, \quad b_1 \neq 0,$$

$g_2^{jk}$  are smooth in polar coordinates  $(r, \theta)$ ,  $g_2^{00} \geq C > 0$ ,  $g_2^{j0} = g_2^{0j} = O(r)$ ,  $1 \leq j \leq 2$ ,  $[g_2^{jk}]_{j,k=1}^2$  is a negative definite matrix when  $|x| < \varepsilon$ ,

$$(3.4) \quad ([g_2^{jk}]_{j,k=1}^2 \alpha, \alpha) \leq -C_0 |\alpha|^2, \quad \forall \alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2.$$

Writing the matrix  $[g^{jk}]_{j,k=0}^2$  in polar coordinates we get

$$(3.5) \quad H\left(r, \theta, \xi_0, \xi_r, \frac{\xi_\theta}{r}\right) = g^{00}\xi_0^2 + 2g^{r0}\xi_0\xi_r + 2g^{\theta 0}\xi_0\frac{\xi_\theta}{r} + g^{rr}\xi_r^2 + 2g^{r\theta}\xi_r\frac{\xi_\theta}{r} + g^{\theta\theta}\frac{\xi_\theta^2}{r^2}.$$

Consider the system of null-bicharacteristics with the Hamiltonian (3.5):

$$(3.6) \quad \frac{dr}{ds} = H_{\xi_r}, \quad \frac{d\theta}{ds} = H_{\xi_\theta}, \quad \frac{dx_0}{ds} = H_{\xi_0}, \quad \frac{d\xi_r}{ds} = -H_r, \quad \frac{d\xi_\theta}{ds} = -H_\theta, \quad \frac{d\xi_0}{ds} = -H_{x_0}.$$

We impose the following initial conditions:

$$(3.7) \quad r(0) = \varepsilon, \quad \theta(0) = \theta_0, \quad \xi_r(0) = \eta_r, \quad \xi_\theta(0) = \eta_\theta, \quad \xi_0(0) = \eta_0.$$



Since  $\frac{\partial H}{\partial x_0} = 0$ , we have that  $\xi_0(s) = \eta_0, \forall s$ , and we choose  $\xi_0 = \eta_0 = 0$ .

There are two family of null-bicharacteristics when  $\xi_0 \equiv 0$ :

$$(3.8) \quad \xi_r^\pm(s) = \frac{-g^{r\theta} \pm \sqrt{(g^{r\theta})^2 - g^{rr}g^{\theta\theta}}}{g^{rr}} \cdot \frac{\xi_\theta^\pm(s)}{r} = \frac{-g^{r\theta} \pm \sqrt{-\Delta}}{g^{rr}} \cdot \frac{\xi_\theta^\pm(s)}{r},$$

assuming that  $g^{rr} \neq 0$ .

The decomposition (3.2) has the following form in polar coordinates

$$(3.9) \quad g^{rr} = \frac{b_1^2}{r^2} + g_2^{rr}, \quad g^{r\theta} = \frac{b_1 b_2}{r^2} + g_2^{r\theta}, \quad g^{\theta\theta} = \frac{b_2^2}{r^2} + g_2^{\theta\theta},$$

$$g^{r0} = \frac{b_1}{r} + g_2^{r0}, \quad g^{\theta 0} = \frac{b_2}{r} + g_2^{\theta 0}.$$

**Theorem 3.1.** *Suppose  $b_1 \neq 0$ . When  $\varepsilon > 0$  is small and  $b_1 < 0$  all null-bicharacteristics (3.6) starting at  $r = \varepsilon$  reach  $r = 0$  as  $x_0$  increases. If  $b_1 > 0$  then all null-bicharacteristics starting at  $r = \varepsilon$  reach  $r = 0$  as  $x_0$  decreases*

**Proof:** Dividing  $\frac{dr}{ds}$  by  $\frac{dx_0}{ds}$  and having  $\xi_0 = 0$  we get from (3.6):

$$(3.10) \quad \frac{dr}{dx_0} = \frac{\left(\frac{b_1^2}{r^2} + g_2^{rr}\right)\xi_r + \left(\frac{b_1 b_2}{r^2} + g_2^{r\theta}\right)\frac{\xi_\theta}{r}}{\left(\frac{b_1}{r} + g_2^{r0}\right)\xi_r + \left(\frac{b_2}{r} + g_2^{\theta 0}\right)\frac{\xi_\theta}{r}}$$

Substituting (3.8) and (3.9) in (3.10) and cancelling  $\frac{\xi_\theta^\pm}{r}$ , we get

$$\begin{aligned} \frac{dr^\pm}{dx_0} &= \frac{\left(\frac{b_1^2}{r^2} + O(1)\right)\left(-\frac{b_2}{b_1} \pm C_1 r + O(r^2)\right) + \frac{b_1 b_2}{r^2} + O(1)}{\left(\frac{b_1}{r} + O(r)\right)\left(-\frac{b_2}{b_1} \pm C_1 r + O(r^2)\right) + \frac{b_2}{r} + O(r)} \\ &= \frac{\frac{\pm b_1^2}{r} C_1 + O(1)}{\pm b_1 C_1 + O(r)} = \frac{b_1}{r} + O(1). \end{aligned}$$

Since  $b_1(\theta) \leq b_0 < 0$  we have  $\frac{dr^\pm}{dx_0} < -\frac{|b_0|}{2r}$ , for small  $r$ . Therefore

$$2r \frac{dr}{dx_0} < -|b_0|, \quad (r^\pm(x_0))^2 - (r^\pm(t))^2 \leq -|b_0|(x_0 - t), \quad r^\pm(t) > 0.$$

When  $x_0$  increases  $r^\pm(x_0) = 0$  for some  $x_0 > t$ . When  $b_1 \geq b_0 > 0$  we have

$$\frac{dr^\pm}{dx_0} \geq \frac{b_0}{2r}, \quad (r^\pm(t))^2 - (r^\pm(x_0))^2 \geq b_0(t - x_0),$$

When  $x_0$  decreases we get  $r^\pm(x_0) = 0$  for some  $x_0 < t$ . □

**3.2. Existence of black or white holes.** The following refinement of Theorem 4.1 in [E1] holds:

**Theorem 3.2.** *Let  $S$  be the ergosphere, i.e.  $S = \{x : \Delta(x) = 0\}$  where  $\Delta(x) = g^{11}g^{22} - (g^{12}(x))^2$ . We assume that  $S$  is a smooth Jordan curve. Let  $\Omega$  be the interior of  $S$  and let  $O \in \Omega$ . Assume that  $\Delta(x) < 0$  for  $x \in \Omega \setminus O$  and assume that  $S$  is not characteristic at any  $x \in S$ . Assume that the metric tensor  $[g^{jk}(x)]_{j,k=0}^2$  has a singularity at  $O$  and conditions (3.2), (3.3), (3.4) are satisfied. If  $b_1 < 0$  then there exists smooth Jordan curves  $S^+$  and  $S^-$  in  $\Omega$  both containing  $O$ ,  $S^-$  is inside the domain with boundary  $S^+$ , such that  $S^+ \times \mathbb{R}$  and  $S^- \times \mathbb{R}$  are black hole horizons and there is no black hole horizons between  $S^+$  and  $S$  and between  $S^-$  and  $O$ .*

*Analogously, when  $b_1 > 0$  there exists white hole horizons  $S^+ \times \mathbb{R}$  and  $S^- \times \mathbb{R}$ . When  $S^- = S^+$  there is only one black hole or white hole horizon between  $S$  and  $O$ .*

**Proof:** Let  $S_\varepsilon = \{x : |x| = \varepsilon\}$ ,  $\varepsilon$  is small, and  $\Omega_\varepsilon = \Omega \setminus \{x : |x| < \varepsilon\}$ . Suppose  $b_1 < 0$  (cf. (3.3)). It follows from Theorem 3.1 that both families of characteristics  $x = x^+(x_0)$  and  $x = x^-(x_0)$  (cf. (4.9) in [E1]) end on  $S_\varepsilon$  when  $x_0$  is increasing. It was shown in [E1] that one of these families, say  $x = x^+(x_0)$ , ends at  $S$  and the second family  $x = x^-(x_0)$  starts at  $S$  when  $x_0$  increases. Therefore if some curve  $x = x_+^{(1)}(x_0)$  touches  $S_\varepsilon$  at some time  $x_0^{(0)}$  it can not reach  $S$  when  $x_0 < x_0^{(0)}$  decreases. Therefore the limit set of the trajectory  $x = x_+^{(1)}(x_0)$  is contained inside  $\Omega_\varepsilon$  and by the Poincare-Bendixson theorem there exists a closed Jordan curve  $x = s_0(x_0)$  such that  $S_0 \times \mathbb{R}$  is the event horizon (i.e. either black hole or white hole horizon), where  $S_0 = \{x = s_0(x_0)\}$  (cf. Theorem 4.1 in [E1]). Consider any curve of the second family  $x = x^-(x_0)$  on  $S_0$ . If it is pointed inside  $S_0$  then  $S_0 \times \mathbb{R}$  is a black hole horizon, and there is no black hole horizon inside  $S_0$ . We denote such  $S_0$  by  $S^-$ . If any curve  $x = x^-(x_0)$  is pointed outward of  $S_0$  then  $S_0 \times \mathbb{R}$  is the white hole event horizon. In this case consider any trajectory  $x = x_-^{(1)}(x_0)$  of  $x = x^-(x_0)$  family that reaches  $S_\varepsilon$  at some time  $x_0^{(1)}$ . When  $x_0 < x_0^{(1)}$  decreases  $x = x_-^{(1)}(x_0)$  can not reach  $S_0$  and again by the Poincare-Bendixson theorem there exists closed Jordan characteristic curve  $x = s_1(x_0)$  such that  $S^{(1)} \times \mathbb{R}$  is a black hole horizon where  $S^{(1)} = \{x = s_1(x_0)\}$  and there are no black holes inside  $S^{(1)}$ . In this case we set  $S^{(1)} = S^-$ . Therefore in both cases  $S^- \times \mathbb{R}$  is the smallest black hole horizon in  $\Omega \setminus O$ .

Consider now a curve  $x = x_+^{(2)}(x_0)$  of the  $x = x^+(x_0)$  family that ends at  $S$  at the time  $x_0 = x_0^{(2)}$ . When  $x_0 < x_0^{(2)}$  decreases this trajectory

can not reach  $S_\varepsilon$  and by the Poincare-Bendixson theorem there is a closed Jordan trajectory  $x = s_2(x_0)$  such that  $S_2 = \{x = s_2(x_0)\} \times \mathbb{R}$  is an event horizon. If  $S_2 \times \mathbb{R}$  is a black hole horizon we denote  $S_2$  by  $S^+$ . If  $S_2 \times \mathbb{R}$  is a white hole horizon then we have, as above, that there is a closed characteristic curve  $S_3 = \{x = s_3(x_0)\}$  belonging to  $x = x^-(x_0)$  family such that  $S_3 \times \mathbb{R}$  is a black hole horizon and there are no black hole horizons between  $S_3$  and  $S$ . In this case we denote  $S_3$  by  $S^+$ .

This concludes the proof of Theorem 3.2 in the case  $b_1 < 0$ . If  $b_1 > 0$  then the same reasoning leads to the proof that there are two white hole horizons  $S^+ \times \mathbb{R}$  and  $S^- \times \mathbb{R}$  such that there is no white hole horizons between  $S^+$  and  $S$  and between  $S^-$  and  $O$ . In the case when  $S^- = S^+$  we have a unique outer black or white hole in  $\Omega \setminus O$ .  $\square$

Note the following property of the black hole with the boundary  $S^+ \times \mathbb{R}$ : disturbance at any point outside  $S^+ \times \mathbb{R}$  propagate to the spatial infinity. The black hole horizon  $S^- \times \mathbb{R}$  has the property that any point inside  $S^- \times \mathbb{R}$  ends on  $O$ .

In Example 4.2 in [E1] the case when there are many black hole horizons and white hole horizons was considered.

**Remark 3.1** In the paper [EH] by Michael Hall and the author (see also Michael Hall, PhD dissertation, UCLA, 2013) the condition that the ergosphere is not characteristic at any point is removed.

#### 4. SPHERICALLY SYMMETRIC BLACK HOLES FOR THE NONSTATIONARY METRICS

**4.1. Spherically symmetric black holes.** Let the nonstationary metric tensor  $[g_{jk}(x_0, x)]_{j,k=0}^n$  be the same as in (1.1). We assume that the metric has a singularity at  $O \times \mathbb{R}$  that satisfies (3.2), (3.3), (3.4) uniformly in  $x_0$ . We shall introduce spherical coordinates  $(r, \theta)$  in  $\mathbb{R}^n$  and we shall write the tensor  $[g^{jk}]_{j,k=0}^n$  in such coordinates (cf. (3.5) for the case of  $n = 2$ ). We shall not assume that the metric is spherically symmetric but we shall require that  $g^{00}(x_0, r)$ ,  $g^{0r} = g^{r0}(x_0, r)$  and  $g^{rr}(x_0, r)$  be dependent of  $x_0, r = |x|$  only.

We are looking for the characteristic surface of the form  $S(x_0, r) = 0$ . Then the characteristic equation has the form

$$(4.1) \quad g^{00}(x_0, r)S_{x_0}^2 + 2g^{r0}(x_0, r)S_{x_0}S_r + g^{rr}(x_0, r)S_r^2 = 0$$

when  $S(x_0, r) = 0$ .

Note that  $S(x_0, r) = 0$  is a curve in  $(x_0, r)$  space but it is a  $(n-1)+1$  dimensional surface in  $\mathbb{R}^n \times \mathbb{R}$ . We shall study when  $S(x_0, r) = 0$  is a black hole horizon starting with the case of nonstationary acoustic metric.

**4.2. Nonstationary acoustic metric.** Consider a time-dependent acoustic metric (3.1) in  $(\mathbb{R}^2 \setminus O) \times \mathbb{R}$  with the velocity flow of the form

$$(4.2) \quad v = \frac{A(x_0)}{r} \hat{r} + \frac{B(x_0, r, \theta)}{r} \hat{\theta}.$$

The characteristic equation for  $S(x_0, r) = 0$  is

$$\left( S_{x_0}(x_0, r) + \frac{A(x_0)}{r} S_r \right)^2 - S_r^2 = 0,$$

or

$$(4.3) \quad S_{x_0}^\pm = \left( -\frac{A(x_0)}{r} \pm 1 \right) S_r^\pm \quad \text{when } S(x_0, r) = 0.$$

Assuming that  $S_r \neq 0$  when  $S = 0$  we can, without loss of generality, look for  $S$  in the form  $S = r - r(x_0) = 0$ . Then  $r = r(x_0)$  satisfies the ordinary differential equation

$$(4.4) \quad r_{x_0}(x_0) = \frac{A(x_0)}{r} \mp 1.$$

There are two family of solutions  $r = r^+(x_0)$  and  $r = r^-(x_0)$  satisfying the equations  $r_{x_0}^+ = \frac{A(x_0)}{r} + 1$  and  $r_{x_0}^- = \frac{A(x_0)}{r} - 1$ , respectively. We assume  $A(x_0) \rightarrow A(\pm\infty)$  when  $x_0 \rightarrow \pm\infty$ . We shall show that there exists a unique (outer) black hole horizon if  $A(x_0) \leq A_0 < 0$  and there exists a unique (outer) white hole horizon if  $A(x_0) \geq A_0 > 0$ . When  $A(x_0) = A_0$  there is an explicit solution of  $\frac{dr^\pm}{dx_0} = \frac{A_0}{r} \mp 1$ . In particular, when  $A_0 < 0$  and  $\frac{dr}{dx_0} = -\frac{|A_0|}{r} + 1$ . We have

$$(4.5) \quad \frac{r}{r - |A_0|} dr = dx_0, \quad r + |A_0| \ln |r - |A_0|| = x_0 + C,$$

$r^+(x_0) \rightarrow |A_0|$  when  $x_0 \rightarrow -\infty$ . If  $r^+(0) = r_0 < |A_0|$  then  $r^+(x_0)$  decreases to 0 when  $x_0$  increases. If  $r^+(0) = r_0 > |A_0|$  then  $r^+(x_0) \rightarrow +\infty$  when  $x_0 \rightarrow +\infty$  (cf. Fig. 1a). The (outer) black hole horizon is  $\{r = |A_0|\} \times \mathbb{R}$ . The solution  $r^+ = |A_0|$  is the separatrix separating the solutions tending to  $+\infty$  when  $x_0 \rightarrow +\infty$  and the solutions ending on  $r = 0$  at some  $x_0 = x_0^{(0)}$ .

Since  $\frac{dr^-(x_0)}{dx_0} = -\frac{|A_0|}{r} - 1 < -1$  all trajectories  $r = r^-(x_0)$  end on  $r = 0$  when  $x_0$  increases (see Fig. 1b). Analogously, if  $A(x_0) = A_0 > 0$  then  $\{r < A_0\} \times \mathbb{R}$  is a white hole. All trajectories of  $\frac{dr^+(x_0)}{dx_0} = \frac{A_0}{r} + 1 > 1$  start at  $r = 0$  and tend to  $+\infty$  when  $x_0$  increases (see Fig. 2a). For the trajectories of  $\frac{dr^-}{dx_0} = \frac{A_0}{r} - 1$  we have that the solution  $r = A_0$  separates the solutions that tend to 0 and to  $+\infty$  when  $x_0$  decreases.




Figure (1a)




Figure (1b)

Figure 1.

- (1a) The family  $r = r^+(x_0)$ ,  $A(x_0) = A_0 < 0$ .  $\{r < |A_0|\} \times \mathbb{R}$  is a black hole.  
(1b) The family  $r = r^-(x_0)$ ,  $A(x_0) = A_0 < 0$ .




Figure (2a)




Figure (2b)

Figure 2.

- (2a) Trajectories  $r = r^+(x_0)$  when  $A_0 > 0$ .  
(2b) Trajectories  $r = r^-(x_0)$  when  $A_0 > 0$ .  $\{r < A_0\} \times \mathbb{R}$  is a white hole.

Now consider the case of  $A(x_0)$  depending on  $x_0$ . Let  $A(x_0) \leq -A_0 < 0$ . Since  $\frac{dr^-}{dx_0} = -\frac{|A(x_0)|}{r} - 1 < -1$  all trajectories  $r = r^-(x_0)$  end on

$r = 0$  (cf. Fig 1b.). Consider

$$(4.6) \quad \frac{dr^+(x_0)}{dx_0} = \frac{A(x_0)}{r} + 1 = \frac{r + A(x_0)}{r}.$$

Make change of variables  $r^+ + A(x_0) = v(x_0)$ , where  $v(x_0)$  will be chosen later. Then  $\frac{dr^+}{dx_0} = \frac{dv}{dx_0} - A'(x_0)$  and  $v(x_0)$  satisfies the equation

$$(4.7) \quad \frac{dv}{dx_0} - \frac{v}{|A(x_0)| + v} = A'(x_0).$$

Let  $w = v \exp(-b(x_0))$ , where  $b(x_0) = \int_0^{x_0} \frac{dy_0}{|A(y_0)| + v(y_0)}$ .

Then  $\frac{dw}{dx_0} \exp b(x_0) = A'(x_0)$  and  $w(x_0) = w(0) + \int_0^{x_0} A'(t) \exp(-b(t)) dt$ .

Therefore  $v = \left( w(0) + \int_0^{x_0} A'(t) \exp(-b(t)) dt \right) \exp b(x_0)$ . Choosing  $w(0) = -\int_0^\infty A'(t) \exp(-b(t)) dt$  we have the following representation for particular solution  $v(x_0)$  of (4.7)

$$(4.8) \quad v(x_0) = - \int_{x_0}^\infty A'(t) \exp \left( - \int_{x_0}^t \frac{dy_0}{|A(y_0)| + v(y_0)} \right) dt.$$

Note that  $v(x_0) \rightarrow 0$  when  $x_0 \rightarrow +\infty$  since  $t > x_0$  in (4.8). Denote by  $B$  the Banach space with norm  $\|v\|_T = \sup_{x_0 \geq T} |v(x_0)|$ ,  $T$  will be large enough. Denote by  $F(v)$  the operator

$$(4.9) \quad F(v) = - \int_{x_0}^\infty A'(t) \exp \left( - \int_{x_0}^t \frac{dy_0}{|A(y_0)| + v(y_0)} \right) dt.$$

We have when,  $\|v\|_T < \frac{1}{2}|A_0|$ ,

$$(4.10) \quad \|F(v)\|_T \leq \int_T^\infty |A'(t)| dt.$$

Also we get, when  $\|v_j\|_T < \frac{|A_0|}{2}, j = 1, 2$ ,

$$(4.11) \quad \|F(v_1) - F(v_0)\|_T \leq C \int_T^\infty t |A'(t)| dt \|v_1 - v_0\|_T.$$

Therefore  $F(v)$  is a contraction map of a ball  $\|v\|_T \leq \varepsilon$  if  $T$  is sufficiently large. Hence there exists a unique bounded solution  $r_0^+(x_0) = -A(x_0) + v$  of (4.6) for  $x_0 > T$ . Taking in (4.8) the limit when  $x_0 \rightarrow +\infty$  we get  $\lim_{x_0 \rightarrow +\infty} r_0^+(x_0) = -\lim_{x_0 \rightarrow +\infty} A(x_0) + \lim_{x_0 \rightarrow +\infty} v(x_0) = -A(+\infty)$ . Since  $A(x_0) \leq -A_0 < 0$  for all  $-\infty < x_0 < +\infty$  and  $\frac{dr_0^+(x_0)}{dx_0} = \frac{A(x_0)}{r} + 1$  we can extend  $r = r_0^+(x_0)$  from  $(T, +\infty)$  to  $(-\infty, +\infty)$ :

For small  $r$  we have  $\frac{dr_0^+}{dx_0} < -\frac{|A_0|}{2r_0}$ . Thus when  $r_0^+(x_0)$  is small it increases when  $x_0$  decreases and hence it cannot hit  $r = 0$  for any  $x_0 \in \mathbb{R}$ . If  $r$  is large then  $\frac{dr_0^+}{dx_0} > \frac{1}{2}$  and  $r_0^+(x_0)$  decreases when  $x_0$  decreases.

Therefore  $r = r_0^+(x_0)$  will remain bounded and nonzero for all  $x_0 \in \mathbb{R}$ . Since  $b(x_0) < 0$  when  $x_0 < 0$  and  $|v(x_0)| < \frac{1}{2}|A_0|$  all solutions of (4.7) tend to zero when  $x_0 \rightarrow -\infty$ . Therefore  $\lim_{x_0 \rightarrow -\infty} r_0^+(x_0) = \lim_{x_0 \rightarrow -\infty} -A(x_0) + \lim_{x_0 \rightarrow -\infty} v(x_0) = -A(-\infty)$ .

Therefore we proved the following theorem.

**Theorem 4.1.** *Consider the acoustic metric with the velocity flow (4.2). Assume that  $A(x_0) \in C^\infty(\mathbb{R})$ ,  $A(x_0) \leq A_0 < 0$ ,  $\lim_{x_0 \rightarrow \pm\infty} A(x_0) = A(\pm\infty)$ . Then there exists a unique (outer) black hole horizon  $r = r_0^+(x_0)$  and  $\lim_{x_0 \rightarrow \pm\infty} r_0^+(x_0) = |A(\pm\infty)|$ . The solution  $r = r_0^+(x_0)$  of (4.6) separates the solutions  $r = r^+(x_0)$  that reach  $r = 0$  at some time  $x_0$  and the solutions  $r = r^+(x_0)$  that tend to the infinity when  $x_0 \rightarrow +\infty$ . Analogously when  $A(x_0) \geq A_0 > 0$  there exists a unique (outer) white hole horizon  $r = r_0^-(x_0)$  such that  $\lim_{x_0 \rightarrow \pm\infty} r_0^-(x_0) = A(\pm\infty)$ .*

The solution  $r = r^-(x_0)$  separates the solutions  $r = r^-(x_0)$  that tend to  $\infty$  when  $x_0 \rightarrow -\infty$  and the solutions  $r = r^-(x_0)$  that end on  $r = 0$  when  $x_0$  decreases.

**4.3. Appearance of black holes at finite time.** Suppose  $A(x_0)$  in (4.6) is such that  $A(x_0) < 0$  when  $x_0 > 0$  and  $A(x_0) \geq 0$  for  $x_0 < 0$ . It was proven above that there exists a solution  $r = r_0^+(x_0)$  of (4.6) for  $x_0 > T$ ,  $T$  is large. We shall show that this solution will vanish at some point  $x_0^{(1)} < 0$ . Let  $x_0 < 0$ . Then  $\frac{dr_0^+(x_0)}{dx_0} = \frac{A(r_0)}{r} + 1 > 1$  since  $A(x_0) \geq 0$  for  $x_0 < 0$ . Thus  $r_0^+(x_0)$  decreases when  $x_0$  decreases and we have, at some time  $x_0 = x_0^{(1)}$ , that  $r_0^+(x_0^{(1)}) = 0$ . Therefore the black hole  $\{r < r_0^+(x_0)\} \times (x_0^{(1)} < x_0 < +\infty)$  starts at the time  $x_0^{(1)}$  and there is no black hole for  $x_0 < x_0^{(1)}$ .

Analogously suppose in (4.6) we have  $A(x_0) > 0$  for  $x_0 < 0$  and  $A(x_0) < 0$  for  $x_0 > 0$ . Let  $r = r_0^-(x_0)$  be the white hole horizon on  $(-\infty, -T)$ . When  $x_0 > 0$  we have  $\frac{dr_0^-(x_0)}{dx_0} = \frac{A(x_0)}{r} - 1 \leq -1$  since  $A(x_0) < 0$  for  $x_0 > 0$ . Therefore  $r^-(x_0^{(2)}) = 0$  for some  $x_0^{(2)}$ . This means that the white hole  $\{r < r_0^-(x_0)\} \times (-\infty, x_0^{(2)})$  ends at the time  $x_0 = x_0^{(2)}$ .

**4.4. The general case.** Consider the metric such that  $g^{00}, g^{r0}, g^{rr}$  depend on  $(x_0, r)$  only so that the characteristic equation has the form (4.1). We assume that  $g^{00}(x_0, r) \geq C_0 > 0$ . We have

$$(4.12) \quad S_{x_0}^\pm = \frac{-g^{r0}(x_0, r) \pm \sqrt{(g^{r0})^2 - g^{00}g^{rr}}}{g^{00}} S_r^\pm.$$

Note that  $S_{x_0}^+$  is the larger root in (4.12) and  $S_{x_0}^-$  is the smaller root.

The strict hyperbolicity implies that

$$(4.13) \quad (g^{r0})^2 - g^{00}g^{rr} \geq C_0 > 0 \quad \forall x_0, r.$$

We shall assume that

$$(4.14) \quad g^{00} = 1 + O\left(\frac{1}{r}\right), \quad g^{r0} = \left(\frac{1}{r}\right), \quad g^{rr} = -1 + O\left(\frac{1}{r}\right)$$

when  $r$  is large. Near  $r = 0$  we assume (cf. (3.2), (3.3), (3.4))

$$(4.15) \quad g^{r0} = \frac{b_1(x_0)}{r} + g_2^{r0}, \quad g^{rr} = \frac{b_1^2(x_0)}{r^2} + g_2^{rr}, \quad g^{00} = 1 + g_2^{00},$$

where  $g_2^{r0} = O(r^2)$ ,  $g_2^{00} = O(r^3)$ ,  $g_2^{rr} \leq -C_0 < 0$ ,  $b_1(x_0) \leq b_0 < 0$ . Then

$$(g^{r0})^2 - g^{00}g^{rr} = -g_2^{rr} + O(r) \geq \frac{C_0}{2}$$

for  $r$  small uniformly in  $x_0$ . Therefore near  $r = 0$  we have

$$(4.16) \quad S_{x_0}^\pm = \left( -\frac{b_1(x_0)}{r} \pm g_0(x_0, r) \right) S_r^\pm,$$

where  $g_0 \geq C > 0$ .

Looking for  $S^\pm = 0$  in the form  $S^\pm = r - r^\mp(x_0) = 0$  we get ordinary differential equations for  $r = r^\pm(x_0)$

$$(4.17) \quad r_{x_0}^+ = \frac{g^{r0}(x_0, r)}{g^{00}(x_0, r)} + d(x_0, r), \quad r_{x_0}^- = \frac{g^{r0}(x_0, r)}{g^{00}(x_0, r)} - d(x_0, r),$$

where  $d(x_0, r) \geq C_0$ . Since  $S^\pm = r - r^\mp(x_0)$  we have that  $r^+$  corresponds to the smaller root in (4.12) and  $r^-(x_0)$  corresponds to the larger root in (4.12).

When  $r$  is close to 0 we have

$$(4.18) \quad r_{x_0}^\pm = \frac{b_1(x_0)}{r} \pm d_1(x_0, r), \quad d_1 \geq C_1 > 0.$$

When  $r$  is large we have

$$(4.19) \quad r_{x_0}^\pm = \pm 1 + O\left(\frac{1}{r}\right).$$

Suppose  $b_1(x_0) < b_0 < 0$  for all  $x_0 \in \mathbb{R}$ . It follows from (4.18) that

$$(4.20) \quad \frac{dr^\pm(x_0)}{dx_0} < -\frac{1}{2}|b_0| \quad \text{for small } r.$$

Therefore if  $r^\pm(x_0^{(1)}) = r_0$  is small then  $(r^\pm(x_0))^\pm - (r^\pm(x_0^{(1)}))^\pm \leq -\frac{1}{2}|b_0|(x_0 - x_0^{(1)})$  and  $r^\pm(x_0^{(2)}) = 0$  for some  $x_0^{(2)} > x_0^{(1)}$ . If  $r^+(x_0^{(1)}) = r_0$  is large it follows from (4.19) that  $r^+(x_0) \rightarrow +\infty$  since

$$(4.21) \quad \frac{dr^+(x_0)}{dx_0} > \frac{1}{2} \quad \text{for } r^+(x_0) > r_0.$$



Denote by  $\mathcal{F}^+$  the set of all  $r^+(x_0)$  that tend to  $+\infty$  when  $x_0 \rightarrow +\infty$ . For each  $t \in \mathbb{R}$  define by  $R^+(t)$  the infimum of  $r^+(t)$  over all  $r^+(x_0) \in \mathcal{F}^+$ :  $R^+(t) = \inf_{\mathcal{F}^+} r^+(t)$ .

Note that  $R^+(t) > 0$ : if  $r^+(t)$  is small then  $r^+(t) \notin \mathcal{F}^+$  since it will end at  $r = 0$ . Let  $R^+(x_0)$  be the solution of (4.17) passing through  $R^+(t)$  at  $x_0 = t$ . Since  $R^+(t) \leq r^+(t)$ ,  $\forall r^+ \in \mathcal{F}^+$ , we have, for any  $x_0$ , that  $R^+(x_0) \leq r^+(x_0)$ ,  $\forall r^+ \in \mathcal{F}^+$ . The solution  $R^+(x_0)$  is bounded when  $x_0 \rightarrow +\infty$  since if  $R^+(x_0)$  is unbounded then we will have  $\frac{dR^+}{dx_0} > \frac{1}{2}$  for  $x_0 \geq x_0^{(0)}$ . Then, choosing  $r_0$  such that  $R^+(x_0^{(0)}) - \varepsilon < r_0 < R^+(x_0^{(0)})$ , where  $\varepsilon$  small, we get that the solution  $r_0^+(x_0)$ , where  $r_0^+(x_0^{(0)}) = r_0$ , also tends to  $+\infty$  when  $x_0 \rightarrow +\infty$ , i.e.  $r_0^+ \in \mathcal{F}^+$ . Since  $r_0^+(x_0^{(0)}) < R^+(x_0^{(0)})$ , we have that  $r_0^+(x_0) < R^+(x_0)$  for all  $x_0$ . In particular,  $r_0^+(t) < R^+(t)$  and this is a contradiction, since  $R^+(t) = \inf_{\mathcal{F}^+} r^+(t)$ . Note that the solution  $R^+(x_0)$  exists for all  $x_0 \in \mathbb{R}$ : it can not end at  $r = 0$  since  $b_1(x_0) < 0$ ,  $\forall x_0$ , and for small  $r$ ,  $r^+(x_0)$  increases when  $x_0$  decreases (cf. (4.20)), and it can not escape to the infinity when  $x_0 \rightarrow -\infty$  since, for large  $r$ ,  $r^+(x_0)$  is decreasing when  $x_0$  is decreasing (see (4.21)). Note that  $R^+(t_1) = \inf_{r^+ \in \mathcal{F}^+} r^+(t_1)$  for any  $t_1$ : Suppose  $R_1^+(t_1) = \inf_{\mathcal{F}^+} r^+(t_1) > R^+(t_1)$ . Denote by  $R_1^+(x_0)$  the solution of (4.17) passing through  $R_1^+(t_1)$  when  $x_0 = t_1$ . Since  $R_1^+(t_1) > R^+(t_1)$  we have that  $R_1^+(t) > R^+(t)$  and this contradicts that  $R^+(t) = \inf_{\mathcal{F}^+} r^+(t)$ , since  $r^+(t) \geq R_1^+(t) > R^+(t)$  for  $\forall r^+ \in \mathcal{F}^+$ .

Since  $R^+(x_0)$  corresponds to the smaller root  $S^-$  in (4.12) the solution  $r = R^+(x_0)$  is an outer black hole horizon (cf. Definition 1.2). Now denote by  $\mathcal{F}^-$  the set of solutions  $r = r^+(x_0)$  that end at  $r = 0$  at some time  $x_0 = x_0^{(0)}$ . We extend  $r = r^+(x_0)$  by zero for  $x_0 \geq x_0^{(0)}$ . For fixed  $t \in \mathbb{R}$  denote by  $R_+(t)$  the supremum of  $r^+(t)$ , where  $r^+ \in \mathcal{F}^-$ :  $R_+(t) = \sup_{\mathcal{F}^-} r^+(t)$ . Let  $R_+(x_0)$  be a solution of (4.17) passing through  $R_+(t)$  at  $x_0 = t$ . Note that  $R_+(t) < +\infty$  since any  $r^+(x_0) \in \mathcal{F}^+$  is larger than any  $r^+(x_0) \in \mathcal{F}^-$ . The same arguments as for  $R^+(x_0)$  show that  $R_+(t_1) = \sup_{\mathcal{F}^-} r^+(t)$  for any  $t_1$ .

As for  $R^+(x_0)$  we have that  $R_+(x_0)$  exists for all  $x_0 \in \mathbb{R}$  providing that  $b(x_0) \leq b_0 < 0$  for all  $x_0 \in \mathbb{R}$ . Therefore  $R_+(x_0)$  is a black hole horizon. Note that  $R_+(x_0) \leq R^+(x_0)$ ,  $\forall x_0$ .

We proved the following theorem:

**Theorem 4.2.** *Consider a nonstationary metric such that  $g^{00}, g^{r0}, g^{rr}$  depend on  $(x_0, r)$  only. Let  $g^{00}(x_0, r) \geq C_0 > 0$ . Assume that conditions (4.13), (4.14), (4.15) are satisfied. Suppose  $b_1(x_0) \leq b_0 < 0$  in (4.15),  $\forall x_0 \in \mathbb{R}$ . Then there exists an outer black hole horizon  $r = R^+(x_0)$  such that  $R^+(x_0) > 0$ ,  $\forall x_0 \in \mathbb{R}$ ,  $R^+(x_0)$  is bounded on  $\mathbb{R}$ .*

Moreover,  $r = R^+(x_0)$  is the boundary of the set of all solutions  $r = r^+(x_0)$  of (4.17) that tend to the infinity when  $x_0 \rightarrow +\infty$ . Also there exists a black hole horizon  $r = R_+(x_0)$  such that  $R_+(x_0) > 0$  on  $\mathbb{R}$ ,  $R_+(x_0)$  is bounded on  $\mathbb{R}$ ,  $R_+(x_0) \leq R^+(x_0)$  and  $R_+(x_0)$  is the boundary of all solutions of (4.17) that end at  $r = 0$  for some  $x_0 \in \mathbb{R}$ . If  $R_+(x_0) = R^+(x_0)$  there is a unique black hole horizon.

**Remark 4.1** Analogous results hold for white hole horizons assuming that  $b_1(x_0) \geq b_0 > 0$  for all  $x_0$ . In particular, there exists white hole horizons  $r = R^-(x_0)$  and  $r = R_-(x_0)$  such that  $R^-(x_0)$  is the boundary of the set of all solutions  $r = r^-(x_0)$  such that  $r^-(x_0) \rightarrow +\infty$  when  $x_0 \rightarrow -\infty$  and  $r = R_-(x_0)$  is the boundary of the set of all  $r = r^-(x_0)$  such that  $r^-(x_0)$  ends at  $r = 0$  when  $x_0$  decreases.

## 5. INVERSE PROBLEMS AND BLACK OR WHITE HOLES

Consider nonstationary spherically symmetric metric

$$(5.1) \quad g_{00}(x_0, r)dx_0^2 + 2g_{0r}(x_0, r)dx_0dr + g'_{rr}(x_0, r)dr^2 - dr^2 - r^2(d\theta)^2,$$

where  $(d\theta)^2$  is a standard metric on  $S^{n-1}$ .

Let

$$\square_g u(x, t) = 0$$

be the wave equation of the form (1.2) corresponding to the metric (5.1). Suppose  $g_{00}, g_{0r}$  and  $g_{rr} = -1 + g'_{rr}$  satisfy the condition of the Theorem 4.2.

Let  $r = R^+(x_0)$  be the boundary of the outer black hole, i.e.  $b_1(x_0) < 0$  in the condition (4.15). Let  $C$  be a cylinder  $C = \{r < a, -\infty < x_0 < +\infty\}$  such that  $r = R^+(x_0)$  is inside  $C$ , i.e.  $a$  is large enough.

Consider the following initial-boundary value problem

$$(5.2) \quad \square_g u = 0 \quad \text{in } C,$$

$$(5.3) \quad u = 0 \quad \text{for } x_0 \ll 0,$$

$$(5.4) \quad u|_{\partial C} = f, \quad f \in C_0^\infty(\partial C).$$

Let  $\Lambda f$  be the Dirichlet-to-Neumann operator, i.e.  $\Lambda f = \frac{\partial u}{\partial \nu}|_{\partial C}$  where  $\nu$  is the outward unit normal to  $\partial C$ ,  $u$  is the solution of (5.2), (5.3), (5.4). The inverse hyperbolic boundary value problem consists of recovery of the metric knowing  $\Lambda f$  on  $\partial C$  for all  $f \in C_0^\infty(\partial C)$ , i.e. knowing the Cauchy data of  $u$  on  $\partial C$ . It is impossible to recover the metric inside the black hole since no information from the interior of the black hole can reach  $\partial C$ . Therefore changes of the metric inside  $r < R^+(x_0)$  does not effect boundary data on  $\partial C$ . However, we shall prove that knowing

the Cauchy data on  $\partial C$  one can recover the metric outside the black hole, i.e. when  $r > R^+(x_0)$ . Assuming that  $f$  does not depend on  $\theta$  on  $\partial C$  we shall consider the solution of the wave equation depending on  $(x_0, r)$  only

(5.5)

$$\begin{aligned} & \frac{1}{\sqrt{-g(x_0, r)}} \frac{\partial}{\partial x_0} \left( \sqrt{-g} g^{00}(x_0, r) \frac{\partial u}{\partial x_0} \right) + \frac{1}{\sqrt{-g}} \frac{\partial}{\partial x_0} \left( \sqrt{-g} g^{0r} \frac{\partial u}{\partial r} \right) \\ & + \frac{1}{\sqrt{-g}} \frac{\partial}{\partial r} \left( \sqrt{g} g^{0r} \frac{\partial u}{\partial x_0} \right) + \frac{1}{\sqrt{-g}} \frac{\partial}{\partial r} \left( \sqrt{-g} g^{rr} \frac{\partial u}{\partial r} \right) = 0, \quad r > 0, \quad x_0 \in \mathbb{R}, \end{aligned}$$

where  $g^{00}, g^{0r} = g^{r0}, g^{rr}$  are the same as in (4.1). Let  $\varphi(x_0, r)$  be a solution of

$$(5.6) \quad g^{00} \varphi_{x_0}^2 + 2g^{r0} \varphi_{x_0} \varphi_r + g^{rr} \varphi_r^2 = 0, \quad r > 0.$$

We assume, as in §4, that  $g^{00} \geq C > 0$  and conditions (4.13), (4.14), (4.15) are satisfied.

Note that we can factor (5.6):

$$(5.7) \quad g^{00} \varphi_{x_0} + g^{0r} \varphi_r \mp \sqrt{(g^{0r})^2 - g^{00} g^{rr}} \varphi_r = 0.$$

Denote by  $\varphi_1(x_0, r), \varphi_2(x_0, r)$  two solutions of (5.7)

$$(5.8) \quad g^{00} \varphi_{1x_0} + b_-(x_0, r) \varphi_{1r} = 0,$$

$$(5.9) \quad g^{00} \varphi_{2x_0} + b_+(x_0, r) \varphi_{2r} = 0,$$

where

$$(5.10) \quad b_{\pm}(x_0, r) = g^{0r} \pm \sqrt{(g^{0r})^2 - g^{00} g^{rr}}.$$

Note that  $b_{\pm}(x_0, r) \neq 0$  near  $r = a$ . Make change of variables

$$(5.11) \quad s = \varphi_1(x_0, r), \quad \tau = \varphi_2(x_0, r).$$

Then (5.5) has the following form in  $(s, \tau)$  coordinates:

$$(5.12) \quad \hat{g}^{s\tau} \frac{\partial^2 \hat{u}}{\partial s \partial \tau} = 0,$$

where

$$(5.13) \quad \hat{g}^{s\tau} = g^{00} \varphi_{1x_0} \varphi_{2x_0} + g^{r0} (\varphi_{1x_0} \varphi_{2r} + \varphi_{1r} \varphi_{2x_0}) + g^{rr} \varphi_{1r} \varphi_{2r}.$$

We used that (5.8), (5.9) imply that  $\hat{g}^{ss} = \hat{g}^{\tau\tau} = 0$  and that  $\sqrt{-\hat{g}} = (\hat{g}^{s\tau})^{-1}$ .

We impose the initial conditions on  $\varphi_1, \varphi_2$ :

$$(5.14) \quad \varphi_1|_{r=a} = x_0 + a, \quad \varphi_2|_{r=a} = -x_0 + a.$$

Let

(5.15)

$$y_0 = \frac{s - \tau}{2} = \frac{\varphi_1(x_0, r) - \varphi_2(x_0, r)}{2}, \quad y_1 = \frac{s + \tau}{2} = \frac{\varphi_1(x_0, r) + \varphi_2(x_0, r)}{2}$$

Then

$$(5.16) \quad y_0|_{r=a} = \frac{x_0 - (-x_0)}{2} = x_0, \quad y_1|_{r=a} = \frac{x_0 + a + (-x_0 + a)}{2} = a$$

Therefore the map  $(x_0, r) \rightarrow (y_0, y_1)$  is the identity when  $r = a$ .

Making the changes of variables (5.15) we get from (5.12)

$$(5.17) \quad \frac{\partial^2 \hat{u}(y_0, y_1)}{\partial y_0^2} - \frac{\partial^2 \hat{u}(y_0, y_1)}{\partial y_1^2} = 0,$$

where

$$(5.18) \quad \hat{u}(y_0, y_1) = u(x_0, r),$$

$(y_0, y_1)$  and  $(x_0, r)$  are related by the equations (5.15).

The characteristics of (5.5), crossing the line  $r = a$ , have the form

$$(5.19) \quad r - r^+(x_0, C_1) = 0, \quad r - r^-(x_0, C_2) = 0,$$

where  $\frac{dr^\pm}{dx_0} = \frac{b_\mp(x_0, r)}{g^{00}(x_0, r)}$ .

As in §4 solutions  $r = r^+(x_0, C_1)$ ,  $\forall C_1$ , tend to  $+\infty$  when  $x_0 \rightarrow +\infty$ , and  $r = r^-(x_0, C_2)$  end at  $r = 0$  when  $x_0$  increases,  $\forall C_2$ . It was shown in Theorem 4.2 that  $r = R^+(t) = \inf_{C_1} r^+(x_0, C_1)$  is the black hole horizon.

The images of the characteristics (5.19) under the map (5.15) are the characteristics  $s = C_1$ ,  $\tau = C_2$  of the equation (5.17).

Denote by  $D_g$  the subdomain in  $[0, a] \times \mathbb{R}^1$  bounded by the equation  $r = R^+(x_0)$  and the line  $\{r = a, x_0 \in \mathbb{R}\}$ . Then (5.15) maps  $D_g$  onto the half-plane  $\{-\infty < y_1 < a, y_0 \in \mathbb{R}\}$ . Since the map (5.15) is the identity on  $\{r = a, x_0 \in \mathbb{R}\}$ , the DN operators  $\Lambda$  and  $\hat{\Lambda}$  for (5.5) and (5.17) are equal on  $\{r = a, x_0 \in \mathbb{R}\}$ , i.e.

$$(5.20) \quad \Lambda f = \hat{\Lambda} f \quad \text{for all } f(x_0) \in C_0^\infty(\mathbb{R}).$$

Consider now any metric  $g_1$  of the form (5.1) assuming that conditions of the form (4.13), (4.14), (4.15) are satisfied. Let

(5.21)

$$\begin{aligned} & \frac{1}{\sqrt{-g_1(x_0, r)}} \frac{\partial}{\partial x_0} \left( \sqrt{-g_1} g_1^{00}(x_0, r) \frac{\partial u'(x_0, r)}{\partial x_0} \right) + \frac{1}{\sqrt{-g_1}} \frac{\partial}{\partial x_0} \left( \sqrt{-g_1} g_1^{0r} \frac{\partial u'}{\partial r} \right) \\ & + \frac{1}{\sqrt{-g_1}} \frac{\partial}{\partial r} \left( \sqrt{-g_1} g_1^{0r} \frac{\partial u'}{\partial x_0} \right) + \frac{1}{\sqrt{-g_1}} \frac{\partial}{\partial r} \left( \sqrt{-g_1} g_1^{rr} \frac{\partial u'}{\partial r} \right) = 0, \quad r > 0, \quad x_0 \in \mathbb{R}, \end{aligned}$$

Let  $r = R_1^+(x_0)$  be the black hole horizon for the metric  $g_1$ .

Denote by  $D_{g_1}$  the domain in  $[0, a) \times \mathbb{R}^1$  bounded by  $r = R_1^+(x_0)$  and  $\{r = a, x_0 \in \mathbb{R}\}$ .

Let  $\varphi'_1(x, r), \varphi'_2(x_0, r)$  be similar to (5.8), (5.9),

$$g_1^{00}\varphi'_{1x_0}(x, r) + b'_-\varphi'_{1r} = 0, \quad g_1^{00}\varphi'_{2x_0}(x_0, r) + b'_+\varphi_{2r} = 0,$$

where  $b'_\pm = g_1^{0r} \pm \sqrt{(g_1^{0r})^2 - g_1^{00}g_1^{rr}}$ .

Make the changes of variables

$$(5.22) \quad y'_0 = \frac{\varphi'_1(x_0, r) - \varphi'_2(x_0, r)}{2}, \quad y'_1 = \frac{\varphi'_1 + \varphi'_2}{2}$$

similar to (5.15), where  $\varphi'_1, \varphi'_2$  satisfy (5.14). Thus we transform the equation (5.21) to the equation

$$(5.23) \quad \frac{\partial^2 \hat{u}'(y'_0, y'_1)}{\partial (y'_0)^2} - \frac{\partial^2 \hat{u}'(y'_0, y'_1)}{\partial (y'_1)^2} = 0,$$

in the domain  $\{-\infty < y'_1 < a, y'_0 \in \mathbb{R}\}$ . Note that

$$(5.24) \quad \hat{u}'(y'_0, y'_1) = u'(x_0, r),$$

where  $(y'_0, y'_1)$  and  $(x_0, r)$  are related by (5.22).

Assume that DN operators  $\Lambda$  and  $\Lambda'$  corresponding to (5.5) and (5.21) coincide on  $\{r = a, x_0 \in \mathbb{R}\}$ , i.e.

$$(5.25) \quad \Lambda f = \Lambda' f \quad \text{for all } f \in C_0^\infty(\mathbb{R}).$$

Let  $\hat{\Lambda}'$  be the DN operator corresponding to (5.23). Since the map (5.22) is an identity on  $\{r = a, x_0 \in \mathbb{R}\}$  we have, as in (5.20) that

$$(5.26) \quad \Lambda' f = \hat{\Lambda}' f, \quad \forall f \in C_0^\infty(\mathbb{R}).$$

Therefore (5.20), (5.25), (5.26) imply that

$$(5.27) \quad \hat{\Lambda} f = \hat{\Lambda}' f, \quad \forall f \in C_0^\infty(\mathbb{R}).$$

Since  $\hat{u}'(y_0, y_1)$  and  $\hat{u}(y_0, y_1)$  satisfy the same equations (5.17) and (5.23), respectively, in  $\{-\infty < y_1 < a, x_0 \in \mathbb{R}\}$  and have the same Cauchy data (5.27), we get that

$$(5.28) \quad \hat{u}(y_0, y_1) = \hat{u}'(y_0, y_1).$$

Denote by  $(x'_0, r') = \sigma(x_0, r)$  the map of  $D_g$  onto  $D_{g'}$  that is a composition of the map (5.15) and the inverse of the map (5.22). Then (5.17), (5.24) and (5.28) imply that

$$(5.29) \quad u'(\sigma(x_0, r)) = u(x_0, r) \quad \text{in } D_g.$$

The equality (5.29) means that metrics  $g$  and  $g'$  are isometric. Note that  $\sigma(x_0, r)$  sends the characteristics of (5.5) in  $D_g$  to the characteristics of (5.21) in  $D_{g'}$ . Thus the boundary  $r = R^+(x_0)$  of  $D_g$  is mapped

to the boundary  $r = R_1^+(x_0)$  of  $D_{g'}$  (cf. Theorem 4.2). Therefore the DN operator of (5.5) on  $\{r = a, x \in \mathbb{R}\}$  determines the boundary  $r = R^+(x_0)$  of the black hole up to an isometry.

Consider the case of the white hole, i.e. when  $b_1(x_0) > 0$  in (4.15). Exactly, as in the case of black holes, one can prove that the DN operator on  $\{r = a, x_0 \in \mathbb{R}\}$  determines the boundary of the white hole  $r = R^-(x_0)$  up to an isometry.

Let  $D_g^-$  be the subdomain of  $\{0 < r < a, x_0 \in \mathbb{R}\}$  bounded by  $\{r = a, x_0 \in \mathbb{R}\}$  and the  $r = R^-(x_0)$ . One can see that any solution  $u(x_0, r)$  of (5.2), (5.3), (5.4) has the support in  $D_g^-$ , i.e. it is equal to zero when  $0 < r < R^-(x_0), x_0 \in \mathbb{R}$ . Thus boundary measurements on  $\partial C = \{r = a, x_0 \in \mathbb{R}\}$  can not recover any information about the metric in  $\{0 < r < R^-(x_0), x_0 \in \mathbb{R}\}$ .

### Acknowledgment

The author is deeply indebted to the referees for correcting mistakes and giving many valuable suggestions.

### REFERENCES

- [BLSV] Barcelo, C., Liberati, S., Sonego, S., Visser, M., Casual structure of acoustic spacetimes, New. J. Phys. 6 (2004), 186
- [BLV] Barcelo, C., Liberati, S., Visser, M., Analogue Gravity, ArXiv:gr-qc/0505065
- [E1] Eskin, G., Inverse hyperbolic problems and optical black holes, Commun. Math. Phys. 297, 817-839 (2010)
- [E2] Eskin, G., Perturbations of the Kerr black hole and the boundness of linear waves, Journ. of Math. Phys. 51, 112501 (2010)
- [E3] Eskin, G., Artificial black holes, Contemporary Mathematics, Vol. 535, 2011, 43-53
- [E4] Eskin, G., Uniqueness and Nonuniqueness in Inverse Hyperbolic Problems and the Black hole Phenomenon, Around research of Vladimir Maz'ya, III, Dodrecht: Springer, 2010
- [EH] Eskin, G. and Hall, M., Existence of black holes for stationary metrics in two space dimensions, arXiv:1402.6381
- [FN] Frolov, V. and Novikov, I., Black hole physics, Kluwer Academic Publishers, London (1998)
- [G] Gordon, W., 1923, Ann. Phys. (Leipzig) 72, 421
- [H] Hall, M., PhD dissertation, UCLA, May 2013
- [LP] Leonhardt, U., Piwnicki, P., 1999, Phys. Rev. A60, 4301
- [NVV] Novello, M., Visser, M., Volovik, G. (editors), Artificial black holes, 2002, World Scientific, Singapore.
- [U] Unruh, W., 1981, Phys. Rv. Lett., 46, 1351
- [V1] Visser, M., 1998, Acoustic black holes, horizons, ergospheres and Hawking radiation, Classical quantum gravity 15, No. 6, 1767-1791.
- [V2] Visser, M., Survey of analogue spacetimes, Lecture Notes in Physics, vol. 870, 31-50 (2013)
- [W] Wald, R., General relativity, Univercity of Chicago Press, Chicago (1988)