

BLOW UP OF SOLUTIONS OF SEMILINEAR HEAT EQUATIONS IN GENERAL DOMAINS

VALERIA MARINO, FILOMENA PACELLA, AND BERARDINO SCIUNZI

ABSTRACT. Consider the nonlinear heat equation $v_t - \Delta v = |v|^{p-1}v$ in a bounded smooth domain $\Omega \subset \mathbb{R}^n$ with $n > 2$ and Dirichlet boundary condition. Given u_p a sign-changing stationary solution fulfilling suitable assumptions, we prove that the solution with initial value ϑu_p blows up in finite time if $|\vartheta - 1| > 0$ is sufficiently small and if p is sufficiently close to the critical exponent.

Since for $\vartheta = 1$ the solution is global, this shows that, in general, the set of the initial data for which the solution is global is not star-shaped. This phenomenon had been previously observed in the case when the domain is a ball and the stationary solution is radially symmetric.

1. INTRODUCTION

We consider a nonlinear heat equation of the type

$$(1.1) \quad \begin{cases} v_t - \Delta v = |v|^{p-1}v & \text{in } \Omega \times (0, T) \\ v = 0 & \text{on } \partial\Omega \times (0, T) \\ v(0) = v_0(x) & \text{in } \Omega \end{cases}$$

where $\Omega \subset \mathbb{R}^n$, $n \in \mathbb{N}$, is a bounded domain, $p > 1$, $T \in (0, +\infty]$ and

$$v_0 \in C_0(\Omega) = \{v \in C(\overline{\Omega}), v(x) = 0 \text{ for } x \in \partial\Omega\}.$$

It is well known that the initial value problem (1.1) is locally well posed in $C_0(\Omega)$. Denoting with T_{v_0} the maximal existence time of the solution of (1.1) with initial datum v_0 , we consider the set of the initial data for which the corresponding solution is global, namely:

$$\mathcal{G} = \{v_0 \in C_0(\Omega), T_{v_0} = \infty\}.$$

It is interesting to understand the geometrical properties of the set \mathcal{G} . If we consider $v_0 = \vartheta w$, with $w \in C_0(\Omega)$ and $\vartheta \in \mathbb{R}$, it is well known that if $|\vartheta|$ is small enough the solution of (1.1) with initial datum ϑw , exists globally. Moreover, if $|\vartheta|$ is sufficiently large, it is easy to see that the solution blows up in finite time as a consequence of the fact that it has negative energy (see [6] and [1]). It is interesting to understand what happens for intermediate values of ϑ . The case when w is positive is completely clear, as a matter of fact from the maximum principle for the heat equation it follows that there exists $\tilde{\vartheta} > 0$ such that if $0 < \vartheta < \tilde{\vartheta}$ then the solution with initial value ϑw is

Key words and phrases. Semilinear heat equation, finite-time blowup, sign-changing stationary solutions, linearized operator, asymptotic behavior.

F. P. was partially supported by PRIN 2009-WRJ3W7 grant(Italy).

B. S. was partially supported by ERC-2011-grant: *Epsilon* and PRIN-2011: *Var. and Top. Met.*

2010 Mathematics Subject Classification: 35K91, 35B35, 35B44, 35J91.

globally defined, while if $\vartheta > \tilde{\vartheta}$ it blows up in finite time. In the borderline case both global existence or blow up in finite time can occur.

Thus, if we define $\mathcal{G}^+ = \{v_0 \in \mathcal{G}, v_0 \geq 0\}$, we can assert that \mathcal{G}^+ is star-shaped with respect to 0 (indeed it is a convex set). When the initial value changes sign the situation is different and, in general, the set \mathcal{G} may be not star-shaped. In fact, if we define by u_p a radial sign changing solution of the stationary problem

$$(1.2) \quad \begin{cases} -\Delta u_p = |u_p|^{p-1} u_p & \text{in } \Omega \\ u_p = 0 & \text{on } \partial\Omega \end{cases}$$

where Ω is the unit ball in \mathbb{R}^n , with $n > 2$ and $p > 1$, it has been shown in [4] that there exists $p^* < p_S$, with $p_S = \frac{n+2}{n-2}$ and there exists $\epsilon > 0$ such that if $p^* < p < p_S$ and $0 < |1 - \vartheta| < \epsilon$ then $\vartheta u_p \notin \mathcal{G}$ i.e. the solution of (1.1), with initial datum ϑu_p , blows up in finite time both for ϑ slightly greater and slightly smaller than 1. Hence \mathcal{G} is not star-shaped since $u_p \in \mathcal{G}$.

Recently a similar result has been proved in [5] in the case when the dimension is two and the exponent p is sufficiently large.

Such a result does not hold in the case $n = 1$ (always considering $p > 1$). As a matter of fact in the one-dimensional case we have that for $|\vartheta| < 1$, $v_{\vartheta,p}$ (the solution with initial value ϑu_p) is global and converges uniformly to zero, while it blows up in finite time if $|\vartheta| > 1$.

The proofs of the results of [4] and [5] exploit strongly the radial symmetry of the stationary solutions. Hence it is natural to ask whether a similar result holds also in general domains and what kind of sign changing stationary solutions give rise to this phenomenon. Note that this cannot be true for any sign changing stationary solution as it is easy to see considering, for example, a nodal solution in the ball which is odd with respect to a symmetry hyperplane and has only two nodal domains.

Here we show that, in the case when $n > 2$ and for exponents close to the critical one, the same blow up phenomenon occurs in any bounded domain considering a suitable class of sign changing solutions u_p of (1.2).

More precisely we deal with solutions u_p of (1.2) with the following properties:

- (a) $\int_{\Omega} |\nabla u_p|^2 dx \rightarrow 2S^{\frac{n}{2}}$ as $p \rightarrow p_S$,
- (b) $\frac{\max u_p}{\min u_p} \rightarrow -\infty$ as $p \rightarrow p_S$,

where S is the best Sobolev constant for the embedding of $H_0^1(\Omega)$ into $L^{2^*}(\Omega)$. It has been proved in [9] that such solutions exist, assuming that Ω is a bounded smooth domain in \mathbb{R}^n with $n > 2$, symmetric with respect to the x_i -coordinates ($i = 1, \dots, n$). Later in [8] the authors extend the same result to any general bounded and smooth domain in \mathbb{R}^n , with $n > 2$. Moreover in [3] it has been proved that condition (a) implies that $\Omega \setminus \{x \in \Omega \mid u_p(x) = 0\}$ has exactly two connected components while, when $n \geq 4$, (b) implies that the nodal surface of u_p does not intersect the boundary $\partial\Omega$ and the positive part u_p^+ and the negative part u_p^- concentrate at the same point. One could easily verify that (a) is equivalent to

$$E_p(u_p) = \frac{1}{2} \int_{\Omega} |\nabla u_p|^2 dx - \frac{1}{p+1} \int_{\Omega} |u_p|^{p+1} dx \rightarrow \frac{2}{n} S^{\frac{n}{2}} \quad \text{as } p \rightarrow p_S.$$

We refer to [3] for further properties of such solutions.

Our goal is to prove the following theorem.

Theorem 1.1. *Given problem (1.1) with $n > 2$, $1 < p < p_S = \frac{n+2}{n-2}$, and Ω a bounded smooth domain in \mathbb{R}^n , there exists $p^* < p_S$ with the following property:*

if $p^ < p < p_S$ and u_p is a sign changing solution of the stationary problem (1.2) satisfying (a) and (b) then there exist $0 < \underline{\vartheta} < 1 < \overline{\vartheta}$ such that if $\underline{\vartheta} < \vartheta < \overline{\vartheta}$ and $\vartheta \neq 1$ then $v_{\vartheta,p}$, solution of (1.1) with initial value ϑu_p , blows up in finite time.*

To prove Theorem 1.1 we use the following result which has been proved in [4] for general domains.

Proposition 1.1. *Let u_p be a sign changing solution of (1.2) and let $\varphi_{1,p}$ be a first eigenfunction of the linearized operator L_p at u_p . Assume that*

$$\int_{\Omega} u_p \varphi_{1,p} \neq 0.$$

Then there exists $\varepsilon > 0$ such that if $0 < |1 - \vartheta| < \varepsilon$, then $v_{\vartheta,p}$, solution of (1.1) with initial value ϑu_p , blows up in finite time.

Thus Theorem 1.1 will be a consequence of the following

Theorem 1.2. *Let $n > 2$, $1 < p < p_S$, $\Omega \subset \mathbb{R}^n$ a bounded smooth domain and u_p a sign changing solution of (1.2) satisfying conditions (a) and (b). Then there exists $p^* < p_S$ such that for $p^* < p < p_S$*

$$(1.3) \quad \int_{\Omega} u_p \varphi_{1,p} dx > 0,$$

where $\varphi_{1,p}$ is the first positive eigenfunction of the linearized operator L_p at u_p .

Let us point out that for the proof of Theorem 1.2 the property (b) of our stationary solutions is crucial. Note that both properties (a) and (b) are actually satisfied in the special case of radial sign changing solutions of (1.2) (in the ball) with two nodal regions.

So this clarifies that it is neither the symmetry nor the one-dimensional character of the solution which leads to the blow up result obtained in [4] but rather these properties of the stationary solution that can hold in any bounded domain. Therefore we believe that also for other semilinear problems where such solutions exist, the same blow up result should be true.

The proof of Theorem 1.2 is based on a rescaling argument about the maximum point of u_p . Indeed, analyzing the asymptotic behavior of the rescaled solutions and of the rescaled first eigenfunctions, we are able to prove (1.3) by using the properties of the solutions of the limit problem.

The same result of Theorem 1.2 can be easily extended to the case when the initial datum is a nodal solution $u_{p,\mathcal{K}}$ of (1.2) with a fixed number $\mathcal{K} > 2$ of nodal regions satisfying:

$$(a)_{\mathcal{K}} \quad \int_{\Omega} |\nabla u_{p,\mathcal{K}}|^2 dx \leq C,$$

(b) $_{\mathcal{K}}$ \exists a nodal region Ω_p^1 such that, setting

$$u_{p,\mathcal{K}}^1 := u_{p,\mathcal{K}} \cdot \chi_{\Omega_p^1} \quad \text{and} \quad \hat{u}_{p,\mathcal{K}} := u_{p,\mathcal{K}} \cdot \chi_{\Omega \setminus \Omega_p^1}$$

then

$$\int_{\Omega_p^1} |\nabla u_{p,\mathcal{K}}^1|^2 dx \rightarrow S^{\frac{n}{2}} \quad \text{as } p \rightarrow p_S$$

and

$$\frac{\|u_{p,\mathcal{K}}^1\|_{\infty}}{\|\hat{u}_{p,\mathcal{K}}^1\|_{\infty}} \rightarrow \infty \quad \text{as } p \rightarrow p_S.$$

Solutions of this type have been found in [8, 9] but other kind of solutions could be considered.

The outline of the proof is the following. In Section 2 we prove some preliminary results, while in Section 3 we study the asymptotic behavior of the first eigenvalue and of the first eigenfunction of the linearized operator at u_p . Finally in Section 4 we prove Theorem 1.2.

2. PRELIMINARIES

Let us start by recalling some properties of our solutions.

Lemma 2.1. *Let (u_p) be a family of sign-changing solutions of (1.2) satisfying (a). Then*

$$\begin{aligned} (i) \quad & \int_{\Omega} |\nabla u_p^+|^2 dx \xrightarrow{p \rightarrow p_S} S^{\frac{n}{2}}, \quad \int_{\Omega} |\nabla u_p^-|^2 dx \xrightarrow{p \rightarrow p_S} S^{\frac{n}{2}}, \\ (ii) \quad & \int_{\Omega} (u_p^+)^{\frac{2n}{n-2}} dx \xrightarrow{p \rightarrow p_S} S^{\frac{n}{2}}, \quad \int_{\Omega} (u_p^-)^{\frac{2n}{n-2}} dx \xrightarrow{p \rightarrow p_S} S^{\frac{n}{2}}, \\ (iii) \quad & u_p \rightharpoonup 0 \text{ as } p \rightarrow p_S, \\ (iv) \quad & M_{p,+} := \max_{\Omega} u_p^+ \xrightarrow{p \rightarrow p_S} +\infty, \quad M_{p,-} := \max_{\Omega} u_p^- \xrightarrow{p \rightarrow p_S} +\infty, \end{aligned}$$

with $u_p^+ = \max_{\Omega}(u_p, 0)$ and $u_p^- = \max_{\Omega}(-u_p, 0)$.

Proof. We refer the reader to [2, Lemma 2.1]. \square

We now describe the rescaled problem. Let us define

$$(2.1) \quad \tilde{u}_p(x) := \frac{1}{M_p} u_p \left(a_p + \frac{x}{M_p^{\frac{p-1}{2}}} \right), \quad \text{for } x \in \tilde{\Omega}_p := M_p^{\frac{p-1}{2}} (\Omega - a_p).$$

where a_p and M_p are such that $|u_p(a_p)| = \|u_p\|_{L^{\infty}(\Omega)} =: M_p$. Without loss of generality, we can assume that $u_p(a_p) > 0$.

Let us consider the limit problem in \mathbb{R}^n , that is

$$(2.2) \quad \begin{cases} -\Delta u = |u|^{p_S-1} u = |u|^{\frac{4}{n-2}} u & \text{in } \mathbb{R}^n \\ u(0) = 1. \end{cases}$$

It is well known that the unique regular positive solution is radial and is given by

$$U(x) = \left(\frac{n(n-2)}{n(n-2) + |x|^2} \right)^{\frac{n-2}{2}} \quad \text{for } x \in \mathbb{R}^n.$$

Moreover any sign changing solution of (2.2) has energy larger than $2S^{\frac{n}{2}}$. We have

Lemma 2.2. *For $p \rightarrow p_S$*

$$\tilde{u}_p \longrightarrow U \text{ in } C_{loc}^2(\mathbb{R}^n).$$

Proof. The proof is the same (with obvious changes) as the one of the similar statement in Theorem 1.1 of [2] (see page 777). \square

Now we study the linearization of the limit problem (2.2), so we define the operator

$$L^*(v) := -\Delta v - p_S |U|^{p_S-1} v, \quad v \in H^2(\mathbb{R}^n)$$

where U is the solution of (2.2). The Rayleigh functional associated to L^* is

$$\mathcal{R}(v) = \int_{\mathbb{R}^n} |\nabla v|^2 - p_S |U|^{p_S-1} v^2 dx$$

and we define

$$(2.3) \quad \lambda_1^* := \inf_{\substack{v \in H^1(\mathbb{R}^n), \\ \|v\|_{L^2(\mathbb{R}^n)}=1}} \mathcal{R}(v).$$

We observe that $\lambda_1^* > -\infty$, since U is bounded.

Remark 2.1. *It can be shown, with standard arguments, that there exists a unique positive minimizer φ_1^* to (2.3) which is radial and radially nonincreasing; moreover λ_1^* is an eigenvalue of L^* and φ_1^* is an eigenvector associated to λ_1^* . For further details see [7].*

Proposition 2.2. *We have the following.*

- (i) $\lambda_1^* < 0$,
- (ii) *every minimizing sequence of (2.3) has a subsequence which strongly converges in $L^2(\mathbb{R}^n)$.*

Proof. Let us compute \mathcal{R} on $U \in H^1(\mathbb{R}^n)$, solution of the limit problem (2.2). We have

$$\begin{aligned} \mathcal{R}(U) &= \int_{\mathbb{R}^n} |\nabla U|^2 - p_S |U|^{p_S+1} dx \\ &= (1 - p_S) \int_{\mathbb{R}^n} |\nabla U|^2 dx < 0 \end{aligned}$$

since $p_S > 1$. By definition (2.3) this implies that $\lambda_1^* < 0$. To prove (ii) let us consider a sequence $w_n \in H^1(\mathbb{R}^n)$, with $\|w_n\|_{L^2(\mathbb{R}^n)} = 1$, which minimizes (2.3). It is easy to see that w_n is bounded in $H^1(\mathbb{R}^n)$; therefore, up to a subsequence, it converges weakly to some $w \in H^1(\mathbb{R}^n)$ and strongly in $L^2(\{|x| \leq R\})$ for every $R > 0$.

By the lower semicontinuity of the norm we have

$$\int_{\mathbb{R}^n} |\nabla w|^2 dx \leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^n} |\nabla w_n|^2 dx \quad \text{and} \quad \|w\|_{L^2(\mathbb{R}^n)} \leq 1.$$

Moreover, for every $\epsilon > 0$, we have

$$\begin{aligned}
& \left| \int_{\mathbb{R}^n} U^{ps-1}(w_n^2 - w^2) dx \right| \leq \int_{\mathbb{R}^n} U^{ps-1} |w_n^2 - w^2| dx \\
&= \int_{|x| \leq R} U^{ps-1} |w_n^2 - w^2| dx + \int_{|x| > R} U^{ps-1} |w_n^2 - w^2| dx \\
&\leq c \int_{|x| \leq R} |w_n^2 - w^2| dx + \frac{c}{R^4} \int_{|x| > R} |w_n^2 - w^2| dx \\
&\leq c \|w_n - w\|_{L^2(|x| \leq R)} + \frac{c}{R^4} \leq \epsilon
\end{aligned}$$

where the last estimate is possible if we fix R large enough and then we take n sufficiently large. Thus

$$\int_{\mathbb{R}^n} U^{ps-1} w_n^2 dx \rightarrow \int_{\mathbb{R}^n} U^{ps-1} w^2 dx, \quad \text{as } n \rightarrow \infty$$

and so

$$\begin{aligned}
\mathcal{R}(w) &= \int_{\mathbb{R}^n} |\nabla w|^2 - p_S |U|^{ps-1} w^2 dx \\
&\leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^n} |\nabla w_n|^2 - p_S |U|^{ps-1} w_n^2 dx = \lambda_1^*.
\end{aligned}$$

This implies that $w \neq 0$ and we can define:

$$\hat{w} = \frac{w}{\|w\|_{L^2(\mathbb{R}^n)}}.$$

If we assume now by contradiction that $\|w\|_{L^2(\mathbb{R}^n)}^2 < 1$, it follows that

$$(2.4) \quad \lambda_1^* \leq \mathcal{R}(\hat{w}) = \frac{\mathcal{R}(w)}{\|w\|_{L^2(\mathbb{R}^n)}^2} \leq \frac{\lambda_1^*}{\|w\|_{L^2(\mathbb{R}^n)}^2} < \lambda_1^*,$$

since $\lambda_1^* < 0$. By (2.4) we deduce therefore that $\|w\|_{L^2(\mathbb{R}^n)}^2 = 1$ and so w is a minimizer. This also allows us to deduce that w_n strongly converges to w in $L^2(\mathbb{R}^n)$, and this concludes the proof of (ii). \square

3. ASYMPTOTIC SPECTRAL ANALYSIS

We consider the linearized operator at u_p , that is:

$$L_p = -\Delta - p|u_p|^{p-1}I.$$

We denote by $\lambda_{1,p}$ the first eigenvalue of L_p in Ω and by $\varphi_{1,p}$ the corresponding positive eigenfunction such that $\varphi_{1,p} > 0$ and $\|\varphi_{1,p}\|_{L^2(\Omega)} = 1$. We have

$$(3.1) \quad -\Delta \varphi_{1,p} - p|u_p|^{p-1} \varphi_{1,p} = \lambda_{1,p} \varphi_{1,p} \quad \text{in } \Omega.$$

Let us define $\tilde{\varphi}_{1,p}$ by

$$\tilde{\varphi}_{1,p}(x) = \left(\frac{1}{M_p^{\frac{p-1}{2}}} \right)^{\frac{n}{2}} \varphi_{1,p} \left(a_p + \frac{x}{M_p^{\frac{p-1}{2}}} \right) \quad \text{in } \tilde{\Omega}_p,$$

and $\tilde{\varphi}_{1,p} = 0$ outside $\tilde{\Omega}_p$. It is easy to see that $\|\tilde{\varphi}_{1,p}\|_{L^2(\mathbb{R}^n)} = 1$ and $\tilde{\varphi}_{1,p}$ satisfies

$$-\Delta \tilde{\varphi}_{1,p} - V_p \tilde{\varphi}_{1,p} = \tilde{\lambda}_{1,p} \tilde{\varphi}_{1,p}$$

where

$$V_p(x) = p \frac{1}{M_p^{p-1}} \left| u_p \left(a_p + \frac{x}{M_p^{\frac{p-1}{2}}} \right) \right|^{p-1} = p |\tilde{u}_p(x)|^{p-1}$$

and

$$\tilde{\lambda}_{1,p} = \frac{\lambda_{1,p}}{M_p^{p-1}}.$$

This means that $\tilde{\varphi}_{1,p}$ is a first eigenfunction of the operator

$$\tilde{L}_p = -\Delta - p |\tilde{u}_p|^{p-1} I$$

and $\tilde{\lambda}_{1,p}$ is the corresponding first eigenvalue.

Lemma 3.1. *The set $\{\tilde{\varphi}_{1,p}, 1 < p < p_S\}$ is bounded in $H^1(\mathbb{R}^n)$.*

Proof. As we have already remarked $\|\tilde{\varphi}_{1,p}\|_{L^2(\mathbb{R}^n)} = 1$. Moreover, since $\lambda_{1,p} < 0$ and $p < p_S$, we get

$$\begin{aligned} \int_{\mathbb{R}^n} |\nabla \tilde{\varphi}_{1,p}(x)|^2 dx &= \frac{1}{M_p^{p-1}} \int_{\tilde{\Omega}_p} \left(\frac{1}{M_p^{\frac{p-1}{2}}} \right)^n \left| \nabla \varphi_{1,p} \left(a_p + \frac{x}{M_p^{\frac{p-1}{2}}} \right) \right|^2 dx \\ &= \frac{1}{M_p^{p-1}} \int_{\Omega} |\nabla \varphi_{1,p}(x)|^2 dx \\ &= \frac{1}{M_p^{p-1}} \int_{\Omega} p |u_p|^{p-1} \varphi_{1,p}^2 dx + \frac{\lambda_{1,p}}{M_p^{p-1}} \int_{\Omega} \varphi_{1,p}^2 dx \\ &\leq \int_{\Omega} p \left(\frac{|u_p|}{M_p} \right)^{p-1} \varphi_{1,p}^2 dx \\ &\leq p \int_{\Omega} \varphi_{1,p}^2 dx < p_S, \end{aligned}$$

i.e. the assertion. □

Theorem 3.2. *We have*

$$(3.2) \quad \tilde{\lambda}_{1,p} \rightarrow \lambda_1^* \quad \text{as } p \rightarrow p_S.$$

Proof. We divide the proof in two steps:

Step 1.: We show that for $\epsilon > 0$ we have

$$(3.3) \quad \lambda_1^* \leq \tilde{\lambda}_{1,p} + \epsilon \quad \text{for } p \text{ sufficiently close to } p_S.$$

By (2.3), we have $\lambda_1^* \leq \mathcal{R}(\tilde{\varphi}_{1,p})$. Thus

$$\begin{aligned} \lambda_1^* &\leq \int_{\mathbb{R}^n} |\nabla \tilde{\varphi}_{1,p}|^2 - p_S |U|^{p_S-1} \tilde{\varphi}_{1,p}^2 dx \\ &= \int_{\tilde{\Omega}_p} |\nabla \tilde{\varphi}_{1,p}|^2 - p |\tilde{u}_p|^{p-1} \tilde{\varphi}_{1,p}^2 dx - \int_{\tilde{\Omega}_p} (p_S |U|^{p_S-1} - p |\tilde{u}_p|^{p-1}) \tilde{\varphi}_{1,p}^2 dx \\ &= \tilde{\lambda}_{1,p} - \int_{\tilde{\Omega}_p} (p_S |U|^{p_S-1} - p |\tilde{u}_p|^{p-1}) \tilde{\varphi}_{1,p}^2 dx \\ &= \tilde{\lambda}_{1,p} - \int_{\tilde{\Omega}_p \cap |x| \leq R} (p_S |U|^{p_S-1} - p |\tilde{u}_p|^{p-1}) \tilde{\varphi}_{1,p}^2 dx + \\ &\quad - \int_{\tilde{\Omega}_p \cap |x| > R} (p_S |U|^{p_S-1} - p |\tilde{u}_p|^{p-1}) \tilde{\varphi}_{1,p}^2 dx \end{aligned}$$

where $R > 0$. Let us first consider the last integral. We want to show that it can be made arbitrarily small. We have

$$(3.4) \quad \begin{aligned} \left| \int_{\tilde{\Omega}_p \cap |x| > R} p_S |U|^{p_S-1} \tilde{\varphi}_{1,p}^2 dx \right| &\leq p_S \int_{\tilde{\Omega}_p \cap |x| > R} |U|^{p_S-1} \tilde{\varphi}_{1,p}^2 dx \\ &\leq \frac{C_1}{R^4} \int_{\mathbb{R}^n} \tilde{\varphi}_{1,p}^2 dx \leq \frac{C_1}{R^4} \end{aligned}$$

for some constant $C_1 > 0$. Therefore we can choose R so large that

$$\left| \int_{\tilde{\Omega}_p \cap |x| > R} p_S |U|^{p_S-1} \tilde{\varphi}_{1,p}^2 dx \right| \leq \epsilon.$$

To estimate the term

$$\left| \int_{\tilde{\Omega}_p \cap |x| > R} p |\tilde{u}_p|^{p-1} \tilde{\varphi}_{1,p}^2 dx \right|$$

note that we can split the integral on $\tilde{\Omega}_p$ in the integral on

$$(3.5) \quad \tilde{\Omega}_p^+ = \{x \in \tilde{\Omega}_p : \tilde{u}_p(x) \geq 0\}$$

and the one on

$$(3.6) \quad \tilde{\Omega}_p^- = \{x \in \tilde{\Omega}_p : \tilde{u}_p(x) < 0\}.$$

Therefore we get

$$(3.7) \quad \begin{aligned} &\left| \int_{\tilde{\Omega}_p \cap |x| > R} p |\tilde{u}_p|^{p-1} \tilde{\varphi}_{1,p}^2 dx \right| \leq \\ &\int_{\tilde{\Omega}_p^+ \cap |x| > R} p |\tilde{u}_p|^{p-1} \tilde{\varphi}_{1,p}^2 dx + \int_{\tilde{\Omega}_p^- \cap |x| > R} p |\tilde{u}_p|^{p-1} \tilde{\varphi}_{1,p}^2 dx. \end{aligned}$$

As for the first term of (3.7) we have

$$(3.8) \quad \begin{aligned} &\int_{\tilde{\Omega}_p^+ \cap |x| > R} p |\tilde{u}_p|^{p-1} \tilde{\varphi}_{1,p}^2 dx \\ &\leq p \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx \right)^{\frac{2}{n}} \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} \tilde{\varphi}_{1,p}^{\frac{2n}{n-2}} dx \right)^{\frac{n-2}{n}} \\ &\leq p \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx \right)^{\frac{2}{n}} \|\tilde{\varphi}_{1,p}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}^n)}^2 \\ &\leq C_2 \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx \right)^{\frac{2}{n}} \end{aligned}$$

where we have used Hölder's inequality (with exponents $\frac{n}{2}$ and $\frac{n}{n-2}$) for the first estimate and the fact that, as a consequence of Lemma 3.1, $\tilde{\varphi}_{1,p}$ is bounded in $L^{\frac{2n}{n-2}}(\mathbb{R}^n)$ to obtain the last inequality.

In order to estimate the last term in (3.8), we use (ii) of Lemma 2.1

to get

$$\begin{aligned} & \int_{\tilde{\Omega}_p^+ \cap |x| \leq R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx + \int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx = \\ & = \int_{\tilde{\Omega}_p^+} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx \xrightarrow{p \rightarrow p_S} S^{\frac{n}{2}} = \int_{\mathbb{R}^n} |U|^{\frac{2n}{n-2}} dx = \\ & \int_{|x| \leq R} |U|^{\frac{2n}{n-2}} dx + \int_{|x| > R} |U|^{\frac{2n}{n-2}} dx \end{aligned}$$

As $\tilde{u}_p \xrightarrow{p \rightarrow p_S} U$ in $C_{loc}^2(\mathbb{R}^n)$, we have

$$\int_{\tilde{\Omega}_p^+ \cap |x| \leq R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx \xrightarrow{p \rightarrow p_S} \int_{|x| \leq R} |U|^{\frac{2n}{n-2}} dx$$

and so

$$(3.9) \quad \int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx \xrightarrow{p \rightarrow p_S} \int_{|x| > R} |U|^{\frac{2n}{n-2}} dx$$

but, as $U \in L^{\frac{2n}{n-2}}(\mathbb{R}^n)$, the term on the right hand side of (3.9) can be made as small as we like, choosing R sufficiently large. Thus we have that, chosen R large enough, we can take p sufficiently close to p_S so that

$$(3.10) \quad \int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{n(p-1)}{2}} dx \leq \epsilon.$$

Let us now estimate the second term of (3.7)

$$\begin{aligned} & \int_{\tilde{\Omega}_p^- \cap |x| > R} p |\tilde{u}_p|^{p-1} \tilde{\varphi}_{1,p}^2 dx \\ & \leq p \left(\frac{\|u_p^-\|_{L^\infty(\Omega)}}{\|u_p^+\|_{L^\infty(\Omega)}} \right)^{p-1} \left(\int_{\tilde{\Omega}_p^- \cap |x| > R} \tilde{\varphi}_{1,p}^2 dx \right) \\ & \leq p \left(\frac{\|u_p^-\|_{L^\infty(\Omega)}}{\|u_p^+\|_{L^\infty(\Omega)}} \right)^{p-1} \xrightarrow{p \rightarrow p_S} 0 \end{aligned}$$

where we used the fact that $\|\tilde{\varphi}_{1,p}\|_{L^2(\mathbb{R}^n)} = 1$ and condition (b) satisfied by our solutions.

Recalling that $\tilde{u}_p \xrightarrow{p \rightarrow p_S} U$ in $C_{loc}^2(\mathbb{R}^n)$, for R fixed as above and p sufficiently close to p_S , we have

$$(3.11) \quad \int_{\tilde{\Omega}_p \cap |x| \leq R} (p_S |U|^{p_S-1} - p |\tilde{u}_p|^{p-1}) \tilde{\varphi}_{1,p}^2 dx \leq \epsilon.$$

Thus (3.3) follows from (3.4)-(3.11).

Step 2.: Now we show that for $\epsilon > 0$ we have

$$(3.12) \quad \tilde{\lambda}_{1,p} \leq \lambda_1^* + \epsilon \quad \text{for } p \text{ sufficiently close to } p_S.$$

Let us consider a regular cut-off function $\psi_R(x) = \psi_R(r)$, for $R > 0$, such that

- $0 \leq \psi_R \leq 1$ and $\psi_R(r) = 1$ for $r \leq R$, $\psi_R(r) = 0$ for $r \geq 2R$,
- $|\nabla \psi_R| \leq \frac{2}{R}$

and let us set

$$w_R := \frac{\psi_R \varphi_1^*}{\|\psi_R \varphi_1^*\|_{L^2(\mathbb{R}^n)}}.$$

Thus

$$\begin{aligned} (3.13) \quad \tilde{\lambda}_{1,p} &\leq \int_{\mathbb{R}^n} |\nabla w_R|^2 - p|\tilde{u}_p|^{p-1} w_R^2 dx \\ &= \int_{\mathbb{R}^n} |\nabla w_R|^2 - p_S |U|^{p_S-1} w_R^2 dx \\ &\quad + \int_{\mathbb{R}^n} (p_S |U|^{p_S-1} - p|\tilde{u}_p|^{p-1}) w_R^2 dx. \end{aligned}$$

It is easy to see that $w_R \rightarrow \varphi_1^*$ in $H^1(\mathbb{R}^n)$ as $R \rightarrow \infty$. Therefore, by (2.3), we have that given $\epsilon > 0$ we can fix $R > 0$ such that

$$\int_{\mathbb{R}^n} |\nabla w_R|^2 - p_S |U|^{p_S-1} w_R^2 dx \leq \lambda_1^* + \epsilon.$$

For such a fixed value of R , arguing as in Step 1, we obtain that

$$(3.14) \quad \int_{\mathbb{R}^n} (p_S |U|^{p_S-1} - p|\tilde{u}_p|^{p-1}) w_R^2 dx \leq \epsilon$$

for p close enough to p_S . Then (3.12) follows from (3.13)-(3.14).

By (3.3) and (3.12) we deduce (3.2). □

Corollary 3.1. $\tilde{\varphi}_{1,p}$ strongly converges to φ_1^* in $L^2(\mathbb{R}^n)$.

Proof. By the definition of $\tilde{\lambda}_{1,p}$, and what is stated in Theorem 3.2, we have

$$\int_{\tilde{\Omega}_p} |\nabla \tilde{\varphi}_{1,p}|^2 - p|U|^{p-1} \tilde{\varphi}_{1,p}^2 dx = \tilde{\lambda}_{1,p} \rightarrow \lambda_1^* \quad \text{as } p \rightarrow p_S.$$

This implies that $\tilde{\varphi}_{1,p}$ is a minimizing sequence for (2.3), and so the assertion follows by Proposition 2.2 (see also Remark 2.1). □

4. PROOF OF THEOREM 1.2

We now proceed proving Theorem 1.2.

Proof of Theorem 1.2. Using $\varphi_{1,p} \in H_0^1(\Omega)$ as a test function in (1.2) we have

$$(4.1) \quad \int_{\Omega} \nabla u_p \cdot \nabla \varphi_{1,p} dx = \int_{\Omega} |u_p|^{p-1} u_p \varphi_{1,p} dx,$$

while using u_p as a test function in (3.1) we obtain

$$(4.2) \quad \int_{\Omega} \nabla u_p \cdot \nabla \varphi_{1,p} dx = \int_{\Omega} p|u_p|^{p-1} u_p \varphi_{1,p} dx + \lambda_{1,p} \int_{\Omega} u_p \varphi_{1,p} dx.$$

Subtracting (4.1) from (4.2) we get

$$-\frac{p-1}{\lambda_{1,p}} \int_{\Omega} |u_p|^{p-1} u_p \varphi_{1,p} dx = \int_{\Omega} u_p \varphi_{1,p} dx.$$

Taking into account that $\lambda_{1,p}$ is negative, we have that, to determine the sign of $\int_{\Omega} u_p \varphi_{1,p} dx$, we can study the sign of

$$(4.3) \quad \int_{\Omega} |u_p|^{p-1} u_p \varphi_{1,p} dx.$$

For convenience we consider

$$M_p^{(\frac{p-1}{2})\frac{n}{2}-p} \int_{\Omega} |u_p|^{p-1} u_p \varphi_{1,p} dx$$

which has the same sign of (4.3). Now we prove that

$$(4.4) \quad \begin{aligned} M_p^{(\frac{p-1}{2})\frac{n}{2}-p} \int_{\Omega} |u_p|^{p-1} u_p \varphi_{1,p} dx &\xrightarrow{p \rightarrow p_S} \int_{\mathbb{R}^n} |U|^{p_S-1} U \varphi_1^* dx \\ &= \int_{\mathbb{R}^n} U^{p_S} \varphi_1^* dx. \end{aligned}$$

Since the term on the right hand side of (4.4) is positive, this will leads to the assertion of Theorem 1.2.

By a simple change of variables it follows that:

$$(4.5) \quad \begin{aligned} &\left| M_p^{(\frac{p-1}{2})\frac{n}{2}-p} \int_{\Omega} |u_p|^{p-1} u_p \varphi_{1,p} dx - \int_{\mathbb{R}^n} |U|^{p_S-1} U \varphi_1^* dx \right| \\ &= \left| \int_{\tilde{\Omega}_p} |\tilde{u}_p|^{p-1} \tilde{u}_p \tilde{\varphi}_{1,p} dx - \int_{\mathbb{R}^n} |U|^{p_S-1} U \varphi_1^* dx \right|. \end{aligned}$$

We take $\epsilon > 0$ and choose $R > 0$ such that

$$\int_{|x|>R} |U|^{p_S-1} U \varphi_1^* dx = \int_{|x|>R} U^{p_S} \varphi_1^* dx \leq \epsilon,$$

this is possible arguing as we did in the proof of (3.4).

We rewrite (4.5) in the following way

$$\begin{aligned} &\left| \int_{\tilde{\Omega}_p \cap |x|>R} |\tilde{u}_p|^{p-1} \tilde{u}_p \tilde{\varphi}_{1,p} dx + \int_{\tilde{\Omega}_p \cap |x|\leq R} |\tilde{u}_p|^{p-1} \tilde{u}_p \tilde{\varphi}_{1,p} dx \right. \\ &- \left. \int_{|x|\leq R} |U|^{p_S-1} U \varphi_1^* dx - \int_{|x|>R} |U|^{p_S-1} U \varphi_1^* dx \right| \\ &\leq \left| \int_{\tilde{\Omega}_p \cap |x|>R} |\tilde{u}_p|^{p-1} \tilde{u}_p \tilde{\varphi}_{1,p} dx \right| + \left| \int_{|x|>R} |U|^{p_S-1} U \varphi_1^* dx \right| \\ &+ \left| \int_{\tilde{\Omega}_p \cap |x|\leq R} |\tilde{u}_p|^{p-1} \tilde{u}_p \tilde{\varphi}_{1,p} dx - \int_{|x|\leq R} |U|^{p_S-1} U \varphi_1^* dx \right|. \end{aligned}$$

Now we analyze each term in the previous inequality. Splitting the integral on $\tilde{\Omega}_p^+$ and on $\tilde{\Omega}_p^-$ (see (3.5) and (3.6) for the definitions of such sets) we have

$$(4.6) \quad \begin{aligned} &\left| \int_{\tilde{\Omega}_p \cap |x|>R} |\tilde{u}_p|^{p-1} \tilde{u}_p \tilde{\varphi}_{1,p} dx \right| \\ &\leq \int_{\tilde{\Omega}_p^+ \cap |x|>R} |\tilde{u}_p|^p \tilde{\varphi}_{1,p} dx + \int_{\tilde{\Omega}_p^- \cap |x|>R} |\tilde{u}_p|^p \tilde{\varphi}_{1,p} dx. \end{aligned}$$

As for the first term of (4.6) we have

$$\begin{aligned}
\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^p \tilde{\varphi}_{1,p} dx &\leq \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{2np}{n+2}} dx \right)^{\frac{n+2}{2n}} \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} \tilde{\varphi}_{1,p}^{\frac{2n}{n-2}} dx \right)^{\frac{n-2}{2n}} \\
&\leq \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{2np}{n+2}} dx \right)^{\frac{n+2}{2n}} \|\tilde{\varphi}_{1,p}\|_{L^{\frac{2n}{n-2}}(\mathbb{R}^n)} \\
&\leq C_4 \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{2np}{n+2}} dx \right)^{\frac{n+2}{2n}}
\end{aligned}$$

where we have used Hölder's inequality (with exponents $\frac{2n}{n+2}$ and $\frac{2n}{n-2}$) for the first estimate and the fact that, as a consequence of Lemma 3.1, $\tilde{\varphi}_{1,p}$ is bounded in $L^{\frac{2n}{n-2}}(\mathbb{R}^n)$.

Thus, with the same argument used to obtain (3.10), we can state that, for every $\epsilon > 0$, having chosen R large enough and taking p close enough to p_S , we have

$$C_4 \left(\int_{\tilde{\Omega}_p^+ \cap |x| > R} |\tilde{u}_p|^{\frac{2np}{n+2}} dx \right)^{\frac{n+2}{2n}} < \epsilon.$$

Next we estimate the second term of (4.6). We have

$$\begin{aligned}
&\int_{\tilde{\Omega}_p^- \cap |x| > R} |\tilde{u}_p|^p \tilde{\varphi}_{1,p} dx \\
&\leq \left(\int_{\tilde{\Omega}_p^- \cap |x| > R} |\tilde{u}_p|^{2p} dx \right)^{\frac{1}{2}} \left(\int_{\tilde{\Omega}_p^- \cap |x| > R} \tilde{\varphi}_{1,p}^2 dx \right)^{\frac{1}{2}} \\
&= \left(\int_{\tilde{\Omega}_p^- \cap |x| > R} |\tilde{u}_p|^{2p - \frac{2n}{n-2}} |\tilde{u}_p|^{\frac{2n}{n-2}} dx \right)^{\frac{1}{2}} \left(\int_{\tilde{\Omega}_p^- \cap |x| > R} \tilde{\varphi}_{1,p}^2 dx \right)^{\frac{1}{2}} \\
&\leq \left(\frac{\|u_p^-\|_{L^\infty(\Omega)}}{\|u_p^+\|_{L^\infty(\Omega)}} \right)^{p - \frac{n}{n-2}} \left(\int_{\tilde{\Omega}_p^- \cap |x| > R} |\tilde{u}_p|^{\frac{2n}{n-2}} dx \right)^{\frac{1}{2}} \\
&\leq C_5 \left(\frac{\|u_p^-\|_{L^\infty(\Omega)}}{\|u_p^+\|_{L^\infty(\Omega)}} \right)^{p - \frac{n}{n-2}} \xrightarrow{p \rightarrow p_S} 0
\end{aligned}$$

where we have used Hölder's inequality (with exponent 2) for the first estimate, the fact that $\|\tilde{\varphi}_{1,p}\|_{L^2(\mathbb{R}^n)} = 1$ for the second and condition (b) satisfied by our solution. Note in particular that, for p close to p_S , we may and do assume that $p > \frac{n}{n-2}$.

Moreover, recalling once again that $\tilde{u}_p \xrightarrow{p \rightarrow p_S} U$ in $C_{loc}^2(\mathbb{R}^n)$, we deduce that:

$$(4.7) \quad \left| \int_{\tilde{\Omega}_p \cap |x| \leq R} |\tilde{u}_p|^{p-1} \tilde{u}_p \tilde{\varphi}_{1,p} dx - \int_{|x| \leq R} |U|^{p_S-1} U \varphi_1^* dx \right| < \epsilon,$$

for R fixed as above and p sufficiently close to p_S .

Finally, for R sufficiently large, the term

$$\int_{|x| > R} |U|^{p_S-1} U \varphi_1^* dx$$

can be made arbitrary small since $U \in L^{\frac{2n}{n-2}}(\mathbb{R}^n)$ and φ_1^* is bounded.

Thus (4.5)-(4.7) and the arbitrary choice of ϵ imply (4.4) concluding the proof. □

REFERENCES

- [1] J. Ball. *Remarks on blow-up and nonexistence theorems for nonlinear evolution equations*, Quarterly Journal of Mathematics 28 (1977) 473-486.
- [2] M. Ben Ayed, K. El Mehdi, F. Pacella. *Blow-up and symmetry of sign-changing solutions to some critical elliptic equations*, Journal of Differential Equations 230 (2006) 771-795.
- [3] M. Ben Ayed, K. El Mehdi, F. Pacella. *Classification of low energy sign-changing solutions of an almost critical problem*, Journal of Functional Analysis 250 (2007) 347-373.
- [4] T. Cazenave, F. Dickstein, F.B. Weissler. *Sign-changing stationary solutions and blow up for the nonlinear heat equation in a ball*, Mathematische Annalen 344(2) (2009) 431-449.
- [5] F. Dickstein, F. Pacella, B. Sciunzi. *Sign-changing stationary solutions and blow up for the nonlinear heat equation in dimension two*, <http://arxiv.org/abs/1304.2571>.
- [6] H.A. Levine. *Some nonexistence and instability theorems for formally parabolic equations of the form $Pu_t = -Au + f(u)$* , Archive for Rational Mechanics and Analysis 51 (1973) 371-386.
- [7] E.H. Lieb, M.Loss. *Analysis*, Graduate Studies in Mathematics, 14, American Mathematical Society, 2001.
- [8] M. Musso, A. Pistoia. *Tower of Bubbles for almost critical problems in general domains*, Journal de Mathématiques Pures et Appliquées 93 (2010), 1-30.
- [9] A. Pistoia, T. Weth. *Sign changing bubble tower solutions in a slightly subcritical semilinear Dirichlet problem*, Annales de l'Institut Henri Poincaré 24 (2007) 325-340.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA "LA SAPIENZA", P.LE A. MORO 2, 00185 ROMA, ITALY
E-mail address: `valeria.marino.m@gmail.com`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA "LA SAPIENZA", P.LE A. MORO 2, 00185 ROMA, ITALY
E-mail address: `pacella@mat.uniroma1.it`

DIPARTIMENTO DI MATEMATICA, UNICAL, PONTE PIETRO BUCCI 31B, 87036 ARCAVACATA DI RENDE, COSENZA, ITALY
E-mail address: `sciunzi@mat.unical.it`