

A new A_4 model for lepton mixingP.M. Ferreira,^{(1,2)*} L. Lavoura,^{(3)†} and P.O. Ludl^{(4)‡}⁽¹⁾ Instituto Superior de Engenharia de Lisboa
1959-007 Lisbon, Portugal⁽²⁾ Centre for Theoretical and Computational Physics, University of Lisbon
1649-003 Lisbon, Portugal⁽³⁾ Universidade de Lisboa, Instituto Superior Técnico, CFTP
1049-001 Lisbon, Portugal⁽⁴⁾ University of Vienna, Faculty of Physics,
Boltzmanngasse 5, A-1090 Vienna, Austria

14 October 2013

Abstract

We present a new model of the lepton sector that uses a family symmetry A_4 to make predictions for lepton mixing which are invariant under any permutation of the three flavours. We show that those predictions broadly agree with the experimental data, leading to a largish $\sin^2 \theta_{12} \gtrsim 0.34$, to $|\cos \delta| \gtrsim 0.7$, and to $|0.5 - \sin^2 \theta_{23}| \gtrsim 0.08$; $\cos \delta$ and $0.5 - \sin^2 \theta_{23}$ are predicted to have identical signs.

*E-mail: ferreira@cii.fc.ul.pt

†E-mail: balio@cftp.ist.utl.pt

‡E-mail: patrick.ludl@univie.ac.at

The experimental discovery that the lepton mixing angle θ_{13} is nonzero [1] caused a profound change in the subject of flavour models for the lepton mass matrices. Many older models ceased to be valid. New models had to be built; some recent examples utilizing the horizontal symmetry group A_4 are collected in refs. [2, 3]. Many of those models use ‘flavons’ and non-renormalizable Lagrangians [2]; in most remaining models [3] there are Higgs doublets at the Fermi scale placed in triplets of the horizontal symmetry.¹ In this paper we present a model that contains only renormalizable terms and only Higgs doublets which are singlets of A_4 .

For any $n \times n$ non-singular matrix $M = [M_{\alpha\beta}]$, one may define a matrix $A = [A_{\alpha\beta}]$ through

$$A_{\alpha\beta} = M_{\alpha\beta} (M^{-1})_{\beta\alpha}, \quad (1)$$

where no sum over either α or β is implied. It is obvious from its definition that A satisfies

$$\sum_{\alpha=1}^n A_{\alpha\beta} = \sum_{\beta=1}^n A_{\alpha\beta} = 1. \quad (2)$$

The matrix A is invariant under

$$M \rightarrow XMY, \quad (3)$$

where X and Y are diagonal non-singular matrices.

For our purposes, M is the (effective) Majorana mass matrix of the three light neutrinos in the weak basis where the charged-lepton mass matrix is diagonal. Therefore, $n = 3$, the indices α and β are in the range $\{e, \mu, \tau\}$, M and A are symmetric, and $X = Y$ in the transformation (3).² The model in this paper predicts³

$$A_{e\mu} = A_{e\tau} = A_{\mu\tau}, \quad (5)$$

hence

$$A = \begin{pmatrix} 1-2t & t & t \\ t & 1-2t & t \\ t & t & 1-2t \end{pmatrix}, \quad (6)$$

¹An exception is the Babu–Ma–Valle model [4], in which the Higgs doublets are A_4 -invariant. That model depends on renormalization to produce realistic neutrino masses and mixings.

²The matrix A was also used in this context in ref. [5].

³Equations (5) may alternatively be stated as

$$M_{ee}(M_{\mu\tau})^2 = M_{\mu\mu}(M_{e\tau})^2 = M_{\tau\tau}(M_{e\mu})^2. \quad (4)$$

where t is in general a complex number. Moreover, through the imposition of an additional CP symmetry on our model, t may be made to be real. We shall show that the conditions (5) fit the experimental data rather well.⁴

Our model has the usual Standard-Model leptonic multiplets $\alpha_R : (\mathbf{1}, -1)$ and $D_{\alpha L} : (\mathbf{2}, -1/2)$.⁵ It has, besides, three right-handed neutrinos $\nu_{\alpha R} : (\mathbf{1}, 0)$. The scalar sector is composed of three Higgs doublets $\phi_k : (\mathbf{2}, 1/2)$, where $k \in \{1, 2, 3\}$. Their conjugate doublets are $\tilde{\phi}_k \equiv i\tau_2 \phi_k^* : (\mathbf{2}, -1/2)$. In our model there are, besides, three *real* scalars $\sigma_\alpha : (\mathbf{1}, 0)$.

The model is based on the well-known discrete symmetry group A_4 possessing the irreducible representations

$$\begin{aligned} \mathbf{1} : & \quad S \rightarrow 1, \quad T \rightarrow 1, \\ \mathbf{1}' : & \quad S \rightarrow 1, \quad T \rightarrow \omega, \\ \mathbf{1}'' : & \quad S \rightarrow 1, \quad T \rightarrow \omega^2, \\ \mathbf{3} : & \quad S \rightarrow \tilde{S}, \quad T \rightarrow \tilde{T}, \end{aligned} \tag{7}$$

where

$$\tilde{S} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \tilde{T} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \tag{8}$$

and $\omega = \exp(i2\pi/3)$. Assigning the fields to the representations as

$$\begin{pmatrix} D_{eL} \\ D_{\mu L} \\ D_{\tau L} \end{pmatrix}, \begin{pmatrix} e_R \\ \mu_R \\ \tau_R \end{pmatrix}, \begin{pmatrix} \nu_{eR} \\ \nu_{\mu R} \\ \nu_{\tau R} \end{pmatrix}, \begin{pmatrix} \sigma_e \\ \sigma_\mu \\ \sigma_\tau \end{pmatrix} : \mathbf{3} \quad \text{and} \quad \begin{aligned} \phi_1 : & \mathbf{1} \\ \phi_2 : & \mathbf{1}' \\ \phi_3 : & \mathbf{1}'' \end{aligned} \tag{9}$$

leads to the Yukawa Lagrangian

$$\begin{aligned} \mathcal{L}_{\text{Yukawa}} = & -y_1 (\bar{D}_{eL} e_R + \bar{D}_{\mu L} \mu_R + \bar{D}_{\tau L} \tau_R) \phi_1 \\ & -y_2 (\bar{D}_{eL} e_R + \omega \bar{D}_{\mu L} \mu_R + \omega^2 \bar{D}_{\tau L} \tau_R) \phi_2 \\ & -y_3 (\bar{D}_{eL} e_R + \omega^2 \bar{D}_{\mu L} \mu_R + \omega \bar{D}_{\tau L} \tau_R) \phi_3 \\ & -y_4 (\bar{D}_{eL} \nu_{eR} + \bar{D}_{\mu L} \nu_{\mu R} + \bar{D}_{\tau L} \nu_{\tau R}) \tilde{\phi}_1 \\ & -y_5 (\bar{D}_{eL} \nu_{eR} + \omega^2 \bar{D}_{\mu L} \nu_{\mu R} + \omega \bar{D}_{\tau L} \nu_{\tau R}) \tilde{\phi}_2 \\ & -y_6 (\bar{D}_{eL} \nu_{eR} + \omega \bar{D}_{\mu L} \nu_{\mu R} + \omega^2 \bar{D}_{\tau L} \nu_{\tau R}) \tilde{\phi}_3 \\ & -y_7 (\bar{\nu}_{eR} C \bar{\nu}_{\mu R}^T \sigma_\tau + \bar{\nu}_{\mu R} C \bar{\nu}_{\tau R}^T \sigma_e + \bar{\nu}_{\tau R} C \bar{\nu}_{eR}^T \sigma_\mu) + \text{H.c.}, \end{aligned} \tag{10}$$

⁴The two conditions (5) represent a total of *four* constraints (two from the real parts and two from the imaginary parts) on the neutrino masses and on lepton mixing. However, because those conditions implicitly involve the Majorana phases of the neutrinos, which are unobservable in oscillation experiments, the predictive power of our model is less than these four constraints might suggest.

⁵The boldface number inside each parentheses is the dimension of the gauge- $SU(2)$ representation; the second number is the weak hypercharge.

where C is the charge-conjugation matrix in Dirac space. The couplings y_{1-7} are dimensionless. There are also bare Majorana mass terms

$$\mathcal{L}_{\text{Majorana}} = -\frac{m}{2} (\bar{\nu}_{eR} C \bar{\nu}_{eR}^T + \bar{\nu}_{\mu R} C \bar{\nu}_{\mu R}^T + \bar{\nu}_{\tau R} C \bar{\nu}_{\tau R}^T) + \text{H.c.}, \quad (11)$$

where m has mass dimension.

When the neutral components of $\phi_{1,2,3}$ get vacuum expectation values (VEVs) $v_k = \langle 0 | \phi_k^0 | 0 \rangle$, the charged leptons acquire masses given by

$$\begin{aligned} m_e &= |y_1 v_1 + y_2 v_2 + y_3 v_3|, \\ m_\mu &= |y_1 v_1 + \omega y_2 v_2 + \omega^2 y_3 v_3|, \\ m_\tau &= |y_1 v_1 + \omega^2 y_2 v_2 + \omega y_3 v_3|. \end{aligned} \quad (12)$$

The three quantities $y_k v_k$ must be finetuned in order that $m_e \ll m_\mu \ll m_\tau$.⁶

We assume the VEVs of the three σ_α to be equal (see appendix A): $\langle 0 | \sigma_e | 0 \rangle = \langle 0 | \sigma_\mu | 0 \rangle = \langle 0 | \sigma_\tau | 0 \rangle \equiv s$. We furthermore assume that s is of the same order of magnitude as m , and that this order of magnitude is very large, *viz.* it is the seesaw scale. Thus, the subgroup $\mathbb{Z}_3$ of A_4 generated by T is preserved at the high (seesaw) scale and only gets spontaneously broken at the low (Fermi⁷) scale, when ϕ_2 and ϕ_3 acquire VEVs.

The neutrino mass matrices M_D and M_R are defined by

$$\mathcal{L}_{\nu \text{ mass}} = -\bar{\nu}_R M_D \nu_L - \frac{1}{2} \bar{\nu}_R M_R C \bar{\nu}_R^T + \text{H.c.} \quad (13)$$

In our model,

$$M_D = \text{diag}(a, b, c), \quad (14)$$

$$M_R = \begin{pmatrix} m & m' & m' \\ m' & m & m' \\ m' & m' & m \end{pmatrix}, \quad (15)$$

⁶Most models require a finetuning in order to obtain $m_e \ll m_\mu \ll m_\tau$. Possible exceptions are models based on the Froggatt–Nielsen paradigm [6] and models with extra dimensions.

⁷Another possibility is that $v_{1,2,3}$ are much lower than the Fermi scale, if the quarks do not have Yukawa couplings to $\phi_{1,2,3}$ and only couple to an extra doublet ϕ_0 which is invariant under A_4 . In that scheme, the VEV of the neutral component of ϕ_0 would be dominant in giving mass both to the gauge bosons and to the quarks, while the VEVs of the neutral components of $\phi_{1,2,3}$ would lie much below the Fermi scale. The masses of the components of the doublets $\phi_{1,2,3}$ would in this picture lie much *above* the Fermi scale, due to a type-II seesaw mechanism for Higgs doublets [7].

where

$$\begin{aligned} a &= y_4^* v_1 + y_5^* v_2 + y_6^* v_3, \\ b &= y_4^* v_1 + \omega y_5^* v_2 + \omega^2 y_6^* v_3, \\ c &= y_4^* v_1 + \omega^2 y_5^* v_2 + \omega y_6^* v_3, \end{aligned} \quad (16)$$

and $m' = y_7 s$. A seesaw mechanism takes place, whereupon an effective mass matrix for the light neutrinos

$$M = -M_D^T M_R^{-1} M_D \quad (17)$$

is generated. The matrix A is then of the form (6), with

$$t = \frac{m'^2}{(m' - m)(2m' + m)}. \quad (18)$$

The values of a , b , and c are irrelevant for A .

One may, if one wants, furnish our model with an extra CP symmetry,

$$CP : \quad \begin{cases} D_{eL}(x) \rightarrow \gamma_0 C \bar{D}_{eL}^T(\bar{x}), \\ D_{\mu L}(x) \rightarrow \gamma_0 C \bar{D}_{\mu L}^T(\bar{x}), \\ D_{\tau L}(x) \rightarrow \gamma_0 C \bar{D}_{\tau L}^T(\bar{x}), \end{cases} \quad \begin{cases} e_R(x) \rightarrow \gamma_0 C \bar{e}_R^T(\bar{x}), \\ \mu_R(x) \rightarrow \gamma_0 C \bar{\mu}_R^T(\bar{x}), \\ \tau_R(x) \rightarrow \gamma_0 C \bar{\tau}_R^T(\bar{x}), \end{cases} \quad (19)$$

$$\begin{cases} \nu_{eR}(x) \rightarrow \gamma_0 C \bar{\nu}_{eR}^T(\bar{x}), \\ \nu_{\mu R}(x) \rightarrow \gamma_0 C \bar{\nu}_{\mu R}^T(\bar{x}), \\ \nu_{\tau R}(x) \rightarrow \gamma_0 C \bar{\nu}_{\tau R}^T(\bar{x}), \end{cases} \quad \begin{cases} \sigma_e(x) \rightarrow \sigma_e(\bar{x}), \\ \sigma_\mu(x) \rightarrow \sigma_\mu(\bar{x}), \\ \sigma_\tau(x) \rightarrow \sigma_\tau(\bar{x}), \\ \phi_k(x) \rightarrow \phi_k^*(\bar{x}) \quad \forall k, \end{cases}$$

where $x = (t, \vec{r})$ and $\bar{x} = (t, -\vec{r})$. This CP symmetry renders $y_1, y_2, \dots, y_7$, and m real. The mass m' will then be real, because the scalars σ_α are Hermitian fields, hence their VEV s is real. Even if the CP symmetry is spontaneously broken by (relatively) complex v_1 , v_2 , and v_3 , the ensuing phases of a , b , and c may be withdrawn from M through appropriate rephasings of the light-neutrino fields. Thus, *there exists a restriction of our model in which M is real*.

We proceed to fit the predictions of our model, *viz.* eqs. (5), to the phenomenological data. The neutrino masses are $m_{1,2,3}$. We use the standard parameterization of lepton mixing in ref. [8], through three mixing angles θ_{12} , θ_{13} , and θ_{23} and one CP -violating phase δ . We have used the following

allowed ranges for the various observables:

$$\begin{aligned}
6.99 \times 10^{-5} \text{ eV}^2 &\leq m_2^2 - m_1^2 \leq 8.20 \times 10^{-5} \text{ eV}^2, \\
2.16 \times 10^{-3} \text{ eV}^2 &\leq |m_3^2 - m_1^2| \leq 2.74 \times 10^{-3} \text{ eV}^2, \\
0.25 &\leq \sin^2 \theta_{12} \leq 0.37, \\
0.016 &\leq \sin^2 \theta_{13} \leq 0.033, \\
0.33 &\leq \sin^2 \theta_{23} \leq 0.68.
\end{aligned} \tag{20}$$

These ranges simultaneously encompass all the corresponding 3σ ranges furnished by the relevant phenomenological analyses [9, 10, 11]. We stress that, even though the fit presented here uses these quite ample ranges, we have also found that *most observables easily fall within their respective 1σ ranges* given in, for instance, ref. [9]; the exception is the mixing angle θ_{12} , which is in our model rather large. We have left δ free, even though refs. [9, 11] provide some bounds on it, which are, however, valid only at the 1σ level.

Our first finding is that in our model the phase δ must be close to either 0 or π ;⁸ if $\delta \approx 0$ then θ_{23} is in the first octant while θ_{23} is in the second octant when $\delta \approx \pi$.⁹ This can be seen in the scatter plot of fig. 1. Therefrom one gathers that in our model $|\cos \delta| \gtrsim 0.7$ (0.8 if the neutrino mass spectrum is inverted) and $|\sin^2 \theta_{23} - 0.5| \gtrsim 0.08$.

In fig. 2 one sees that our model is unable to predict θ_{13} but neatly predicts $\sin^2 \theta_{12} \gtrsim 0.34$. This is a rather large value, which is allowed by the phenomenological data only at the 2σ (or even 3σ) level. It can moreover be seen in fig. 2 (and also in fig. 4) that the restricted real version of our model does not have much more predictive power than the general version, even though it has one degree of freedom less.

In the right panel of fig. 2 one observes that the *lower bound* on θ_{12} in our model coincides with the *prediction* for θ_{12} in a model with trimaximal mixing (TM₂ in the nomenclature of ref. [12]). Trimaximal mixing is defined to be the situation where $|U_{\alpha 2}| = 3^{-1/2} \ \forall \alpha \in \{e, \mu, \tau\}$. A model with TM₂ has been suggested a few years ago [13]. In TM₂ $|U_{e2}|^2 = \sin^2 \theta_{12} \cos^2 \theta_{13} = 1/3$ and therefore $\sin^2 \theta_{12} \approx 0.34$, which is a bit large but has not deterred several authors—see for instance ref. [14]—from having recently suggested models and *Ansätze* featuring TM₂. Note that TM₂, just as our model, is

⁸The predictions of our model are symmetric under $\mu \leftrightarrow \tau$. In the parameterization of the lepton mixing matrix that we use, the $\mu \leftrightarrow \tau$ interchange corresponds to $\cos \delta \rightarrow -\cos \delta$ and $\sin^2 \theta_{23} \rightarrow 0.5 - \sin^2 \theta_{23}$. This symmetry is easily observable in figs. 1 and 3 and in the left panel of fig. 4.

⁹This contradicts the phenomenological findings (at the 1σ level) of ref. [9], according to which θ_{23} lies in the first octant and δ is close to π . However, those findings are not in agreement with the ones of refs. [10, 11].




Figure 1: Scatter plot of $\cos \delta$ against $\sin^2 \theta_{23}$. The green (light grey) points are for an inverted neutrino mass spectrum ($m_3 < m_{1,2}$), the blue (black) points for a normal one ($m_3 > m_{1,2}$); this convention is used in all the figures of this paper.

characterized by predictions for lepton mixing which are invariant under any permutation of the lepton flavours.

One further prediction of TM_2 is¹⁰

$$(\cos \delta) \tan(2\theta_{23}) = \frac{\cos(2\theta_{13})}{(\sin \theta_{13}) \sqrt{3 \cos^2 \theta_{13} - 1}} > 0. \quad (21)$$

Therefore, in TM_2 , just as in our model, δ is in the first (or fourth) quadrant when θ_{23} is in the first octant, and δ is in the second (or third) quadrant when θ_{23} is in the second octant. Moreover, in TM_2 θ_{23} becomes closer to $\pi/4$ when $|\cos \delta|$ becomes smaller.

Figure 3 shows that, if we want a lower θ_{12} in our model, then we must accept $|\cos \delta|$ to be ever closer to 1, *i.e.* a more stringent phenomenological upper bound on $\sin^2 \theta_{12}$ translates in our model into a more stringent lower bound on $|\cos \delta|$.

In fig. 4 one sees that our model's bound on θ_{23} depends only faintly on θ_{13} . One also sees that our model's *lower bound* on $|0.5 - \sin^2 \theta_{23}|$ coincides with the *prediction* of TM_2 for $|0.5 - \sin^2 \theta_{23}|$ when $|\cos \delta|$ is *maximal*.

Figure 5 shows the scale of the neutrino masses in our model.

¹⁰Reference [15] contains relations that generalize eq. (21).




Figure 2: Scatter plots of $\sin^2 \theta_{12}$ against $\sin^2 \theta_{13}$. The left plot is for the general (complex) version of the model, the right plot is for the real version with $\cos \delta = -1$. The black line in the right plot displays the prediction [13] $\sin^2 \theta_{12} (1 - \sin^2 \theta_{13}) = 1/3$ of trimaximal mixing (TM_2). The crosses (blue (black) for a normal neutrino mass spectrum, green (light grey) for an inverted spectrum) indicate the phenomenological best-fit points of ref. [9].

We next speculate on possible experimental indications for (or against) our model. As is clear in the first three lines of eq. (10), in our model the Yukawa couplings of the neutral scalars to the charged leptons conserve flavour. Therefore, flavour-changing neutral Yukawa interactions only arise at loop level and are suppressed by a loop factor $(16\pi^2)^{-1}$ and by two additional Yukawa couplings. One may, moreover, show [16] that the (loop induced) flavour changing neutral couplings of the charged leptons to the photon and to the Z^0 are suppressed by inverse powers of the seesaw scale and are therefore, in general, unobservably small. Therefore, no decays like $\mu \rightarrow e\gamma$ or $Z^0 \rightarrow e^+\mu^-$ are expected, but decays like $h \rightarrow \tau^+\mu^-$ might be observable at the LHC (h is the observed scalar particle with mass 125 GeV).

When extending our model to the quark sector, one may either add to it further Higgs doublets or—a more economic possibility—make the quarks have Yukawa couplings to any one of the Higgs doublets ϕ_1 , ϕ_2 , or ϕ_3 (or else the up-type quarks may couple to one of them and the down-type quarks to another one). The fact that our Higgs doublets are singlets of the flavour group allows for this inviting possibility. Depending on the specific Yukawa couplings used, the signatures of the model at the LHC will vary. It is worth pointing out that, in any multi-Higgs-doublet model (MHDM), a physical scalar couples to $Z^0 Z^0$ with a coupling at most as strong as the one of




Figure 3: Scatter plot of $\cos \delta$ against $\sin^2 \theta_{12}$.




Figure 4: Scatter plots of $\sin^2 \theta_{23}$ against $\sin^2 \theta_{13}$. The left plot is for the general version and the right plot is for the real one with $\cos \delta = -1$. The black lines correspond to eq. (21), predicted by TM_2 [13], when $\cos \delta = \pm 1$ (full lines), $\cos \delta = \pm 0.9$ (dashed lines), and $\cos \delta = \pm 0.8$ (dotted lines). The crosses indicate the phenomenological best-fit points of ref. [9].




Figure 5: Ranges of allowed masses of the neutrinos in our model. The left scatter plot shows the neutrino mass relevant for neutrinoless double beta decay as a function of $\cos \delta$; the right plot correlates that quantity with the sum of the three light neutrino masses. We have imposed the cut $m_1 + m_2 + m_3 \leq 1$ eV on the sum of the neutrino masses.

the Higgs particle of the Standard Model (SM) [17]. This naively suggests that the observed decay $h \rightarrow Z^0 Z^0$ could be used to strongly constrain the parameter space of a MHDM, but this is not the case. Indeed, in a MHDM the couplings of any particular scalar to the top and bottom quarks might be either (much) stronger or (much) weaker than in the SM, and therefore in a MHDM both the production cross section and the total decay width of h will in general be at variance with those expected in the SM.

To summarize, we have presented in this paper a seesaw model featuring a simple application of the A_4 symmetry and which makes the predictions (5) for lepton mixing. We have shown that our model is compatible with experiment provided θ_{12} is in the upper part of its allowed range. Our model predicts non-maximal θ_{23} and also makes the prediction that $|\cos \delta|$ is very close to 1.

Acknowledgements:

LL thanks João P. Silva for a useful discussion. The work of PMF is supported by the Portuguese Foundation for Science and Technology (FCT) through the projects PEst-OE/FIS/UI0618/2011, PTDC/FIS/117951/2010, and the FP7 Reintegration Grant PERG08-GA-2010-277025. The work of LL

is supported through the Marie Curie Initial Training Network “UNILHC” PITN-GA-2009-237920 and through the projects PEst-OE/FIS/UI0777/2013, PTDC/FIS-NUC/0548/2012, and CERN/FP/123580/2011 of FCT. The work of POL is supported by the Austrian Science Fund (FWF) through the project P 24161-N16.

A Appendix: the VEVs of the σ_α

The potential for the σ_α is

$$V_\sigma = \mu (\sigma_e^2 + \sigma_\mu^2 + \sigma_\tau^2) + \lambda_1 (\sigma_e^2 + \sigma_\mu^2 + \sigma_\tau^2)^2 + \tilde{m} \sigma_e \sigma_\mu \sigma_\tau + \lambda_2 (\sigma_e^2 \sigma_\mu^2 + \sigma_\mu^2 \sigma_\tau^2 + \sigma_\tau^2 \sigma_e^2). \quad (\text{A1})$$

We have neglected terms which include both the σ_α and the ϕ_k since the VEVs of the ϕ_k should be much smaller than the VEVs of the σ_α and therefore those terms should have a negligible influence on the equations which determine the VEVs of the σ_α .

One may parameterize the VEVs of the σ_α as

$$\langle 0 | \sigma_e | 0 \rangle = U \cos \vartheta, \quad \langle 0 | \sigma_\mu | 0 \rangle = U \sin \vartheta \cos \varphi, \quad \langle 0 | \sigma_\tau | 0 \rangle = U \sin \vartheta \sin \varphi, \quad (\text{A2})$$

with $U \geq 0$, $\vartheta \in [0, \pi]$, and $\varphi \in [0, 2\pi[$. Then,

$$V_{\sigma 0} \equiv \langle 0 | V_\sigma | 0 \rangle = \mu U^2 + \lambda_1 U^4 + \frac{\tilde{m} U^3}{2} \sin^2 \vartheta \cos \vartheta \sin 2\varphi + \lambda_2 U^4 \sin^2 \vartheta \left(\cos^2 \vartheta + \frac{1}{4} \sin^2 \vartheta \sin^2 2\varphi \right). \quad (\text{A3})$$

There is a range of $\tilde{m}$ and λ_2 for which the minimum of $V_{\sigma 0}$ occurs when $\sin 2\varphi$ is at the boundary of its range, *viz.* when $\sin 2\varphi = 1$. There,

$$V_{\sigma 0} = \mu U^2 + \lambda_1 U^4 + \frac{\tilde{m} U^3}{2} \sin^2 \vartheta \cos \vartheta + \frac{\lambda_2 U^4}{4} \sin^2 \vartheta (1 + 3 \cos^2 \vartheta). \quad (\text{A4})$$

Then,

$$\frac{\partial V_{\sigma 0}}{\partial \cos \vartheta} = U^3 (1 - 3 \cos^2 \vartheta) \left(\frac{\tilde{m}}{2} + \lambda_2 U \cos \vartheta \right). \quad (\text{A5})$$

Within a range of $\tilde{m}$ and λ_2 the minimum of $V_{\sigma 0}$ occurs when $\cos^2 \vartheta = 1/3$ and, indeed, $\cos \vartheta = 1/\sqrt{3}$. In this way one obtains $\langle 0 | \sigma_e | 0 \rangle = \langle 0 | \sigma_\mu | 0 \rangle = \langle 0 | \sigma_\tau | 0 \rangle = U/\sqrt{3}$ as desired.

References

- [1] Y. Abe *et al.* (Double Chooz Coll.), *Indication for the disappearance of reactor electron antineutrinos in the Double Chooz experiment*, *Phys. Rev. Lett.* **108** (2012) 131801;
 F.P. An *et al.* (Daya Bay Coll.), *Observation of electron-antineutrino disappearance at Daya Bay*, *Phys. Rev. Lett.* **108** (2012) 171803;
 J.K. Ahn *et al.* (RENO Coll.), *Observation of reactor electron antineutrino disappearance in the RENO experiment*, *Phys. Rev. Lett.* **108** (2012) 191802.
- [2] F. Feruglio, C. Hagedorn, Y. Lin, and L. Merlo, *Lepton flavour violation in a supersymmetric model with A_4 flavour symmetry*, *Nucl. Phys. B* **832** (2010) 251;
 S.F. King and C. Luhn, *A_4 models of tri-bimaximal-reactor mixing*, *J. High Energy Phys.* **1203** (2012) 036;
 M.-C. Chen, J. Huang, J.-M. O'Bryan, A.M. Wijangco, and F. Yu, *Compatibility of θ_{13} and the type I seesaw model with A_4 symmetry*, *J. High Energy Phys.* **1302** (2013) 021;
 M. Holthausen, M. Lindner, and M.A. Schmidt, *Lepton flavor at the electroweak scale: A complete A_4 model*, *Phys. Rev. D* **87** (2013) 033006;
 S.F. King, S. Morisi, E. Peinado, and J.W.F. Valle, *Quark-lepton mass relation in a realistic A_4 extension of the Standard Model*, *Phys. Lett. B* **724** (2013) 68;
 S. Morisi, M. Nebot, K.M. Patel, E. Peinado, and J.W.F. Valle, *Quark-lepton mass relation and CKM mixing in an A_4 extension of the Minimal Supersymmetric Standard Model*, *Phys. Rev. D* **88**, 036001 (2013).
- [3] D. Meloni, S. Morisi, and E. Peinado, *Neutrino phenomenology and stable dark matter with A_4* , *Phys. Lett. B* **697** (2011) 339;
 S. Gupta, A.S. Joshipura and K.M. Patel, *Minimal extension of tri-bimaximal mixing and generalized $Z_2 \times Z_2$ symmetries*, *Phys. Rev. D* **85** (2012) 031903;
 H. Ishimori and E. Ma, *New simple A_4 neutrino model for nonzero θ_{13} and large δ_{CP}* , *Phys. Rev. D* **86** (2012) 045030;
 E. Ma, *Self-organizing neutrino mixing matrix*, *Phys. Rev. D* **86** (2012) 117301;
 E. Ma, A. Natale, and A. Rashed, *Scotogenic A_4 neutrino model for nonzero θ_{13} and large δ_{CP}* , *Int. J. Mod. Phys. A* **27** (2012) 1250134;
 Y. Ben Tov, X.-G. He, and A. Zee, *An $A_4 \times Z_4$ model for neutrino mixing*, *J. High Energy Phys.* **1212** (2012) 093;
 Y.H. Ahn, S.K. Kang, and C.S. Kim, *Spontaneous CP violation in A_4*

- flavor symmetry and leptogenesis*, *Phys. Rev. D* **87** (2013) 113012;
A.E. Carcamo Hernandez, I.d.M. Varzielas, S.G. Kovalenko, H. Päs and
I. Schmidt, *Lepton masses and mixings in a A_4 multi-Higgs model with
radiative seesaw*, arXiv:1307.6499 [hep-ph].
- [4] K.S. Babu, E. Ma, and J.W.F. Valle, *Underlying A_4 symmetry for the
neutrino mass matrix and the quark mixing matrix*, *Phys. Lett. B* **552**
(2003) 207;
For a modern version, see S. Morisi, D.V. Forero, J.C. Romão, and
J.W.F. Valle, *Neutrino mixing with revamped A_4 flavour symmetry*,
Phys. Rev. D **88** (2013) 016003.
 - [5] P.M. Ferreira, L. Lavoura, and P.O. Ludl, *Five models for lepton mixing*,
J. High Energy Phys. **1308** (2013) 113.
 - [6] C.D. Froggatt and H.B. Nielsen, *Hierarchy of quark masses, Cabibbo
angles and CP violation*, *Nucl. Phys. B* **147** (1979) 277.
 - [7] W. Grimus, L. Lavoura, and B. Radovčić, *Type II seesaw mechanism for
Higgs doublets and the scale of new physics*, *Phys. Lett. B* **674** (2009)
117.
 - [8] J. Beringer *et al.* (Particle Data Group), *Review of particle physics*,
Phys. Rev. D **86** (2012) 010001.
 - [9] G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, and A.M. Ro-
tunno, *Global analysis of neutrino masses, mixings and phases: Enter-
ing the era of leptonic CP violation searches*, *Phys. Rev. D* **86** (2012)
013012.
 - [10] D.V. Forero, M. Tórtola, and J.W.F. Valle, *Global status of neutrino
oscillation parameters after Neutrino-2012*, *Phys. Rev. D* **86** (2012)
073012.
 - [11] M.C. Gonzalez-Garcia, M. Maltoni, J. Salvado, and T. Schwetz, *Global
fit to three neutrino mixing: Critical look at present precision*, *J. High
Energy Phys.* **1212** (2012) 123.
 - [12] C.H. Albright, A. Dueck, and W. Rodejohann, *Possible alternatives to
tri-bimaximal mixing*, *Eur. Phys. J. C* **70** (2010) 1099.
 - [13] W. Grimus and L. Lavoura, *A model for trimaximal lepton mixing*, *J.
High Energy Phys.* **0809** (2008) 106.

- [14] N. Memenga, W. Rodejohann, and H. Zhang, *A_4 flavor symmetry model for Dirac neutrinos and sizable U_{e3}* , *Phys. Rev. D* **87** (2013) 053021;
 S.F. King, T. Neder, and A.J. Stuart, *Lepton mixing predictions from $\Delta(6n^2)$ family symmetry*, arXiv:1305.3200 [hep-ph];
 H. Qu and B.-Q. Ma, *New mixing pattern for neutrinos*, *Phys. Rev. D* **88** (2013) 037301.
- [15] S.-F. Ge, D.A. Dicus, and W.W. Repko, *$\mathbb{Z}_2$ symmetry prediction for the leptonic Dirac CP phase*, *Phys. Lett. B* **702** (2011) 220;
 S.-F. Ge, D.A. Dicus, and W.W. Repko, *Residual symmetries for neutrino mixing with a large θ_{13} and nearly maximal δ_D* , *Phys. Rev. Lett.* **108** (2012) 041801.
- [16] W. Grimus and L. Lavoura, *Soft lepton flavor violation in a multi-Higgs-doublet seesaw model*, *Phys. Rev. D* **66** (2002) 014016;
 R. Alonso, M. Dhen, M.B. Gavela, and T. Hambye, *Muon conversion to electron in nuclei in type-I seesaw models*, *J. High Energy Phys.* **1301** (2013) 118.
- [17] See for instance W. Grimus, L. Lavoura, O.M. Ogreid, and P. Osland, *A precision constraint on multi-Higgs-doublet models*, *J. Phys. G* **35** (2008) 075001;
 G.C. Branco, P.M. Ferreira, L. Lavoura, M.N. Rebelo, M. Sher, and J.P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rep.* **516** (2012) 1.