

# The Cauchy problem for $\mathscr{D}$ -modules on Ran spaces.

Giuseppe Bonavolontà

## Abstract

We will adopt an elementary approach to  $\mathscr{D}$ -modules on Ran spaces in terms of two-limits; the aim here is to define the category of coherent  $\mathscr{D}$ -modules, characteristic varieties and non-characteristic maps. An application will be the proof of the Cauchy-Kowaleski-Kashiwara theorem in this setting.

## 1 Introduction

The pioneering work [BD04] on the geometric foundations of Quantum Field Theory and Conformal Field Theory stressed the importance played by so-called Ran spaces. For example chiral, Lie<sup>\*</sup> and factorization algebras can be considered as geometric objects living on such spaces (see [BD04]). The most appropriate approach to the theory of  $\mathscr{D}$ -modules on Ran spaces (as presented in [FG11]) requires many sophisticated tools from Lurie's work on  $(\infty, 1)$ -categories. The main goal of this paper is to explore some definitions and properties of  $\mathscr{D}$ -modules on Ran spaces employing the classical theory of prestacks and two limits ([SGA4]). This simplified approach makes the theory accessible and sheds light on some definitions: the notions of characteristic varieties and of coherent  $\mathscr{D}$ -modules are really natural in this language. In paragraph §2 we recall some basics on the classical theory of  $\mathscr{D}$ -modules and microlocal geometry; in §3 we recall some definitions relative to prestacks, two-limits, etc.; in paragraph §4 we introduce the theory of modules on Ran manifolds by the use of §3; as an application we prove the Cauchy-Kowaleski-Kashiwara theorem for Ran manifolds.

**Acknowledgments.** I am deeply grateful to Pierre Schapira for having called my attention on the study of coherent  $\mathscr{D}$ -modules over Ran spaces. I have much benefited from the reading of the unpublished manuscript [GS06] and I wish to kindly thank the authors. Moreover I thank the Luxembourgian National Research Fund for support via AFR grant PhD 09-072.

## 2 Preliminaries

### Sheaves

We shall mainly follow the notations of [KS90] for sheaves, that of [KS06] for categories and that of [Ka03] for  $\mathscr{D}$ -modules.

Let  $X$  be a real manifold and let  $\mathbf{k}$  be a field. One denotes by  $\mathbf{k}_X$  the constant sheaf on  $X$  with stalk  $\mathbf{k}$  and by  $\text{Mod}(\mathbf{k}_X)$  the abelian category of sheaves of  $\mathbf{k}$ -modules on  $X$ . We denote by  $D^b(\mathbf{k}_X)$  the bounded derived category of  $\text{Mod}(\mathbf{k}_X)$ . One simply calls an object of this category, “a sheaf”.

We shall use the classical six operations on sheaves, the internal hom  $\mathcal{H}om$ , the tensor product  $\otimes$ , the direct image  $f_*$ , the proper direct image  $f_!$ , the inverse image  $f^{-1}$  and the derived functors  $R\mathcal{H}om$ ,  $^L\otimes$ ,  $Rf_*$ ,  $Rf_!$ ,  $f^{-1}$ , as well as the extraordinary inverse image  $f^!$ , right adjoint to  $Rf_!$ .

We also use the notation  $\boxtimes$  for the external product: if  $X$  and  $Y$  are two manifolds and where  $q_1$  and  $q_2$  denote the first and second projection on  $X \times Y$ , we set for  $F \in \text{Mod}(\mathbf{k}_X)$  and  $G \in \text{Mod}(\mathbf{k}_Y)$ , one sets  $F \boxtimes G := q_1^{-1}F \otimes q_2^{-1}G$ .

## Microlocal geometry

For a real or complex manifold  $X$ , we denote by  $\tau: TX \rightarrow X$  its tangent vector bundle and by  $\pi: T^*X \rightarrow X$  its cotangent vector bundle. If  $E \rightarrow X$  is a vector bundle, we identify  $X$  with the zero-section.

Let  $f: X \rightarrow Y$  be a morphism of real or complex manifolds. To  $f$  are associated the tangent morphisms

$$(2.1) \quad \begin{array}{ccccc} TX & \xrightarrow{f'} & X \times_Y TY & \xrightarrow{f_\tau} & TY \\ \downarrow \tau & & \downarrow \tau & & \downarrow \tau \\ X & \xlongequal{\quad} & X & \xrightarrow{f} & Y. \end{array}$$

By duality, we deduce the diagram:

$$(2.2) \quad \begin{array}{ccccc} T^*X & \xleftarrow{f_d} & X \times_Y T^*Y & \xrightarrow{f_\pi} & T^*Y \\ \downarrow \pi & & \downarrow \pi & & \downarrow \pi \\ X & \xlongequal{\quad} & X & \xrightarrow{f} & Y. \end{array}$$

One sets

$$T_X^*Y := \ker f_d = f_d^{-1}(T_X^*X).$$

**Definition 2.1.** Let  $\Lambda \subset T^*Y$  be a closed  $\mathbb{R}^+$ -conic subset. One says that  $f$  is non-characteristic for  $\Lambda$ , or else,  $\Lambda$  is non-characteristic for  $f$ , if

$$f_\pi^{-1}(\Lambda) \cap T_X^*Y \subset X \times_Y T_Y^*Y.$$

## $\mathcal{O}$ -modules

Now assume that  $(X, \mathcal{O}_X)$  is a complex manifold of complex dimension  $d_X$ . We denote by  $\text{Mod}(\mathcal{O}_X)$  the abelian category of sheaves of  $\mathcal{O}_X$ -modules and by  $\mathcal{H}om_{\mathcal{O}_X}$  and  $\otimes_{\mathcal{O}_X}$  the internal hom and tensor product in this category. The sheaf of rings  $\mathcal{O}_X$  is Noetherian and we denote by  $\text{Mod}_{\text{coh}}(\mathcal{O}_X)$  the thick abelian subcategory of  $\text{Mod}(\mathcal{O}_X)$  consisting of coherent sheaves.

We denote by  $D^b(\mathcal{O}_X)$  the bounded derived category of  $\text{Mod}(\mathcal{O}_X)$  and by  $D_{\text{coh}}^b(\mathcal{O}_X)$  the full triangulated subcategory consisting of objects with coherent cohomology.

Let  $f: X \rightarrow Y$  be a morphism of complex manifolds. We keep the notation  $f_*$  and  $f_!$  for the two direct image functors for  $\mathcal{O}$ -modules and we denote by  $f^*$  the inverse image functor for  $\mathcal{O}$ -modules:

$$f^*\mathcal{G} := \mathcal{O}_X \otimes_{f^{-1}\mathcal{O}_Y} f^{-1}\mathcal{G}.$$

Hence we have the pair of adjoint functors  $(f^*, f_*)$

$$\text{Mod}(\mathcal{O}_X) \begin{matrix} \xrightarrow{f_*} \\ \xleftarrow{f^*} \end{matrix} \text{Mod}(\mathcal{O}_Y)$$

It follows that  $f_*$  is left exact and  $f^*$  is right exact. We denote by  $Lf^*$  the left derived functor of  $f^*$ .

Given two manifolds  $X$  and  $Y$ , we denote by  $\boxtimes$  the external product for  $\mathcal{O}$ -modules. Hence,

$$\mathcal{F} \boxtimes \mathcal{G} := \mathcal{O}_{X \times Y} \otimes_{\mathcal{O}_X \boxtimes \mathcal{O}_Y} (\mathcal{F} \boxtimes \mathcal{G}).$$

### $\mathcal{D}$ -modules

We denote by  $\mathcal{D}_X$  the sheaf of rings of holomorphic differential operators, the subalgebra of  $\mathcal{E}nd(\mathcal{O}_X)$  generated by  $\mathcal{O}_X$  and  $\Theta_X$ , the sheaf of holomorphic vector fields.

Unless otherwise specified, a  $\mathcal{D}_X$ -module is a left- $\mathcal{D}_X$ -module. Hence a right  $\mathcal{D}_X$ -module is a  $(\mathcal{D}_X^{\text{op}})$ -module. We denote by  $\text{Mod}(\mathcal{D}_X)$  the abelian category of  $\mathcal{D}_X$ -modules and by  $D^b(\mathcal{D}_X)$  its bounded derived category.

Recall the operations

$$\begin{aligned} \mathcal{H}om_{\mathcal{O}_X} &: \text{Mod}(\mathcal{D}_X)^{\text{op}} \times \text{Mod}(\mathcal{D}_X) \rightarrow \text{Mod}(\mathcal{D}_X), \\ \mathcal{H}om_{\mathcal{O}_X} &: \text{Mod}(\mathcal{D}_X^{\text{op}})^{\text{op}} \times \text{Mod}(\mathcal{D}_X^{\text{op}}) \rightarrow \text{Mod}(\mathcal{D}_X), \\ \otimes_{\mathcal{O}_X} &: \text{Mod}(\mathcal{D}_X) \times \text{Mod}(\mathcal{D}_X) \rightarrow \text{Mod}(\mathcal{D}_X), \\ \otimes_{\mathcal{O}_X} &: \text{Mod}(\mathcal{D}_X^{\text{op}}) \times \text{Mod}(\mathcal{D}_X) \rightarrow \text{Mod}(\mathcal{D}_X^{\text{op}}). \end{aligned}$$

We denote by  $\boxtimes$  the external product for  $\mathcal{D}$ -modules:

$$\mathcal{M} \boxtimes \mathcal{N} := \mathcal{D}_{X \times Y} \otimes_{\mathcal{D}_X \boxtimes \mathcal{D}_Y} (\mathcal{M} \boxtimes \mathcal{N}).$$

For a morphism  $f: X \rightarrow Y$  of complex manifolds, we denote as usual by  $\mathcal{D}_{X \rightarrow Y}$  the transfert bimodule, a  $(\mathcal{D}_X, f^{-1}\mathcal{D}_Y)$ -bimodule

$$\mathcal{D}_{X \rightarrow Y} := \mathcal{O}_X \otimes_{f^{-1}\mathcal{O}_Y} f^{-1}\mathcal{D}_Y.$$

One should be aware that the left  $\mathcal{D}_X$  structure of  $\mathcal{D}_{X \rightarrow Y}$  is not the one induced by that of  $\mathcal{O}_X$ . We then have the inverse and direct image functors

$$\begin{aligned} f^D &: D^b(\mathcal{D}_Y) \rightarrow D^b(\mathcal{D}_X), \quad f^D \mathcal{N} = \mathcal{D}_{X \rightarrow Y} \overset{\text{L}}{\otimes}_{f^{-1}\mathcal{D}_Y} f^{-1}\mathcal{N}, \\ f_D &: D^b(\mathcal{D}_X^{\text{op}}) \rightarrow D^b(\mathcal{D}_Y^{\text{op}}), \quad f_D(\mathcal{M}) = \text{R}f_*(\mathcal{M} \overset{\text{L}}{\otimes}_{\mathcal{D}_X} \mathcal{D}_{X \rightarrow Y}). \end{aligned}$$

We denote by  $\text{Mod}_{\text{coh}}(\mathcal{D}_X)$  the thick abelian subcategory of  $\text{Mod}(\mathcal{D}_X)$  consisting of coherent  $\mathcal{D}_X$ -modules and by  $\text{D}_{\text{coh}}^b(\mathcal{D}_X)$  the full triangulated subcategory consisting of objects with coherent cohomology.

For  $\mathcal{M} \in \text{D}_{\text{coh}}^b(\mathcal{D}_X)$ , we denote by  $\text{char}(\mathcal{M})$  its characteristic variety, a closed  $\mathbb{C}^\times$ -conic complex analytic subvariety of  $T^*X$ .

## The Cauchy-Kowaleski-Kashiwara theorem

**Definition 2.2.** Let  $f : X \rightarrow Y$  be a morphism of manifolds and  $\mathcal{N}$  be a coherent  $\mathcal{D}_Y$ -module. One says that  $f$  is non-characteristic for  $\mathcal{N}$  if  $f$  is non-characteristic for  $\text{char}(\mathcal{N})$ .

**Proposition 2.3.** ([Ka70]) *Let  $\mathcal{N}$  be a coherent  $\mathcal{D}_Y$ -module. Suppose that  $f : X \rightarrow Y$  is non-characteristic for  $\mathcal{N}$ . Then*

- (a)  $H^k(f^D \mathcal{N}) = 0$  for  $k \neq 0$ ,
- (b)  $H^0(f^D \mathcal{N})$  is a coherent  $\mathcal{D}_X$ -module,
- (c)  $\text{char}(f^D \mathcal{N}) = f_* f_\pi^{-1} \text{char} \mathcal{N}$ .

For  $\mathcal{N}_1$  and  $\mathcal{N}_2$  in  $\text{D}^b(\mathcal{D}_Y)$  we have a natural morphism

$$f^{-1} \text{R}\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{N}_1, \mathcal{N}_2) \rightarrow \text{R}\mathcal{H}om_{\mathcal{D}_X}(f^D \mathcal{N}_1, f^D \mathcal{N}_2).$$

Note that  $f^D \mathcal{O}_X \simeq \mathcal{O}_Y$ .

The following theorem is known as the *Cauchy-Kowalesky-Kashiwara* theorem:

**Theorem 2.4.** ([Ka70]) *Let  $f : X \rightarrow Y$  be a morphism of complex manifolds and  $\mathcal{N}$  be a coherent  $\mathcal{D}_Y$ -module. If  $f$  is non-characteristic for  $\mathcal{N}$ , then*

$$(2.3) \quad f^{-1} \text{R}\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{N}, \mathcal{O}_Y) \rightarrow \text{R}\mathcal{H}om_{\mathcal{D}_X}(f^D \mathcal{N}, \mathcal{O}_X)$$

*is an isomorphism.*

## 3 Grothendieck stacks

References for this section are made to [SGA4] (see also [KS06] for an exposition). The results of this section are well-known to the specialists. We will work in a given universe  $\mathcal{U}$ , a category means a  $\mathcal{U}$ -category and a set means a small set. Here **Cat** denotes the big 2-category of all  $\mathcal{U}$ -categories.

### Prestack

Let  $\mathfrak{S}$  be a 2-projective system of categories indexed by a small category  $\mathcal{I}$ , that is, a functor  $\mathfrak{S} : \mathcal{I}^{\text{op}} \rightarrow \mathbf{Cat}$ . For short, we consider  $\mathcal{I}$  as a presite and call  $\mathfrak{S}$  a prestack on  $\mathcal{I}$ . Hence

- (a) for each  $i \in \mathcal{I}$ ,  $\mathfrak{S}(i)$  is a category;
- (b) for each morphism  $s : i_1 \rightarrow i_2$  in  $\mathcal{I}$ ,  $\mathfrak{S}(s)$  is a functor  $\mathfrak{S}(i_2) \rightarrow \mathfrak{S}(i_1)$ , called the restriction functor (for convenience, we shall write in the sequel  $\rho_s$  instead of  $\mathfrak{S}(s)$ );

- (c) for  $s: i_1 \rightarrow i_2$  and  $t: i_2 \rightarrow i_3$ , we have an isomorphism of functor  $c_{s,t}: \rho_s \circ \rho_t \xrightarrow{\sim} \rho_{t \circ s}$ , making the diagram below commutative for each  $u: i_3 \rightarrow i_4$ :

$$\begin{array}{ccc} \rho_s \circ \rho_t \circ \rho_u & \xrightarrow{c_{t,u}} & \rho_s \circ \rho_{u \circ t} \\ c_{s,t} \downarrow & & \downarrow c_{s,u \circ t} \\ \rho_{t \circ s} \circ \rho_u & \xrightarrow{c_{t \circ s, u}} & \rho_{u \circ t \circ s}. \end{array}$$

- (d) finally  $\rho_{\text{id}_i} = \text{id}_{\mathfrak{S}(i)}$  and  $c_{\text{id}_i, \text{id}_i} = \text{id}_{\text{id}_{\mathfrak{S}(i)}}$  for any  $i \in \mathcal{I}$ .

## Functor of prestacks

Let  $\mathfrak{S}_\nu$  ( $\nu = 1, 2$ ) be prestacks on  $\mathcal{I}$  with the restrictions  $\rho'_s$  and the composition isomorphisms  $c'_{s,t}$ .

Recall that a functor of prestacks  $\Phi: \mathfrak{S}_1 \rightarrow \mathfrak{S}_2$  on  $\mathcal{I}$  is the data of:

- (a) for any  $i \in \mathcal{I}$ , a functor  $\Phi(i): \mathfrak{S}_1(i) \rightarrow \mathfrak{S}_2(i)$ ;
- (b) for any morphism  $s: i_1 \rightarrow i_2$ , an isomorphism  $\Phi_s$  of functors from  $\mathfrak{S}_1(i_2)$  to  $\mathfrak{S}_2(i_1)$

$$\Phi_s: \Phi(i_1) \circ \rho_s^1 \xrightarrow{\sim} \rho_s^2 \circ \Phi(i_2);$$

these data satisfying: for any sequence of morphisms  $i_1 \xrightarrow{s} i_2 \xrightarrow{t} i_3$  the following diagram commutes

$$(3.1) \quad \begin{array}{ccccc} \Phi(i_1) \circ \rho_s^1 \circ \rho_t^1 & \xrightarrow{\Phi_s} & \rho_s^2 \circ \Phi(i_2) \circ \rho_t^1 & \xrightarrow{\Phi_t} & \rho_s^2 \circ \rho_t^2 \circ \Phi(i_3) \\ \downarrow c_{s,t}^1 & & & & \downarrow c_{s,t}^2 \\ \Phi(i_1) \circ \rho_{t \circ s}^1 & \xrightarrow{\Phi_{t \circ s}} & & & \rho_{t \circ s}^2 \circ \Phi(i_3). \end{array}$$

## Morphism of functors of prestacks

We will need the following definition

**Definition 3.1.** Let  $\Phi_\nu: \mathfrak{S}_1 \rightarrow \mathfrak{S}_2$  ( $\nu = 1, 2$ ) be two functors of prestacks on  $\mathcal{I}$ . A morphism of functors of prestacks  $\theta: \Phi_1 \rightarrow \Phi_2$  is the data for any  $i \in \mathcal{I}$  of a morphism of functors  $\theta(i): \Phi_1(i) \rightarrow \Phi_2(i)$  such that for any morphism  $s: i_1 \rightarrow i_2$  in  $\mathcal{I}$ , the following diagram commutes

$$(3.2) \quad \begin{array}{ccc} \Phi_1(i_1) \circ \rho_s^1 & \xrightarrow{\theta(i_1)} & \Phi_2(i_1) \circ \rho_s^1 \\ \downarrow \Phi_s^1 & & \downarrow \Phi_s^2 \\ \rho_s^2 \circ \Phi_1(i_2) & \xrightarrow{\theta(i_2)} & \rho_s^2 \circ \Phi_2(i_2). \end{array}$$

## 2-projective limits

For a prestack  $\mathfrak{S}$  on  $\mathcal{I}$ , one defines the category  $\mathfrak{S}(\mathcal{I})$  as the 2-projective limit of the functor  $\mathfrak{S}: \mathcal{I}^{\text{op}} \rightarrow \mathbf{Cat}$ :

$$\mathfrak{S}(\mathcal{I}) = 2 \varprojlim_{i, s} \mathfrak{S}(i).$$

More explicitly  $\mathfrak{S}(\mathcal{I})$  is given as follows.

**Definition 3.2.** (a) An object  $F$  of  $\mathfrak{S}(\mathcal{J})$  is a family  $\{(F_i, \varphi_s)\}_{i,s}$  ( $i \in \mathcal{J}$ ,  $s \in \text{Mor}(\mathcal{J})$ ) where

- (i) for any  $i \in \mathcal{J}$ ,  $F_i$  is an object of  $\mathfrak{S}(i)$ ,
- (ii) for any morphism  $s: i_1 \rightarrow i_2$  in  $\mathcal{J}$ ,  $\varphi_s: \rho_s F_{i_2} \xrightarrow{\sim} F_{i_1}$  is an isomorphism such that
  - for all  $i \in \mathcal{J}$ ,  $\varphi_{\text{id}_i} = \text{id}_{F_i}$ ,
  - for any sequence  $i_1 \xrightarrow{s} i_2 \xrightarrow{t} i_3$  of morphisms in  $\mathcal{J}$ , the following diagram commutes

$$(3.3) \quad \begin{array}{ccc} \rho_s \rho_t F_{i_3} & \xrightarrow[\sim]{\rho_s(\varphi_t)} & \rho_s F_{i_2} \\ c_{s,t} \downarrow \wr & & \varphi_s \downarrow \wr \\ \rho_{tos} F_{i_3} & \xrightarrow[\sim]{\varphi_{tos}} & F_{i_1}. \end{array}$$

- (b) A morphism  $f: \{(F_i, \varphi_s)\}_{i,s} \rightarrow \{(F'_i, \varphi'_s)\}_{i,s}$  in  $\mathfrak{S}(\mathcal{J})$  is a family of morphisms  $\{f_i: F_i \rightarrow F'_i\}_{i \in \mathcal{J}}$  such that for any  $s: i_1 \rightarrow i_2$ , the diagram below commutes:

$$(3.4) \quad \begin{array}{ccc} \rho_s(F_{i_2}) & \xrightarrow{\rho_s(f_{i_2})} & \rho_s(F'_{i_2}) \\ \varphi_s \downarrow \wr & & \wr \downarrow \varphi'_s \\ F_{i_1} & \xrightarrow{f_{i_1}} & F'_{i_1}. \end{array}$$

**Remark 3.3.** Denote by  $\text{Mor}(\mathcal{J})$  the category whose objects are the morphisms of  $\mathcal{J}$ , a morphism  $(s: i \rightarrow j) \rightarrow (s': i' \rightarrow j')$  being visualized by the commutative diagram

$$\begin{array}{ccc} i & \xrightarrow{s} & j \\ t \downarrow & & \uparrow t' \\ i' & \xrightarrow{s'} & j'. \end{array}$$

For two objects  $F = \{(F_i, \varphi_s)\}_{i,s}$  and  $F' = \{(F'_i, \varphi'_s)\}_{i,s}$  in  $\mathfrak{S}(\mathcal{J})$ , consider the functor  $\Psi: \text{Mor}(\mathcal{J})^{\text{op}} \rightarrow \mathbf{Set}$  defined as follows. For  $(s: i \rightarrow j) \in \text{Mor}(\mathcal{J})$ , set

$$\Psi(s) = \text{Hom}_{\mathfrak{S}(i)}(F_i, \rho_s F'_j).$$

For  $(s': i' \rightarrow j') \in \text{Mor}(\mathcal{J})$  and for a morphism  $(t, t'): s \rightarrow s'$  in  $\text{Mor}(\mathcal{J})$ , define  $\Psi(s') \rightarrow \Psi(s)$  as the composition

$$\begin{aligned} \text{Hom}_{\mathfrak{S}(i')}(F_{i'}, \rho_{s'} F'_{j'}) &\rightarrow \text{Hom}_{\mathfrak{S}(i')}(F_{i'}, \rho_t F'_{i'}, \rho_t \rho_{s'} F'_{j'}) \\ &\rightarrow \text{Hom}_{\mathfrak{S}(i)}(F_i, \rho_s F'_j) \end{aligned}$$

where the last map is associated with the morphisms  $F_i \rightarrow \rho_t F'_{i'}$  and  $\rho_t \rho_{s'} F'_{j'} \rightarrow \rho_t \rho_{s'} \rho_{t'} F'_j \simeq \rho_s F'_j$ .

Then

$$\text{Hom}_{\mathfrak{S}(\mathcal{J})}(F, F') \simeq \varprojlim_{s \in \text{Mor}(\mathcal{J})^{\text{op}}} \Psi(s).$$

**Proposition 3.4.** *Let  $\mathfrak{S}$  be a prestack on  $\mathcal{I}$ . Assume that, for each object  $i \in \mathcal{I}$ , the category  $\mathfrak{S}(i)$  admits inductive (resp. projective) limits indexed by a small category  $A$  and that, for each morphism  $s: i \rightarrow j$  in  $\mathcal{I}$ , the restriction functor  $\rho_s$  commutes with such limits. Then the category  $\mathfrak{S}(\mathcal{I})$  admits inductive (resp. projective) limits indexed by  $A$  and the restriction functors  $\rho_i: \mathfrak{S}(\mathcal{I}) \rightarrow \mathfrak{S}(i)$  commute with such limits.*

### Some Lemmas

**Lemma 3.5.** *Every functor of prestacks  $\Psi: \mathfrak{S}_1 \rightarrow \mathfrak{S}_2$  on  $\mathcal{I}$  induces a functor  $\Psi_\infty: \mathfrak{S}_1(\mathcal{I}) \rightarrow \mathfrak{S}_2(\mathcal{I})$  on the corresponding two-limit categories.*

*Proof.* To each object  $\{(F_i, \varphi_s)\}_{i,s} \in \mathfrak{S}_1(\mathcal{I})$ ,  $\Psi$  associates  $\{(\Psi(i)F_i, \tilde{\varphi}_s)\}_{i,s} \in \mathfrak{S}_2(\mathcal{I})$ , with  $\tilde{\varphi}_s$  defined as the composition  $\rho_s^2 \Psi(j)F_j \xrightarrow{\sim} \Psi(i)\rho_s^1 F_j \xrightarrow{\sim} \Psi(i)F_i$  where the first isomorphism is the inverse of  $\Psi_s$  and the second is  $\Psi(i)(\varphi_s)$ . Notice that for any sequence  $i_1 \xrightarrow{s} i_2 \xrightarrow{t} i_3$  of morphisms in  $\mathcal{I}$ , the following diagram commutes

$$(3.5) \quad \begin{array}{ccc} \rho_s^2 \rho_t^2 \Psi(i_3)F_{i_3} & \xrightarrow[\sim]{\rho_s^2(\tilde{\varphi}_t)} & \rho_s^2 \Psi(i_2)F_{i_2} \\ c_{s,t}^2 \downarrow \wr & & \tilde{\varphi}_s \downarrow \wr \\ \rho_{tos}^2 \Psi(i_3)F_{i_3} & \xrightarrow[\sim]{\tilde{\varphi}_{tos}} & \Psi(i_1)F_{i_1}; \end{array}$$

in fact it is enough to apply  $\Psi(i_1)$  to the corresponding diagram in def (3.2) and then use property (3.1) of the morphism  $\Psi$ .

Finally  $\Psi_\infty$  associates to each morphism  $f: \{(F_i, \varphi_s)\}_{i,s} \rightarrow \{(F'_i, \varphi'_s)\}_{i,s}$ ,  $\Psi_\infty f = \{\Psi(i)f_i\}: \mathfrak{S}_1(\mathcal{I}) \rightarrow \mathfrak{S}_2(\mathcal{I})$  with

$$(3.6) \quad \begin{array}{ccc} \rho_s(\Psi(i_2)F_{i_2}) & \xrightarrow{\rho_s \Psi(i_2)(f_{i_2})} & \rho_s(\Psi(i_2)F'_{i_2}) \\ \tilde{\varphi}_s \downarrow \wr & & \wr \downarrow \tilde{\varphi}'_s \\ \Psi(i_1)F_{i_1} & \xrightarrow{\Psi(i_1)f_{i_1}} & \Psi(i_1)F'_{i_1} \end{array}$$

(it is enough to apply the functor  $\Psi(i_1)$  to the diagram (3.4))

□

The next lemma is obvious

**Lemma 3.6.** *Let  $\Psi_1: \mathfrak{S}_1 \rightarrow \mathfrak{S}_2$  and  $\Psi_2: \mathfrak{S}_2 \rightarrow \mathfrak{S}_3$  be two morphisms of prestacks on  $\mathcal{I}$ . There exists an isomorphism of functors  $\Psi_1 \circ \Psi_2 \circ \infty \xrightarrow{\sim} (\Psi_1 \circ \Psi_2)_\infty: \mathfrak{S}_1(\mathcal{I}) \rightarrow \mathfrak{S}_3(\mathcal{I})$ .*

Define

$$(3.7) \quad \pi: \mathfrak{S}(\mathcal{I}) \rightarrow \prod_{i \in \mathcal{I}} \mathfrak{S}(i)$$

as the functor that associates to each  $\{(F_i, \varphi_s)\}_{i,s}$ , the collection  $\{F_i\}_i$  and acts on each morphism  $\{f_i\}_{i \in I}$  forgetting the property (3.4).

**Lemma 3.7.** *The functor  $\pi$  is conservative.*

*Proof.* Consider  $\{f_i\}_{i \in I}$  a morphism in  $\mathfrak{S}(\mathcal{J})$  between  $\{F_i, \varphi_s\}_{i,s}$  and  $\{F'_i, \varphi'_s\}_{i,s}$ . Assume  $\{f_i\}_{i \in I}$  is an isomorphism in  $\prod_{i \in \mathcal{J}} \mathfrak{S}(i)$ . For each  $f_i$  denote by  $g_i$  the corresponding inverse morphism. Consider the diagram

$$(3.8) \quad \begin{array}{ccccc} \rho_s(F'_{i_2}) & \xrightarrow{\rho_s(g_{i_2})} & \rho_s(F_{i_2}) & \xrightarrow{\rho_s(f_{i_2})} & \rho_s(F'_{i_2}) \\ \varphi'_s \downarrow \wr & & \wr \downarrow \varphi_s & & \varphi'_s \downarrow \wr \\ F'_{i_1} & \xrightarrow{g_{i_1}} & F_{i_1} & \xrightarrow{f_{i_1}} & F'_{i_1}; \end{array}$$

the right square is commutative by hypothesis and the large rectangle is trivially commutative. Then the equality  $f_{i_1} \circ \varphi_s \circ \rho_s(g_{i_2}) = f_{i_1} \circ g_{i_1} \circ \varphi'_s$  implies  $\varphi_s \circ \rho_s(g_{i_2}) = g_{i_1} \circ \varphi'_s$  since each  $f_{i_1}$  is an isomorphism.

**Lemma 3.8.** *Let  $\Psi_\nu: \mathfrak{S}_1 \rightarrow \mathfrak{S}_2$  ( $\nu = 1, 2$ ) be two functors of prestacks on  $\mathcal{J}$ , and  $\theta: \Psi_1 \rightarrow \Psi_2$  a morphisms of functors.*

- (a)  *$\theta$  defines a morphism of functors  $\theta_\infty: \Psi_1 \infty \rightarrow \Psi_2 \infty$ ,*
- (b) *if  $\theta$  is an isomorphism, then  $\theta_\infty$  is an isomorphism as well.*

*Proof.* Obvious.

## Grothendieck prestacks

Recall that a  $\mathbf{k}$ -abelian prestack  $\mathfrak{S}$  on a presite  $\mathcal{J}$  is a prestack such that  $\mathfrak{S}(i)$  is a  $\mathbf{k}$ -abelian category for each  $i \in \mathcal{J}$  and the restriction functors  $\rho_s$  are exact. For a site  $\mathcal{J}$ , a  $\mathbf{k}$ -abelian stack is a  $\mathbf{k}$ -abelian prestack which is a stack. If there is no risk of confusion, we shall not mention the field  $\mathbf{k}$ .

**Definition 3.9.** (see [GS06])

- (i) A Grothendieck prestack  $\mathfrak{S}$  over  $\mathbf{k}$  on a presite  $\mathcal{J}$  is a  $\mathbf{k}$ -abelian prestack such that the abelian category  $\mathfrak{S}(i)$  is a Grothendieck category for each object  $i \in \mathcal{J}$  and the restriction functor  $\rho_s$  commutes with small inductive limits for each morphism  $s: i \rightarrow j$ . (Recall that  $\rho_s$  is exact.)
- (ii) For a site  $\mathcal{J}$ , a Grothendieck stack is a Grothendieck prestack which is a stack.

The next result is a deep results of [SGA4] (see Exposé I, Theorem 9.22).

**Theorem 3.10.** *Let  $\mathfrak{S}$  be a Grothendieck prestack on  $\mathcal{J}$ . Then  $\mathfrak{S}(\mathcal{J})$  is a Grothendieck category.*

If  $\mathcal{C}$  is an abelian category, we denote as usual by  $D^b(\mathcal{C})$  (resp.  $D^+(\mathcal{C})$ ) its bounded (resp. bounded from below) derived category. Hence, for each  $i \in \mathcal{J}$ , the functor  $\rho_i: \mathfrak{S}(\mathcal{J}) \rightarrow \mathfrak{S}(i)$  extends as a functor  $\tilde{\rho}_i: D^+(\mathfrak{S}^*(\mathcal{J})) \rightarrow D^+(\mathfrak{S}^*(i))$  and one checks easily that these functors define a functor

$$(3.9) \quad \tilde{\rho}: D^+(\mathfrak{S}(\mathcal{J})) \rightarrow 2 \varprojlim_{i,s} D^+(\mathfrak{S}(i)).$$



The  $2\varprojlim$  of triangulated categories is no more triangulated in general; the functor  $\tilde{\rho}$  is neither full nor faithful. However it would be interesting to check if this functor is conservative.

## 4 Ran manifolds

We shall specialize the general theory of 2-limits presented in the previous section. Let  $\mathcal{J}$  be a small category.

- Definition 4.1.** (a) A real (resp. complex) Ran-manifold indexed by  $\mathcal{J}$  is a projective system  $X = \{X_i\}_{i \in \mathcal{J}}$  of real (resp. complex) manifolds indexed by  $\mathcal{J}$  such that for every morphism  $s \in \text{Hom}_{\mathcal{J}}(j, i)$  the corresponding morphism  $\Delta_s : X_i \rightarrow X_j$  is a closed embedding.
- (b) Let  $X = \{X_i\}_{i \in \mathcal{J}}$  and  $Y = \{Y_i\}_{i \in \mathcal{J}}$  be two Ran-manifolds. A morphism  $f : X \rightarrow Y$  is defined as a collection of morphisms  $\{f_i\}_{i \in \mathcal{J}}$  such that all the diagrams below commute

$$(4.1) \quad \begin{array}{ccc} X_i & \xrightarrow{f_i} & Y_i \\ \Delta_s \downarrow & & \downarrow \Delta_s \\ X_j & \xrightarrow{f_j} & Y_j \end{array}$$

For each morphism  $s : j \rightarrow i$ , with the corresponding embedding  $\Delta_s : X_i \hookrightarrow X_j$ , we consider the following diagram

$$(4.2) \quad \begin{array}{ccccc} T^*X_i & \xleftarrow{\Delta_{sd}} & X_i \times_{X_j} T^*X_j & \xrightarrow{\Delta_{s\pi}} & T^*X_j \\ \pi \downarrow & & \downarrow \pi & & \downarrow \pi \\ X_i & \xlongequal{\quad} & X_i & \xrightarrow{\Delta_s} & X_j \end{array}$$

We denote by  $T_s^*X_j$  the kernel of  $\Delta_{sd}$ , that is,  $\Delta_{sd}^{-1}T_{X_i}^*X_i$ .

**Definition 4.2.** (a) We define  $T^*X$  as the collection of all the diagrams (4.2).

(b) A family  $\{\Lambda_i\}_{i \in \mathcal{J}}$  of closed  $\mathbb{R}^+$ -conic subsets  $\Lambda_i \subset T^*X_i$  satisfying

$$(4.3) \quad \Lambda_i \subset \Delta_{sd}(\Delta_{s\pi}^{-1}(\Lambda_j))$$

for all  $s : j \rightarrow i$  is said to be a closed  $\mathbb{R}^+$ -conic subset  $\Lambda \subset T^*X$ .

(c) A closed  $\mathbb{R}^+$ -conic subset  $\Lambda$  of  $T^*X$  is said to be transversal to the identity, if for each  $s : j \rightarrow i$ , the map  $\Delta_s$  is non characteristic with respect to  $\Lambda_j$ , that is,  $\Delta_{sd}^{-1}T_{X_i}^*X_i \cap \Delta_{s\pi}^{-1}\Lambda_j \subset X_i \times_{X_j} T_{X_j}^*X_j$ .

Consider a morphism  $f: X \rightarrow Y$  of Ran-manifolds. It gives rise to the commutative diagrams associated to  $s: j \rightarrow i$ :

$$(4.4) \quad \begin{array}{ccccc} T^*X_i & \xleftarrow{\quad} & X_i \times_{X_j} T^*X_j & \xhookrightarrow{\quad} & T^*X_j \\ \uparrow f_{id} & \searrow \Delta_{f_s,d} & \uparrow & & \uparrow f_{jd} \\ X_i \times_{Y_i} T^*Y_i & \xleftarrow{\quad} & X_i \times_{Y_j} T^*Y_j & \xhookrightarrow{\quad} & X_j \times_{Y_j} T^*Y_j \\ \downarrow f_{i\pi} & & \downarrow & \searrow \Delta_{f_s,\pi} & \downarrow f_{j\pi} \\ T^*Y_i & \xleftarrow{\quad} & Y_i \times_{Y_j} T^*Y_j & \xhookrightarrow{\quad} & T^*Y_j. \end{array}$$

Notice that in Diagram 4.4, we have defined the maps

$$\Delta_{f_s,d}: X_i \times_{Y_j} T^*Y_j \rightarrow T^*X_i, \quad \Delta_{f_s,\pi}: X_i \times_{Y_j} T^*Y_j \rightarrow T^*Y_j.$$

where  $f_s: X_i \rightarrow Y_j$  is given by the sequence of morphisms of manifolds  $X_i \xrightarrow{f_i} Y_i \xrightarrow{\Delta_s^{(Y)}} Y_j$ .

**Definition 4.3.** Let us fix a closed  $\mathbb{R}^+$ -conic subset  $\Lambda \subset T^*Y$ .

We say that  $\Lambda$  is transversal to  $f$  if for each  $s: j \rightarrow i$ , the map  $f_s: X_i \rightarrow Y_j$  is non-characteristic with respect to  $\Lambda_j$ , that is  $\Delta_{f_s,d}^{-1}(T_{X_i}^*X_i) \cap \Delta_{f_s,\pi}^{-1}(\Lambda_j) \subset X_i \times_{Y_j} T_{Y_j}^*Y_j$ .

Note that to be transversal is stronger than to be non characteristic. When  $f: X \rightarrow Y = X$  is the identity, we recover the notion of Definition 4.2 (c).

**Notation 4.4.** We will denote by  $\Delta_s^{(Y)}$ ,  $\Delta_s^{(X)}$ ,  $\forall s \in \text{Hom}(\mathcal{I})$ , the closed embeddings of  $Y$  and  $X$  respectively. However when it is clear from the context we will omit the superscript.

**Lemma 4.5.** Let  $\Lambda$  be a closed  $\mathbb{R}^+$ -conic subset of  $T^*Y$ ; moreover assume  $\Lambda$  is transversal to  $f$ . Then for every  $s: j \rightarrow i$

- (a)  $\Delta_s^{(Y)}$  is non-characteristic for  $\Lambda_j$  in a neighborhood of  $f_i(X_i)$ ;
- (b)  $\Delta_s^{(X)}$  is non-characteristic for  $f_{jd}f_{j\pi}^{-1}(\Lambda_j)$ .

*Proof.* (a) By hypothesis the morphism  $f_s$  is non-characteristic for  $\Lambda_j$ . The result follows from Lemma 4.10 pag.65 in [Ka03] applied to the following commutative diagram:

$$(4.5) \quad \begin{array}{ccccc} T^*X_i & \xleftarrow{f_{id}} & X_i \times_{Y_i} TY_i & \xleftarrow{\varphi} & X_i \times_{Y_j} T^*Y_j \\ \downarrow f_{i\pi} & & \downarrow & & \downarrow \psi \\ T^*Y_i & \xleftarrow{\Delta_{sd}} & Y_i \times_{Y_j} T^*Y_j & & \\ & & \downarrow \Delta_{s\pi} & & \\ & & T^*Y_j. & & \end{array}$$

(b) Same argument applied to  $f_s = f_j \circ \Delta_s^{(X)}$ .

□

## 5 Sheaves on Ran-manifolds

Let  $X$  be a Ran-manifold and  $\mathcal{J}$  a small category. Consider  $D^b(\mathbf{k}_{X_i})$  the bounded derived category of  $\text{Mod}(\mathbf{k}_{X_i})$ . Define the functor  $\mathfrak{S}: \mathcal{J}^{op} \rightarrow \mathbf{Cat}$  as follows:

- (a) to any  $i \in \mathcal{J}$  set  $D^b(\mathbf{k}_{X_i})$ ;
- (b) for any morphism  $s: i_1 \rightarrow i_2$  set  $\rho_s: D^b(\mathbf{k}_{X_{i_1}}) \rightarrow D^b(\mathbf{k}_{X_{i_2}})$ , with  $\rho_s := \Delta_s^{-1}$ ;
- (c) for  $s: i_1 \rightarrow i_2$ ,  $t: i_2 \rightarrow i_3$  the isomorphism of functors  $c_{s,t}$  is  $\Delta_t^{-1} \circ \Delta_s^{-1} \simeq \Delta_{t \circ s}^{-1}$ .

**Definition 5.1.** We set

$$D^b(\mathbf{k}_X) := 2 \varprojlim_{i,s} (D^b(\mathbf{k}_{X_i}), \Delta_s^{-1}).$$

Hence, an object  $D^b(\mathbf{k}_X)$  is the data of  $\{(F_i, \varphi_s)\}_{i,s}$  ( $i \in \mathcal{J}, s \in \text{Mor}(\mathcal{J})$ ) with  $F_i \in D^b(\mathbf{k}_{X_i})$  and  $\varphi_s: \Delta_s^{-1} F_j \xrightarrow{\sim} F_i$  satisfying the compatibility conditions in def 3.2.

### Inverse images

Let  $f: X \rightarrow Y$  be a morphism of Ran manifolds.

**Definition 5.2.** The inverse image functor  $f^{-1}: D^b(\mathbf{k}_Y) \rightarrow D^b(\mathbf{k}_X)$  is defined as follows. Given  $G \in D^b(\mathbf{k}_Y)$ ,  $G = \{(G_i, \psi_s)\}_{i,s}$  with  $i \in \mathcal{J}, s \in \text{Hom}_{\mathcal{J}}(j, i)$ , we set  $f^{-1}G = \{(F_i, \varphi_s)\}_{i,s}$  where  $F_i = f_i^{-1}G_i$  and  $\varphi_s: \Delta_s^{-1} F_j \simeq F_i$  is given by the isomorphisms

$$\Delta_s^{-1} F_j = \Delta_s^{-1} f_j^{-1} G_j \simeq f_i^{-1} \Delta_s^{-1} G_j \simeq f_i^{-1} G_i = F_i.$$

## 6 $\mathcal{D}$ -modules on complex Ran-manifolds

The  $\infty$ -category of  $\mathcal{D}$ -modules on Ran spaces has been introduced and studied in [FG11]. We propose here a more elementary approach.

First we shall adapt Definition 5.1 in the  $\mathcal{D}$ -modules setting. Let  $X$  be a Ran-manifold and  $\mathcal{J}$  a small category. Define the functor  $\mathfrak{S}: \mathcal{J}^{op} \rightarrow \mathbf{Cat}$  by

- (a) to  $i \in \mathcal{J}$ ,  $\mathfrak{S}(i) = D^b(\mathcal{D}_{X_i})$ ;
- (b) for any morphism  $s: i_1 \rightarrow i_2$ ,  $\rho_s: D_{\text{coh}}^b(\mathcal{D}_{X_{i_1}}) \rightarrow D_{\text{coh}}^b(\mathcal{D}_{X_{i_2}})$  is the functor  $\Delta_s^D$ ;
- (c) for  $s: i_1 \rightarrow i_2$ ,  $t: i_2 \rightarrow i_3$ , the isomorphism of functors  $c_{s,t}$  is the obvious one  $\Delta_t^D \circ \Delta_s^D \simeq \Delta_{t \circ s}^D$ .

**Definition 6.1.** We set

$$D^b(\mathcal{D}_X) = 2 \varprojlim_{i,s} (D^b(\mathcal{D}_{X_i}), \Delta_s^D).$$

**Definition 6.2.** Denote by  $D_{\text{coh nc}}^b(\mathcal{D}_{X_j})$  the full triangulated subcategory of  $D_{\text{coh}}^b(\mathcal{D}_{X_j})$  whose objects  $\mathcal{M}_j$  are such that for every  $s \in \text{Hom}_{\mathcal{J}}(j, i)$  the corresponding morphism  $\Delta_s: X_i \rightarrow X_j$  is non-characteristic with respect to  $\mathcal{M}_j$ .

**Definition 6.3.** We set

$$D_{\text{coh nc}}^b(\mathcal{D}_X) = 2 \varprojlim_{i, s} (D_{\text{coh nc}}^b(\mathcal{D}_{X_i}), \Delta_s^D).$$

Let  $X$  and  $Y$  be two complex Ran-manifolds and  $f: X \rightarrow Y$ . We shall mimic the previous definitions

**Definition 6.4.** Denote with  $D_{\text{coh } f\text{-nc}}^b(\mathcal{D}_{Y_j})$  the full triangulated subcategory of  $D_{\text{coh}}^b(\mathcal{D}_{Y_j})$  whose objects  $\mathcal{N}_j$  are such that for every  $s \in \text{Hom}_{\mathcal{J}}(j, i)$  the corresponding morphism  $f_s = f_j \circ \Delta_s^{(X)}: X_i \rightarrow Y_j$  is non-characteristic with respect to  $\mathcal{N}_j$ .

**Definition 6.5.** We set  $D_{\text{coh } f\text{-nc}}^b(\mathcal{D}_Y)$  as the 2-limit category

$$2 \varprojlim_{i, s} (D_{\text{coh } f\text{-nc}}^b(\mathcal{D}_{Y_i}), \Delta_s^D).$$

Of course the two definitions coincide in the case  $f = id$ .

**Example 6.6.** Let  $X$  be a complex Ran-manifold. Since  $\Delta_s^D \mathcal{O}_{X_j} \simeq \mathcal{O}_{X_i}$ , the family  $\{\mathcal{O}_{X_i}\}_i$  defines an object in  $D_{\text{coh}}^b(\mathcal{D}_X)$ .

**Lemma 6.7.** Any object  $\mathcal{N} \in D_{\text{coh } f\text{-nc}}^b(\mathcal{D}_Y)$  belongs to the subcategory of  $D_{\text{coh}}^b(\mathcal{D}_Y)$  consisting of objects s.t. for every  $s: j \rightarrow i$ ,  $\Delta_s$  is non-characteristic for  $\mathcal{N}_j$  in a neighborhood of  $f_i(X_i)$ .

*Proof.* It is a consequence of Lemma 4.5. □

### The inverse image functor $f^D$

Let  $f: X \rightarrow Y$  be a morphism of Ran-manifolds. We will need the following result (which holds in view of Lemma 4.5).

**Lemma 6.8.** For all  $s: j \rightarrow i$  the corresponding map  $\Delta_s: X_i \rightarrow X_j$  is non characteristic for the corresponding  $f_j^D \mathcal{N}_j$ .

**Proposition 6.9.** The map  $f$  induces a functor  $f^D: D_{\text{coh } f\text{-nc}}^b(\mathcal{D}_Y) \rightarrow D_{\text{coh nc}}^b(\mathcal{D}_X)$

*Proof.* We set  $f^D := \{f_i^D\}_{i \in \mathcal{J}}$ . This inverse image functor is well-defined due to Lemma 6.8. In view of Lemma 3.5 it is enough to prove that  $\{f_i^D\}_{i \in \mathcal{J}}$  is a functor of prestack. In particular this means that each functor  $f_i^D$  commutes with the restriction morphisms. By Definition 4.1 of morphism of Ran manifolds we have  $\Delta_s \circ f_i = f_j \circ \Delta_s$  and applying the inverse image functor we obtain the desired property  $f_i^D \circ \Delta_s^D \simeq \Delta_s^D \circ f_j^D$ . □

**The functor**  $\mathrm{R}\mathcal{H}om_{\mathcal{D}_Y}(-, \mathcal{O}_Y)$

**Lemma 6.10.** *The functor  $\mathrm{R}\mathcal{H}om_{\mathcal{D}_Y}(-, \mathcal{O}_Y) := \{\mathrm{R}\mathcal{H}om_{\mathcal{D}_{Y_i}}(-, \mathcal{O}_{Y_i})\}_{i \in I}$ ,*

$$\mathrm{R}\mathcal{H}om_{\mathcal{D}_Y}(-, \mathcal{O}_Y) : \mathrm{D}_{\mathrm{coh}}^b(\mathcal{D}_Y) \rightarrow \mathrm{D}_{\mathrm{coh}}^b(\mathbf{k}_X)$$

*is well defined.*

*Proof.* Consider an object  $\{\mathcal{N}_i, \varphi_s\}_{i,s} \in \mathrm{D}_{\mathrm{coh}}^b(\mathcal{D}_Y)$ ; by the CKK theorem (Theorem 2.4), there is the isomorphism  $\Delta_s^{-1} \mathrm{R}\mathcal{H}om_{\mathcal{D}_{Y_i}}(\mathcal{N}_i, \mathcal{O}_{Y_i}) \simeq \mathrm{R}\mathcal{H}om_{\mathcal{D}_{Y_j}}(\Delta_s^D \mathcal{N}_i, \mathcal{O}_{Y_j})$ .

By the composition of the last isomorphism with  $\varphi_s : \Delta_s^D \mathcal{N}_i \xrightarrow{\sim} \mathcal{N}_j$  we obtain the isomorphism  $\Delta_s^{-1} \mathrm{R}\mathcal{H}om_{\mathcal{D}_{Y_i}}(\mathcal{N}_i, \mathcal{O}_{Y_i}) \xrightarrow{\sim} \mathrm{R}\mathcal{H}om_{\mathcal{D}_{Y_j}}(\mathcal{N}_j, \mathcal{O}_{Y_j})$ . Lemma 3.5 completes the proof.  $\square$

### The Cauchy-Kowaleski-Kashiwara theorem for Ran spaces

**Theorem 6.11.** *There is an isomorphism of functors of prestacks*

$$\theta_\infty : f^{-1} \circ \mathrm{R}\mathcal{H}om_{\mathcal{D}_Y}(-, \mathcal{O}_Y) \xrightarrow{\sim} \mathrm{R}\mathcal{H}om_{\mathcal{D}_Y}(-, \mathcal{O}_Y) \circ f^D.$$

*In other words for any  $\mathcal{N} \in \mathrm{D}_{\mathrm{coh} \, f\text{-nc}}^b(\mathcal{D}_Y)$  we have the isomorphism*

$$f^{-1} \mathrm{R}\mathcal{H}om_{\mathcal{D}_Y}(\mathcal{N}, \mathcal{O}_Y) \xrightarrow{\sim} \mathrm{R}\mathcal{H}om_{\mathcal{D}_X}(f^D \mathcal{N}, \mathcal{O}_X)$$

*functorial in  $\mathcal{N} \in \mathrm{D}_{\mathrm{coh} \, f\text{-nc}}^b(\mathcal{D}_Y)$ .*

The theorem is visualized by the following quasi commutative diagram:

$$(6.1) \quad \begin{array}{ccc} \mathrm{D}_{\mathrm{coh} \, f\text{-nc}}^b(\mathcal{D}_Y) & \xrightarrow{f^D} & \mathrm{D}_{\mathrm{coh} \, \mathrm{nc}}^b(\mathcal{D}_X) \\ \mathrm{R}\mathcal{H}om_{\mathcal{D}_Y}(-, \mathcal{O}_Y) \downarrow & & \downarrow \mathrm{R}\mathcal{H}om_{\mathcal{D}_X}(-, \mathcal{O}_X) \\ \mathrm{D}^b(\mathbf{k}_Y) & \xrightarrow{f^{-1}} & \mathrm{D}^b(\mathbf{k}_X). \end{array}$$

*Proof.* In view of Lemmas 3.6, 3.6, 3.8, it is enough to observe that each morphism

$$\theta_i : f_i^{-1} \circ \mathrm{R}\mathcal{H}om_{\mathcal{D}_{Y_i}}(-, \mathcal{O}_{Y_i}) \xrightarrow{\sim} \mathrm{R}\mathcal{H}om_{\mathcal{D}_{Y_i}}(-, \mathcal{O}_{Y_i}) \circ f_i^D$$

is an isomorphism due to Theorem 2.4.  $\square$

**Example 6.12.** Let  $\mathcal{J}$  be the category of finite non-empty sets and surjective maps. Let  $X$  be a complex manifold. The Ran space  $\mathrm{Ran} \, X$  in [BD04] is defined as follows: to  $I \in \mathcal{J}$  one associates the product manifold  $X^I$  and to a surjection  $s : J \rightarrow I$  is associated the diagonal embedding

$$(6.2) \quad \delta_s : X^I \hookrightarrow X^J$$

which maps  $\{x_i\}_{i \in I} \in X^I$  to  $\{x_j\}_{j \in J} \in X^J$  with  $x_j = x_i$  if  $s(j) = i$ .

## Appendix

There are two other natural categories,  $\mathfrak{S}^+(\mathcal{J})$  and  $\mathfrak{S}^-(\mathcal{J})$ .

**Definition 6.13.** (a) An object  $F$  of  $\mathfrak{S}^+(\mathcal{J})$  (resp.  $\mathfrak{S}^-(\mathcal{J})$ ) is a family  $\{(F_i, \phi_s)\}_{i,s}$  ( $i \in \mathcal{J}, s \in \text{Mor}(\mathcal{J})$ ) where

- (i) for any  $i \in \mathcal{J}$ ,  $F_i$  is an object of  $\mathfrak{S}(i)$ ,
- (ii) for any morphism  $s: i_1 \rightarrow i_2$  in  $\mathcal{J}$ ,  $\phi_s: F_{i_1} \rightarrow \rho_s(F_{i_2})$  (resp.  $\phi_s: \rho_s(F_{i_2}) \rightarrow F_{i_1}$ ) is a morphism such that
  - for all  $i \in \mathcal{J}$ ,  $\phi_{\text{id}_i} = \text{id}_{F_i}$ ,
  - for any sequence  $i_1 \xrightarrow{s} i_2 \xrightarrow{t} i_3$  of morphisms in  $\mathcal{J}$ , the following diagram commutes

$$(6.3) \quad \begin{array}{ccc} \rho_s \rho_t(F_{i_3}) & \xleftarrow{\rho_s(\phi_t)} & \rho_s(F_{i_2}) \\ c_{s,t} \Big| \wr & & \uparrow \phi_s \\ \rho_{t \circ s}(F_{i_3}) & \xleftarrow{\phi_{t \circ s}} & F_{i_1} \end{array}$$

(resp. the following diagram commutes

$$(6.4) \quad \begin{array}{ccc} \rho_s \rho_t(F_{i_3}) & \xrightarrow{\rho_s(\phi_t)} & \rho_s(F_{i_2}) \\ c_{s,t} \Big| \wr & & \downarrow \phi_s \\ \rho_{t \circ s}(F_{i_3}) & \xrightarrow{\phi_{t \circ s}} & F_{i_1} \end{array}$$

- (b) A morphism  $f: \{(F_i, \phi_s)\}_{i,s} \rightarrow \{(F'_i, \phi'_s)\}_{i,s}$  in  $\mathfrak{S}^+(\mathcal{J})$  (resp.  $\mathfrak{S}^-(\mathcal{J})$ ) is a family of morphisms  $f_i: F_i \rightarrow F'_i$  such that for any  $s: i_1 \rightarrow i_2$ , the diagram below commutes:

$$(6.5) \quad \begin{array}{ccc} \rho_s(F_{i_2}) & \xrightarrow{\rho_s(f_{i_2})} & \rho_s(F'_{i_2}) \\ \uparrow \phi_s & & \uparrow \phi'_s \\ F_{i_1} & \xrightarrow{f_{i_1}} & F'_{i_1} \end{array}$$

(resp. the diagram below commutes:

$$(6.6) \quad \begin{array}{ccc} \rho_s(F_{i_2}) & \xrightarrow{\rho_s(f_{i_2})} & \rho_s(F'_{i_2}) \\ \downarrow \phi_s & & \downarrow \phi'_s \\ F_{i_1} & \xrightarrow{f_{i_1}} & F'_{i_1} \end{array}$$

- (c) We consider  $\mathfrak{S}(\mathcal{J})$  as the full subcategory of  $\mathfrak{S}^+(\mathcal{J})$  or  $\mathfrak{S}^-(\mathcal{J})$  consisting of objects  $\{(F_i, \phi_s)\}_{i \in \mathcal{J}, s \in \text{Mor}(\mathcal{J})}$  such that for all  $s \in \text{Mor}(\mathcal{J})$ , the morphisms  $\phi_s$  are isomorphisms and we denote by  $\iota_{\mathcal{J}}^+: \mathfrak{S}(\mathcal{J}) \rightarrow \mathfrak{S}^+(\mathcal{J})$  and  $\iota_{\mathcal{J}}^-: \mathfrak{S}(\mathcal{J}) \rightarrow \mathfrak{S}^-(\mathcal{J})$  the natural faithful functors.

## References

- [BD04] A. Beilinson, V. Drinfeld, *Chiral algebras*, volume 51 of A. M. S. Colloquium Publications. American Mathematical Society, Providence, RI, 2004.
- [FG11] J. Francis and D. Gaitsgory, *Chiral Koszul duality*, arXiv:1103.5803.
- [GS06] S. Guillermou and P. Schapira, *Operations on projective limits of categories of sheaves*, unpublished (2006).
- [Ka70] M. Kashiwara, *Algebraic study of systems of partial differential equations*, Thesis, Tokyo Univ. (1970), translated by A. D'Agnolo and J-P. Schneiders, Mémoires Soc. Math. France **63** (1995).
- [Ka75] M. Kashiwara, *On the maximally overdetermined system of linear differential equations I*, Publ. Res. Inst. Math. Sci. **10** (1974/75) 563–579.
- [Ka03] ———, *D-modules and microlocal calculus*, volume 217 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 2003.
- [KS90] M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Grundlehren der Mathematischen Wissenschaften, **292**. Springer-Verlag, 1990, x+512 pp.
- [KS06] ———, *Categories and Sheaves*, volume 332 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2006.
- [SGA4] S-G-A 4, *Théorie des topos et cohomologie étale des schémas*, Sémin. Géom. Algébrique (1963-64) by M. Artin, A. Grothendieck and J.-L. Verdier, Lecture Notes in Math. **260, 270, 305** Springer Verlag (1972/73).

Giuseppe Bonavolontà,  
Campus Kirchberg, Mathematics Research Unit  
6, rue R. Coudenhove-Kalergi, L-1359 Luxembourg City