

Global Well-posedness for the Generalized Navier-Stokes System

Zeng Zhang¹ Zhaoyang Yin²

Department of Mathematics, Sun Yat-sen University,
510275, Guangzhou, P. R. China.

Abstract

In this paper we investigate well-posedness of the Cauchy problem of the three dimensional generalized Navier-Stokes system. We first establish local well-posedness of the GNS system for any initial data in the Fourier-Herz space χ^{-1} . Then we show that if the χ^{-1} norm of the initial data is smaller than $C\nu$ in the GNS system where ν is the viscosity coefficient, the corresponding solution exists globally in time. Moreover, we prove global well-posedness of the Navier-Stokes system without norm restrictions on the corresponding solutions provided the χ^{-1} norm of the initial data is less than ν . Our obtained results cover and improve recent results in [2, 11].

2010 Mathematics Subject Classification: 35Q30, 76D05, 35A01, 35A02

Keywords: Generalized Navier-Stokes system; Navier-Stokes system; Local well-posedness; Global well-posedness; Fourier-Herz space.

Contents

1 Introduction

2

¹Corresponding author. Email: zhangzeng534534@163.com; Tel.: +8613725201644;
Address: No. 135, Xingang Xi Road, Guangzhou, 510275, P. R. China

²Email: mcsyzy@mail.sysu.com.cn

2 Preliminaries	5
3 Proofs of main theorems	6
References	14

1 Introduction

In this paper we consider the generalized Navier-Stokes system

$$(GNS_\nu) \quad \begin{cases} \partial_t u - \nu \Delta u = Q(u, u), & t > 0, x \in \mathbb{R}^3, \\ u(x, 0) = u_0, \end{cases}$$

with ν being a positive constant and the bilinear operator Q defined as

$$(1.1) \quad Q^j(u, v) = \sum_{k,l,m=1}^3 q_{k,l}^{j,m} \partial_m (u^k v^l), \quad j = 1, 2, 3,$$

where $q_{k,l}^{j,m}(a) = \sum_{n,p=1}^3 a_{k,l}^{j,m,p,n} \mathcal{F}^{-1}(\frac{\xi_n \xi_p}{|\xi|^2} \widehat{a}(\xi))$, and $a_{k,l}^{j,m,p,n}$ are real numbers.

It is obvious that the incompressible Navier-Stokes system

$$(NS_\nu) \quad \begin{cases} \partial_t u - \nu \Delta u = \mathcal{P} \nabla (u \otimes u), & t > 0, x \in \mathbb{R}^3, \\ \operatorname{div} u = 0, \\ u(x, 0) = u_0, \end{cases}$$

is a particular case of the system (GNS_ν) . Here u stands for the velocity field of the fluid, ν is the viscosity and \mathcal{P} is the Leray projection operator defined by the formula:

$$(1.2) \quad \mathcal{F}(\mathcal{P}f)^j(\xi) = \mathcal{F}(f)^j(\xi) - \frac{1}{|\xi|^2} \sum_{k=1}^3 \xi_j \xi_k \mathcal{F}(f)^k(\xi), \quad j = 1, 2, 3.$$

From now on we always assume that the initial data u_0 is divergence free and C denotes a generic constant.

It is well known that the space BMO^{-1} is the largest space which is included in the tempered distribution and enjoys the property of translation and scaling invariant (see [4, 6] for instance). The global well-posedness for the Navier-Stokes system in the space BMO^{-1} was studied by Koch and Tataru [6]. Many works in subspaces of the space BMO^{-1} , such as $\dot{H}^{\frac{3}{2}-1}(\mathbb{R}^3)$, $L^3(\mathbb{R}^3)$,

and $\dot{B}_{p,\infty}^{-1+\frac{3}{p}}(\mathbb{R}^3)$ also have been down: the Navier-Stokes system is known to be globally well-posed for sufficiently small data $u_0 \in \dot{H}^{\frac{3}{2}-1}(\mathbb{R}^3)$, and locally well-posed for any $u_0 \in \dot{H}^{\frac{3}{2}-1}(\mathbb{R}^3)$, as proved by Fujita and Kato [1]; the global well-posedness for small data is due to Kato [7] in $L^3(\mathbb{R}^3)$, Cannone, Meyer and Planchon [9] in $\dot{B}_{p,\infty}^{-1+\frac{3}{p}}(\mathbb{R}^3)$, respectively. For more studies in these spaces, the reader may refer to [5, 8, 10]. Let us mention that the well-posedness for the Navier-Stokes system in those above spaces can be extended to the generalized Navier-Stokes system [3].

Recently Lei and Lin [2] studied a new space

$$\chi^{-1} = \{f \in \mathcal{D}'(\mathbb{R}^3); \int_{\mathbb{R}^3} |\xi|^{-1} |\widehat{f}| d\xi < \infty\},$$

which is contained in BMO^{-1} [2] and is equivalent to the Fourier-Herz space \dot{B}_1^{-1} [11]. They pointed out that $H^s(\mathbb{R}^3)$ ($s > \frac{1}{2}$) $\subseteq \chi^{-1}$, and they also presented an example to show that $H^{\frac{3}{2}-1}(\mathbb{R}^3) \not\subseteq \chi^{-1}$. Here we give an example to show that $\chi^{-1} \not\subseteq \dot{H}^{\frac{3}{2}-1}(\mathbb{R}^3)$. In fact, let $f(x) = \mathcal{F}^{-1}(\frac{1}{|\xi|} h(|\xi|))$, where $h(r)$ is defined by

$$h(r) = \begin{cases} 2^{\frac{j+1}{2}}, & 1 - 2^{-j} \leq r < 1 - 2^{-(j+1)}, j = 0, 1, \dots, \\ 0, & r \geq 1. \end{cases}$$

It is easy to deduce that

$$\int_{\mathbb{R}^3} |\xi|^{-1} |\widehat{f}(\xi)| d\xi \leq C \int_0^\infty h(r) dr = C \sum_{j=0}^\infty 2^{\frac{j+1}{2}} 2^{-(j+1)} < \infty.$$

Similarly, we can also get

$$\int_{\mathbb{R}^3} |\xi| |\widehat{f}(\xi)|^2 d\xi = C \int_0^\infty r h(r)^2 dr \geq C \sum_{j=0}^\infty (1 - 2^{-j}) 2^{j+1} 2^{-(j+1)} = \infty.$$

Combing the above two inequalities yields that $f \in \chi^{-1}$, but $f \notin \dot{H}^{\frac{3}{2}-1}(\mathbb{R}^3)$. Thus, we conclude that χ^{-1} and $\dot{H}^{\frac{3}{2}-1}(\mathbb{R}^3)$ do not contain each other.

Lei and Lin [2] proved that if the initial data u_0 in χ^{-1} satisfying $\|u_0\|_{\chi^{-1}} < \nu$, the system (NS_ν) admits a global mild solution. They also proved that this global mild solution is unique under the condition $\|u\|_{L^\infty(\mathbb{R}^+; \chi^{-1})} < \nu$. Cannone and Wu [11] gave a global well-posedness result for small initial data in a family of critical Fourier-Herz spaces \dot{B}_q^{-1} ($q \in [1, 2]$). They also showed this global solution is unique under the condition $\|u\|_{L^\infty(\mathbb{R}^+; \mathcal{B}_q^{-1}) \cap L^1(\mathbb{R}^+; \mathcal{B}_q^1)} \leq \frac{50}{9} \|u_0\|_{\mathcal{B}_q^{-1}}$.

However, it is not clear whether there exists a solution to the system (NS_ν) for large initial data. Moreover, without norm restrictions on the solutions, is the uniqueness of solutions to the system (NS_ν) still valid?

In the paper we will give definite answers to these two questions. In Section 3, we will solve the system (GNS_ν) by means of a contraction mapping argument (see Lemma 1.1 below). Thus, we can obtain a unique local mild solution to the system (GNS_ν) for any initial data in χ^{-1} and prove the corresponding solution will be global if the initial data is sufficient small. Especially for the Navier-Stokes system, we show that if $\|u_0\|_{\chi^{-1}} < \nu$ then the solution to the system (NS_ν) will be unique and global without norm restrictions on the solutions.

Lemma 1.1. ([3]) *Let E be a Banach space, \mathcal{B} a continuous bilinear map from $E \times E \rightarrow E$, and a positive real number such that $\alpha < \frac{1}{4\|\mathcal{B}\|}$, with*

$$\|\mathcal{B}\| = \sup_{\|u\| \leq 1, \|v\| \leq 1} \|\mathcal{B}(u, v)\|.$$

For any a in the ball $B(0, \alpha)$ in E , then there exists a unique x in $B(0, 2\alpha)$ such that

$$x = a + \mathcal{B}(x, x).$$

We will also use the spaces

$$(1.3) \quad \chi^i = \{f \in \mathcal{S}'(\mathbb{R}^3); \int_{\mathbb{R}^3} |\xi|^i |\hat{f}| d\xi < \infty\}, \quad i = -1, 0, 1.$$

The norm of χ^i is denoted by $\|\cdot\|_{\chi^i}$. Let $T \in (0, \infty]$. Note that the spaces $L^2([0, T]; \chi^0)$ and $L^\infty([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1)$ are Banach spaces and are translation and shift invariant.

Now we are in the position to state our main results:

Theorem 1.2. *Let u_0 be in χ^{-1} . There exists a positive time T such that the system (GNS_ν) has a unique solution u in $L^2([0, T]; \chi^0)$ which also belongs to*

$$C([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1) \cap L^\infty([0, T]; \chi^{-1}).$$

Let T_{u_0} denote the maximal time of existence of such a solution. Then:

(i) There exists a constant C such that if $\|u_0\|_{\chi^{-1}} \leq C\nu$, then

$$T_{u_0} = \infty.$$

(ii) If T_{u_0} is finite, then

$$\int_0^{T_{u_0}} \|u(t)\|_{\chi^0}^2 dt = \infty.$$

Theorem 1.3. *Let u_0 be in χ^{-1} . There exists a positive time T such that the system (NS_ν) has a unique solution u in $L^2([0, T]; \chi^0)$ which also belongs to*

$$C([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1) \cap L^\infty([0, T]; \chi^{-1}).$$

Let T_{u_0} denote the maximal time of existence of such a solution. Then:

(i) There exists a constant C such that if $\|u_0\|_{\chi^{-1}} < \nu$, then

$$T_{u_0} = \infty.$$

(ii) If T_{u_0} is finite, then

$$\int_0^{T_{u_0}} \|u(t)\|_{\chi^0}^2 dt = \infty.$$

Remark 1.4. *Although the result (i) in Theorem 1.2 is the same as Theorem (1.1) in [2], our method here is different from their method in [2]. In particular, our proof relies on the obtained local well-posedness result and the blow-up criterion (ii), but not on additional norm restrictions on the corresponding solutions.*

2 Preliminaries

Let $B(u, v)$ be the solution to the heat equation

$$(2.1) \quad \begin{cases} \partial_t B(u, v) - \nu \Delta B(u, v) = Q(u, v), \\ B(u, v)|_{t=0} = 0, \end{cases}$$

with the bilinear operator Q defined as in (1.1).

Solving (GNS_ν) amounts to finding a fixed point for the map

$$u \mapsto e^{t\nu\Delta} u_0 + B(u, u).$$

By Duhamel's formula in Fourier space and (1.1), we have

$$(2.2) \quad \begin{aligned} |\widehat{B(u, v)}(t, \xi)| &= \left| \int_0^t e^{-\nu(t-s)|\xi|^2} \widehat{Q(u, v)}(s, \xi) ds \right| \\ &\leq C \int_0^t e^{-\nu(t-s)|\xi|^2} |\xi| (|\widehat{u}| * |\widehat{v}|)(s, \xi) ds. \end{aligned}$$

We now give two useful propositions which will be used in the sequel.

Proposition 2.1.

$$(2.3) \quad \|u\|_{L^2([0,T];\chi^0)}^2 \leq \|u\|_{L^\infty([0,T];\chi^{-1})} \|u\|_{L^1([0,T];\chi^1)}.$$

Proof. It is easy to check that

$$\begin{aligned} \|u\|_{L^2([0,T];\chi^0)}^2 &= \int_0^T \left(\int_{\mathbb{R}^3} |\widehat{u}(s, \xi) d\xi \right)^2 dt \\ &\leq \int_0^T \left(\int_{\mathbb{R}^3} |\xi|^{-1} |\widehat{u}(s, \xi) d\xi \right) \left(\int_{\mathbb{R}^3} |\xi| |\widehat{u}(s, \xi) d\xi \right) dt \\ &\leq \|u\|_{L^\infty([0,T];\chi^{-1})} \|u\|_{L^1([0,T];\chi^1)}. \end{aligned}$$

Proposition 2.2. *A constant C exists such that*

$$(2.4) \quad \|B(u, v)\|_{L^2([0,T];\chi^0)} \leq \frac{C}{\nu^{\frac{1}{2}}} \|u\|_{L^2([0,T];\chi^0)} \|v\|_{L^2([0,T];\chi^0)}.$$

Proof. Thanks to the inequality (2.2) and Minkowski's inequality, we have

$$\begin{aligned} \|B(u, v)\|_{L^2([0,T];\chi^0)} &\leq C \left\| \int_{\mathbb{R}^3} \int_0^T I_{[0,t]}(s) e^{-\nu(t-s)|\xi|^2} |\xi| (|\widehat{u}| * |\widehat{v}|)(s, d\xi) ds d\xi \right\|_{L^2(0,T)} \\ &\leq C \int_{\mathbb{R}^3} \int_0^T \left(\int_0^T I_{[0,t]}(s) e^{-2\nu(t-s)|\xi|^2} |\xi|^2 dt \right)^{\frac{1}{2}} (|\widehat{u}| * |\widehat{v}|)(s, d\xi) ds d\xi \\ &\leq C \int_{\mathbb{R}^3} \int_0^T \frac{1}{\nu^{\frac{1}{2}}} (|\widehat{u}| * |\widehat{v}|)(s, d\xi) ds d\xi \\ &\leq \frac{C}{\nu^{\frac{1}{2}}} \int_0^T \|\widehat{u}(s)\|_{L^1} \|\widehat{v}(s)\|_{L^1} ds \\ &\leq \frac{C}{\nu^{\frac{1}{2}}} \|u\|_{L^2([0,T];\chi^0)} \|v\|_{L^2([0,T];\chi^0)}. \end{aligned}$$

3 Proofs of main theorems

To prove the first part of Theorem 1.2. we shall use Lemma 1.1. Given some $u_0 \in \chi^{-1}$, thanks to Minkowski's inequality, we have

$$\begin{aligned} (3.1) \quad \|e^{\nu t \Delta} u_0\|_{L^2([0,T];\chi^0)} &= \left(\int_0^T \left(\int_{\mathbb{R}^3} e^{-\nu t |\xi|^2} |\widehat{u_0}(\xi) d\xi \right)^2 dt \right)^{\frac{1}{2}} \\ &\leq \int_{\mathbb{R}^3} \left(\int_0^T e^{-2\nu t |\xi|^2} |\widehat{u_0}|^2(\xi) ds \right)^{\frac{1}{2}} d\xi \end{aligned}$$

$$\begin{aligned} &\leq \int_{R^3} \frac{1}{(2\nu|\xi|^2)^{\frac{1}{2}}} |\widehat{u_0}|(\xi) d\xi \\ &\leq \frac{1}{(2\nu)^{\frac{1}{2}}} \|u_0\|_{\chi^{-1}}. \end{aligned}$$

Thus, combining Proposition 2.2 and the inequality (3.1) gives that if $\|u_0\|_{\chi^{-1}} \leq \frac{\nu}{2^{\frac{3}{2}}C_0}$, with $C_0 > C$, then

$$\|e^{\nu t \Delta} u_0\|_{L^2([0,T];\chi^0)} \leq \frac{1}{4\frac{C_0}{\nu^{\frac{1}{2}}}} < \frac{1}{4\|B\|}.$$

According to Lemma 1.1, there exists a unique solution of the system (GNS_ν) in the ball with center 0 and radius $\frac{\nu^{\frac{1}{2}}}{2C_0}$ in the space $L^2([0,T];\chi^0)$.

We now consider the case of a large initial data $u_0 \in \chi^{-1}$. We shall split u_0 into a small part in χ^{-1} and a large part with compactly supported Fourier transform. For that, we fix some positive real number ρ_{u_0} such that

$$(3.2) \quad \int_{|\xi| \geq \rho_{u_0}} |\xi|^{-1} |\widehat{u_0}|(\xi) d\xi \leq \frac{\nu}{2^{\frac{5}{2}}C_0}.$$

Using the inequality (3.1) again and defining $u_0^b = \mathcal{F}^{-1}(I_{B(0,\rho_{u_0})}(\xi)\widehat{u_0}(\xi))$, we get

$$\|e^{\nu t \Delta} u_0\|_{L^2([0,T];\chi^0)} \leq \frac{\nu^{\frac{1}{2}}}{8C_0} + \|e^{\nu t \Delta} u_0^b\|_{L^2([0,T];\chi^0)}.$$

From which we can deduce that

$$\begin{aligned} \|e^{\nu t \Delta} u_0^b\|_{L^2([0,T];\chi^{-1})} &= \left(\int_0^T \left(\int_{|\xi| \leq \rho_{u_0}} e^{-\nu t |\xi|^2} |\widehat{u_0}|(\xi) d\xi \right)^2 dt \right)^{\frac{1}{2}} \\ &= \left(\int_0^T \left(\int_{|\xi| \leq \rho_{u_0}} e^{-\nu t |\xi|^2} |\xi| |\xi|^{-1} |\widehat{u_0}|(\xi) d\xi \right)^2 dt \right)^{\frac{1}{2}} \\ &\leq \rho_{u_0} T^{\frac{1}{2}} \|u_0\|_{\chi^{-1}}. \end{aligned}$$

Thus if

$$(3.3) \quad T \leq \left(\frac{\nu^{\frac{1}{2}}}{8\rho_{u_0}C_0\|u_0\|_{\chi^{-1}}} \right)^2,$$

then we conclude the existence of a unique solution in the ball with center 0 and radius $\frac{\nu^{\frac{1}{2}}}{2C_0}$ in the space $L^2([0,T];\chi^0)$.

Next we claim that if u is a solution of the system (GNS_ν) in $L^2([0, T]; \chi^0)$, then u also belongs to $C([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1) \cap L^\infty([0, T]; \chi^{-1})$. In fact, it is easy to deduce that

$$\begin{aligned} \|B(u, v)(t)\|_{\chi^{-1}} &\leq C \int_{\mathbb{R}^3} \int_0^t |\xi|^{-1} e^{-\nu(t-s)|\xi|^2} |\xi| (|\hat{u}| * |\hat{v}|)(s, \xi) ds d\xi \\ &\leq C \int_{\mathbb{R}^3} \int_0^t e^{-\nu(t-s)|\xi|^2} (|\hat{u}| * |\hat{v}|)(s, \xi) ds d\xi \\ &\leq C \int_0^t \int_{\mathbb{R}^3} (|\hat{u}| * |\hat{v}|)(s, \xi) d\xi ds \\ &\leq C \|u\|_{L^2([0, T]; \chi^0)} \|v\|_{L^2([0, T]; \chi^0)}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \|B(u, v)\|_{L^1([0, T]; \chi^1)} &\leq C \int_0^T \int_{\mathbb{R}^3} \int_0^t |\xi| I_{[0, t]}(s) e^{-\nu(t-s)|\xi|^2} |\xi| (|\hat{u}| * |\hat{v}|)(s, \xi) ds d\xi dt \\ &\leq C \int_{\mathbb{R}^3} \int_0^T \left(\int_0^t I_{[0, t]}(s) e^{-\nu(t-s)|\xi|^2} |\xi|^2 dt \right) (|\hat{u}| * |\hat{v}|)(s, \xi) ds d\xi \\ &\leq C \int_{\mathbb{R}^3} \int_0^T \frac{1}{\nu} (|\hat{u}| * |\hat{v}|)(s, \xi) ds d\xi \\ &\leq \frac{C}{\nu} \|u\|_{L^2([0, T]; \chi^0)} \|v\|_{L^2([0, T]; \chi^0)}. \end{aligned}$$

Combing the above two inequalities yields that

$$B(u, u) \in L^\infty([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1).$$

Noticing the following two facts:

$$\|e^{\nu t \Delta} u_0\|_{\chi^{-1}} \leq \int_{\mathbb{R}^3} e^{-\nu t |\xi|^2} |\xi|^{-1} |\widehat{u_0}|(\xi) d\xi \leq \|u_0\|_{\chi^{-1}},$$

$$\|e^{\nu t \Delta} u_0\|_{L^1([0, T]; \chi^{-1})} \leq \int_0^T \int_{\mathbb{R}^3} e^{-\nu t |\xi|^2} |\xi| |\widehat{u_0}|(\xi) d\xi dt \leq \frac{1}{\nu} \|u_0\|_{\chi^{-1}},$$

we have

$$e^{\nu t \Delta} u_0 \in L^\infty([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1).$$

We can thus conclude that

$$u \in L^\infty([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1).$$

To get further the regularity of $u(t, x)$ with respect to t , we come back to the system (GNS_ν) . It is obvious that Δu is in $L^1([0, T]; \chi^{-1})$. Thanks to Proposition 2.1 we have

$$\begin{aligned}
 (3.4) \quad & \|Q(u, v)\|_{L^1([0, T]; \chi^{-1})} \\
 &= \int_0^T \int_{\mathbb{R}^3} |\xi|^{-1} |\widehat{Q(u, v)}|(s, \xi) d\xi ds \\
 &\leq C \int_0^T \int_{\mathbb{R}^3} (|\hat{u}| * |\hat{v}|)(s, \xi) d\xi ds \\
 &\leq C \|u\|_{L^2([0, T]; \chi^0)} \|v\|_{L^2([0, T]; \chi^0)} \\
 &\leq C \|u\|_{L^\infty([0, T]; \chi^{-1})}^{\frac{1}{2}} \|u\|_{L^1([0, T]; \chi^1)}^{\frac{1}{2}} \|v\|_{L^\infty([0, T]; \chi^{-1})}^{\frac{1}{2}} \|v\|_{L^1([0, T]; \chi^1)}^{\frac{1}{2}}.
 \end{aligned}$$

Thus, $Q(u, u) \in L^1([0, T]; \chi^{-1})$. Then we conclude that $\partial_t u \in L^1([0, T]; \chi^{-1})$. We hence prove the announced properties:

$$u \in C([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1) \cap L^\infty([0, T]; \chi^{-1}).$$

The uniqueness part relies on the following lemma.

Lemma 3.1. *Let v be a solution in $C([0, T]; \mathcal{S}'(R^3))$ of the Cauchy problem*

$$(3.5) \quad \begin{cases} \partial_t v - \nu \Delta v = f, \\ v|_{t=0} = v_0, \end{cases}$$

with $f \in L^1([0, T]; \chi^{-1})$ and $v_0 \in \chi^{-1}$. Then for any $0 \leq t_0 \leq T$, we have

$$(3.6) \quad \int_{\mathbb{R}^3} \sup_{0 \leq t \leq t_0} |\hat{v}|(t, \xi) |\xi|^{-1} d\xi \leq \|v_0\|_{\chi^{-1}} + \|f\|_{L^1([0, t_0]; \chi^{-1})},$$

$$(3.7) \quad \|v\|_{L^\infty([0, t_0]; \chi^{-1})} + \nu \|v\|_{L^1([0, t_0]; \chi^1)} \leq 2(\|v_0\|_{\chi^{-1}} + \|f\|_{L^1([0, t_0]; \chi^{-1})}).$$

Proof. Due to Duhamel's formula in Fourier space, we can write that

$$\widehat{v}(t, \xi) = e^{-\nu t |\xi|^2} \widehat{v}_0(\xi) + \int_0^t e^{-\nu(t-s) |\xi|^2} \widehat{f}(s, \xi) ds.$$

Then we have

$$|\widehat{v}(t, \xi)| \leq e^{-\nu t |\xi|^2} |\widehat{v}_0(\xi)| + \int_0^t e^{-\nu(t-s) |\xi|^2} |\widehat{f}(s, \xi)| ds.$$

For any $0 \leq t_0 \leq T$, we get

$$\sup_{0 \leq t \leq t_0} |\widehat{v}(t, \xi)| \leq |\widehat{v}_0|(\xi) + \int_0^{t_0} |\widehat{f}|(s, \xi) ds.$$

Taking the L^1 norm with respect to $|\xi|^{-1} d\xi$ allows us to conclude that

$$\begin{aligned} \int_{\mathbb{R}^3} \sup_{0 \leq t \leq t_0} |\widehat{v}(t, \xi)| |\xi|^{-1} d\xi &\leq \int_{\mathbb{R}^3} |\widehat{v}_0|(\xi) |\xi|^{-1} d\xi + \int_{\mathbb{R}^3} \int_0^{t_0} |\widehat{f}|(s, \xi) |\xi|^{-1} ds d\xi \\ &\leq \|v_0\|_{\chi^{-1}} + \|f\|_{L^1([0, t_0]; \chi^{-1})}. \end{aligned}$$

The first result (3.6) is thus proved.

Similarly, taking the L^1 norm with respect to $|\xi|^1 d\xi dt$, one have,

$$\begin{aligned} &\int_0^{t_0} \int_{\mathbb{R}^3} |\widehat{v}|(s, \xi) |\xi| d\xi ds \\ &\leq \int_0^{t_0} \int_{\mathbb{R}^3} e^{-\nu t |\xi|^2} |\widehat{v}_0|(\xi) |\xi| d\xi dt + \int_0^{t_0} \int_{\mathbb{R}^3} \int_0^t e^{-\nu(t-s) |\xi|^2} |\widehat{f}|(s, \xi) |\xi| ds d\xi dt \\ &\leq \int_{\mathbb{R}^3} \left(\int_0^{t_0} e^{-\nu t |\xi|^2} |\xi|^2 dt \right) |\widehat{v}_0|(\xi) |\xi|^{-1} d\xi \\ &\quad + \int_0^{t_0} \int_{\mathbb{R}^3} \left(\int_s^{t_0} e^{-\nu(t-s) |\xi|^2} |\xi|^2 dt \right) |\widehat{f}|(s, \xi) |\xi|^{-1} ds d\xi \\ &\leq \frac{1}{\nu} \left(\int_{\mathbb{R}^3} |\widehat{v}_0|(\xi) |\xi|^{-1} d\xi + \|f\|_{L^1([0, t_0]; \chi^{-1})} \right). \end{aligned}$$

This implies that

$$(3.8) \quad \nu \|v\|_{L^1([0, t_0]; \chi^1)} \leq \|v_0\|_{\chi^{-1}} + \|f\|_{L^1([0, t_0]; \chi^{-1})}.$$

The inequalities (3.6) and (3.8) lead to (3.7).

Now consider two solutions u_1 and u_2 with the same initial data u_0 , and assume that

$$u_i \in C([0, T]; \chi^{-1}) \cap L^1([0, T]; \chi^1) \cap L^\infty([0, T]; \chi^{-1}), \quad i = 1, 2.$$

Let $w = u_1 - u_2$, we note that w satisfies

$$(3.9) \quad \begin{cases} \partial_t w - \nu \Delta w = Q(w, u_1) + Q(u_2, w), \\ w|_{t=0} = 0. \end{cases}$$

Thanks to the inequality (3.4), we have that

$$Q(w, u_1) + Q(u_2, w) \in L^1([0, T]; \chi^{-1}),$$

and that

$$\begin{aligned}
 (3.10) \quad & \|Q(w, u_1) + Q(u_2, w)\|_{L^1([0, t_0]; \chi^{-1})} \\
 & \leq C \|w\|_{L^\infty([0, t_0]; \chi^{-1})}^{\frac{1}{2}} (\|u_1\|_{L^\infty([0, t_0]; \chi^{-1})} + \|u_2\|_{L^\infty([0, t_0]; \chi^{-1})})^{\frac{1}{2}} \\
 & \quad \times \|w\|_{L^1([0, t_0]; \chi^1)}^{\frac{1}{2}} (\|u_2\|_{L^1([0, t_0]; \chi^1)} + \|u_2\|_{L^1([0, t_0]; \chi^1)})^{\frac{1}{2}}.
 \end{aligned}$$

By virtue of Lemma 3.1, we infer that for any $0 \leq t_0 \leq T$,

$$\begin{aligned}
 (3.11) \quad & \|w\|_{L^\infty([0, t_0]; \chi^{-1})} + \nu \|w\|_{L^1([0, t_0]; \chi^1)} \\
 & \leq 2 \|Q(w, u_1) + Q(u_2, w)\|_{L^1([0, t_0]; \chi^{-1})} \\
 & \leq C \|w\|_{L^\infty([0, t_0]; \chi^{-1})}^{\frac{1}{2}} (\|u_1\|_{L^\infty([0, t_0]; \chi^{-1})} + \|u_2\|_{L^\infty([0, t_0]; \chi^{-1})})^{\frac{1}{2}} \\
 & \quad \times \|w\|_{L^1([0, t_0]; \chi^1)}^{\frac{1}{2}} (\|u_2\|_{L^1([0, t_0]; \chi^1)} + \|u_2\|_{L^1([0, t_0]; \chi^1)})^{\frac{1}{2}} \\
 & \leq \varepsilon (\|u_1\|_{L^\infty([0, t_0]; \chi^{-1})} + \|u_2\|_{L^\infty([0, t_0]; \chi^{-1})}) \|w\|_{L^\infty([0, t_0]; \chi^{-1})} \\
 & \quad + C_\varepsilon (\|u_1\|_{L^1([0, t_0]; \chi^1)} + \|u_2\|_{L^1([0, t_0]; \chi^1)}) \|w\|_{L^1([0, t_0]; \chi^1)}.
 \end{aligned}$$

Choosing $\varepsilon > 0$ such that

$$\varepsilon (\|u_1\|_{L^\infty([0, T]; \chi^{-1})} + \|u_2\|_{L^\infty([0, T]; \chi^{-1})}) \leq \frac{1}{2},$$

then there exists a positive number δ satisfying $0 < \delta \leq T$ and

$$C_\varepsilon (\|u_1\|_{L^1([0, \delta]; \chi^1)} + \|u_2\|_{L^1([0, \delta]; \chi^1)}) \leq \frac{\nu}{2}.$$

We then infer that

$$\begin{aligned}
 & \|w\|_{L^\infty([0, \delta]; \chi^{-1})} + \nu \|w\|_{L^1([0, \delta]; \chi^1)} \\
 & \leq \varepsilon (\|u_1\|_{L^\infty([0, \delta]; \chi^{-1})} + \|u_2\|_{L^\infty([0, \delta]; \chi^{-1})}) \|w\|_{L^\infty([0, \delta]; \chi^{-1})} \\
 & \quad + C_\varepsilon (\|u_1\|_{L^1([0, \delta]; \chi^1)} + \|u_2\|_{L^1([0, \delta]; \chi^1)}) \|w\|_{L^1([0, \delta]; \chi^1)} \\
 & \leq \frac{1}{2} (\|w\|_{L^\infty([0, \delta]; \chi^{-1})} + \nu \|w\|_{L^1([0, \delta]; \chi^1)}).
 \end{aligned}$$

This implies that $w(t) = 0$, $0 \leq t \leq \delta$. Basic connective argument then yields uniqueness on $[0, T]$.

Theorem 1.2 is thus proved up to the blow-up criterion. Assume that we have a solution of the system (GNS_ν) on a time interval $[0, T]$ ($T < \infty$) such that

$$(3.12) \quad \int_0^T \|u\|_{\chi^0}^2 dt < \infty.$$

We claim that the lifespan T_{u_0} of u is greater than T . Indeed, thanks to Lemma 3.1, the inequalities (3.4) and (3.12), we have

$$\begin{aligned} \int_{\mathbb{R}^3} \sup_{0 \leq t \leq T} |\widehat{v}(t, \xi)| |\xi|^{-1} d\xi &\leq \|u_0\|_{\chi^{-1}} + \|Q(u, u)\|_{L^1([0, T]; \chi^{-1})} \\ &\leq \|u_0\|_{\chi^{-1}} + C \|u\|_{L^2[0, T]; \chi^0}^2 \\ &< \infty. \end{aligned}$$

Thus, a positive number ρ exists such that

$$\forall t \in [0, T], \int_{|\xi| \geq \rho} |\xi|^{-1} |\widehat{u}(t, \xi)| d\xi \leq \frac{\nu}{2^{\frac{5}{2}} C_0}.$$

The condition (3.3) now implies that for any $t \in [0, T]$, the lifespan for a solution of (GNS_ν) with initial data $u(t)$ is bounded from below by a positive real number C which is independent of t . Thus $T_{u_0} > T$, and the whole of Theorem 1.2 is now proved.

As the system (NS_ν) is a particular case of the system (GNS_ν) , we only need to show that the Navier-Stokes system is well-posed globally in time for $\|u_0\|_{\chi^{-1}} < \nu$. This result has been obtained in [2] by mollifying initial data. Here, however, due to Theorem 1.2, for any $u_0 \in \chi^{-1}$ there already exists a unique local solution $u(x, t)$ on some interval $[0, T^*)$, thus we can directly obtain estimates of the solution u instead of approximate solutions [2].

Taking the Fourier transform of the system (NS_ν) , one has

$$\begin{aligned} (3.13) \quad \partial_t \widehat{u}(t, \xi) + \nu |\xi|^2 \widehat{u}(t, \xi) \\ = i \int_{\mathbb{R}^3} \widehat{u}(\eta) \otimes \widehat{u}(\xi - \eta) d\eta \cdot \xi - i \left(\frac{1}{|\xi|^2} \xi^T \cdot \int_{\mathbb{R}^3} \widehat{u}(\eta) \otimes \widehat{u}(\xi - \eta) d\eta \cdot \xi \right) \xi, \end{aligned}$$

The condition $\operatorname{div} u = 0$ implies that $\xi \cdot \widehat{u} = 0$.

Letting $(3.13) \cdot \overline{\widehat{u}} + \widehat{u} \cdot \overline{(3.13)}$, we get

$$\begin{aligned} \partial_t |\widehat{u}|^2(t, \xi) + 2\nu |\xi|^2 |\widehat{u}|^2(t, \xi) \\ = i \left[\xi^T \cdot \int_{\mathbb{R}^3} \widehat{u}(\eta) \otimes \widehat{u}(\xi - \eta) d\eta \cdot \widehat{\bar{u}} - \widehat{u}^T \cdot \int_{\mathbb{R}^3} \widehat{\bar{u}}(\eta) \otimes \widehat{\bar{u}}(\xi - \eta) d\eta \cdot \xi \right]. \end{aligned}$$

For any positive ε , we have

$$\partial_t(|\widehat{u}|^2 + \varepsilon)^{\frac{1}{2}} = \frac{\partial_t |\widehat{u}|^2}{2(|\widehat{u}|^2 + \varepsilon)^{\frac{1}{2}}}.$$

Thus, integrating with respect to t gives

$$\begin{aligned} & (|\widehat{u}|^2(t, \xi) + \varepsilon)^{\frac{1}{2}} + \nu \int_0^t \frac{|\xi|^2 |\widehat{u}|^2(s, \xi)}{(|\widehat{u}|^2(s, \xi) + \varepsilon)^{\frac{1}{2}}} ds \\ & \leq (|\widehat{u}_0|^2(\xi) + \varepsilon)^{\frac{1}{2}} + \int_0^t \int_{\mathbb{R}^3} \frac{|\widehat{u}(s, \xi)| |\xi| |\widehat{u}(s, \eta)| |\widehat{v}(s, \xi - \eta)|}{(|\widehat{u}|^2(s, \xi) + \varepsilon)^{\frac{1}{2}}} d\eta ds. \end{aligned}$$

Letting ε tend to zero gives

$$|\widehat{u}|(t, \xi) + \nu \int_0^t |\xi|^2 |\widehat{u}|(s, \xi) ds \leq |\widehat{u}_0|(\xi) + \int_0^t \int_{\mathbb{R}^3} |\xi| |\widehat{u}|(s, \eta) |\widehat{u}|(s, \xi - \eta) d\eta ds.$$

Taking the L^1 norm with respect to $|\xi|^{-1} d\xi$, we have

$$\begin{aligned} & \int_{\mathbb{R}^3} |\widehat{u}|(t, \xi) |\xi|^{-1} d\xi + \nu \int_0^t \int_{\mathbb{R}^3} |\widehat{u}|(s, \xi) |\xi| d\xi ds \\ & \leq \int_{\mathbb{R}^3} |\widehat{u}_0|(\xi) |\xi|^{-1} d\xi + \|u\|_{L^2([0, t]; \chi^0)}^2 \\ & \leq \|u_0\|_{\chi^{-1}} + \|u\|_{L^\infty([0, t]; \chi^{-1})} \|u\|_{L^1([0, t]; \chi^1)}. \end{aligned}$$

If $\|u_0\|_{\chi^{-1}} < \nu$, then $\|u(t)\|_{\chi^{-1}} < \nu$ at least for a very short time interval $[0, \delta]$.

Consequently, on such a time interval, we have

$$\|u(t)\|_{\chi^{-1}} \leq \|u(0)\|_{\chi^{-1}} < \nu.$$

The basic continuity argument yields that

$$\|u(t)\|_{\chi^{-1}} \leq \|u(0)\|_{\chi^{-1}} < \nu,$$

for all $t \in [0, T^*)$ (see the inequalities (2.5)-(2.6) in [2]).

We then derive that

$$\|u(t)\|_{\chi^{-1}} + (\nu - \|u(0)\|_{\chi^{-1}}) \int_0^t \|u(s)\|_{\chi^1} ds \leq \|u_0\|_{\chi^{-1}},$$

for all $t \in [0, T^*)$.

Thanks to Proposition 2.1, one has

$$\int_0^{T^*} \|u(t)\|_{\chi^0}^2 dt \leq \frac{\|u_0\|_{\chi^{-1}}^2}{\nu - \|u(0)\|_{\chi^{-1}}} < \infty.$$

According to Theorem 1.2 again, this implies that $T^* = \infty$, and the whole Theorem 1.3 is proved.

Acknowledgements. This work was partially supported by NNSFC (No. 11271382), RFDP (No. 20120171110014), and the key project of Sun Yat-sen University (No. c1185).

References

- [1] T. Kato and H. Fujita, *On the Navier-Stokes initial value problem I*, Arch. Ration. Mech. Anal., **16** (1964), 269-351.
- [2] Z. Lei and F. Lin, *Global mild solutions of Navier-Stokes equations*, Comm. Pure Appl. Math., **64**:9 (2011), 1297C1304
- [3] H. Bahouri, J.-Y. Chemin and R. Danchin, *Fourier Analysis and Nonlinear Partial Differential Equations*, Grundlehren der Mathematischen Wissenschaften, Vol. 343. New York: Springer-Verlag (2011).
- [4] J.-Y. Chemin and I. Gallagher, *Wellposedness and stability results for the Navier-Stokes equations in \mathbb{R}^3* , Ann. I. H. Poincaré-AN, **26**:2 (2009), 599C624.
- [5] T. Kato, *Nonstationary flows of viscous and ideal fluids in \mathbb{R}^3* , J. Funct. Anal., **9** (1972), 296-305.
- [6] H. Koch and D. Tataru, *Well-posedness for the Navier-Stokes equations*, Adv. Math., **157**:1 (2001), 22C35.
- [7] T. Kato, *Strong L^p solutions of the Navier-Stokes equations in \mathbb{R}^m , with applications to weak solutions*, Math. Z., **187**:4 (1984), 471C480.
- [8] J.-Y. Chemin, *Théorèmes d'unicité pour le système de Navier-Stokes tridimensionnel*, J. Anal. Math., **77** (1999), 27C50.
- [9] M. Cannone, Y. Meyer and F. Planchon, *Solutions autosimilaires des équations de Navier-Stokes*, Séminaire Équations aux Dérivées Partielles de l'École Polytechnique, 1993C1994.

- [10] I. Gallagher, D. Iftimie and F. Planchon, *Asymptotics and stability for global solutions to the Navier-Stokes equations*, Ann. Inst. Fourier., **53** (2003), 1387C1424.
- [11] M. Cannone and G. Wu, *Global well-posedness for Navier-Stokes equations in critical Fourier-Herz spaces*, Nonlinear Anal., **75** (2012), 3754 C3760.