

Is elastic scattering at the LHC absorptive or geometric?

S.M. Troshin, N.E. Tyurin

*Institute for High Energy Physics,
Protvino, Moscow Region, 142281, Russia*

Abstract

We discuss the role of the impact-parameter dependent quantities. Their use along with the available experimental data on elastic scattering at the LHC could serve for an additional insight into the asymptotics.

There is a general opinion that the upper (Froissart-Martin) bound for the total cross-sections should be saturated at asymptotics. The hopes to understand asymptotic behavior of strongly interacting particles always arise when a new particle machine starts to operate. But, after not so long time, it becomes evident again that asymptotic limit is still elusive. Cosmic rays experiments probe much higher energies, but those have much lower statistics. However, their use even allowed to produce a claim that the solution of the asymptotic puzzle has been found [1]. There are two reasons preventing acceptance of this conclusion as an only relevant reflection of the physical reality, i.e. those results are, in fact, model-dependent ones and are limited to the use of the experimental data for the scattering in the forward region only.

Despite that the functional energy dependence of the on-shell hadron total cross-section is considered in most cases to follow $\ln^2 s$ -dependence, the value of the numerical factor in front of $\ln^2 s$ remains obscure. This issue is closely related to the selection of the upper limit for the partial amplitude, namely, should this limit correspond to the maximum of the inelastic channel contribution to the elastic unitarity with asymptotics

$$\sigma_{el}(s)/\sigma_{tot}(s) \rightarrow 1/2 \quad (1)$$

or it corresponds to a maximal value of the partial amplitudes allowed by unitarity and results in the asymptotics

$$\sigma_{el}(s)/\sigma_{tot}(s) \rightarrow 1. \quad (2)$$

Under assumption of the limit (1) the original Froissart-Martin bound for the total cross-sections has been improved [2]. The bound for the total inelastic cross-section reduced by factor of 4 has also been derived [2]. The ratio of the elastic to total cross-section (2) corresponds to energy increase of the total inelastic cross-section slower than $\ln^2 s$. It should be noted that this ratio is, in fact, a factor standing in front of $\ln^2 s$ in the bound for the total cross-section[3]. Various asymptotic limits have been considered in [4] in almost model-independent way, but also for the forward scattering data only.

We argue that the inclusion into the analysed data set of the elastic differential cross-section can provide new information on the asymptotics in hadron scattering. The main tool here is reconstruction of the impact-parameter dependent quantities from this experimental dataset.

It has been known for a long time (cf. [5]) that position operator (impact parameter) deprives of its unpleasant features related to its non-commutativity with the Hamiltonian with the energy increase. At the moment the highest

available particle collision energy is provided at the LHC. It seems that at such high energies the impact parameter representation for the spinless particles can be used. One of the attractive features of this representation is diagonalization of the unitarity equation for the elastic scattering amplitude $f(s, b)$, i.e. at high energies

$$\text{Im}f(s, b) = h_{el}(s, b) + h_{inel}(s, b)$$

with accuracy up to $\mathcal{O}(1/s)$ [6]. The function $h_{el}(s, b) \equiv |f(s, b)|^2$ is the contribution of the elastic channel, while $h_{inel}(s, b)$ takes into account contribution of all the inelastic intermediate channels, b is an impact parameter of the colliding hadrons. This equation is instrumental for the reconstruction of $h_{inel}(s, b)$ ¹ from the elastic scattering data².

It should be noted that unitarity implies an existence of the two scattering modes - absorptive and geometric ones. Namely, the elastic scattering S -matrix (related to the elastic scattering amplitude as $S(s, b) = 1 + 2if(s, b)$) can be presented in the form

$$S(s, b) = \kappa(s, b) \exp[2i\delta(s, b)]$$

with the two real functions $\kappa(s, b)$ and $\delta(s, b)$. The function κ ($0 \leq \kappa \leq 1$) is an absorption factor³, its value $\kappa = 0$ corresponds to a complete absorption. At the LHC energies the real part of the scattering amplitude would be neglected, i.e. we can perform replacement $f \rightarrow if$. This assumption is widely used. Selection of the elastic scattering mode, namely, absorptive or geometric is governed then by the phase $\delta(s, b)$. The standard assumption is $S(s, b) \rightarrow 0$ when the impact parameter b is fixed and $s \rightarrow \infty$. It corresponds to a black disk limit and the elastic scattering is then completely absorptive, i.e. it is just a shadow of all the inelastic processes.

There is another possibility, namely, the function $S(s, b) \rightarrow -1$ when b is fixed and $s \rightarrow \infty$, i.e. $\kappa \rightarrow 1$ and $\delta \rightarrow \pi/2$. This case corresponds to a pure geometric scattering [9]. The principal point is that the phase is non-zero, i.e. δ is equal to $\pi/2$.

We discuss now the observable effects sensitive to the presence of the non-zero phase. One of the useful quantities for that purposes is the ratio of elastic to total impact-parameter dependent cross-sections, i.e.

$$\mathcal{R}(s, b) = \sigma_{el}(s, b)/\sigma_{tot}(s, b).$$

¹The function $h_{inel}(s, b)$ is not well suited for the studies of asymptotics due to absence of the one-to-one correspondence between the functions $f(s, b)$ and $h_{inel}(s, b)$.

²Cf. e.g. [7] for an earlier analysis of $h_{inel}(s, b)$ and [8] for the most recent one.

³Its meaning is different in the reflection region, as it will be discussed further.

The impact-parameter dependent quantities $\sigma_{el}(s, b)$, $\sigma_{inel}(s, b)$ and $\sigma_{tot}(s, b)$ can be extracted from the experimental data on the $d\sigma/dt$. The function $\mathcal{R}(s, b)$ at the energies $s > s_0$ (the value of s_0 is determined by the solution of the equation $f(s, b = 0) = 1/2$) has two different forms depending on the value of impact parameter, namely at $b > R(s)$ this function has the form

$$\mathcal{R}(s, b) = [1 - \kappa(s, b)]/2 < 1/2,$$

while in the region $b < R(s)$ the presence of the non-zero phase (i.e. $\cos 2\delta = -1$) changes this dependence to

$$\mathcal{R}(s, b) = [1 + \kappa(s, b)]/2 > 1/2.$$

It means that the hadron scattering becomes predominantly geometric one at small impact parameters and high energies $s > s_0$. Thus, the border of the region where reflective scattering is presented, is determined by the function $R(s)$. This is solution of the equation

$$f(s, b = R(s)) = 1/2$$

at $s > s_0$. The scattering in this case has a dominant absorptive contribution in the peripheral region of the impact parameters only. The extraction from the experimental data of the function $\mathcal{R}(s, b)$ at the finite energies can provide a hint on the asymptotics of the soft strong interactions, namely if the inequality

$$\mathcal{R}(s, b) > 1/2$$

does result from the experimental data analysis at small impact parameters and high values of the collision energy, one can arrive to conclusion that the black disk limit will be violated at the asymptotic energies and in this region the scattering would have a geometric origin since reflective scattering mode dominates [9]. This would testify in favor of asymptotic limit (2). Of course, this claim is based on the assumed monotonic energy dependence of the function \mathcal{R} at high energies.

The following asymptotic limit takes place, namely, $\mathcal{R}(s, b) \rightarrow 1$ when impact parameter b is fixed and $s \rightarrow \infty$ (i.e. geometric elastic scattering saturates the total cross-section in this limit), while $\mathcal{R}(s, b) \rightarrow 0$ at fixed energy s and $b \rightarrow \infty$, i.e. at large impact parameters elastic scattering cross-section decreases faster than the cross-sections of the all inelastic processes, i.e.

$$\mathcal{R}(s, b) = \bar{\mathcal{R}}(s, b)/[1 + \bar{\mathcal{R}}(s, b)],$$

where

$$\bar{\mathcal{R}}(s, b) \equiv \sigma_{el}(s, b)/\sigma_{inel}(s, b) \rightarrow 0$$

the following relation takes place

$$\bar{\mathcal{R}}(s, b) = f(s, b)/[1 - f(s, b)]$$

at fixed s and $b \rightarrow \infty$ and $f(s, b) \rightarrow 0$ in this limit since .

Here the new ratio $\bar{\mathcal{R}}(s, b)$ has been used. It is clear that

$$\bar{\mathcal{R}}(s, b) \rightarrow \infty$$

at fixed b and $s \rightarrow \infty$ under transition to the non-zero phase $\cos 2\delta = -1$. The function $\bar{\mathcal{R}}(s, b)$ is another quantity sensitive to the presence of the reflective scattering mode and it determines the generalized reaction matrix dependence. Namely, in the U -matrix approach for the unitarization (rational form of unitarization) the elastic scattering matrix element in the impact parameter representation is a linear fractional transform:

$$S(s, b) = \frac{1 - U(s, b)}{1 + U(s, b)}. \quad (3)$$

$U(s, b)$ is the generalized reaction matrix, which is assumed to be input dynamical quantity. This relation (3) is a one-to-one transform and can be easily inverted. The ratio $\bar{\mathcal{R}}(s, b)$ determines the function $U(s, b)$, i.e.

$$U(s, b) = \bar{\mathcal{R}}(s, b)$$

and this relation can be used as a tool for the reconstruction of the function $U(s, b)$ from the experimental data.

In the models the function $U(s, b)$ passes through unity with increasing energy providing that way a gradual transition to the geometric scattering mode. This transition implies crossing zero value by the function $S(s, b)$ and acquiring the phase $\delta = \pi/2$ when the function $U(s, b)$ passes through unity. The solution of the equation $U(s, b) = 1$ separates the regions of absorptive and geometric scattering and corresponds to the maximum value of $h_{inel}(s, b) = 1/4$ since the derivative of $h_{inel}(s, b)$ has the form

$$\frac{\partial h_{inel}(s, b)}{\partial b} = S(s, b) \frac{\partial f(s, b)}{\partial b}$$

and equals to zero at $U(s, b) = 1$. The derivative of the inelastic overlap function has the sign opposite to the sign of $\partial f(s, b)/\partial b$ in the region where $U(s, b) > 1$ and the non-zero phase is, therefore, responsible for the transformation of the central impact-parameter profile of the function $f(s, b)$ into a peripheral one of the inelastic overlap function $h_{inel}(s, b)$. The appearance of

the non-zero phase results (cf. Eq. 3) from transition into the region where $U(s, b) > 1$. Since the function $U(s, b)$ (its imaginary part in fact, but here we consider a pure imaginary case) gets contributions from the inelastic intermediate channels only, its increasing behavior with energy can be associated with increasing contribution of the successive opening new inelastic channels. This can be considered as a dynamical origin for the relation $\cos 2\delta = -1$.

It should be noted that the derivative of the elastic overlap function has no sign-changing factor in front of $\partial f(s, b)/\partial b$, namely

$$\frac{\partial h_{el}(s, b)}{\partial b} = [1 - S(s, b)] \frac{\partial f(s, b)}{\partial b}$$

with $1 - S(s, b)$ being a non-negative at all values of s and b .

Thus, the role of the non-zero phase in the high energy scattering is essential. In the presence of the non-zero phase at the LHC energies the geometric scattering dominates at small impact parameters while inelastic processes are peripheral. The albedo (coefficient of reflection) increases with energy at $s > s_0$ [9]. The factor $\kappa(s, b)$ plays the role of albedo at $s > s_0$ and $b < R(s)$ and hence should be considered as a reflective factor rather than absorption factor when this region is taken into consideration.

This effect results in a number of consequences. The peripherality of the inelastic amplitudes could be a dominating mechanism of the ridge and double-ridge effects observed in the two-particle correlation functions in proton-proton collisions [10, 11]. The appearance of the non-zero phase at $s > s_0$ leading to the dominance of the geometric scattering, provides change in the slope ("the knee") of the energy spectrum of the cosmic particles at the ground level [9].

At this point comments on the present experimental situation with elastic scattering at the LHC are to be done. Most recent review of the data for elastic scattering with the analysis of the theoretical models has been given in [12]. As it was noted, the elastic scattering from the experimental viewpoint is consistent with BEL picture when the protons becomes blacker, edgier and larger [13]. This is indicated by the analysis of the TOTEM data [14, 15]. However, at the moment one cannot exclude the possibility of a gradual transition to the picture which name can be abbreviated as a REL one, i.e. when the scattering region of the protons starts to become reflective at the center and simultaneously edgier and larger at its peripherality. The TOTEM data do not provide a straightforward clue for the asymptotic behavior, however the most recent luminosity-independent measurements at $\sqrt{s} = 8$ TeV [16] have confirmed steady increase of the ratio σ_{el}/σ_{tot} providing the value 0.266 ± 0.006 for this quantity.

The processes at much higher collision energies can be observed in the cosmic rays. As it was noted in the beginning of the present paper, the cosmic rays data have significant error bars and therefore are not extremely conclusive. The highest-energy cosmic data for the total inelastic cross-sections are provided now at the Pierre Auger Observatory [17]. Estimations on the base of these data [12] might be considered in favor of the REL scattering picture since a decrease of the ratio $\sigma_{inel}/\sigma_{tot}$ from 0.75 at $\sqrt{s} = 7$ TeV to 0.67 at $\sqrt{s} = 57 \pm 6$ TeV could take place. This result, however, should be taken with caution and several reservations related to the low statistics of the cosmic data points and model dependence of the extrapolations.

Indirect information on the possible asymptotics can also be extracted from the deep-elastic scattering which is sensitive to the region of small impact parameters and from the studies of the correlations of elastic scattering process with the particle production processes [18].

To get a more straightforward further information on which scattering mode takes place at asymptotic energies, namely, absorptive or geometric, one can use a full reconstruction of the impact-parameter dependent scattering amplitude from the experimental data set. This set should include differential cross-section of the elastic scattering. Despite that such an analysis assumes using the models for the phase of the scattering amplitude it has a full potential to extract the impact-parameter dependent quantities. We would like to point out that the recent analysis [19] implies that the amplitude $f(s, b)$ can reach and even already crossed the black-disk limit at $b = 0$ and $\sqrt{s} = 7$ TeV⁴. If it is so, it would testify in favor of a gradual transition to the geometric elastic scattering starting at the LHC energies.

The model-independent analysis of the LHC data aiming to the reconstruction of the impact parameter amplitude $f(s, b)$ is desirable and would be instrumental for the clarification of the hadron asymptotics.

Acknowledgement

We are grateful to E. Martynov for letting us know the results of the impact parameter analysis of the elastic scattering.

References

- [1] M.M. Block, F. Halzen, Phys. Rev. Lett. **107**, 212002 (2011).

⁴It should be noted here that the value of $\text{Im}f(s, b = 0)$ has increased from 0.36 (CERN ISR) to 0.492 ± 0.008 (Tevatron) and is very close to the black disk limit in this energy domain[20].

- [2] A. Martin, Phys. Rev. D **80**, 065013 (2009).
- [3] V. Singh, S.M. Roy, Ann. of Phys. **57**, 461 (1970).
- [4] D.A. Fagundes, M.J. Menon, Nucl. Phys. A **880**, 1 (2012).
- [5] B.R. Webber, Nucl. Phys. B **87**, 269 (1975).
- [6] M.L. Goldberger, K.M. Watson, *Collision Theory*, John Wiley and Sons, Inc., New-York—London—Sydney 1964.
- [7] U. Amaldi, K.S. Schubert, Nucl. Phys. B **166**, 301 (1980).
- [8] I.M. Dremin, V.A. Nechitailo, arXiv:1306.5384 [hep-ph].
- [9] S.M. Troshin, N.E. Tyurin, Int. J. Mod. Phys. A **22**, 4437 (2007).
- [10] S.M. Troshin, N.E. Tyurin, Int. J. Mod. Phys. A **26**, 4703 (2011).
- [11] S.M. Troshin, N.E. Tyurin, Mod. Phys. Lett. A **28**, 1350031 (2013).
- [12] I.M. Dremin, Physics-Uspekhi, **56**, 3 (2013).
- [13] R. Henzi, P. Valin, Phys. Lett. B, **132**, 443, (1983).
- [14] G. Antchev et al. (TOTEM Collaboration), Europhys. Lett. **95**, 41001 (2011).
- [15] G. Antchev et al. (TOTEM Collaboration), Europhys. Lett. **96**, 21002 (2011).
- [16] G. Antchev et al. (TOTEM Collaboration), Phys. Rev. Lett. **111**, 012001 (2013).
- [17] P. Aubreu et al. (Pierre Auger Collaboration), Phys. Rev. Lett. **109**, 062002 (2012)
- [18] S.M. Troshin, N.E. Tyurin, Phys. Lett. B, **707**, 558, (2012).
- [19] A. Alkin, O. Kovalenko, E. Martynov, Europhys. Lett. **102**, 31001 (2013).
- [20] P. Giromini, Proc. of the Vth BLOIS Workshop, Elastic and Diffractive Scattering, Eds. H.M. Fried, K. Kang, C-I Tan, World Scientific, Singapore, 1994, p. 30.