

PREPARED FOR SUBMISSION TO JHEP

Masses, mixing angles and phases of general Majorana neutrino mass matrix

Biswajit Adhikary^a, Mainak Chakraborty^b, Ambar Ghosal^b

a)Department of Physics, Gurudas College, Narkeldanga, Kolkata-700054, India

b) Saha Institute of Nuclear Physics, 1/AF Bidhannagar, Kolkata 700064, India

E-mail: biswajitadhikary@gmail.com, mainak.chakraborty@saha.ac.in,
ambar.ghosal@saha.ac.in

KEYWORDS: Neutrino Physics, Beyond Standard Model

Contents

1	Introduction	2
2	General Solution	3
2.1	Mass Eigenvalues	3
2.2	Mixing Angles and Dirac CP phase	5
3	Majorana Phases	6
4	Cyclic Symmetry	9
4.1	Basic Formalism	9
4.2	Parametrization and Diagonalization	10
5	Breaking of cyclic symmetry	11
5.1	Numerical results and phenomenological discussions	12
6	Summary	15
A	Appendix	16
A.1	Elements of h in terms of elements of m_ν (a_i, b_i)	16
A.2	Coefficients of the cubic equation in terms of elements of h	16

General Majorana neutrino mass matrix is complex symmetric and for three generations of neutrinos it contains 12 real parameters. We diagonalize this general neutrino mass matrix and express the three neutrino masses, three mixing angles, one Dirac CP phase and two Majorana phases (removing three unphysical phases) in terms of the neutrino mass matrix elements. We apply the results in the context of a neutrino mass matrix derived from a broken cyclic symmetry invoking type-I seesaw mechanism. Phenomenological study of the above mass matrix allows enough parameter space to satisfy the neutrino oscillation data with only 10% breaking of this symmetry. In this model only normal mass hierarchy is allowed. In addition, the Dirac CP phase and the Majorana phases are numerically estimated. Σm_i and $|m_{\nu_{ee}}|$ are also calculated.

1 Introduction

It is very useful to have a straightforward framework to find the masses and mixing angles of a generalized neutrino mass matrix. In this work special emphasis is given on the diagonalization procedure of the most general 3×3 complex symmetric effective neutrino mass matrix (m_ν). Starting from a most general m_ν we calculate three masses directly (without any approximation) in terms of the elements of m_ν . Knowing the mass eigenvalues, three mixing angles and the Dirac CP phase are also obtained. Apart from the Dirac CP phase the total diagonalization matrix consists of three unphysical phases and two Majorana phases. Eliminating the unphysical phases, extraction of the Majorana phases (for generalized m_ν) are also done. We would like to emphasize that those expressions are readily applicable in case of any symmetric or broken symmetric mass matrix. More importantly, the diagonalization is exact and the corresponding neutrino observables are calculated in an exact form without assuming any approximate procedure regarding diagonalization. To illustrate, we employ the obtained expressions in the context of a neutrino mass matrix derived from a broken symmetry.

In the field of neutrino physics, it is now a challenging task to build a suitable model which can accommodate neutrino oscillation experimental data comprising solar[1, 2], atmospheric[3] and recent reactor neutrino [4–7] experiments as well as the constraint on the sum of the three neutrino masses arising from cosmological data[8, 9]. Furthermore, for Majorana type neutrino, an additional constraint on the $|m_{\nu ee}|$ element of the neutrino mass matrix [10–12] is also necessary to take into account. Popular paradigm is to invoke some symmetries or ansatz[13–15], *viz.* A_4 [16], $\mu\tau$ symmetry[17]–[44], scaling ansatz[45]–[54], to generate nondegenerate mass eigenvalues[55] and $\theta_{23}, \theta_{12} \neq 0$ with $\theta_{13} = 0$ at the leading order and nonzero θ_{13} [56]–[73] is generated by further breaking of such symmetries or ansatz. Contrary to the above idea, in the present work, we explore a typical symmetry, cyclic permutation symmetry[74–76], in which it is possible to generate all three mixing angles nonzero at the leading order, however, the mass eigenvalues become degenerate. To circumvent this loop hole, we break the symmetry in such a way that the degeneracy in mass eigenvalues is lifted but the mixing angles are still compatible with the extant data.

We consider standard $SU(2)_L \times U(1)_Y$ model with three right handed neutrinos N_{eR} , $N_{\mu R}$, $N_{\tau R}$ and invoke type-I seesaw mechanism to generate light neutrino masses. We further impose a cyclic permutation symmetry on both left and right chiral neutrino fields as

$$\begin{aligned} \nu_{eL} &\rightarrow \nu_{\mu L} \rightarrow \nu_{\tau L} \rightarrow \nu_{eL}, \\ N_{eR} &\rightarrow N_{\mu R} \rightarrow N_{\tau R} \rightarrow N_{eR}. \end{aligned} \tag{1.1}$$

Cyclic permutation symmetry is a subgroup of S_3 permutation symmetry[77] with three of its elements as $\{P_0, P_{123}, P_{132}\}$ ¹. One of the motivation to study the S_3 symmetry is to realize the well known Tribimaximal (TBM) mixing pattern.

¹Permutation of three objects $\{a, b, c\}$ form S_3 group. There are six elements: $P_0, P_{12}, P_{13}, P_{23}, P_{123}, P_{132}$. Their operations are as follows: $P_0(a, b, c) \rightarrow (a, b, c)$, $P_{12}(a, b, c) \rightarrow (b, a, c)$, $P_{13}(a, b, c) \rightarrow (c, b, a)$, $P_{23}(a, b, c) \rightarrow (a, c, b)$, $P_{123}(a, b, c) \rightarrow (c, a, b)$, $P_{132}(a, b, c) \rightarrow (b, c, a)$.

The paper is organized as follows: In section 2, we present the most general solution of a complex 3×3 symmetric mass matrix to obtain three masses, three mixing angles and the Dirac CP phase. Expressions for the Majorana phases are given in section 3. Section 4 deals with a convenient parametrization and diagonalization of the proposed cyclic symmetry invariant Majorana neutrino mass matrix. Expression of m_ν in parametric form due to broken cyclic symmetry and corresponding numerical results and phenomenological discussions on the allowed parameter ranges are presented in Section 5. Section 6 contains a summary of the present work.

2 General Solution

In this section we calculate the exact algebraic expressions for the masses and mixing angles of the most general complex symmetric neutrino mass matrix (m_ν) which is written in terms of real (a_i) and imaginary (b_i) parts as

$$m_\nu = \begin{pmatrix} a_1 + ib_1 & a_2 + ib_2 & a_3 + ib_3 \\ a_2 + ib_2 & a_4 + ib_4 & a_5 + ib_5 \\ a_3 + ib_3 & a_5 + ib_5 & a_6 + ib_6 \end{pmatrix}. \quad (2.1)$$

2.1 Mass Eigenvalues

It is well known that any complex symmetric mass matrix can be diagonalized by a unitary transformation as

$$U^\dagger m_\nu U^* = \text{diag}(m_1, m_2, m_3) \quad (2.2)$$

where U is a unitary matrix and m_i 's ($i = 1, 2, 3$) are real positive masses. However, the columns of U can not be the eigenvectors of m_ν because

$$m_\nu U^* = U \text{diag}(m_1, m_2, m_3) \quad (2.3)$$

is essentially in the form

$$m_\nu |m_i\rangle^* = m_i |m_i\rangle \quad (2.4)$$

by considering $|m_i\rangle$ as columns of U . Since, the states in the l.h.s and r.h.s of eq.(2.4) are different, it is not possible to utilize the equation of the type $\text{Det}(m_\nu - \lambda \mathbf{I}) = 0$ to obtain the masses m_i . It is therefore necessary to construct a hermitian matrix h as $h = m_\nu m_\nu^\dagger$. Explicit expressions of the elements of h matrix in terms of mass matrix parameters a_i and b_i are provided in Appendix A.1. The squared mass eigenvalues are obtained by direct diagonalization of h matrix as

$$U^\dagger h U = \text{diag}(m_1^2, m_2^2, m_3^2) \quad (2.5)$$

where the matrix U is constructed with the eigenvectors of h . It is now straightforward to write down the characteristic equation as $\text{Det}(h - \lambda I) = 0$ to find the eigenvalues. This gives a cubic equation

$$a\lambda^3 + b\lambda^2 + c\lambda + d = 0 \quad (2.6)$$

where the coefficients a, b, c, d are expressed in terms of the elements of h matrix and spelt out in Appendix A.2. The nature of the roots in eq.(2.6) depend on the sign of the discriminant Δ where

$$\Delta = 18abcd - 4b^3d + b^2c^2 - 4ac^3 - 27a^2d^2. \quad (2.7)$$

Depending on the sign of Δ two cases arise as

Case I: $\Delta \geq 0 \Rightarrow$ All roots are real. The roots are distinct for $\Delta > 0$ and degenerate roots occur for $\Delta = 0$.

Case II: $\Delta < 0 \Rightarrow$ One of the root is real and the other two are complex conjugate to each other.

Since hermitian matrix has real roots we stick to the condition $\Delta \geq 0$. The general expressions of the three roots of eq.(2.6) are given by

$$\begin{aligned} \lambda_1 = & -\frac{b}{3a} - \frac{1}{3a} \sqrt[3]{\frac{1}{2}(2b^3 - 9abc + 27a^2d + \sqrt{-27a^2\Delta})} \\ & - \frac{1}{3a} \sqrt[3]{\frac{1}{2}(2b^3 - 9abc + 27a^2d - \sqrt{-27a^2\Delta})} \end{aligned} \quad (2.8)$$

$$\begin{aligned} \lambda_2 = & -\frac{b}{3a} - \frac{1 + i\sqrt{3}}{6a} \sqrt[3]{\frac{1}{2}(2b^3 - 9abc + 27a^2d + \sqrt{-27a^2\Delta})} \\ & - \frac{1 - i\sqrt{3}}{6a} \sqrt[3]{\frac{1}{2}(2b^3 - 9abc + 27a^2d - \sqrt{-27a^2\Delta})} \end{aligned} \quad (2.9)$$

$$\begin{aligned} \lambda_3 = & -\frac{b}{3a} - \frac{1 - i\sqrt{3}}{6a} \sqrt[3]{\frac{1}{2}(2b^3 - 9abc + 27a^2d + \sqrt{-27a^2\Delta})} \\ & - \frac{1 + i\sqrt{3}}{6a} \sqrt[3]{\frac{1}{2}(2b^3 - 9abc + 27a^2d - \sqrt{-27a^2\Delta})}. \end{aligned} \quad (2.10)$$

Subject to the condition $\Delta \geq 0$ eq.(2.8) is simplified as

$$\lambda_1 = -\frac{b}{3a} - \frac{1}{3\sqrt[3]{2a}} (\sqrt[3]{x + iy} + \sqrt[3]{x - iy}) \quad (2.11)$$

where $x = 2b^3 - 9abc + 27a^2d$, $y = 3\sqrt{3}a\sqrt{\Delta}$.

Substituting $x = r \cos 3\theta$, $y = r \sin 3\theta$ in eq.(2.11) the complex part cancels out and λ_1 is simplified to

$$\lambda_1 = -\frac{b}{3a} - \frac{2\sqrt[3]{r}}{3\sqrt[3]{2a}} \cos \theta. \quad (2.12)$$

Following similar substitutions in eq.(2.9) and eq.(2.10) we get the simplified roots as

$$\lambda_2 = -\frac{b}{3a} + \frac{\sqrt[3]{r}}{3\sqrt[3]{2a}} (\cos \theta - \sqrt{3} \sin \theta) \quad (2.13)$$

$$\lambda_3 = -\frac{b}{3a} + \frac{\sqrt[3]{r}}{3\sqrt[3]{2a}} (\cos \theta + \sqrt{3} \sin \theta). \quad (2.14)$$

The mapping of $(\lambda_1, \lambda_2, \lambda_3)$ to (m_1^2, m_2^2, m_3^2) is done through utilization of the experimental data.

2.2 Mixing Angles and Dirac CP phase

In the above section we have calculated the mass eigenvalues by directly solving the characteristic equation. In other words, the matrix h is diagonalized through a rotation by a unitary matrix U , which is known as mixing matrix, as

$$U^\dagger h U = \text{diag}(m_1^2, m_2^2, m_3^2) = D \quad (2.15)$$

or,

$$h U = U D. \quad (2.16)$$

Eq.(2.16) is our key equation to get generalized expression of U_{ij} . Comparing l.h.s and r.h.s of eq.(2.16) we get 9 equations, and these 9 equations are clubbed in three equations in the following way

$$(h_{11} - m_i^2)U_{1i} + h_{12}U_{2i} + h_{13}U_{3i} = 0 \quad (2.17)$$

$$h_{12}^*U_{1i} + (h_{22} - m_i^2)U_{2i} + h_{23}U_{3i} = 0 \quad (2.18)$$

$$h_{13}^*U_{1i} + h_{23}^*U_{2i} + (h_{33} - m_i^2)U_{3i} = 0 \quad (2.19)$$

where $i = 1, 2, 3$. The unitary property of the U matrix further constrains the elements as

$$|U_{1i}|^2 + |U_{2i}|^2 + |U_{3i}|^2 = 1. \quad (2.20)$$

Thus utilizing eq.(2.17) to eq.(2.20) we get rowwise elements of U as

$$\begin{aligned} U_{1i} &= \frac{(h_{22} - m_i^2)h_{13} - h_{12}h_{23}}{N_i} \\ U_{2i} &= \frac{(h_{11} - m_i^2)h_{23} - h_{12}^*h_{13}}{N_i} \\ U_{3i} &= \frac{|h_{12}|^2 - (h_{11} - m_i^2)(h_{22} - m_i^2)}{N_i} \end{aligned} \quad (2.21)$$

where N_i is the normalization constant given by

$$\begin{aligned} |N_i|^2 &= |(h_{22} - m_i^2)h_{13} - h_{12}h_{23}|^2 + |(h_{11} - m_i^2)h_{23} - h_{12}^*h_{13}|^2 + \\ &\quad (|h_{12}|^2 - (h_{11} - m_i^2)(h_{22} - m_i^2))^2. \end{aligned} \quad (2.22)$$

The U matrix obtained here in general can have three phases and three mixing angles. This can be understood easily by looking at the h matrix. The h matrix has six moduli and three phases in three off diagonal elements. After diagonalization we have three real positive eigenvalues and a unitary matrix U in which remaining six parameters (three angles and three phases) are contained. Rotating the h matrix by a diagonal phase matrix P : $h' = P^\dagger h P$ we can absorb atmost two phases from two off diagonal elements and the survived phase in rest off diagonal elements will be same as the phase of $h_{12}h_{23}h_{31}$ ², term.

²With $P = \text{diag}(e^{i\alpha_1}, e^{i\alpha_2}, e^{i\alpha_3})$ we have $h'_{12} = e^{i(\alpha_2 - \alpha_1)}h_{12}$, $h'_{13} = e^{i(\alpha_3 - \alpha_1)}h_{13}$, $h'_{23} = e^{i(\alpha_3 - \alpha_2)}h_{23}$. h'_{12} , h'_{13} can be made real with the choice $\alpha_2 - \alpha_1 = -\arg h_{12}$, $\alpha_3 - \alpha_1 = -\arg h_{13}$ which in turn fixes $\alpha_3 - \alpha_2 = \arg h_{12} - \arg h_{13}$ and stops further absorption of phase. Hence survived phase in h'_{23} will be $\arg h_{12} + \arg h_{23} - \arg h_{13} \equiv \arg h_{12}h_{23}h_{31}$.

Phase of the quantity $h_{12}h_{23}h_{31}$ is independent of phase rotation i.e, phase of $h'_{12}h'_{23}h'_{31}$ is same as the phase of $h_{12}h_{23}h_{31}$. Now, unitary matrix with three angles and single phase in CKM type parametrization (following PDG[78] convention) is

$$U^{\text{CKM}} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \quad (2.23)$$

with $c_{ij} = \cos \theta_{ij}$, $s_{ij} = \sin \theta_{ij}$ and δ is the Dirac CP phase. Obtained solution of U_{ij} elements in (2.16) may contain unwanted phases which only can appear as the overall phase factor in elements of the U matrix. Hence, we can directly compare their modulus with the modulus of U_{ij}^{CKM} : $|U_{ij}^{\text{CKM}}| = |U_{ij}|$. This gives the expressions of three mixing angles as

$$\tan \theta_{23} = \frac{|U_{23}|}{|U_{33}|} \quad (2.24)$$

$$\tan \theta_{12} = \frac{|U_{12}|}{|U_{11}|} \quad (2.25)$$

$$\sin \theta_{13} = |U_{13}|. \quad (2.26)$$

To obtain the δ phase we utilize the phase rotation independent quantity $h_{12}h_{23}h_{31}$. Obviously, absence of phase factor in $h_{12}h_{23}h_{31}$ makes the h matrix real symmetric under phase rotation. Therefore, $\text{Im}(h_{12}h_{23}h_{31})$ must be proportional to $\sin \delta$:

$$\text{Im}(h_{12}h_{23}h_{31}) = \frac{(m_2^2 - m_1^2)(m_3^2 - m_2^2)(m_3^2 - m_1^2) \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \sin \delta}{8} \quad (2.27)$$

which can be easily inverted to obtain the phase δ . Thus, from h we are able to find out three masses, three mixing angles and the Dirac CP phase in terms of the elements of neutrino mass matrix. Our next goal is to find out remaining two Majorana phases which we will explore in the next section.

3 Majorana Phases

In this section we explicitly calculate the Majorana phases assuming the three neutrino masses, three mixing angles and the Dirac CP phase are calculable in terms of the elements of neutrino mass matrix. For a complex symmetric m_ν matrix there are twelve independent parameters arising from six complex elements. These twelve parameters are counted as (i) three masses, (ii) three mixing angles, (iii) one Dirac CP phase, (iv) two Majorana phases and (v) three unphysical phases. These three unphysical phases take crucial part in diagonalization. Now, the unitary matrix with three angles and six phases can be parametrized as:

$$U_{\text{tot}} = P_\phi U^{\text{PMNS}} \quad (3.1)$$

where

$$U^{\text{PMNS}} = U^{\text{CKM}} \begin{pmatrix} e^{\frac{i\alpha_M}{2}} & 0 & 0 \\ 0 & e^{\frac{i\beta_M}{2}} & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (3.2)$$

and

$$P_\phi = \begin{pmatrix} e^{i\phi_1} & 0 & 0 \\ 0 & e^{i\phi_2} & 0 \\ 0 & 0 & e^{i\phi_3} \end{pmatrix}. \quad (3.3)$$

P_ϕ is the unphysical phase matrix with unphysical phases $\phi_{1,2,3}$. Phase matrix in extreme right of the U^{PMNS} matrix contains two Majorana phases α_M and β_M . Now m_ν can be diagonalized as

$$U_{tot}^\dagger m_\nu U_{tot}^* = \text{diag}(m_1, m_2, m_3) \quad (3.4)$$

which can be inverted as

$$m_\nu = U_{tot} \text{diag}(m_1, m_2, m_3) U_{tot}^T. \quad (3.5)$$

Equating both sides of eq.(3.5) elements of m_ν matrix can be written in terms of masses, mixing angles and phases as

$$(m_\nu)_{11} = e^{2i\phi_1} (c_{12}^2 c_{13}^2 m_1 e^{i\alpha_M} + s_{12}^2 c_{13}^2 m_2 e^{i\beta_M} + m_3 s_{13}^2 e^{-2i\delta}) \quad (3.6)$$

$$(m_\nu)_{12} = e^{i(\phi_1+\phi_2)} c_{13} \{ -m_1 e^{i\alpha_M} (c_{12} s_{12} c_{23} + c_{12}^2 s_{13} s_{23} e^{i\delta}) + m_2 e^{i\beta_M} (c_{12} s_{12} c_{23} - s_{12}^2 s_{13} s_{23} e^{i\delta}) + m_3 s_{13} s_{23} e^{-i\delta} \} \quad (3.7)$$

$$(m_\nu)_{13} = e^{i(\phi_1+\phi_3)} c_{13} \{ m_1 e^{i\alpha_M} (-c_{12}^2 c_{23} s_{13} e^{i\delta} + c_{12} s_{12} s_{23}) - m_2 e^{i\beta_M} (c_{12} s_{12} s_{23} + s_{12}^2 s_{13} c_{23} e^{i\delta}) + m_3 s_{13} c_{23} e^{-i\delta} \} \quad (3.8)$$

$$(m_\nu)_{22} = e^{2i\phi_2} \{ m_1 e^{i\alpha_M} (s_{12} c_{23} + c_{12} s_{23} s_{13} e^{i\delta})^2 + m_2 e^{i\beta_M} (c_{12} c_{23} - s_{12} s_{13} s_{23} e^{i\delta})^2 + m_3 c_{13}^2 s_{23}^2 \} \quad (3.9)$$

$$(m_\nu)_{23} = e^{i(\phi_2+\phi_3)} [m_1 e^{i\alpha_M} \{ c_{12} s_{12} s_{13} (c_{23}^2 - s_{23}^2) e^{i\delta} + c_{12}^2 c_{23} s_{23} s_{13}^2 e^{2i\delta} - s_{12}^2 s_{23} c_{23} \} - m_2 e^{i\beta_M} \{ c_{12} s_{12} (c_{23}^2 - s_{23}^2) s_{13} e^{i\delta} + c_{12}^2 s_{23} c_{23} - s_{12}^2 s_{13}^2 s_{23} c_{23} e^{2i\delta} \} + m_3 c_{13}^2 c_{23} s_{23}] \quad (3.10)$$

$$(m_\nu)_{33} = e^{2i\phi_3} \{ m_1 e^{i\alpha_M} (c_{12} c_{23} s_{13} e^{i\delta} - s_{12} s_{23})^2 + m_2 e^{i\beta_M} (s_{12} c_{23} s_{13} e^{i\delta} + c_{12} s_{23})^2 + m_3 c_{13}^2 c_{23}^2 \}. \quad (3.11)$$

We now extract α_M and β_M eliminating unwanted ϕ_i phases. Modulus $|(m_\nu)_{ij}|$ of all elements are free from ϕ_i phases. The combinations such as $\frac{|(m_\nu)_{ij}|^2}{(m_\nu)_{ii}(m_\nu)_{jj}}$ ($i \neq j$) are also independent of those ϕ_i phases. Neglecting terms of $O(s_{13}^2)$ and higher order, we find, among all the $|(m_\nu)_{ij}|$ terms, the term $|(m_\nu)_{11}|$ has the simplest structure and independent of ϕ_i . We can easily extract $\beta_M - \alpha_M$ from this term as

$$\cos(\beta_M - \alpha_M) = \frac{|(m_\nu)_{11}|^2 - c_{12}^4 m_1^4 - s_{12}^4 m_2^2}{2c_{12}^2 s_{12}^2 m_1 m_2}. \quad (3.11)$$

To find the individual value of Majorana phases, we consider the term $\frac{[(m_\nu)_{23}]^2}{(m_\nu)_{22}(m_\nu)_{33}}$ which looks simpler by neglecting terms like $(c_{23}^2 - s_{23}^2)s_{13}$, s_{13}^2 and their higher power. Substituting Majorana phase difference $\beta_M - \alpha_M$ in the term $\frac{[(m_\nu)_{23}]^2}{(m_\nu)_{22}(m_\nu)_{33}}$ we can construct two different complex equations only with α_M and β_M respectively. It is straightforward to find out two Majorana phases with the chain of expressions in a generic form as

$$\tan \theta_j = \frac{Y'_j W_j - W'_j Y_j}{X_j W'_j - W_j X'_j} \quad (3.12)$$

where $j = 1, 2$ and $\theta_1 = \alpha_M$, $\theta_2 = \beta_M$ with

$$\begin{aligned} X_1 &= A_i - \{D_r \sin(\beta_M - \alpha_M) + D_i \cos(\beta_M - \alpha_M) + F_r \sin 2(\beta_M - \alpha_M) + \\ &\quad F_i \cos 2(\beta_M - \alpha_M) + E_i\} \\ X'_1 &= \{D_r \cos(\beta_M - \alpha_M) - D_i \sin(\beta_M - \alpha_M) + F_r \cos 2(\beta_M - \alpha_M) - \\ &\quad F_i \sin 2(\beta_M - \alpha_M) + E_r\} - A_r \\ Y_1 &= A_r + \{D_r \cos(\beta_M - \alpha_M) - D_i \sin(\beta_M - \alpha_M) + F_r \cos 2(\beta_M - \alpha_M) - \\ &\quad F_i \sin 2(\beta_M - \alpha_M) + E_r\} \\ Y'_1 &= A_i + \{D_r \sin(\beta_M - \alpha_M) + D_i \cos(\beta_M - \alpha_M) + F_r \sin 2(\beta_M - \alpha_M) + \\ &\quad F_i \cos 2(\beta_M - \alpha_M) + E_i\} \\ W_1 &= B_r + C_r \cos(\beta_M - \alpha_M) - C_i \sin(\beta_M - \alpha_M) \\ W'_1 &= B_i + C_r \sin(\beta_M - \alpha_M) + C_i \cos(\beta_M - \alpha_M) \end{aligned} \quad (3.13)$$

and

$$\begin{aligned} X_2 &= A_i - \{D_i \cos(\beta_M - \alpha_M) - D_r \sin(\beta_M - \alpha_M) + E_i \cos 2(\beta_M - \alpha_M) - \\ &\quad E_r \sin 2(\beta_M - \alpha_M) + F_i\} \\ X'_2 &= \{D_r \cos(\beta_M - \alpha_M) + D_i \sin(\beta_M - \alpha_M) + E_r \cos 2(\beta_M - \alpha_M) + \\ &\quad E_i \sin 2(\beta_M - \alpha_M) + F_r\} - A_r \\ Y_2 &= A_r + \{D_r \cos(\beta_M - \alpha_M) + D_i \sin(\beta_M - \alpha_M) + E_r \cos 2(\beta_M - \alpha_M) + \\ &\quad E_i \sin 2(\beta_M - \alpha_M) + F_r\} \\ Y'_2 &= A_i + \{D_i \cos(\beta_M - \alpha_M) - D_r \sin(\beta_M - \alpha_M) + E_i \cos 2(\beta_M - \alpha_M) - \\ &\quad E_r \sin 2(\beta_M - \alpha_M) + F_i\} \\ W_2 &= C_r + B_r \cos(\beta_M - \alpha_M) + B_i \sin(\beta_M - \alpha_M) \\ W'_2 &= C_i + B_i \cos(\beta_M - \alpha_M) - B_r \sin(\beta_M - \alpha_M) \end{aligned} \quad (3.14)$$

where suffix i and r stand for imaginary and real part respectively. The complex quantities A, B, C, D, E and F are defined as follows

$$\begin{aligned}
A &= m_3^2[Z - 1] \\
B &= m_3 m_1 \left[Z s_{12}^2 \frac{1 + t_{23}^4}{t_{23}^2} - Z \sin 2\theta_{12} s_{13} e^{i\delta} \frac{t_{23}^2 - 1}{t_{23}} + 2s_{12}^2 \right] \\
C &= m_3 m_2 \left[Z c_{12}^2 \frac{1 + t_{23}^4}{t_{23}^2} + Z \sin 2\theta_{12} s_{13} e^{i\delta} \frac{t_{23}^2 - 1}{t_{23}} + 2c_{12}^2 \right] \\
D &= m_1 m_2 \left[2Z c_{12}^2 s_{12}^2 + \sin 2\theta_{12} \cos 2\theta_{12} s_{13} e^{i\delta} \frac{t_{23}^2 - 1}{t_{23}} - 2s_{12}^2 c_{12}^2 \right] \\
E &= m_1^2 \left[Z s_{12}^4 + Z s_{12}^2 \sin 2\theta_{12} s_{13} e^{i\delta} \frac{t_{23}^2 - 1}{t_{23}} - s_{12}^4 \right] \\
F &= m_2^2 \left[Z c_{12}^4 - Z c_{12}^2 \sin 2\theta_{12} s_{13} e^{i\delta} \frac{t_{23}^2 - 1}{t_{23}} - c_{12}^4 \right]
\end{aligned} \tag{3.15}$$

with $t_{23} = \tan \theta_{23}$ and $Z = \frac{[(M_\nu)_{23}]^2}{(M_\nu)_{22}(M_\nu)_{33}}$. Again in the expressions of B, C, D, E and F terms containing $s_{13}(t_{23}^2 - 1)e^{i\delta}$ is propotional to $s_{13}(c_{23}^2 - s_{23}^2)$. Dropping those terms one can further simplify the expressions of B, C, D, E and F keeping other dominating terms. This simplification makes expressions of A to F free from the Dirac phase and their complex nature is only due to Z parameter.

Thus, apart from the masses, finally, we gather complete information of the U^{PMNS} matrix containing mixing angles and physical phases from a general three generation Majorana neutrino mass matrix.

4 Cyclic Symmetry

4.1 Basic Formalism

The most general leptonic mass term of the Lagrangian in the present model is

$$- \mathcal{L}_{\text{mass}} = (m_\ell)_{ll'} \bar{l}_L l'_R + m_{D_{ll'}} \bar{\nu}_{lL} N_{l'R} + M_{R_{ll'}} \bar{N}_{l'L}^c N_{l'R} \tag{4.1}$$

where $l, l' = e, \mu, \tau$. We demand that the neutrino part of the Lagrangian is invariant under the cyclic permutation symmetry as given in eq.(1.1). The symmetry invariant Dirac neutrino mass matrix m_D takes the form

$$m_D = \begin{pmatrix} y_1 & y_2 & y_3 \\ y_3 & y_1 & y_2 \\ y_2 & y_3 & y_1 \end{pmatrix} \tag{4.2}$$

where in general all the entries are complex. Without loss of generality, we consider a basis in which the right handed neutrino mass matrix M_R and charged lepton mass matrix m_ℓ are diagonal. Further, imposition of cyclic symmetry dictates the texture of M_R as

$$M_R = \begin{pmatrix} m & 0 & 0 \\ 0 & m & 0 \\ 0 & 0 & m \end{pmatrix}. \tag{4.3}$$

Now, within the framework of type-I seesaw mechanism the effective neutrino mass matrix m_ν ,

$$m_\nu = -m_D M_R^{-1} m_D^T \quad (4.4)$$

takes the following form with cyclic symmetric m_D (eq.(4.2)) and M_R (eq.(4.3)) as

$$m_\nu = -\frac{1}{m} \begin{pmatrix} y_1^2 + y_2^2 + y_3^2 & y_1 y_2 + y_2 y_3 + y_3 y_1 & y_1 y_2 + y_2 y_3 + y_3 y_1 \\ y_1 y_2 + y_2 y_3 + y_3 y_1 & y_1^2 + y_2^2 + y_3^2 & y_1 y_2 + y_2 y_3 + y_3 y_1 \\ y_1 y_2 + y_2 y_3 + y_3 y_1 & y_1 y_2 + y_2 y_3 + y_3 y_1 & y_1^2 + y_2^2 + y_3^2 \end{pmatrix}. \quad (4.5)$$

4.2 Parametrization and Diagonalization

With a suitable choice of parametrization the effective neutrino mass matrix given in eq.(4.5) can be rewritten as

$$m_\nu = m_0 \begin{pmatrix} 1 + p^2 e^{2i\alpha} + q^2 e^{2i\beta} & p e^{i\alpha} + q e^{i\beta} + p q e^{i(\alpha+\beta)} & p e^{i\alpha} + q e^{i\beta} + p q e^{i(\alpha+\beta)} \\ p e^{i\alpha} + q e^{i\beta} + p q e^{i(\alpha+\beta)} & 1 + p^2 e^{2i\alpha} + q^2 e^{2i\beta} & p e^{i\alpha} + q e^{i\beta} + p q e^{i(\alpha+\beta)} \\ p e^{i\alpha} + q e^{i\beta} + p q e^{i(\alpha+\beta)} & p e^{i\alpha} + q e^{i\beta} + p q e^{i(\alpha+\beta)} & 1 + p^2 e^{2i\alpha} + q^2 e^{2i\beta} \end{pmatrix} \quad (4.6)$$

where we have parametrized the different elements (y_1, y_2, y_3) of m_ν in terms of p, q and two phases α, β accordingly

$$m_0 = -\frac{y_3^2}{m}, \quad p e^{i\alpha} = \frac{y_1}{y_3}, \quad q e^{i\beta} = \frac{y_2}{y_3}. \quad (4.7)$$

Denoting

$$\begin{aligned} P &= 1 + p^2 e^{2i\alpha} + q^2 e^{2i\beta} \\ Q &= p e^{i\alpha} + q e^{i\beta} + p q e^{i(\alpha+\beta)} \end{aligned} \quad (4.8)$$

m_ν is written in a convenient form as

$$m_\nu = m_0 \begin{pmatrix} P & Q & Q \\ Q & P & Q \\ Q & Q & P \end{pmatrix}. \quad (4.9)$$

We construct the matrix $h(= m_\nu m_\nu^\dagger)$ to calculate the mixing angles and mass eigenvalues. Expression of h obtained as

$$h = m_\nu m_\nu^\dagger = m_0^2 \begin{pmatrix} A & B & B \\ B & A & B \\ B & B & A \end{pmatrix} \quad (4.10)$$

where

$$\begin{aligned} A &= |P|^2 + 2|Q|^2 \\ B &= |Q|^2 + P Q^* + P^* Q. \end{aligned} \quad (4.11)$$

Diagonalizing the matrix h given in eq.(4.10) through $U^\dagger h U = \text{diag}(m_1^2, m_2^2, m_3^2)$ we get the mass squared eigenvalues as

$$\begin{aligned} m_1^2 &= m_0^2(A - B) \\ m_2^2 &= m_0^2(A + 2B) \\ m_3^2 &= m_0^2(A - B). \end{aligned} \quad (4.12)$$

However, there is a problem of unique determination of the diagonalization matrix U due to the degeneracy in the eigenvalues ($m_1^2 = m_3^2 \neq m_2^2$). Any vector in the plane orthogonal to the unique eigenvector of eigenvalue m_2^2 can be an eigen vector of m_1^2 or m_3^2 . One can choose two mutually orthogonal eigenvectors on that plane for the eigenvalues m_1^2 and m_3^2 . So, in effect, we can have the U matrix of the above case with these three eigenvectors. But, choice of eigenvectors for m_1^2 and m_3^2 on the degenerate plane is arbitrary. Any other two orthogonal combinations of these two eigenvectors are equally good for construction of the U matrix for the same eigenvalues. So, the diagonalization matrix can not be unique and hence the derived mixing angles are also not unique.

Here, one observation is that the eigenvector of m_2^2 : $(1/\sqrt{3}, 1/\sqrt{3}, 1/\sqrt{3})$ coincides with the 2nd column of TBM mixing matrix. Due to degeneracy $m_1^2 = m_3^2$, one of the possible choice of diagonalization matrix could be the well known TBM mixing matrix. However, it is also possible to generate all three mixing angles nonzero by proper combination of eigenvectors corresponding to the degenerate eigenvalues. Furthermore, in order to accommodate solar and atmospheric neutrino mass squared differences it is necessary to break the symmetry to remove the degeneracy between the mass eigenvalues.

5 Breaking of cyclic symmetry

In this scheme, we break the cyclic symmetry in the right chiral neutrino sector only. Retaining the flavour diagonal texture of M_R , we introduce only two symmetry breaking parameters ϵ_1 and ϵ_2 in any two diagonal entries. (It is sufficient to incorporate two symmetry breaking parameters to achieve all the eigenvalues of M_R different). This can be done in three ways as

$$\begin{aligned} \text{(i)} & M_R = \text{diag} \left(m, m + \epsilon_1, m + \epsilon_2 \right), \\ \text{(ii)} & M_R = \text{diag} \left(m + \epsilon_1, m + \epsilon_2, m \right), \\ \text{(iii)} & M_R = \text{diag} \left(m + \epsilon_1, m, m + \epsilon_2 \right). \end{aligned}$$

It is to be noted that instead of perturbative approach, we directly diagonalize the broken symmetric mass matrix with the help of the results obtained in section 2. Let us first consider case (i) where symmetry breaking occurs at ‘22’ and ‘33’ elements. Using the expression of M_R given in (i) and m_D as given in eq.(4.2), the effective neutrino mass matrix is obtained due to type-I seesaw mechanism as

$$m_\nu = -\frac{y_3^2}{m} \begin{pmatrix} \frac{y_1^2}{y_3} + \frac{y_2^2}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{1}{(1+\epsilon'_2)} & \frac{y_1}{y_3} + \frac{y_1 y_2}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{y_2}{y_3} \frac{1}{(1+\epsilon'_2)} & \frac{y_1 y_2}{y_3} + \frac{y_2}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{y_1}{y_3} \frac{1}{(1+\epsilon'_2)} \\ \frac{y_1}{y_3} + \frac{y_1 y_2}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{y_2}{y_3} \frac{1}{(1+\epsilon'_2)} & 1 + \frac{y_1^2}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{y_2^2}{y_3} \frac{1}{(1+\epsilon'_2)} & \frac{y_2}{y_3} + \frac{y_1}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{y_1 y_2}{y_3} \frac{1}{(1+\epsilon'_2)} \\ \frac{y_1 y_2}{y_3} + \frac{y_2}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{y_1}{y_3} \frac{1}{(1+\epsilon'_2)} & \frac{y_2}{y_3} + \frac{y_1}{y_3} \frac{1}{(1+\epsilon'_1)} + \frac{y_1 y_2}{y_3} \frac{1}{(1+\epsilon'_2)} & \frac{y_2^2}{y_3} + \frac{y_1^2}{y_3} \frac{1}{(1+\epsilon'_2)} + \frac{1}{(1+\epsilon'_1)} \end{pmatrix} \quad (5.1)$$

where we have defined $\epsilon'_1 = \frac{\epsilon_1}{m}, \epsilon'_2 = \frac{\epsilon_2}{m}$. We rewrite m_ν as

$$m_\nu = m_0 \begin{pmatrix} p^2 e^{2i\alpha} + \frac{q^2 e^{2i\beta}}{(1+\epsilon'_1)} + \frac{1}{(1+\epsilon'_2)} & p e^{i\alpha} + \frac{q e^{i\beta}}{(1+\epsilon'_2)} + \frac{p q e^{i(\alpha+\beta)}}{(1+\epsilon'_1)} & \frac{p e^{i\alpha}}{(1+\epsilon'_2)} + \frac{q e^{i\beta}}{(1+\epsilon'_1)} + p q e^{i(\alpha+\beta)} \\ p e^{i\alpha} + \frac{q e^{i\beta}}{(1+\epsilon'_2)} + \frac{p q e^{i(\alpha+\beta)}}{(1+\epsilon'_1)} & 1 + \frac{p^2 e^{2i\alpha}}{(1+\epsilon'_1)} + \frac{q^2 e^{2i\beta}}{(1+\epsilon'_2)} & \frac{p e^{i\alpha}}{(1+\epsilon'_1)} + q e^{i\beta} + \frac{p q e^{i(\alpha+\beta)}}{(1+\epsilon'_2)} \\ \frac{p e^{i\alpha}}{(1+\epsilon'_2)} + \frac{q e^{i\beta}}{(1+\epsilon'_1)} + p q e^{i(\alpha+\beta)} & \frac{p e^{i\alpha}}{(1+\epsilon'_1)} + q e^{i\beta} + \frac{p q e^{i(\alpha+\beta)}}{(1+\epsilon'_2)} & \frac{p^2 e^{2i\alpha}}{(1+\epsilon'_2)} + q^2 e^{2i\beta} + \frac{1}{(1+\epsilon'_1)} \end{pmatrix} \quad (5.2)$$

where we mimic the parametrization previously shown in eq.(4.7). The other two cases, case (ii) and (iii) also produce the same form of m_ν given in eq.(5.2) with a different set of parametrizations given by

•Case (ii)

$$m_0 = -\frac{y_1^2}{m}, \quad p e^{i\alpha} = \frac{y_2}{y_1}, \quad q e^{i\beta} = \frac{y_3}{y_1} \quad (5.3)$$

•Case(iii)

$$m_0 = -\frac{y_2^2}{m}, \quad p e^{i\alpha} = \frac{y_3}{y_2}, \quad q e^{i\beta} = \frac{y_1}{y_2}. \quad (5.4)$$

5.1 Numerical results and phenomenological discussions

It is now straightforward to calculate the eigenvalues and mixing angles of the above mass matrix m_ν . The coefficients a, b, c and d of the general characteristic equation (eq.(2.6)) can be written in terms of Lagrangian parameters (p, q, α, β) through the substitution of elements of general m_ν (eq.(2.1)) by the corresponding elements of broken symmetric m_ν (eq.(5.2)). Substituting these values in eq.(2.8), (2.9) and (2.10) it is possible to calculate three eigenvalues. The mapping of $(\lambda_1, \lambda_2, \lambda_3)$ to (m_1^2, m_2^2, m_3^2) is done by utilizing neutrino oscillation experimental data shown in Table 5.1.

Table 1. Input experimental values [10]

Quantity	3σ ranges/other constraint
Δm_{21}^2	$7.12 < \Delta m_{21}^2 (10^5 \text{ eV}^{-2}) < 8.20$
$ \Delta m_{31}^2 (N)$	$2.31 < \Delta m_{31}^2 (10^3 \text{ eV}^{-2}) < 2.74$
$ \Delta m_{31}^2 (I)$	$2.21 < \Delta m_{31}^2 (10^3 \text{ eV}^{-2}) < 2.64$
θ_{12}	$31.30^\circ < \theta_{12} < 37.46^\circ$
θ_{23}	$36.86^\circ < \theta_{23} < 55.55^\circ$
θ_{13}	$7.49^\circ < \theta_{13} < 10.46^\circ$
δ	$0 - 2\pi$

Before proceeding to carry out the numerical analysis few remarks are in order :

i) Taking into account different cosmological experiments with recent PLANCK satellite experimental results [8] the upper limit of the sum of the three neutrino masses can vary mostly within the range as $\Sigma m_i (= m_1 + m_2 + m_3) < (0.23 - 1.11) \text{ eV}$ [79]. A combined analysis of PLANCK, WMAP low l polarization, gravitational lensing and results of prior

on the Hubble constant H_0 from Hubble space telescope data corresponds to the higher value of Σm_i whereas inclusion of SDSS DR8 result with the above combination sharply reduce the upper limit of Σm_i at the above mentioned lower edge. However, in our set up individual masses of the neutrinos and sum of the neutrino masses are considered as predictions of this model. We investigate to check the viability of the sum of the three neutrino masses in view of the upper bound provided by the extant cosmological data.

ii) Another constrain arises from $\beta\beta_{0\nu}$ decay experiments [10–12] on the matrix element $|m_{\nu_{ee}}| (= m_{\nu_{11}})$. At present lots of experiments are running/proposed among them EXO-200 Collaboration [80] has quoted a range on the upper limit of $|m_{\nu_{ee}}|$ as $|m_{\nu_{ee}}| < (0.14 - 0.38)\text{eV}$. In the present work, we are not restricting the value of $|m_{\nu_{ee}}|$ rather treat it also as a prediction to testify the present model in the foreseeable future.

We have varied the symmetry breaking parameters ϵ'_1, ϵ'_2 in the range $-0.1 < \epsilon'_1, \epsilon'_2 < 0.1$ to keep the symmetry breaking effect small. With such values of $\epsilon'_{1,2}$ and taking neutrino experimental data [10, 81, 82] given in Table 5.1 as input, we find admissible parameter space of the model. The allowed region of the p vs q parametric plane is shown in left panel of figure 1, wherefrom the allowed ranges of p and q can be read as $0.27 < p < 2.09$, $0.44 < q < 2.21$. The two phase parameters α and β are varied as $-180^\circ < \alpha, \beta < 180^\circ$ and the allowed parameter space in α vs β plane is shown in right panel of figure 1. Two tiny disconnected patches are allowed and one is mirror image to the other. The allowed ranges of α, β obtained as $-161.12^\circ < \alpha < -89.35^\circ$ with $91.09^\circ < \beta < 166.53^\circ$ and $90.80^\circ < \alpha < 161.02^\circ$ with $-166.35^\circ < \beta < -92.11^\circ$. Next in figure 2, in the left panel we plot Σm_i vs $|m_{\nu_{ee}}|$ and the ranges obtained as $0.076\text{eV} < \Sigma m_i < 0.23\text{eV}$, $0.002\text{eV} < |m_{\nu_{ee}}| < 0.069\text{eV}$. The upper limit of Σm_i obtained from figure 2 marginally touches the most optimistic cosmological upper bound 0.23 eV , however, the lower limit is very far to probe in the near future. On the otherhand, both the higher and lower values $|m_{\nu_{ee}}|$ is well within the upper bound of running/proposed experiments (for example KamLAND+Zen, EXO). In the right panel of figure 2, m_1 vs $m_{2,3}$ plot is given and it is clear from the plot that the mass ordering is normal ($m_1 < m_2 < m_3$). The ranges of individual mass eigenvalues obtained as $0.0122\text{eV} < m_1 < 0.0720\text{eV}$, $0.0143\text{eV} < m_2 < 0.0730\text{eV}$, $0.0495\text{eV} < m_3 < 0.09\text{eV}$. Thus, the testability of the present model crucially relies upon the determination of the neutrino mass hierarchy by future neutrino experiments. We have successively plotted the variation of Jarlskog invariant J_{CP} ³ with the Dirac CP phase (δ) in the left panel of figure 3 and Majorana phases α_M vs β_M in the right panel of figure 3. We see that $-0.044 < J_{\text{CP}} < 0.044$ and all values of δ lies within the range -90° to 90° whereas Majorana phases admit almost all values in the range $-90^\circ < \alpha_M, \beta_M < 90^\circ$. Before concluding this section we like to comment on the necessity of the two breaking parameters ϵ'_1 and ϵ'_2 . It is seen from the present analysis that in the present model it is possible to explain the neutrino oscillation data with either of the ϵ'_i parameter equal to zero.

$$^3 J_{\text{CP}} = \frac{\text{Im}(h_{12}h_{23}h_{31})}{(m_2^2 - m_1^2)(m_3^2 - m_2^2)(m_3^2 - m_1^2)} = \frac{\sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \sin \delta}{8}$$




Figure 1. (colour online) Plot of the allowed parameter space in p, q (left) plane and α, β (right) plane satisfying input data shown in Table 5.1




Figure 2. (colour online) Plot of Σm_i vs $|m_{\nu ee}|$ (left), m_1 vs $m_{2,3}$ (right) satisfying input data shown in Table 5.1




Figure 3. (colour online) Plot of δ vs J_{CP} (left) and α_M vs β_M (right) satisfying input data shown in Table 5.1

6 Summary

The main objective of this paper is to develop a simple methodology to obtain exact mass eigenvalues, mixing angles, the Majorana phases and the Dirac CP phase of a general complex symmetric Majorana neutrino mass matrix without any approximation. The hermitian matrix h constructed from m_ν ($h = m_\nu m_\nu^\dagger$) is solved to get the squared mass eigenvalues. The elements of the diagonalization matrix U and hence, three mixing angles and the Dirac CP phase δ are calculated by solving the set of eigenvalue equations. Since m_ν has twelve independent parameters, the total diagonalization matrix which diagonalizes m_ν , should contain five more phase parameters apart from the Dirac CP phase (The other six parameters are three mass squared values and three mixing angles.). The above mentioned five phase parameters contain three unphysical phases and two Majorana phases. General expressions for the Majorana phases are obtained by eliminating those unphysical ones.

We demonstrate this general and exact methodology in the context of a neutrino mass matrix obtained from a cyclic symmetry transformation invoking type-I seesaw mechanism. The symmetry invariant structure of the effective neutrino mass matrix leads to degeneracy in the mass eigenvalues and thereby, prohibited by the experimental data. The symmetry is broken in the right handed neutrino sector only in order to fulfill the phenomenological demands of nonzero mass squared differences and mixing angles. All the physical parameters (three mixing angles, one Dirac CP phase, two Majorana phases) of the total diagonalization matrix (U_{tot}) and the mass eigenvalues of the broken symmetric mass matrix are readily expressed in terms of the Lagrangian parameters through the utilization of the results obtained from general diagonalization procedure. For completeness of the analysis, we explore the parameter space and it is revealed that the mass hierarchy of the neutrinos is normal and inverted hierarchy is completely ruled out. Plots of the allowed parameter space show that this model is capable of producing those observables (mixing angles, solar and atmospheric mass squared differences) within experimentally constrained

ranges. Finally, the exact expressions obtained for physical parameters can be directly applicable in any (symmetry invariant or broken) neutrino mass matrix.

A Appendix

A.1 Elements of h in terms of elements of m_ν (a_i, b_i)

$$h_{11} = a_1^2 + b_1^2 + a_2^2 + b_2^2 + a_3^2 + b_3^2 \quad (\text{A.1})$$

$$h_{22} = a_2^2 + b_2^2 + a_4^2 + b_4^2 + a_5^2 + b_5^2 \quad (\text{A.2})$$

$$h_{33} = a_3^2 + b_3^2 + a_5^2 + b_5^2 + a_6^2 + b_6^2 \quad (\text{A.3})$$

$$h_{12} = (a_1a_2 + b_1b_2 + a_2a_4 + b_2b_4 + a_3a_5 + b_3b_5) \\ + i(b_1a_2 - a_1b_2 + b_2a_4 - a_2b_4 + b_3a_5 - a_3b_5) \quad (\text{A.4})$$

$$h_{13} = (a_1a_3 + b_1b_3 + a_2a_5 + b_2b_5 + a_3a_6 + b_3b_6) \\ + i(b_1a_3 - a_1b_3 + b_2a_5 - a_2b_5 + b_3a_6 - a_3b_6) \quad (\text{A.5})$$

$$h_{23} = (a_2a_3 + b_2b_3 + a_4a_5 + b_4b_5 + a_5a_6 + b_5b_6) \\ + i(b_2a_3 - a_2b_3 + b_4a_5 - a_4b_5 + b_5a_6 - a_5b_6) \quad (\text{A.6})$$

A.2 Coefficients of the cubic equation in terms of elements of h

$$a = 1 \quad (\text{A.7})$$

$$b = -(h_{11} + h_{22} + h_{33}) \quad (\text{A.8})$$

$$c = h_{33}h_{11} + h_{33}h_{22} + h_{11}h_{22} - |h_{12}|^2 - |h_{13}|^2 - |h_{23}|^2 \quad (\text{A.9})$$

$$d = h_{11}|h_{23}|^2 + h_{33}|h_{12}|^2 + h_{22}|h_{13}|^2 - h_{11}h_{22}h_{33} - 2\text{Re}(h_{12}h_{23}h_{13}^*) \quad (\text{A.10})$$

References

- [1] B. Aharmim *et al.* [SNO Collaboration], *An Independent Measurement of the Total Active B-8 Solar Neutrino Flux Using an Array of He-3 Proportional Counters at the Sudbury Neutrino Observatory*, Phys. Rev. Lett. **101** (2008) 111301, [arXiv:0806.0989 [nucl-ex]].
- [2] B. Aharmim *et al.* [SNO Collaboration], *Low Energy Threshold Analysis of the Phase I and Phase II Data Sets of the Sudbury Neutrino Observatory*, Phys. Rev. **C 81** (2010) 055504 [arXiv:0910.2984 [nucl-ex]].
- [3] R. Wendell *et al.* [Super-Kamiokande Collaboration], *Atmospheric neutrino oscillation analysis with sub-leading effects in Super-Kamiokande I, II, and III*, Phys. Rev. **D 81** (2010) 092004 [arXiv:1002.3471 [hep-ex]].
- [4] [T2K Collaboration] K. Abe *et al.*, *Indication of Electron Neutrino Appearance from an Accelerator-produced Off-axis Muon Neutrino Beam*, Phys. Rev. Lett. **107** (2011) 041801 [arXiv:1106.2822 [hep-ex]].
- [5] [RENO Collaboration] J. K. Ahn *et al.*, *Observation of Reactor Electron Antineutrino Disappearance in the RENO Experiment*, Phys.Rev.Lett. **108** (2012) 191802 [arXiv:1204.0626 [hep-ex]].

- [6] [DAYA-BAY Collaboration] F. P. An *et al.*, *Observation of electron-antineutrino disappearance at Daya Bay*, Phys. Rev. Lett. **108** (2012) 171803 [arXiv:1203.1669 [hep-ex]].
- [7] [Double Chooz Collaboration] Y. Abe *et al.*, Phys. Rev. Lett. **108** (2012) 131801.
- [8] [Planck Collaboration] P. A. R. Ade *et al.*, *Planck 2013 results. XVI. Cosmological parameters*, arXiv:1303.5076 [astro-ph.CO].
- [9] C. L. Bennett, D. Larson, J. L. Weiland, N. Jarosik, G. Hinshaw, N. Odegard, K. M. Smith and R. S. Hill *et al.*, *Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results*, arXiv:1212.5225 [astro-ph.CO].
- [10] D. V. Forero, M. Tortola and J. W. F. Valle, *Global status of neutrino oscillation parameters after Neutrino-2012*, Phys. Rev. **D 86** (2012) 073012 [arXiv:1205.4018 [hep-ph]].
- [11] A. Giuliani, *Searches for neutrinoless double beta decay*, Acta Phys. Polon. **B 41** (2010) 1447.
- [12] W. Rodejohann, *Neutrinoless double beta decay and neutrino physics*, J. Phys. **G 39** (2012) 124008 [arXiv:1206.2560 [hep-ph]].
- [13] S. Morisi and J. W. F. Valle, *Neutrino masses and mixing: a flavour symmetry roadmap*, arXiv:1206.6678 [hep-ph].
- [14] S. F. King and C. Luhn, *Neutrino Mass and Mixing with Discrete Symmetry*, Rept. Prog. Phys. **76** (2013) 056201 [arXiv:1301.1340 [hep-ph]].
- [15] A. Y. Smirnov, *Neutrino mass, mixing and discrete symmetries*, arXiv:1305.4827 [hep-ph].
- [16] G. Altarelli, *Status of neutrino masses and mixing in 2009*, [arXiv:0905.3265]; E. Ma and G. Rajasekaran, *Softly broken A_4 symmetry for nearly degenerate neutrino masses*, Phys. Rev. **D 64** (2001) 113012 [hep-ph/0106291]; K.S. Babu, E. Ma and J.W.F. Valle, *Underlying A_4 symmetry for the neutrino mass matrix and the quark mixing matrix*, Phys. Lett. **B 552** (2003) 207 [hep-ph/0206292]; M. Hirsch, J.C. Romao, S. Skadhauge, J.W.F. Valle and A. Villanova del Moral, *Degenerate neutrinos from a supersymmetric A_4 model*, [hep-ph/0312244], *Phenomenological tests of supersymmetric A_4 family symmetry model of neutrino mass*, Phys. Rev. **D 69** (2004) 093006 [hep-ph/0312265]; E. Ma, *A_4 origin of the neutrino mass matrix*, Phys. Rev. **D 70** (2004) 031901 [hep-ph/0404199], *Non-Abelian discrete symmetries and neutrino masses: two examples*, New J. Phys. **6** (2004) 104 [hep-ph/0405152], *Tetrahedral family symmetry and the neutrino mixing matrix*, Mod. Phys. Lett. A **20** (2005) 2601 [hep-ph/0508099], *Aspects of the tetrahedral neutrino mass matrix*, Phys. Rev. **D 72** (2005) 037301 [hep-ph/0505209]; S.-L. Chen, M. Frigerio and E. Ma, *Hybrid seesaw neutrino masses with A_4 family symmetry*, Nucl. Phys. **B 724** (2005) 423 [hep-ph/0504181]; M. Hirsch, A. Villanova del Moral, J.W.F. Valle and E. Ma, *Predicting neutrinoless double beta decay*, Phys. Rev. **D 72** (2005) 091301, [Erratum *ibid* **D 72** (2005) 119904] [hep-ph/0507148], K.S. Babu and X.-G. He, *Model of geometric neutrino mixing*, [hep-ph/0507217]; A. Zee, *Obtaining the neutrino mixing matrix with the tetrahedral group*, Phys. Lett. **B 630** (2005) 58 [hep-ph/0508278]; X.-G. He, Y.-Y. Keum and R.R. Volkas, *A_4 flavour symmetry breaking scheme for understanding quark and neutrino mixing angles* JHEP**04** (2006)039 [hep-ph/0601001]; B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma and M.K. Parida, *A_4 symmetry and prediction of U_{e3} in a modified Altarelli-Feruglio model*, Phys. Lett. **B 638** (2006) 345 [hep-ph/0603059]; E. Ma, *Tribimaximal neutrino mixing from a supersymmetric model with A_4 family symmetry*, Phys. Rev. **D 73** (2006) 057304 [hep-ph/0511133], *Suitability of A_4 as a family symmetry in grand unification*, Mod. Phys.

- Lett. **A 21** (2006) 2931 [hep-ph/0607190], *Supersymmetric $A_4 \times Z_3$ and A_4 realizations of neutrino tribimaximal mixing without and with corrections*, Mod. Phys. Lett. **A 22** (2007) 101 [hep-ph/0610342]; S.F. King and M. Malinsky, *A_4 family symmetry and quark-lepton unification*, Phys. Lett. **B 645** (2007) 351 [hep-ph/0610250]; S. Morisi, M. Picariello and E. Torrente-Lujan, *A model for fermion masses and lepton mixing in $SO(10) \times A_4$* , Phys. Rev. **D 75** (2007) 075015 [hep-ph/0702034]; F. Yin, *Neutrino mixing matrix in the 3-3-1 model with heavy leptons and A_4 symmetry*, Phys. Rev. **D 75** (2007) 073010 [arxiv:0704.3827]; F. Bazzocchi, S. Kaneko and S. Morisi, *A SUSY A_4 model for fermion masses and mixings* JHEP **03** (2008) 063 [arXiv:0707.3032]; F. Bazzocchi, S. Morisi and M. Picariello, *Embedding A_4 into left-right flavor symmetry: tribimaximal neutrino mixing and fermion hierarchy*, Phys. Lett. **B 659** (2008) 628 [arXiv:0710.2928]; M. Honda and M. Tanimoto, *Deviation from tri-bimaximal neutrino mixing in A_4 flavor symmetry*, Prog. Theor. Phys. **119** (2008) 583 [arXiv:0801.0181]; B. Brahmachari, S. Choubey and M. Mitra, *The A_4 flavor symmetry and neutrino phenomenology*, Phys. Rev. **D 77** (2008) 073008 [Erratum *ibid* **D 77** (2008) 119901] [arXiv:0801.3554]; B. Adhikary and A. Ghosal, *Nonzero U_{e3} , CP-violation and leptogenesis in a see-saw type softly broken A_4 symmetric model*, Phys. Rev. **D 78** (2008) 073007 [arXiv:0803.3582]; A. Ghosal, *Recent models of neutrino masses and mixing*, [hep-ph/0612245]; B. Adhikary and A. Ghosal, *Constraining CP violation in a softly broken A_4 symmetric model*, Phys. Rev. **D 75** (2007) 073020 [hep-ph/0609193]; G. Altarelli, F. Feruglio and C. Hagedorn, *A SUSY $SU(5)$ grand unified model of tri-bimaximal mixing from A_4* JHEP **03** (2008) 052 [arXiv:0802.0090]; M. Hirsch, S. Morisi and J.W.F. Valle, *Tri-bimaximal neutrino mixing and neutrinoless double beta decay*, Phys. Rev. **D 78** (2008) 093007 [arXiv:0804.1521]; P.H. Frampton and S. Matsuzaki, *Renormalizable A_4 model for lepton sector*, [arXiv:0806.4592]; C. Cski, C. Delaunay, C. Grojean and Y. Grossman, *A model of lepton masses from a warped extra dimension* JHEP **10** (2008) 055 [arXiv:0806.0356]; F. Bazzocchi, M. Frigerio and S. Morisi, *Fermion masses and mixing in models with $SO(10) \times A_4$ symmetry*, Phys. Rev. **D 78** (2008) 116018 [arXiv:0809.3573]; S. Morisi, *Tri-bimaximal lepton mixing with A_4 semidirect product $Z_2 \times Z_2 \times Z_2$* , Phys. Rev. **D 79** (2009) 033008 [arXiv:0901.1080]; P. Ciafaloni, M. Picariello, E. Torrente-Lujan and A. Urbano, *Neutrino masses and tribimaximal mixing in Minimal renormalizable SUSY $SU(5)$ Grand Unified Model with A_4 flavor symmetry*, Phys. Rev. **D 79** (2009) 116010 [0901.2236]; M.-C. Chen and S.F. King, *A_4 see-saw models and form dominance* JHEP **06** (2009) 072 [arXiv:0903.0125]; G. Altarelli and F. Feruglio, *Tri-bimaximal neutrino mixing from discrete symmetry in extra dimensions*, Nucl. Phys. **B 720** (2005) 64 [hep-ph/0504165]; G. Altarelli and F. Feruglio, *Tri-bimaximal neutrino mixing, A_4 and the modular symmetry*, Nucl. Phys. **B 741** 215 [hep-ph/0512103]; G. Altarelli, F. Feruglio and Y. Lin, *Tri-bimaximal neutrino mixing from orbifolding*, Nucl. Phys. **B 775** (2007) 31 [hep-ph/0610165]; Y. Lin, *A predictive A_4 model, charged lepton hierarchy and tri-bimaximal sum rule*, Nucl. Phys. **B 813** (2009) 91 [arXiv:0804.2867], *A dynamical approach to link low energy phases with leptogenesis*, [arXiv:0903.0831]; G. Altarelli and D. Meloni, *Embedding A_4 into $SU(3) \times U(1)$ flavor symmetry: large neutrino mixing and fermion mass hierarchy in $SO(10)$ GUT*, J. Phys. **G 36** (2009) 085005 [arXiv:0802.1693].
- [17] T. Fukuyama and H. Nishiura, *Mass matrix of Majorana neutrinos*, arXiv:hep-ph/9702253.
- [18] W. Rodejohann, *Broken $\mu - \tau$ symmetry and leptonic CP violation Prepared for 12th International Workshop on Neutrinos Telescopes: Twenty Years after the Supernova 1987A Neutrino Bursts Discovery, Venice, Italy, 6-9 Mar 2007*
- [19] J. C. Gomez-Izquierdo and A. Perez-Lorenzana, *Softly broken $\mu \leftrightarrow \tau$ symmetry in the*

minimal see-saw model Phys. Rev. **D 77**, 113015 (2008).

- [20] T. Baba, *What does mu - tau symmetry imply about leptonic CP violation?*, Int. J. Mod. Phys. **E 16** (2007) 1373.
- [21] N. Nimai Singh, H. Zeen Devi and M. Patgiri, *Phenomenology of neutrino mass matrices obeying μ - τ reflection symmetry*, arXiv:0707.2713 [hep-ph].
- [22] A. S. Joshipura and B. P. Kodrani, *Complex CKM matrix, spontaneous CP violation and generalized μ - τ symmetry*, Phys. Lett. **B 670**, 369 (2009).
- [23] B. Adhikary, *Soft breaking of $L_\mu - L_\tau$ symmetry: Light neutrino spectrum and leptogenesis*, Phys. Rev. **D 74** (2006) 033002 [arXiv:hep-ph/0604009].
- [24] T. Baba and M. Yasue, *Correlation between leptonic CP violation and mu - tau symmetry breaking*, Phys. Rev. **D 75**, 055001 (2007).
- [25] W. Grimus, *Realizations of mu - tau interchange symmetry*, arXiv:hep-ph/0610158.
- [26] Z. z. Xing, H. Zhang and S. Zhou, *Nearly tri-bimaximal neutrino mixing and CP violation from mu - tau symmetry breaking*, Phys. Lett. **B 641** (2006) 189.
- [27] N. Haba and W. Rodejohann, *A Supersymmetric Contribution to the Neutrino Mass Matrix and Breaking of mu-tau Symmetry*, Phys. Rev. **D 74** (2006) 017701.
- [28] R. N. Mohapatra, S. Nasri and H. B. Yu, *Grand unification of mu - tau symmetry* Phys. Lett. **B 636** (2006) 114.
- [29] Y. H. Ahn, S. K. Kang, C. S. Kim and J. Lee, *Phased breaking of mu - tau symmetry and leptogenesis*, Phys. Rev. **D 73** (2006) 093005.
- [30] K. Fuki and M. Yasue, *What does mu - tau symmetry imply in neutrino mixings?*, Phys. Rev. **D 73** (2006) 055014.
- [31] S. Nasri, *Implications of $\mu \leftrightarrow \tau$ symmetry on neutrinos and leptogenesis* Int. J. Mod. Phys. **A 20** (2005) 6258.
- [32] I. Aizawa and M. Yasue, *A new type of complex neutrino mass texture and mu - tau symmetry*, Phys. Rev. **D 73** (2006) 015002.
- [33] R. N. Mohapatra and W. Rodejohann, *Broken mu-tau Symmetry and Leptonic CP Violation*, Phys. Rev. **D 72** (2005) 053001.
- [34] I. Aizawa, M. Ishiguro, M. Yasue and T. Kitabayashi, *$\mu\tau$ permutation symmetry and neutrino mixing for a two-loop radiative mechanism*, J. Korean Phys. Soc. **46** (2005) 597.
- [35] T. Kitabayashi and M. Yasue, *mu-tau symmetry and maximal CP violation* Phys. Lett. **B 621** (2005) 133.
- [36] R. N. Mohapatra, S. Nasri and H. B. Yu, *Leptogenesis, $\mu - \tau$ Symmetry and θ_{13}* , Phys. Lett. **B 615** (2005) 231.
- [37] R. N. Mohapatra and S. Nasri, *Leptogenesis and mu - tau symmetry*, Phys. Rev. **D 71** (2005) 033001.
- [38] R. N. Mohapatra, *θ_{13} as a probe of $\mu \leftrightarrow \tau$ symmetry for leptons*, JHEP **0410** (2004) 027.
- [39] I. Aizawa, M. Ishiguro, T. Kitabayashi and M. Yasue, *Bilarge neutrino mixing and mu-tau permutation symmetry for two-loop radiative mechanism* Phys. Rev. **D 70** (2004) 015011.

- [40] I. de Medeiros Varzielas and G. G. Ross, *SU(3) family symmetry and neutrino bi-tri-maximal mixing*, Nucl. Phys. **B 733** (2006) 31.
- [41] P. F. Harrison and W. G. Scott, *mu-tau reflection symmetry in lepton mixing and neutrino oscillations*, Phys. Lett. **B 547** (2002) 219.
- [42] T. Kitabayashi and M. Yasue, *$S_{(2L)}$ permutation symmetry for left-handed mu and tau families and neutrino oscillations in an $SU(3)_L \times U(1)_N$ gauge model*, Phys. Rev. **D 67** (2003) 015006.
- [43] A. Ghosal, *A neutrino mass model with reflection symmetry*, Mod. Phys. Lett. **A 19** (2004) 2579.
- [44] B. Adhikary, A. Ghosal and P. Roy, *$\mu\tau$ symmetry, tribimaximal mixing and four zero neutrino Yukawa textures* JHEP **0910** (2009) 040 [arXiv:0908.2686 [hep-ph]].
- [45] B. Adhikary, M. Chakraborty and A. Ghosal, *Scaling ansatz, four zero Yukawa textures and large θ_{13}* , Phys. Rev. **D 86** (2012) 013015 [arXiv:1205.1355 [hep-ph]].
- [46] A. S. Joshipura and W. Rodejohann, *Scaling in the Neutrino Mass Matrix, $\mu - \tau$ Symmetry and the See-Saw Mechanism*, Phys. Lett. **B 678** (2009) 276 [arXiv:0905.2126 [hep-ph]].
- [47] R. N. Mohapatra and W. Rodejohann, *Scaling in the neutrino mass matrix* Phys. Lett. **B 644** (2007) 59 [hep-ph/0608111].
- [48] A. Blum, R. N. Mohapatra and W. Rodejohann, *Inverted mass hierarchy from scaling in the neutrino mass matrix: Low and high energy phenomenology*, Phys. Rev. **D 76** (2007) 053003 [arXiv:0706.3801 [hep-ph]].
- [49] M. Obara, *The Possible Textures in the Seesaw Realization of the Strong Scaling Ansatz and the Implications for Thermal Leptogenesis* arXiv:0712.2628 [hep-ph].
- [50] A. Damanik, M. Satriawan, Muslim and P. Anggraita, *Neutrino mass matrix from seesaw mechanism subjected to texture zero and invariant under a cyclic permutation* arXiv:0705.3290 [hep-ph].
- [51] S. Goswami and A. Watanabe, *Minimal Seesaw Textures with Two Heavy Neutrinos*, Phys. Rev. **D 79** (2009) 033004 [arXiv:0807.3438 [hep-ph]].
- [52] W. Grimus and L. Lavoura, *Softly broken lepton number $L_e - L_\mu - L_\tau$ with non-maximal solar neutrino mixing* J. Phys. **G 31** (2005) 683 [hep-ph/0410279].
- [53] M. S. Berger and S. Santana, *Combined flavor symmetry violation and lepton number violation in neutrino physics*, Phys. Rev. **D 74** (2006) 113007 [hep-ph/0609176].
- [54] S. Goswami, S. Khan and W. Rodejohann, *Minimal Textures in Seesaw Mass Matrices and their low and high Energy Phenomenology*, Phys. Lett. **B 680** (2009) 255 [arXiv:0905.2739 [hep-ph]].
- [55] A. Merle and W. Rodejohann, *The Elements of the neutrino mass matrix: Allowed ranges and implications of texture zeros*, Phys. Rev. **D 73** (2006) 073012 [hep-ph/0603111].
- [56] J. Gluza and R. Szafron, *Real and complex random neutrino mass matrices and θ_{13}* , Phys. Rev. **D 85** (2012) 047701 [arXiv:1111.7278 [hep-ph]].
- [57] X. -G. He and S. K. Majee, *Implications of Recent Data on Neutrino Mixing and Lepton Flavour Violating Decays for the Zee Model*, JHEP **1203**, (2012) 023 [arXiv:1111.2293 [hep-ph]].

- [58] G. Mangano, G. Miele, S. Pastor, O. Pisanti and S. Sarikas, *Updated BBN bounds on the cosmological lepton asymmetry for non-zero θ_{13}* , Phys. Lett. **B 708** (2012) 1 [arXiv:1110.4335 [hep-ph]].
- [59] Q. -H. Cao, S. Khalil, E. Ma and H. Okada, *Nonzero θ_{13} for Neutrino Mixing in a Supersymmetric B-L Gauge Model with T_7 Lepton Flavor Symmetry*, Phys. Rev. **D 84** (2011) 071302 [arXiv:1108.0570 [hep-ph]].
- [60] W. Chao and Y. -j. Zheng, *Relatively Large θ_{13} from Modification to the Tri-bimaximal, Bimaximal and Democratic Neutrino Mixing Matrices*, arXiv:1107.0738 [hep-ph].
- [61] D. Meloni, *Bimaximal mixing and large θ_{13} in a SUSY $SU(5)$ model based on S_4* , JHEP **1110** (2011) 010 [arXiv:1107.0221 [hep-ph]].
- [62] N. Haba and R. Takahashi, *Predictions via large θ_{13} from cascades*, Phys. Lett. **B 702** (2011) 388 [arXiv:1106.5926 [hep-ph]].
- [63] A. B. Balantekin, *The Last Neutrino Mixing angle θ_{13}* , J. Phys. Conf. Ser. **337** (2012) 012049 [arXiv:1106.5021 [hep-ph]].
- [64] P. Novella and f. t. D. C. collaboration, *Double Chooz: Searching for θ_{13} with reactor neutrinos*, arXiv:1105.6079 [hep-ex].
- [65] A. B. Balantekin, *Neutrino interactions in astrophysics and the third neutrino mixing angle θ_{13}* , AIP Conf. Proc. **1269** (2010) 195 [arXiv:1006.2836 [nucl-th]].
- [66] M. C. Gonzalez-Garcia, M. Maltoni and J. Salvado, *Updated global fit to three neutrino mixing: status of the hints of $\theta_{13} > 0$* , JHEP **1004**, (2010) 056 [arXiv:1001.4524 [hep-ph]].
- [67] E. E. Jenkins and A. V. Manohar, *Tribimaximal Mixing, Leptogenesis, and θ_{13}* , Phys. Lett. **B 668** (2008) 210 [arXiv:0807.4176 [hep-ph]].
- [68] A. B. Balantekin and D. Yilmaz, *Contrasting solar and reactor neutrinos with a non-zero value of θ_{13}* , J. Phys. **G 35** (2008) 075007 [arXiv:0804.3345 [hep-ph]].
- [69] V. Barger, R. Gandhi, P. Ghoshal, S. Goswami, D. Marfatia, S. Prakash, S. K. Raut and S. U. Sankar, *Neutrino mass hierarchy and octant determination with atmospheric neutrinos*, arXiv:1203.6012 [hep-ph].
- [70] Y. H. Ahn and S. K. Kang, *Non-zero θ_{13} and CP violation in a model with A_4 flavor symmetry*, arXiv:1203.4185 [hep-ph].
- [71] B. Brahmachari and A. Raychaudhuri, *Perturbative generation of θ_{13} from tribimaximal neutrino mixing*, arXiv:1204.5619 [hep-ph].
- [72] H. Ishimori and E. Ma, *New Simple A_4 Neutrino Model for Nonzero θ_{13} and Large δ_{CP}* , Phys. Rev. **D 86** (2012) 045030 [arXiv:1205.0075 [hep-ph]].
- [73] R. Dutta, U. Ch, A. K. Giri and N. Sahu, *Perturbative Bottom-up Approach for Neutrino Mass Matrix in Light of Large θ_{13} and Role of Lightest Neutrino Mass*, arXiv:1303.3357 [hep-ph].
- [74] Y. Koide, *Quark and lepton mass matrices with a cyclic permutation invariant form*, hep-ph/0005137.
- [75] A. Damanik, M. Satriawan, P. Anggraita, A. Hermanto and Muslim, *Neutrino Mass Matrix from Seesaw Mechanism with Heavy Majorana Neutrino Subject to Texture Zero and*

Invariant Under a Cyclic Permutation, J. Theor. Comput. Stud. **8**, (2008) 0102 [arXiv:0710.1742 [hep-ph]].

- [76] A. Damanik, arXiv:1004.1457 [hep-ph].
- [77] C.H. Albright, *Overview of neutrino mixing models and ways to differentiate among them*, [arXiv:0905.0146]; E. Ma, *$S_3 \times Z_3$ model of lepton mass matrices*, Phys. Rev. **D 44** (1991) 587; Y. Koide, *Universal seesaw mass matrix model with an S_3 symmetry*, Phys. Rev. **D 60** (1999) 077301 [hep-ph/9905416]; M. Tanimoto, *Large mixing angle MSW solution in S_3 flavor symmetry*, Phys. Lett. **B 483** (2000) 417 [hep-ph/0001306]; J. Kubo, *Majorana phase in minimal S_3 invariant extension of the standard model*, Phys. Lett. **B 578** (2004) 156 [Erratum ibid **B 619** (2005) 387] [hep-ph/0309167]; F. Caravaglios and S. Morisi, *Neutrino masses and mixings with an S_3 family permutation symmetry*, [hep-ph/0503234]; S. Morisi and M. Picariello, *The flavor physics in unified gauge theory from an S_{3P} discrete symmetry*, Int. J. Theor. Phys. **45** (2006) 1267 [hep-ph/0505113]; P.F. Harrison and W.G. Scott, *Permutation symmetry, tri-bimaximal neutrino mixing and the S_3 group characters*, Phys. Lett. **B 557** (2003) 76 [hep-ph/0302025]; W. Grimus and L. Lavoura, *$S_3 \times Z_2$ model for neutrino mass matrices* JHEP**08** (2005) 013 [hep-ph/0504153]; R.N. Mohapatra, S. Nasri and H.-B. Yu, *S_3 symmetry and tri-bimaximal mixing*, Phys. Lett. **B 639** (2006) 318 [hep-ph/0605020]; N. Haba and K. Yoshioka, *Discrete flavor symmetry, dynamical mass textures and grand unification*, Nucl. Phys. **B 739** (2006) 254 [hep-ph/0511108]; C.-Y. Chen and L. Wolfenstein, *Consequences of approximate S_3 symmetry of the neutrino mass matrix*, Phys. Rev. **D 77** (2008) 093009 [0709.3767]; S. Kaneko, H. Sawanaka, T. Shingai, M. Tanimoto and K. Yoshioka, *New approach to texture-zeros with S_3 symmetry: flavor symmetry and vacuum aligned mass textures*, [hep-ph/0703250]; Y. Koide, *S_3 symmetry and neutrino masses and mixings*, Eur. Phys. J. **C 50** (2007) 809 [hep-ph/0612058]; Y. Koide, *Permutation symmetry S_3 and VEV structure of flavor-triplet Higgs scalars*, Phys. Rev. **D 73** (2006) 057901 [hep-ph/0509214]; T. Teshima, *Flavor mass and mixing and S_3 symmetry: an S_3 invariant model reasonable to all*, Phys. Rev. **D 73** (2006) 045019 [hep-ph/0509094]; L. Lavoura and E. Ma, *Two predictive supersymmetric $S_3 \times Z_2$ models for the quark mass matrices*, Mod. Phys. Lett. **A 20** (2005) 1217 [hep-ph/0502181]; T. Araki, J. Kubo and E.A. Paschos, *S_3 flavor symmetry and leptogenesis*, Eur. Phys. J. **C 45** (2006) 465 [hep-ph/0502164]; N. Haba, A. Watanabe and K. Yoshioka, *Twisted flavors and tri-bi-maximal neutrino mixing*, Phys. Rev. Lett. **97** (2006) 041601 [hep-ph/0603116]; F. Feruglio and Y. Lin, *Fermion mass hierarchies and flavour mixing from a minimal discrete symmetry*, Nucl. Phys. **B 800** (2008) 77 [arXiv:0712.1528]; A.S. Joshipura and S.D. Rindani, *Vacuum solutions of neutrino anomalies through a softly broken $U(1)$ symmetry*, Eur. Phys. J. **C 14** (2000) 85 [hep-ph/9811252]; R.N. Mohapatra, A. Perez-Lorenzana and C.A. de Sousa Pires, *Type II seesaw and a gauge model for the bimaximal mixing explanation of neutrino puzzles*, Phys. Lett. **B 474** (2000) 355 [hep-ph/9911395]; Q. Shafi and Z. Tavartkiladze, *Anomalous flavor $U(1)$: predictive texture for bi-maximal neutrino mixing*, Phys. Lett. **B 482** (2000) 145 [hep-ph/0002150]; L. Lavoura, *New model for the neutrino mass matrix*, Phys. Rev. **D 62** (2000) 093011 [hep-ph/0005321]; W. Grimus and L. Lavoura, *A neutrino mass matrix with seesaw mechanism and two-loop mass splitting*, Phys. Rev. **D 62** (2000) 093012 [hep-ph/0007011]; T. Kitabayashi and M. Yasue, *Radiatively induced neutrino masses and oscillations in an $SU(3)_L \times U(1)_N$ gauge model*, Phys. Rev. **D 63** (2001) 095002 [hep-ph/0010087]; A. Aranda, C.D. Carone and P. Meade, *2002 $U(2)$ -like flavor symmetries and approximate bimaximal neutrino mixing*, Phys. Rev. **D 65** (2002) 013011 [hep-ph/0109120]; K.S. Babu and R.N. Mohapatra, *Predictive schemes*

- for *bimaximal neutrino mixings*, Phys. Lett. **B 532** (2002) 77 [hep-ph/0201176]; H.-J. He, D.A. Dicus and J.N. Ng, *Minimal schemes for large neutrino mixings with inverted hierarchy*, Phys. Lett. **B 536** (2002) 83 [hep-ph/0203237]; H.S. Goh, R.N. Mohapatra and S.-P. Ng, 2002 *Testing neutrino mass matrices with approximate $L_e - L_\mu - L_\tau$ symmetry*, Phys. Lett. **B 542** (2002) 116 [hep-ph/0205131]; G.K. Leontaris, J. Rizos and A. Psallidas, *Majorana neutrino masses from anomalous $U(1)$ symmetries*, Phys. Lett. **B 597** (2004) 182 [hep-ph/0404129] S. Dev, R. R. Gautam and L. Singh, *Broken S_3 Symmetry in the Neutrino Mass Matrix and Non-Zero θ_{13}* , Phys. Lett. **B 708** (2012) 284 [arXiv:1201.3755 [hep-ph]]; S. Dev, S. Gupta and R. R. Gautam, *Broken S_3 Symmetry in the Neutrino Mass Matrix*, Phys. Lett. **B 702** (2011) 28 [arXiv:1106.3873 [hep-ph]]; S. Zhou, *Relatively large θ_{13} and nearly maximal θ_{23} from the approximate S_3 symmetry of lepton mass matrices*, Phys. Lett. **B 704** (2011) 291 [arXiv:1106.4808 [hep-ph]]. [78]
- [78] J. Beringer *et al.* [Particle Data Group Collaboration], *Review of Particle Physics (RPP)*, Phys. Rev. **D 86** (2012) 010001.
- [79] E. Giusarma, R. de Putter, S. Ho and O. Mena, *Constraints on neutrino masses from Planck and Galaxy Clustering data* arXiv:1306.5544 [astro-ph.CO].
- [80] M. Auger *et al.* [EXO Collaboration], *Search for Neutrinoless Double-Beta Decay in ^{136}Xe with EXO-200* Phys. Rev. Lett. **109** (2012) 032505 [arXiv:1205.5608 [hep-ex]].
- [81] G. L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo and A. M. Rotunno, *Global analysis of neutrino masses, mixings and phases: entering the era of leptonic CP violation searches*, Phys. Rev. D **86** (2012) 013012 [arXiv:1205.5254 [hep-ph]].
- [82] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, *Global fit to three neutrino mixing: critical look at present precision*, JHEP **1212** (2012) 123 [arXiv:1209.3023 [hep-ph]].