

Ultraviolet Complete Technicolor and Higgs Physics at LHC

Matti Antola^{*†} and Stefano Di Chiara[‡]
*Helsinki Institute of Physics, P.O.Box 64,
FI-000140, Univ. of Helsinki, Finland*

Kimmo Tuominen[§]
*Department of Physics, P.O.Box 35,
FI-40014, Univ. of Jyväskylä, Finland and
Helsinki Institute of Physics, P.O.Box 64,
FI-000140, Univ. of Helsinki, Finland*

Abstract

We construct a Technicolor model which provides masses for the electroweak gauge bosons and for all the Standard Model matter fields. Starting from an ultraviolet complete supersymmetric technicolor, we propose a scenario where all elementary scalars, gauginos, and higgsinos are decoupled at an energy scale substantially higher than the electroweak scale, therefore avoiding the little hierarchy problem of the minimal supersymmetric standard model. The resulting low energy theory has an $SU(3)$ global symmetry whose breaking to $SO(3)$ leads to electroweak symmetry breaking. We study in detail the phenomenology of this theory and demonstrate that it reproduces the present LHC data at the same level of precision as the Standard Model itself.

^{*} Currently at Valo Research and Trading, Helsinki, Finland

[†]Electronic address: antola.matti@gmail.com

[‡]Electronic address: dichiara@helsinki.fi

[§]Electronic address: kimmo.i.tuominen@jyu.fi

I. INTRODUCTION

The data collected at the LHC experiments during the 7 and 8 TeV runs, with the epochal discovery of the Higgs boson [1, 2] and the measurement of its couplings [3, 4], seem to have provided the experimental verification of the Standard Model (SM) in its entirety. Compatibility of the present LHC data with the SM predictions does not imply that no beyond SM physics exists. Rather, there are several reasons to expect the contrary: The SM Higgs does not explain why the electroweak scale exists, but merely parametrizes its consequences. The masses of the SM fermions are not explained but remain as parameters fitted to their measured values; in particular, SM provides no insight to the flavor patterns or why there exists three generations of matter fields. There is no dark matter candidate in SM nor an explanation for the matter-antimatter asymmetry.

What the LHC data clearly indicates, is that the new physics coupled with the electroweak SM currents must have a typical scale of the order of a TeV or higher. One possibility is that the new physics scale is much above the terascale, and then naturalness as a model building paradigm should be reinterpreted [5, 6]. If the new physics scale on the other hand is near the terascale, then the present LHC data should provide hints of a new spectrum awaiting discovery in the future runs at the LHC. In this paper we investigate a model framework falling into the latter category.

In Technicolor (TC) theories [7, 8] the new physics scale is naturally of the order of a TeV. The electroweak (EW) symmetry is broken by a new strong interaction which generates a fermion condensate, in a way analogous to QCD but with a technipion decay constant approximately equal to the EW scale. The absence of light elementary scalars in TC automatically solves the fine-tuning problem traditionally associated with such massive fields without any protective symmetries. Gauge theories whose beta function stays small over a large range of energies constitute a popular class of TC models called walking TC [9, 10]. The lightest scalar in walking TC is the pseudo Nambu-Goldstone boson associated with the weakly broken conformal symmetry, and is therefore expected to be lighter than the other composite states [11–14]. Recently it has been shown that a further suppression of the naive prediction for the composite Higgs mass, obtained by scaling up the mass of the σ state of QCD [15], arises from the loop corrections from the heavy SM states [16]. These alone might explain the measured value, 125 GeV, of the Higgs mass.

To explain the observed mass patterns of the known matter fields within the TC framework one needs to couple TC fields with the matter fields of the SM. A well-known approach is extended TC (ETC) [17, 18], in which the technicolored fermions (technifermions) and the SM fermions are embedded into a representation of a larger gauge group containing the SM and TC gauge groups. If the ETC gauge group breaks sequentially, such ETC model may explain the observed mass hierarchies of the SM fermions [19, 20].

Another framework for extending TC with couplings to SM matter fields is bosonic TC (BTC). In BTC the couplings between SM fermions and composite scalars are introduced by assuming that both SM fermions and technifermions have Yukawa couplings to elementary heavy Higgs scalars [21–25]. To control fine-tuning at scales above the mass scales of the elementary scalars one can introduce supersymmetry (SUSY) [26, 27].

Recently we constructed a model along these lines and called it Minimal Super Conformal Technicolor (MSCT) [28, 29]. In MSCT the low energy effective theory is Minimal Walking Technicolor (MWT) [11, 30, 31] whose global symmetry is $SU(4)$. In this paper we describe another possibility within this model framework. We show that even though the particle content of the model is large, the low energy mass spectrum required by phenomenological viability can be simple: All the elementary scalars and gauginos are assumed heavy and integrated out. At the TeV scale this leaves, besides the SM fermions and gauge bosons, three Weyl technifermions, a technigluon, and a heavy fourth generation EW doublet of leptons. The TC sector of the mesoscopic Lagrangian obtained in this case features an $SU(3)$ global symmetry; due to its global symmetry and the resemblance to MWT we call this model $SU(3)$ Minimal Walking Technicolor (3MWT).

The 3MWT offers a concrete example of how the flavor physics sector can have a large effect on the low energy phenomenology, when compared to the underlying TC model without the coupling to the flavor sector. The four fermion interactions responsible for this deformation in our low energy theory result from attractive Yukawa interactions. This can be contrasted with the traditional ETC approach where the corresponding operators arise from gauge interactions which can be attractive or repulsive.

In Section II we briefly introduce MSCT as an ultraviolet (UV) complete TC theory and review its main features [29]. In Section III we derive the mesoscopic Lagrangian of 3MWT, by integrating out the heavy states below the SUSY breaking scale, m_{SUSY} , but above the dynamical scale of TC, Λ_{TC} . In Section IV we determine the effective Lagrangian at scales

Superfield	SU(2) _{TC}	SU(2) _L	U(1) _Y
Φ_L	Adj	\square	1/2
Φ_3	Adj	1	-1
V	Adj	1	0
Λ_L	1	\square	-3/2
N	1	1	1
E	1	1	2

TABLE I: Non-MSSM superfield content of MSCT. Here Adj and \square denote the adjoint and fundamental representations, respectively. None of the fields above is charged under SU(3)_c.

below Λ_{TC} in terms of SM fields, composite scalars (including the light Higgs boson), and composite vector degrees of freedom. In Section V we study the phenomenology of the model. Then, in Section VI we perform a goodness of fit analysis for 3MWT based on the current collider data (from both LHC and Tevatron) for Higgs physics, as well as the precision EW parameters S and T [32, 33]. We show that the data favors 3MWT over the SM. In Section VII we present our conclusions.

II. MSCT: A UV COMPLETE TECHNICOLOR

In MSCT the flavor extension (generating the SM fermion masses) corresponds to adding two Higgs scalar EW doublets with opposite hypercharge to MWT and supersymmetrizing the whole theory. The resulting model is gauge invariant under SU(2)_{TC} × SU(3)_c × SU(2)_L × U(1)_Y and can be viewed as the Minimal Supersymmetric Standard Model (MSSM) extended by a fourth lepton family and the TC sector which, due to supersymmetrization, is $\mathcal{N}=4$ Super Yang Mills (4SYM) theory. The non-MSSM superfields appearing in MSCT and their quantum numbers are given in Table I.

The renormalizable lepton and baryon number conserving superpotential for MSCT is

$$P = P_{\text{MSSM}} + P_{\text{TC}}, \quad (1)$$

where P_{MSSM} is the MSSM superpotential, and

$$P_{\text{TC}} = -\frac{g_{\text{TC}}}{\sqrt{2}} \epsilon^{abc} \Phi_L^a \cdot \Phi_L^b \Phi_3^c + y_U \Phi_L^a \cdot H_u \Phi_3^a + y_N \Lambda_L \cdot H_u N + y_E \Lambda_L \cdot H_d E + y_R E \Phi_3^a \Phi_3^a, \quad (2)$$

with H_u (H_d) denoting the $Y = +1/2$ ($-1/2$) Higgs superfield. The dot (\cdot) indicates a contraction between the $SU(2)_L$ doublets with the antisymmetric two-index Levi-Civita tensor ϵ .

To the Lagrangian corresponding to the superpotential in Eq. (1) we add the soft SUSY breaking terms of the MSSM as well as those corresponding to P_{TC} , with the latter expressed by:

$$\begin{aligned} \mathcal{L}_{soft}^{TC} = & - \left[a_{TC} \epsilon^{abc} \hat{U}_L^a \hat{D}_L^b \hat{U}_R^{*c} + a_U \hat{Q}_L^a \cdot \hat{H}_u \hat{U}_R^{*a} + a_N \hat{\Lambda}_L \cdot \hat{H}_u \hat{N}_R^* \right. \\ & + a_E \hat{\Lambda}_L \cdot \hat{H}_d \hat{E}_R^* + a_R \hat{U}_R^{*a} \hat{U}_R^{*a} \hat{E}_R^* + \frac{1}{2} M_D D_R^{\dagger a} D_R^{\dagger a} + c.c. \left. \right] - M_Q^2 \hat{Q}_L^{\dagger a} \hat{Q}_L^a \\ & - M_U^2 \hat{U}_R^{*a} \hat{U}_R^a - M_L^2 \hat{L}_L^\dagger \hat{L}_L - M_N^2 \hat{N}_R^* \hat{N}_R - M_E^2 \hat{E}_R^* \hat{E}_R. \end{aligned} \quad (3)$$

We denote the scalar components of a chiral supermultiplet with a hat.

This model constitutes our fundamental description in terms of the elementary degrees of freedom and forces. The relevant scales of the problem are the SUSY breaking scale, m_{SUSY} , and the EW scale that we identify with the low-energy strongly coupled regime of the TC theory $\Lambda_{TC} \sim 4\pi v_w$, which for $v_w = 246$ GeV implies $\Lambda_{TC} \sim 3$ TeV. We will assume here that the two scales satisfy

$$m_{SUSY} > \Lambda_{TC}. \quad (4)$$

With this ordering the EW symmetry is broken dynamically. Furthermore, we arrange the spectrum in the following way:

- 1) All SUSY breaking masses are taken to be of the order of m_{SUSY} .
- 2) We will argue that in our model the μ parameter, which gives the mass of the Higgsinos, is much larger than Λ_{TC} . Therefore the Higgsinos are ignored in the low energy phenomenology.
- 3) The remaining composite states acquire a mass, which at most is of the order of Λ_{TC} .

After the identification of the relevant scales and their ordering we now start systematically deriving the effective Lagrangian description towards the electroweak scale.

III. MESOSCOPIC LAGRANGIAN

In this section we derive the full Lagrangian of 3MWT, including the four fermion interactions, by integrating out all elementary scalars, all gauginos, and all higgsinos. The key

point of this model is that the technigaugino D_R^\dagger is removed from low energy dynamics, and this modifies the initial technicolor theory, MWT, strongly. Now the particle content of the TC theory is

$$Q_L^a = \begin{pmatrix} U_L^a \\ D_L^a \end{pmatrix}, \quad U_R^a, \quad a = 1, 2, 3, \quad (5)$$

where, in the Weyl basis, U_L and D_L are the left-handed techniup and technidown, respectively, and U_R is the right-handed up-type techniparticle. These are the fermionic states of the supermultiplets denoted by Φ_L and Φ_3 in Section II.

To determine the 3MWT Lagrangian, valid between the TeV-scale and m_{SUSY} , we now consider the Higgs Yukawa couplings in MSCT:

$$\begin{aligned} -\mathcal{L}_{\text{Yukawa}}^{\text{MSCT}} &= \hat{H}_u \cdot F_u + \hat{H}_d \cdot F_d + \text{h.c.}, \\ F_u &= q_{Lu}^i Y_u^i u_R^\dagger + y_U Q_L U_R^\dagger + y_N L_L N_R^\dagger, \\ F_d &= q_{Ld}^i Y_d^i d_R^\dagger + l_L^i Y_l^i e_R^\dagger + y_E L_L E_R^\dagger. \end{aligned} \quad (6)$$

The fields \hat{H}_u and \hat{H}_d are the two Higgs scalar doublets. The flavor index is denoted by $i = 1\dots 3$ and it is summed over. The matrices Y_u , Y_d , and Y_l are diagonal, and the CKM matrix V is hidden in the definitions of the vectors

$$q_{Lu}^{Ti} = (u_L^i, V^{ij} d_L^j) \quad \text{and} \quad q_{Ld}^{Ti} = (V^{\dagger ij} u_L^j, d_L^i). \quad (7)$$

The potential of the MSSM Higgs fields is

$$V_{\text{MSSM}} = (m_{\text{SUSY}}^2 + |\mu|^2) |\hat{H}_u|^2 + (m_{\text{SUSY}}^2 + |\mu|^2) |\hat{H}_d|^2 - (b \hat{H}_u \hat{H}_d + \text{h.c.}) + \dots \quad (8)$$

The Higgs states are diagonalized via:

$$\begin{pmatrix} \hat{H}_u \\ \hat{H}_d^c \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \hat{H}_1 \\ \hat{H}_2 \end{pmatrix}, \quad (9)$$

where $\hat{H}_d^c = i\sigma^2 \hat{H}_d^*$, with σ^2 being a Pauli matrix. The tree-level masses $m_{1,2}^2 = \mu^2 + m_{\text{SUSY}}^2 \pm b$ of $\hat{H}_{1,2}$ are traded for two convenient parameters θ and m_s defined via the following relations:

$$\frac{1}{2} \left(\frac{1}{m_1^2} + \frac{1}{m_2^2} \right) = \frac{c_\theta^2}{m_s^2}, \quad \frac{1}{2} \left(\frac{1}{m_1^2} - \frac{1}{m_2^2} \right) = \frac{c_\theta s_\theta}{m_s^2}. \quad (10)$$

In terms of the parameters appearing in Eq. (8) we have

$$m_s^2 = (\mu^2 + m_{\text{SUSY}}^2) \frac{(\mu^2 + m_{\text{SUSY}}^2)^2 - b^2}{(\mu^2 + m_{\text{SUSY}}^2)^2 + b^2}, \quad t_\theta = \frac{b}{\mu^2 + m_{\text{SUSY}}^2}. \quad (11)$$

In the above equations we have denoted $\sin \theta$, $\cos \theta$, $\tan \theta$ simply with s_θ , c_θ , t_θ . Decoupling all the scalars, as well as the gauginos and higgsinos, leads to the following intermediate scale (or mesoscopic) interaction Lagrangian for the fermions:

$$\begin{aligned} \mathcal{L}_{4-fermion} = & \frac{c_\theta^2}{m_s^2} \left(F_u^\dagger F_u + F_d^\dagger F_d \right) - \frac{c_\theta s_\theta}{m_s^2} (F_u \cdot F_d + \text{h.c.}) \\ & + \frac{g_{\text{TC}}^2}{m_{\text{SUSY}}^2} \epsilon_{abc} \epsilon_{cde} \eta_i^{\alpha a} \eta_{j\alpha}^b \eta_{i\dot{\beta}}^{\dagger d} \eta_j^{\dagger \dot{\beta} e} , \end{aligned} \quad (12)$$

with

$$\eta_\alpha^T = \left(U_{L\alpha}, D_{L\alpha}, -i\sigma_{\alpha\beta}^2 U_R^{\dagger\beta} \right) . \quad (13)$$

We have included the operators of mass dimension less or equal to six. The indices i and j denote SU(3) flavor; the first letters of the alphabet are reserved for the adjoint SU(2) technicolor indices, while the Greek indices label the spin component. The color indices are contracted and suppressed, while the TC indices, running from 1 to 3, are written explicitly only in the last term. We suppress summed spin indices as long as it can be done consistently.

The first line in Eq. (12) derives from decoupling the Higgs scalars, and breaks the global SU(3) symmetry, while the last term originates from the Yukawa interaction between techniquarks and techni-quarks. This interaction stems from the superpotential and gauge interactions of the $\mathcal{N} = 4$ sector. Because of its origin, the last term in Eq.(12) respects the global SU(3) symmetry of the pure TC sector.

IV. EFFECTIVE LAGRANGIAN AT THE ELECTROWEAK SCALE

In this section we derive the effective Lagrangian at scales below Λ_{TC} . The relevant degrees of freedom are the composite states associated to the strong TC interaction, and the form of the effective Lagrangian is constrained by requiring the effective theory to satisfy the approximate global symmetries of the fundamental Lagrangian. In addition to the composite scalar fields, we introduce also composite vector fields in a consistent manner.

The effective theory at the electroweak scale can be seen as composed of two parts:

$$\mathcal{L}_{eff}(M, K, K') = \mathcal{L}_{3\text{MWT}}(M) + \mathcal{L}_F(M, K, K') , \quad (14)$$

where M is the composite scalar matrix. The pure technicolor theory, corresponding to the particle content in Eq. (5) and the effective Lagrangian $\mathcal{L}_{3\text{MWT}}$, respects the full SU(3)

symmetry. The explicit breaking of this global symmetry due to the flavor extension is represented by the effective Lagrangian \mathcal{L}_F containing the spurions K and K' , which are defined in Subsection IV B. We will now define these effective Lagrangian contributions in detail.

A. Technicolor Scalar Sector

The composite scalar matrix field M , singlet under $SU(2)_{\text{TC}}$, has minimal particle content given by the techniquark bilinears:

$$M_{ij} \sim \eta_i^\alpha \eta_j^\beta \varepsilon_{\alpha\beta} = \eta_i \eta_j, \quad \text{with} \quad i, j = 1 \dots 3. \quad (15)$$

The field M transforms under the full $SU(3)$ group according to

$$M \rightarrow \mathbf{u} M \mathbf{u}^T, \quad \text{with} \quad \mathbf{u} \in SU(3). \quad (16)$$

The effective linearly transforming $SU(3)$ invariant Lagrangian reads:

$$\mathcal{L}_M = \frac{1}{2} \text{Tr} [D_\mu M^\dagger D^\mu M] - \mathcal{V}_M, \quad (17)$$

where the covariant derivative is given by

$$D_\mu M = \partial_\mu M - i g_L [G_\mu M + M G_\mu^T],$$

with

$$G_\mu = \tilde{W}_\mu^a \frac{\lambda^a}{2} + t_\xi B_\mu Y_M, \quad a = 1, 2, 3. \quad (18)$$

In the above equation λ^a are the Gell-Mann matrices, $t_\xi = \tan \xi$ with ξ the EW mixing angle, \tilde{W}_μ and B_μ are the SM EW gauge fields, and

$$Y_M = \text{diag} \left(\frac{1}{2}, \frac{1}{2}, -1 \right). \quad (19)$$

The most general $SU(3)$ preserving effective potential, including only up to dimension four operators, is

$$\mathcal{V}_M = -\frac{m^2}{2} \text{Tr} [M^\dagger M] + \frac{\lambda}{4} \text{Tr} [M^\dagger M]^2 + \lambda' \text{Tr} [M^\dagger M M^\dagger M] - 2m' [\det M + \det M^\dagger], \quad (20)$$

which breaks $SU(3)$ spontaneously to $SO(3)$ for positive m^2 , as we show explicitly in Appendix A. The TC gauge interaction is actually invariant under $U(3) \equiv SU(3) \times U(1)_A$, rather

than just $SU(3)$. However the $U(1)_A$ axial symmetry is anomalous, and is therefore broken at the quantum level. This corresponds to the $\det M$ term in Eq. (20).

The components of the matrix $M \sim \eta^T \eta$ can be described in terms of the transformation properties of the composite states under $SU(2)_L \times U(1)_Y$. This notation is introduced in Table II. Using this notation, the matrix M is written in terms of complex scalars as

Field	$SU(2)_L$	$U(1)_Y$
$\Delta \sim Q_L Q_L$	$\square\square$	1
$\sigma \sim Q_L U_R^\dagger$	\square	$-\frac{1}{2}$
$\delta^{--} \sim U_R^\dagger U_R^\dagger$	1	-2

TABLE II: Transformation properties of the component fields of the matrix M under $SU(2)_L \times U(1)_Y$. The complex scalars are grouped, based on their transformation properties under $SU(2)_L$, into one triplet, one doublet, and one singlet.

$$M = \begin{pmatrix} \sqrt{2}\Delta^{++} & \Delta^+ & \sigma^0 \\ \Delta^+ & \sqrt{2}\Delta^0 & \sigma^- \\ \sigma^0 & \sigma^- & \sqrt{2}\delta^{--} \end{pmatrix}. \quad (21)$$

This notation is suitable to study the vacuum, since the flavor extension sector breaks the global symmetry of the potential from $SU(3)$ down to $SU(2) \times U(1)$, (i.e. the EW gauge group). Next, we determine the effective Lagrangian terms of the flavor extension of 3MWT below scale Λ_{TC} .

B. Flavor Extension Sector

The four-fermion theory, Eq. (12), is given just below the SUSY breaking scale and the techniquark condensate needs to be evolved down to the EW scale. This is achieved by multiplying the techniquark Yukawa coupling y_U , renormalized at the SUSY breaking scale, with the dimensionless factor

$$\omega = \frac{\langle U_L U_R^\dagger \rangle_{m_{SUSY}}}{\langle U_L U_R^\dagger \rangle_{\Lambda_{TC}}} = \left(\frac{m_{SUSY}}{\Lambda_{TC}} \right)^\gamma, \quad (22)$$

written under the assumption that the anomalous dimension γ of the techniquark mass operator is constant.

Note that in the following we neglect the contribution of the last term in Eq. (12) because that term respects the global SU(3) symmetry, and therefore its effects should already be parametrized by the quartic couplings in the TC effective Lagrangian, Eq. (20). The masses of the SM fermions and the fourth family leptons arise from the terms on the first line of Eq. (12): more specifically those masses are generated by the following four-fermion operator

$$\eta^T K \eta , \quad (23)$$

with

$$K_{ij} = \frac{y_U c_\theta \omega}{m_s^2} \left[\delta_{ik} c_\theta \left(q_{Lu}^{\dagger k} Y_u^* u_R + y_N^* L_L^{\dagger k} N_R \right) - \epsilon_{ik} s_\theta \left(q_{Ld}^k Y_d d_R^\dagger + l_L^k Y_l e_R^\dagger + y_E L_L^k E_R^\dagger \right) \right] \delta_{3j},$$

$$i, j = 1, \dots, 3; \quad k = 1, 2; \quad \epsilon_{3k} \equiv 0 , \quad (24)$$

upon condensation of the techniquarks. Under SU(3) global symmetry the spurion K transforms as $K \rightarrow u^* K u^\dagger$.

The four-techniquark term on the other hand is

$$\frac{y_U^2 c_\theta^2}{m_s^2} \omega^2 (Q_L U_R^\dagger)(Q_L^\dagger U_R) = K'_{ijkl} \eta_i^\alpha \eta_{j\alpha} \eta_{k\beta}^\dagger \eta_l^{\dagger\beta} , \quad K'_{ijkl} = \frac{y_U^2 c_\theta^2}{m_s^2} \omega^2 (\delta_{ik1} + \delta_{ik2}) \delta_{jl3}, \quad (25)$$

where α and β are spin indices. For this term to be invariant under SU(3), the spurion K' must transform as $K'_{ijkl} \rightarrow u_{im} u_{jn} u_{ko}^* u_{lp}^* K'_{mnop}$, with $u \in \text{SU}(3)$. To estimate the effects of renormalization, we simply assumed factorization, leading to a multiplicative factor of ω^2 .

At the lowest order in the spurions, the SU(3) breaking effective Lagrangian, obtained from Eqs. (24,25) is:

$$\mathcal{L}_F = c_1 \Lambda_{\text{TC}}^2 \text{Tr} [MK] + c_2 \Lambda_{\text{TC}}^4 K'_{ijkl} M_{ij} M_{kl}^* + \text{h.c.} . \quad (26)$$

The factors of Λ_{TC} have been added to make dimensionless the coefficients c_i , which parametrize the uncertainty in the couplings of the effective Lagrangian in terms of those of the underlying theory. We estimate these coefficients using dimensional analysis [34–36] and find

$$c_1 = \mathcal{O}(\Upsilon^{-1}) , \quad c_2 = \mathcal{O}(\Upsilon^{-2}) , \quad \Upsilon \equiv \frac{\Lambda_{\text{TC}}}{v_w} . \quad (27)$$

Finally, we note that, due to the last term in Eq. (26), the global SU(3) symmetry breaking pattern of the TC potential is altered compared to the pure TC scenario presented in Appendix A. We will study the ground state of the full potential in Section V A. To conclude this section, we next discuss how to introduce also composite vector fields consistently.

C. Vector Sector

Similarly to QCD, in TC a tower of composite states is predicted to arise at low energies. The lightest states of the spectrum are constituted by the scalar composite states, already introduced in Subsection IV A, and the vector resonances. A minimal set of composite vector fields transforming homogeneously under SU(3) can be written, in terms of Gell-Mann matrices λ^a , as

$$A_\mu = A_\mu^a \frac{\lambda^a}{2}, \quad (28)$$

which transform under SU(3) according to

$$A_\mu \rightarrow u A_\mu u^\dagger, \quad \text{with} \quad u \in \text{SU}(3). \quad (29)$$

The elementary particle content of A^μ is expressed by the equivalence

$$A_i^{\mu j} \sim \eta_i^\alpha \sigma_{\alpha\dot{\beta}}^\mu \eta^{\dagger\dot{\beta}j} - \frac{1}{3} \delta_i^j \eta_k^\alpha \sigma_{\alpha\dot{\beta}}^\mu \eta^{\dagger\dot{\beta}k} = \bar{\Psi}^j \gamma^\mu \Psi_i - \frac{1}{3} \delta_i^j \bar{\Psi}^k \gamma^\mu \Psi_k, \quad \bar{\Psi} = (\bar{U}_L, \bar{D}_L, \bar{U}_R^c), \quad (30)$$

where the components of Ψ in SU(3) space are Dirac spinors, with the superscript c on the last entry denoting the charge conjugation. The vector and axial-vector charge eigenstates and their elementary particle content are given in Appendix (B).

The effective Lagrangian including composite vector fields, A^μ , has already been derived in [31] by applying the hidden local symmetry principle [37, 38] for MWT, which features an SU(4) global symmetry in the TC sector. Those results can be straightforwardly used for SU(3) symmetric TC by defining the corresponding vector field and the field strength tensor:

$$C_\mu = A_\mu - \epsilon G_\mu, \quad \epsilon = \frac{g_L}{g_{\text{TC}}}, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i g_{\text{TC}} [A_\mu, A_\nu], \quad (31)$$

with G_μ defined in Eq. (18). The vector field C_μ has the same transformation law as A_μ :

$$C_\mu \rightarrow u C_\mu u^\dagger, \quad \text{with} \quad u \in \text{SU}(3). \quad (32)$$

The kinetic and mass terms for the vector fields can then be written as:

$$\mathcal{L}_k \supset -\frac{1}{2} \text{Tr} [\tilde{W}_{\mu\nu} \tilde{W}^{\mu\nu}] - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - \frac{1}{2} \text{Tr} [F_{\mu\nu} F^{\mu\nu}] + m_A^2 \text{Tr} [C_\mu C^\mu], \quad (33)$$

while the scalar-vector field interaction terms up to dimension four operators read:

$$\mathcal{L}_{M-V} = g_{\text{TC}}^2 r_1 \text{Tr} [C_\mu C^\mu M M^\dagger] + g_{\text{TC}}^2 r_2 \text{Tr} [C_\mu M C^{\mu T} M^\dagger] - g_{\text{TC}}^2 \frac{r_3}{4} \text{Tr} [C_\mu C^\mu] \text{Tr} [M M^\dagger], \quad (34)$$

with the r_i constants being of $\mathcal{O}(1)$. A few remarks are in order: First, higher dimensional operators are suppressed by positive powers of Λ_{TC} , and therefore are subleading. Second, terms proportional to y_U , which explicitly break $\text{SU}(3)$ global symmetry, are small compared to those proportional to g_{TC}^2 and therefore negligible at leading order. Third, to simplify the phenomenological analysis of 3MWT, presented in the next section, we neglected also a covariant derivative coupling term (see [31] for its precise definition).¹

V. PHENOMENOLOGY

In this section we determine the physical states arising at low energy in 3MWT, with particular focus on the light composite Higgs boson, and then study their contributions to the EW precision parameters S and T .

A. EW Symmetry Breaking & Mass Eigenstates

The most general ground state which breaks the EW gauge group down to the electromagnetic $\text{U}(1)_Q$ can be parametrized by the following form of the vacuum expectation value (vev) of the matrix field M

$$\langle M \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & v_\sigma \\ 0 & \sqrt{2}v_\Delta & 0 \\ v_\sigma & 0 & 0 \end{pmatrix}. \quad (35)$$

Minimizing the scalar potential given in Eqs. (20,26) leads to the following relations between the parameters:

$$m^2 = \lambda (v_\sigma^2 + v_\Delta^2) + 4\lambda'v_\Delta^2 - 2\lambda''v_\sigma^2, \quad \tilde{m}^2 = 2(\lambda' + \lambda'')(v_\sigma^2 - 2v_\Delta^2), \quad (36)$$

where we have denoted

$$\tilde{m}^2 = c_2 y_U^2 c_\theta^2 \frac{\Lambda_{\text{TC}}^4 \omega^2}{m_s^2}, \quad \lambda'' \equiv -\frac{m'}{v_\Delta}. \quad (37)$$

A sufficient condition for the potential to be bounded from below is $\lambda, \lambda' > 0$. In the limit $v_\sigma = \sqrt{2}v_\Delta$ we obtain $\tilde{m} = 0$, consistently with the pure TC result in Appendix A.

¹ Neglecting this term is simply a restriction on the parameter space: this term could be included in more thorough future analyses.

We notice that the ground state changes because of the four-technifermion interaction in Eq. (25). An analogous change of ground state occurs in ETC theories where some of the chiral symmetries of the pure TC theory are broken by extended gauge interactions. In our model setup the effective four fermion interactions at low energy arise from attractive Yukawa couplings which is different from the usual ETC scenario where the underlying gauge interactions can be either repulsive or attractive.

The mass spectrum of 3MWT, corresponding to the vev in Eq. (35), includes two neutral scalars, h^0 and H^0 , as well as one neutral pseudoscalar, Π^0 , one charged and two doubly charged scalars, H^\pm , $h^{\pm\pm}$, and $H^{\pm\pm}$, respectively, with the corresponding squared mass matrices given in Appendix C. We introduce for later use the mixing angle φ defined by

$$\begin{pmatrix} h^0 \\ H^0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} c_\varphi & -s_\varphi \\ s_\varphi & c_\varphi \end{pmatrix} \begin{pmatrix} \Re(\sigma^0) \\ \Re(\Delta^0) \end{pmatrix}. \quad (38)$$

Assuming that only the third generation SM fermions have non-negligible Yukawa couplings, it follows from Eq. (26) that the masses of the upper component u and the lower component d of a generic SM fermion EW doublet are given, respectively, by

$$m_u = c_1 \frac{c_\theta^2 y_U y_u \omega \Lambda_{\text{TC}}^2}{m_s^2} \frac{v_\sigma}{\sqrt{2}}, \quad m_d = \frac{y_d}{y_u} t_\theta m_u. \quad (39)$$

Using the previous equation, the fact that both E and N have to be heavier than about 100 GeV, and requiring the Yukawa couplings to be perturbative, we deduce that θ cannot be close to either 0 or $\pi/2$. To simplify the study of 3MWT from here on we take $m_s = m_{\text{SUSY}}$, which from Eq. (11) is equivalent to imposing:

$$m_{\text{SUSY}} = m_s \quad \Rightarrow \quad \frac{b}{\mu^2} = \frac{t_\theta + t_\theta^{-1}}{2}. \quad (40)$$

From the fact that neither t_θ nor its inverse are large, it follows that our choice does not introduce any large hierarchy between b and μ^2 , which can be taken both of $\mathcal{O}(m_s^2)$. This implies in particular that the higgsinos are heavy and decoupled from low energy phenomenology.

Finally, the composite axial-vector and vector resonances mix with the SM gauge bosons, while the doubly charged baryon technivector does not. The resulting physical massive vector states are $Z_\mu, Z'_\mu, Z''_\mu, W_\mu^\pm, W''_\mu^\pm, W'''_\mu^\pm, \Omega_\mu^{\pm\pm}$, with the corresponding squared mass matrices given in Appendix C. The masses of W^\pm and Z in the limit of negligible mixing ($\epsilon = 0$) read:

$$m_W^2 \cong \frac{1}{4} g_L^2 v_w^2, \quad m_Z^2 \cong \frac{1}{4} (g_L^2 + g_Y^2) (1 + t_\rho^2) v_w^2, \quad (41)$$

where $t_\rho = \sqrt{2}v_\Delta/v_\sigma$, and the EW scale is given by

$$v_w^2 = (\sqrt{2}G_F)^{-1} = (246 \text{ GeV})^2 = v_\sigma^2 + 2v_\Delta^2 . \quad (42)$$

B. Higgs Mass

The light Higgs deserves some further discussion. In [16] it has been shown that $m_{h^0}^2$ receives a large negative corrections at one loop from the top quark and the W^\pm and Z bosons. Because of these corrections the tree level Higgs mass can be significantly larger than 125 GeV: Adapting the formulas given in [16] to the 3MWT case, while neglecting in first approximation any mixing between the two neutral scalar states, we can write

$$(m_{h^0}^2)_{\text{tree}} \simeq m_{h^0}^2 + \frac{8}{3}\kappa^2 \left[2a_f^2 (3m_t^2 + m_E^2 + m_N^2) - 3a_\pi^2 \left(m_W^2 + \frac{m_Z^2}{2} \right) \right] , \quad (43)$$

where $(m_{h^0}^2)_{\text{tree}}$ is the scalar mass due to pure strong dynamics before coupling with the EW gauge currents and SM matter. Furthermore, κ is an $\mathcal{O}(1)$ renormalization coefficient, while a_f and a_π are rescaling coefficients of the SM Higgs linear coupling to fermions and quadratic coupling to gauge bosons, respectively. In writing the formula above we set the dimension of the TC representation to 3 and the number of technidoublets to 0.75, as it is the case for 3MWT. To illustrate the idea we evaluate the formula above by assuming couplings and renormalization coefficient to be SM like ($a_F = a_\pi = \kappa = 1$), and $m_E = m_N = m_t$. With these choices we find a light Higgs mass equal to 740 GeV. This value is not far from the naive TC estimate for 3MWT, obtained by scaling up the mass of the $f_0(500)$ QCD resonance [39] ($m_{f_0}=400\text{-}550$ GeV):

$$(m_{h^0}^2)_{\text{naive}} \simeq \frac{4}{3} \frac{v_w^2}{f_\pi^2} m_{f_0}^2 = 1200\text{-}1700 \text{ GeV} . \quad (44)$$

The required suppression of the tree level Higgs mass, around 50%, might realistically come from near-conformal dynamics. This should be indeed the case for 3MWT, which has 1.5 adjoint Dirac flavors and is outside but close to the conformal window [40]. Lattice results moreover suggest that for 2 Dirac flavors the lower bound of the conformal window extends below the analytic estimates based on two loop perturbative beta function and ladder approximation of the Schwinger-Dyson equation for the fermion propagator. Therefore the suppression of the tree level Higgs mass in our theory could be more pronounced than what perturbative estimates might suggest.

Taking into account one loop corrections to the mixing of the two neutral scalar states would require us to introduce the one loop effective potential: this is beyond the task of our study, which is to perform an initial investigation of the viability of 3MWT. For this reason we simply take the tree level light Higgs mass to be 125 GeV, by assuming that this value incorporates not only the strong dynamics contribution but receives also non-leading corrections from the global SU(3) breaking sector of 3MWT.

C. Oblique Corrections

The precision EW parameters [32, 33] can be calculated directly from the vector-boson sector of the effective Lagrangian, Eqs. (33,34), by integrating out the heavy charged and neutral states and then using the formulas provided by [41, 42].² At tree level and linear order in the mixing parameter ϵ we find:

$$S_{tree} = 0, \quad \alpha_e T_{tree} = -\frac{2v_\Delta^2}{v_w^2}. \quad (45)$$

For the T -parameter to be consistent with the experiments, the vev component v_Δ clearly has to be small relatively to the EW scale. We note that the S parameter is zero up to corrections of order ϵ^4 while the T parameter obtains further contributions of order ϵ^2 which we neglect since $\epsilon \ll 1$.

The intrinsic TC contribution is usually calculated from the one loop perturbative diagrams of the technifermions, which are assumed to be massive because of dynamical symmetry breaking. The dynamical mass divided by m_Z is usually taken infinite, as this gives a meaningful result with no unknown parameters. These contributions are denoted by S_{naive} and T_{naive} . For our underlying technicolor theory, Eq. (5), the dynamical masses are such that while the up-techniquark, U , gains only a Dirac mass, m_U , the down-techniquark, D_L , acquires also a Majorana mass, m_L . Oblique corrections for this general case have been calculated in [43] in terms of integral functions, which we used to derive the explicit formulas given in Appendix D. In the pure 3MWT limit the Dirac and Majorana masses are equal to

² For this task we adapted the code provided by the authors of [31] to 3MWT.

each other³, $m_L = m_U$, and we find

$$S_{naive} = \frac{3}{4\pi} \quad , \quad T_{naive} = \frac{m_U^2 \log[\frac{m_U}{\Lambda_{NP}}]}{4m_Z^2 s_\xi^2 c_\xi^2 \pi} . \quad (46)$$

The dependence of T on the renormalization scale Λ_{NP} is due to the Majorana mass, and it should be matched onto a renormalizable term in the underlying theory. For the phenomenological purposes of our current analysis, we assume that the renormalization scale Λ_{NP} is close to m_U making the one-loop contribution T_{naive} negligible when compared to the tree-level contribution, T_{tree} .⁴ With this assumption we use the following values of the naive parameters:

$$S_{naive} = \frac{3}{4\pi} \quad , \quad T_{naive} = 0 . \quad (47)$$

Another independent contribution to S and T in our model comes from the fourth family leptons N and E . We denote these contributions by $S_{N,E}$ and $T_{N,E}$. These have been already evaluated in [44]: for $m_E > m_N$, the $S_{N,E}$ contribution is actually negative and can offset the positive S_{naive} . We can therefore summarize the non-negligible contributions to the S and T parameters, with the assumptions described above, as:

$$S = S_{naive} + S_{N,E} \quad , \quad T = T_{tree} + T_{N,E} , \quad (48)$$

where T_{tree} , S_{naive} , and the Higgs contributions are given in Eqs. (45,47).

VI. EXPERIMENTAL VALIDATION

In this section we study the viability of 3MWT by performing a goodness of fit analysis based on the recent LHC and Tevatron data on Higgs physics, as well as the experimental values of the S and T parameters. In Subsection VIA we derive the linear couplings of the light Higgs and relate these to the corresponding SM coupling strengths. In Subsection VIB we implement the direct search lower bounds on the new physics mass spectrum by performing a numerical scan over the parameter space and collecting data points that satisfy these constraints and the EW precisions tests. Finally, in Subsection VIC we define the observables included in the goodness of fit analysis and then present the statistical results on the viability of 3MWT.

³ This follows from the invariance of the mass terms implied by Eq. (21) under SO(3).

⁴ A generic renormalization scale still allows viability since T_{naive} can be canceled by T_{tree} , though this requires fine tuning of v_Δ .

A. Coupling Coefficients

In our model the linear Higgs coupling to charged vector bosons can be written in compact form as

$$\mathcal{L} \supset \frac{2m_A^2}{v_w} \bar{W}_\mu^\dagger \Xi \bar{W}^\mu h^0, \quad \bar{W}_\mu^\dagger = \left(\tilde{W}_\mu^-, V_\mu^-, A_\mu^-, \Omega_\mu^{--} \right), \quad (49)$$

with the non-zero terms of the matrix Ξ given by

$$\begin{aligned} \Xi_{1,1} &= (x^2 + \epsilon^2 z_1) \left(c_\varphi c_\rho - \sqrt{2} s_\varphi s_\rho \right) + \frac{\epsilon^2 z_3}{\sqrt{2}} \left(s_\varphi s_\rho - \sqrt{2} c_\varphi c_\rho \right), \\ \Xi_{2,2} &= z_1 \left(c_\varphi c_\rho - \sqrt{2} s_\varphi s_\rho \right) + z_2 \left(c_\varphi s_\rho - \sqrt{2} c_\rho s_\varphi \right) + \frac{z_3}{\sqrt{2}} \left(s_\varphi s_\rho - \sqrt{2} c_\varphi c_\rho \right), \\ \Xi_{3,3} &= z_1 \left(c_\varphi c_\rho - \sqrt{2} s_\varphi s_\rho \right) - z_2 \left(c_\varphi s_\rho - \sqrt{2} c_\rho s_\varphi \right) + \frac{z_3}{\sqrt{2}} \left(s_\varphi s_\rho - \sqrt{2} c_\varphi c_\rho \right), \\ \Xi_{1,2} &= -\frac{\epsilon}{2\sqrt{2}} \Xi_{2,2}, \quad \Xi_{1,3} = -\frac{\epsilon}{2\sqrt{2}} \Xi_{3,3}, \\ \Xi_{4,4} &= 2c_\varphi c_\rho (z_1 + z_2) + \frac{z_3}{\sqrt{2}} \left(s_\varphi s_\rho - \sqrt{2} c_\varphi c_\rho \right). \end{aligned} \quad (50)$$

Here φ is the mixing angle of the neutral scalars, Eq. (38), and

$$\epsilon = \frac{g_L}{g_{TC}}, \quad x = \frac{g_L v_w}{2m_A}, \quad t_\rho = \frac{\sqrt{2}v_\Delta}{v_\sigma}, \quad z_i = \left(\frac{g_{TC} v_w}{2m_A} \right)^2 r_i, \quad i = 1, 2, 3. \quad (51)$$

To simplify the analysis we select the slice of parameter space where the axial vector coupling to the light Higgs is zero ($\Xi_{1,3} = \Xi_{3,3} = 0$). Moreover, we require the mixing mass term of the axial-vector to arise only from mixing ($(\mathcal{M}_W^2)_{1,3} = -\epsilon m_A^2 / \sqrt{2}$), which is taken to be small. Together these conditions are satisfied by imposing

$$z_1 = \frac{1}{4} (3 + c_{2\rho}^{-1}) z_3, \quad z_2 = \frac{1}{4} t_{2\rho} z_3. \quad (52)$$

If we apply the above substitution together with

$$z_3 \rightarrow \frac{4s^2}{s_{2\rho} t_{2\rho}} \quad (53)$$

to the charged vector boson squared mass matrix in Eq. (C6), we reproduce the corresponding result quoted in [45]:

$$\mathcal{M}_W^2 = \begin{pmatrix} m_W^2 & -\frac{\epsilon m_V^2}{\sqrt{2}} & -\frac{\epsilon m_A^2}{\sqrt{2}} \\ -\frac{\epsilon m_V^2}{\sqrt{2}} & m_V^2 & 0 \\ -\frac{\epsilon m_A^2}{\sqrt{2}} & 0 & m_A^2 \end{pmatrix}, \quad (54)$$

with

$$m_{\bar{W}}^2 = [x^2 + (1 + s^2) \epsilon^2] m_A^2, \quad m_V^2 = (1 + 2s^2) m_A^2. \quad (55)$$

The mass eigenvalues can be expanded in x and ϵ , which in TC are both expected to be small:

$$\begin{aligned} m_W^2 &\cong m_A^2 x^2 [1 - \epsilon^2], \quad m_{W''}^2 \cong m_A^2 \left[1 + \frac{1}{2} (1 + x^2) \epsilon^2 - \frac{1}{8} \left(2 + \frac{1}{s^2} \right) \epsilon^4 \right], \\ m_{W'}^2 &\cong m_A^2 \left[1 + 2s^2 + \frac{1}{2} (1 + 2s^2 + x^2) \epsilon^2 + \frac{1}{8} \left(2 + \frac{1}{s^2} \right) \epsilon^4 \right], \end{aligned} \quad (56)$$

where we kept contributions up to $\mathcal{O}(x^n \epsilon^{4-n})$, with $n = 0, \dots, 4$.

Generally the linear couplings of the light Higgs can be conveniently expressed in terms of coupling coefficients defined by

$$\begin{aligned} \mathcal{L}_{\text{eff}} &= \sum_i a_{W_i} \frac{2m_{W_i}^2}{v_w} h W_{i\mu}^+ W_i^{-\mu} + \sum_j a_{Z_j} \frac{m_{Z_j}^2}{v_w} h Z_{j\mu} Z_j^\mu \\ &- a_f \sum_{\psi=t,b,\tau,N,E} \frac{m_\psi}{v_w} h \bar{\psi} \psi - \sum_k a_{S_k} \frac{2m_{S_k}^2}{v_w} h S_k^+ S_k^-, \end{aligned} \quad (57)$$

where the indices i and k run over all the charged scalars and vector bosons (including the states with double charge). By substituting the charged vector mass eigenstates obtained from Eq. (54) in Eq. (49), we find the linear Higgs couplings to the charged vector bosons. By normalizing these couplings according to Eq. (57), with the masses given by Eqs. (56,C7), we determine

$$\begin{aligned} a_W &= \left(c_\varphi c_\rho - \sqrt{2} s_\varphi s_\rho \right) \left[1 - \frac{x^2 \epsilon^2}{2} \left(1 + \frac{1}{1 + 2s^2} \right) \right] + \frac{x^2 \epsilon^2 s^2}{2(1 + 2s^2)^2} \left(\frac{c_\varphi}{c_\rho} - \sqrt{2} \frac{s_\varphi}{s_\rho} \right), \\ a_W + a_{W'} + a_{W''} &= c_\varphi c_\rho - \sqrt{2} s_\varphi s_\rho + \frac{s^2}{1 + 2s^2} \left(\frac{c_\varphi}{c_\rho} - \sqrt{2} \frac{s_\varphi}{s_\rho} \right), \\ a_\Omega &= \frac{s^2}{s_\rho} \frac{2c_\varphi (1 + t_\rho^{-1}) + \sqrt{2} c_{2\rho} c_\rho^{-2} s_\varphi}{2 - 4s^2 + s^2 (2(t_\rho^{-1} + s_\rho^{-2}) + c_\rho^{-2})}. \end{aligned} \quad (58)$$

The corresponding result for fermions reads simply

$$a_f = \frac{c_\varphi}{c_\rho}. \quad (59)$$

Among the neutral vector resonances only Z is relevant for the LHC observables we include in this study. As we discuss in the next section, the numerical values of a_Z that we find are very close to a_W , and for all practical purposes they can therefore be taken equal to each other.⁵

⁵ The analytic expressions for a_{Z_j} and a_{S_k} are lengthy and we do not reproduce them here.

B. Parameter Space Scan

We perform a scan of the parameter space to find data points that satisfy experimental constraints on the new physics mass spectrum and on the S and T parameters. We require perturbativity of all but the g_{TC} coupling and stability of the potential at large values of the scalar fields. The unknown parameters derived from strong dynamics are estimated using dimensional analysis [34–36]. Finally, we fix the anomalous dimension to $\gamma = 1.5$, and require the supersymmetry breaking scale to be larger than 5 TeV. Putting all this together, the free parameters in our scan acquire values in the domain defined by the following relations:

$$210 \text{ GeV} \leq |v_\sigma| \leq 246 \text{ GeV}, \quad m_A = 1 \text{ TeV}, \quad \gamma = 1.5, \quad \pi \leq \Upsilon \leq 4\pi, \quad 0.5 \leq c_1 \Upsilon^{-1} \leq 5, \\ 0.1 \leq \lambda \leq (2\pi)^2, \quad 0.1 \leq y_t, y_N, y_E, y_U \leq 2\pi, \quad 0.1 \leq \lambda'' \leq 200, \quad 0 \leq z_3 \leq 1, \quad |\epsilon| \leq 2x. \quad (60)$$

The remaining model parameters, $\lambda', \theta, x, v_\Delta$ are determined in terms of the ones above by using, respectively, Eqs. (C1,39,42,56) together with the observed masses of the Higgs boson, top quark, W boson, and the EW vev. The last relation above ensures that the physical W is mostly made of the $SU(2)_L$ gauge field. We scanned over the parameter space defined above and collected 1000 data points, each satisfying the following constraints:

$$m_{\text{SUSY}} > 5 \text{ TeV}, \quad m_{H^0} > 600 \text{ GeV}, \quad m_{H^\pm}, m_{h^{\pm\pm}}, m_{H^{\pm\pm}}, m_E, m_N \geq 100 \text{ GeV}, \quad (61)$$

as well as the experimental limits on S and T [39]. We also checked that the collected data points satisfy the ATLAS lower limit on the mass of a sequential W' boson [46], equal to 2.55 TeV, after an appropriate rescaling of the limit which takes into account the non-SM value of the W' coupling to fermions [45].

C. Goodness of Fit

We performed a goodness of fit analysis by using the observed Higgs decay rates to $\gamma\gamma, ZZ, WW, \tau\tau, bb, \gamma\gamma JJ$ at ATLAS [47–51] and CMS [52–55], and to $\gamma\gamma, WW$, and bb at Tevatron [56], as well as the S and T experimental values [39], for a total of 19 observables. The LHC and Tevatron results are expressed in terms of the signal strengths, defined as

$$\hat{\mu}_{ij} = \frac{\sigma_{\text{tot}} \text{Br}_{ij}}{\sigma_{\text{tot}}^{\text{SM}} \text{Br}_{ij}^{\text{SM}}}, \quad \sigma_{\text{tot}} = \sum_{\Phi'=h, q\bar{q}, \dots} \epsilon_{\Phi'} \sigma_{\Phi \rightarrow \Phi'}, \quad \Phi = pp, p\bar{p}, \quad (62)$$

where $\epsilon_{\Phi'}$ is the efficiency associated with the given final state Φ' in an exclusive search, while for inclusive searches one simply has $\sigma_{\text{tot}} = \sigma_{pp \rightarrow h^0(X)}$, the h^0 production total cross section.

The combined signal strengths from ATLAS, CMS⁶, and Tevatron are given in Table III, while the signal strengths and efficiencies⁷ for dijet associated $\gamma\gamma$ production at ATLAS and

ij	ATLAS	CMS	Tevatron
ZZ	1.50 ± 0.40	0.91 ± 0.27	
$\gamma\gamma$	1.65 ± 0.32	1.11 ± 0.31	6.20 ± 3.30
WW	1.01 ± 0.31	0.76 ± 0.21	0.89 ± 0.89
$\tau\tau$	0.70 ± 0.70	1.10 ± 0.40	
bb	-0.40 ± 1.10	1.30 ± 0.70	1.54 ± 0.77

TABLE III: Combined signal strengths from LHC and Tevatron experiments.

CMS are listed in Table IV.

	ATLAS 7TeV	ATLAS 8TeV	CMS 7TeV	CMS 8TeV
$\gamma\gamma JJ$	2.7 ± 1.9	2.8 ± 1.6	2.9 ± 1.9	0.3 ± 1.3
$pp \rightarrow h$	22.5%	45.0%	26.8%	46.8%
$pp \rightarrow qqh$	76.7%	54.1%	72.5%	51.1%
$pp \rightarrow t\bar{t}h$	0.6%	0.8%	0.6%	1.7%
$pp \rightarrow Vh$	0.1%	0.1%	0%	0.5%

TABLE IV: Signal strengths and efficiencies for Higgs decay to $\gamma\gamma$ associated to a dijet at LHC.

Finally, the observed values for the S and T parameters read [39]:

$$S = 0.04 \pm 0.09, \quad T = 0.07 \pm 0.08, \quad r(S, T) = 88\%, \quad (63)$$

with the last quantity defining the correlation of the two parameters.

For a detailed description of the present fit we refer the reader to [45], where the same statistical analysis has been performed for a different model. Given that no new physics

⁶ We use the mass cut based result for CMS result on the Higgs to diphoton decay.

⁷ We chose to include only the loose categories from the ATLAS and CMS dataset at 8 TeV.

has been detected, only the contributions of new charged particles at one loop to $\Gamma_{h\rightarrow\gamma\gamma}$ become relevant when comparing the 3MWT predictions to the data in Tables III, IV. More explicitly one has [57]

$$\Gamma_{h\rightarrow\gamma\gamma} = \frac{\alpha_e^2 m_h^3}{256\pi^3 v_w^2} \left| \sum_i N_i e_i^2 F_i \right|^2, \quad (64)$$

with i summed over all the charged particles, N_i is the number of colors, e_i the charge in electron units, and F_i a function of the mass m_i and the coupling coefficient defined in [45]. In the limit of new particles being much heavier than the light Higgs, one finds

$$F_{W_i} = 7a_{W_i}, \quad F_E = F_N = -a_f \frac{4}{3}, \quad F_{S_i} = -a_{S_i} \frac{1}{3}, \quad (65)$$

with the coupling coefficients defined by Eq. (57). We can therefore mimic the contribution of the charged non-SM particles in 3MWT to the observables in Tables III, IV by including only the new contribution of a heavy single charged vector boson with coupling coefficient $a_{V'}$ determined by

$$a_{V'} \equiv \frac{1}{7} (F_{W'} + F_{W''} + 4F_\Omega) - \frac{a_S}{21}, \quad a_S \equiv -3(16F_E + 4F_N + F_{H^\pm} + 4F_{h^{\pm\pm}} + 4F_{H^{\pm\pm}}), \quad (66)$$

where the factors of 4 account for the double charge of the corresponding states. Moreover to simplify the analysis consistently with [45] we redefine

$$a_Z \approx a_W \equiv a_V, \quad (67)$$

where the numerical deviations from the first approximate equality above are negligible for the collected data points compared to the uncertainties on the coupling coefficients, which we present at the end of this section. At each collected data point we determine the numerical values of a_f , a_V , and $a_{V'}$ by Eqs. (58,66,67), while we calculate numerically the coupling coefficients of the charged scalars. In Fig. 1 we plot the viable data points on the (a_V, a_f) (left panel) and $(a_V, a_{V'})$ (right panel) planes together with the 68% (green), 90% (blue), and 95% (yellow) confidence level (CL) regions. In both plots, the third parameter is fixed to the optimal value marked with a blue star. It is clear from Fig. 1, left panel, that the W and Z couplings are enhanced, compared to their SM values, while the SM fermion couplings are suppressed. This result for the 3MWT couplings enhances the Higgs decay to diphotons. However, the contribution of the new charged fermions and scalars, as can be seen from Eq. (66), is large and interferes destructively with the W contribution to the same




FIG. 1: Viable data points in the (a_V, a_f) (left panel) and $(a_V, a_{V'})$ (right panel) planes, together with the 68% (green), 90% (blue), and 95% (yellow) CL region. The blue star in each plot marks the optimal coupling coefficients on the respective planes.

process. The probability associated with the data point minimizing χ^2 in the (a_f, a_V, a_S) space, obtained from a 3MWT particle spectrum lacking composite vector resonances at low energy would be ruled out entirely:

$$\begin{aligned}
 a_V &= 1.00, & a_f &= 1.00, & a_S &= 20.5, & S &= 0.04, & T &= 0.07; \\
 \chi_{\min}^2/\text{d.o.f.} &= 3.42, & P(\chi^2 > \chi_{\min}^2) &= 0.0004\%, & \text{d.o.f.} &= 16.
 \end{aligned}
 \tag{68}$$

In calculating $\chi_{\min}^2/\text{d.o.f.}$ in the the above equations we assumed the model to allow for 3 free paramters (a_f, S, T) , since a_V is strongly correlated with a_f near χ_{\min}^2 and a_S is basically constant. Fortunately the goodness of the fit changes dramatically once we include composite vector resonances in the 3MWT low energy spectrum, as it can be guessed from Fig. 1, right panel, and from Fig. 2. The contribution of the new charged vector bosons, and especially that of the vector baryon Ω , to the Higgs decay into diphoton is large, and offsets entirely the negative contribution of E , N , and charged scalars in Eq. (66). Among the 1000 viable data points, the one producing the minimum value for χ^2 gives:

$$\begin{aligned}
 a_V &= 1.01, & a_f &= 0.99, & a_{V'} &= 0.21, & S &= 0.04, & T &= 0.07; \\
 \chi_{\min}^2/\text{d.o.f.} &= 0.83, & P(\chi^2 > \chi_{\min}^2) &= 65\%, & \text{d.o.f.} &= 15,
 \end{aligned}
 \tag{69}$$




FIG. 2: Viable data points in the $(a_{V'}, a_f)$ plane passing through the point with optimal coupling coefficients in the $(a_{V'}, a_f, a_V)$ space, marked by a blue star, together with the 68% (green), 90% (blue), and 95% (yellow) CL region.

where the number of degrees of freedom (d.o.f.) has decreased by one, since $a_{V'}$ is a free parameter. It is interesting to notice that the optimal value of $a_{V'}$ above is equal to the average $a_{V'}$, calculated over the 1000 data points, while the average values of a_f and a_V are, respectively, 0.98 and 1.03, which are very close to the corresponding optimal values given above. This shows that strong dynamics, that we used to determine the scanned range of values of the free parameters, generates rather naturally the coupling strengths favored by LHC data, at least once the direct constraints on the mass spectrum and the EW precision parameters are satisfied.

The 3MWT result in Eq. (69) should be compared to the SM one:

$$\chi_{\min}^2/\text{d.o.f.} = 0.89, \quad P(\chi^2 > \chi_{\min}^2) = 60\%, \quad \text{d.o.f.} = 19. \quad (70)$$

While the SM fit is less satisfactory than the 3MWT one, it clearly shows that the SM is still perfectly viable in light of present collider data. It is instructive to notice that the fit performed with completely free coupling coefficients, therefore not motivated by any specific

underlying theory, produces a worse fit than the 3MWT:

$$\begin{aligned}
 a_V &= 0.97_{-0.11}^{+0.10}, & a_f &= 1.02_{-0.32}^{+0.25}, & a_{V'} &= 0.21_{-0.18}^{+0.16}, \\
 \chi_{\min}^2/\text{d.o.f.} &= 0.85, & P(\chi^2 > \chi_{\min}^2) &= 62\%, & \text{d.o.f.} &= 14.
 \end{aligned}
 \tag{71}$$

This is because the underlying strong dynamics introduces a large correlation between a_f and a_V , hence increasing the number of d.o.f by one, while achieving a χ_{\min}^2 very close to the corresponding result obtained with free coupling coefficients.

VII. CONCLUSIONS

We have constructed a supersymmetric TC model and studied its low energy phenomenology in detail. We started from the ultraviolet complete model obtained by coupling Minimal Walking Technicolor and the SM with two Higgs doublets and supersymmetrizing the entire theory. We then considered a situation where the supersymmetry breaking occurs at scales much higher than the electroweak scale, leading to decoupling of all elementary scalars, gauginos, and higgsinos. What remains at low energies is a TC theory with global $SU(3)$ chiral symmetry and its coupling with the SM fermions. We labeled this model 3MWT.

We constructed the low energy effective Lagrangian of 3MWT and analysed its phenomenological consequences. We discussed the structure of the vacuum expectation values and emphasised the differences with respect to the related MWT model. Then we discussed how the observed light Higgs boson emerges from 3MWT. We computed the oblique corrections and performed a scan over the parameter space of the model to contrast it with the present data. The goodness of fit analysis, which we performed by using the observed Higgs coupling strengths as well as the S and T EW parameters, demonstrated that 3MWT yields a slightly better fit to the data than the SM. In particular, the role played by heavy composite vector resonances is critical, as their contribution to the diphoton decay of the light Higgs is absolutely necessary to bring the corresponding 3MWT prediction within the experimental constraints. These composite vector resonances, having mass of $\mathcal{O}(\text{TeV})$, should be in principle observable at LHC, for example through Drell-Yan processes with two leptons in the final state or through a W' decaying into three leptons plus missing energy.

Acknowledgments

We thank R. Foadi for providing the code to evaluate the EW oblique corrections and for discussions.

Appendix A: EW Symmetry Breaking in Global SU(3) Invariant Technicolor

The SU(3) symmetry of the microscopic TC Lagrangian is spontaneously broken to the maximal diagonal subgroup, SO(3). The symmetry breaking pattern leaves us with five broken generators with associated Goldstone bosons. Such a breaking is driven by the condensate

$$\langle \eta_i^\alpha \eta_j^\beta \epsilon_{\alpha\beta} E^{ij} \rangle = \langle 2U_R^\dagger U_L + D_L D_L \rangle , \quad (\text{A1})$$

where the indices $i, j = 1, \dots, 3$ denote the components of the triplet of η , and the Greek indices indicate the ordinary spin. The matrix E is a 3×3 matrix defined as

$$E = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix} . \quad (\text{A2})$$

The above condensate is invariant under an SO(3) symmetry. It is convenient to separate the eight generators of SU(3) into the three that leave the vacuum invariant, S^a , and the remaining five that do not, X^a . Then the S^a generators of the SO(3) subgroup satisfy the relation

$$S^a E + E S^{aT} = 0 , \quad \text{with} \quad a = 1, \dots, 3 , \quad (\text{A3})$$

so that $\mathbf{u} E \mathbf{u}^T = E$, for $\mathbf{u} \in \text{SO}(3)$. An explicit realization of the generators is shown in Appendix (B).

The scalar and pseudoscalar degrees of freedom, necessary to model the Goldstone bosons and spontaneous symmetry breaking, consist of a composite Higgs and its pseudoscalar partner, as well as five pseudoscalar Goldstone bosons and their scalar partners. These can be assembled in the matrix

$$M = \left[\frac{\sigma + i\Theta}{\sqrt{3}} I_3 + \sqrt{2}(i\Pi^a + \tilde{\Pi}^a) X^a \right] E , \quad (\text{A4})$$

which transforms under the full SU(3) group according to

$$M \rightarrow \mathbf{u} M \mathbf{u}^T , \quad \text{with} \quad \mathbf{u} \in \text{SU}(3) . \quad (\text{A5})$$

The X^a 's, $a = 1, \dots, 5$ are the generators of the SU(3) group which do not leave the vacuum expectation value (VEV) of M invariant

$$\langle M \rangle = \frac{v}{\sqrt{3}} E . \quad (\text{A6})$$

Appendix B: SU(3) Generators

The generators S^i of SO(3) satisfy $S^i E + E S^{iT} = 0$. The other generators of SU(3) are written as X^i . The generators are normalized as

$$\text{Tr}[S^i S^j] = \delta^{ij}/2 \quad \text{Tr}[X^i X^j] = \delta^{ij}/2 \quad \text{Tr}[X^i S^j] = 0 \quad (\text{B1})$$

and given in terms of the Gell-Mann matrices λ^i by

$$S^1 = \frac{1}{2\sqrt{2}} (\lambda^1 - \lambda^6) \quad (\text{B2})$$

$$S^2 = \frac{1}{2\sqrt{2}} (\lambda^2 - \lambda^7) \quad (\text{B3})$$

$$S^3 = \frac{1}{4} (\lambda^3 + \sqrt{3}\lambda^8) \quad (\text{B4})$$

$$X^1 = \frac{1}{2\sqrt{2}} (\lambda^1 + \lambda^6) \quad (\text{B5})$$

$$X^2 = \frac{1}{2\sqrt{2}} (\lambda^2 + \lambda^7) \quad (\text{B6})$$

$$X^3 = \frac{1}{4} (\sqrt{3}\lambda^3 - \lambda^8) \quad (\text{B7})$$

$$X^4 = \frac{1}{2}\lambda^4 \quad (\text{B8})$$

$$X^5 = \frac{1}{2}\lambda^5 \quad (\text{B9})$$

Using the generators above, it is straightforward to obtain the vector and axial-vector charge eigenstates and their elementary particle content from Eqs. (28,30). First note that the charge operator is $Q = S^3$. We find first the linear combinations of the generators corresponding to charge eigenvalues 0, ± 1 and ± 2 . Then we project the corresponding vector

states, e.g. $v_\mu^0 = 2\text{Tr}(S^3 A_\mu)$, and obtain:

$$\begin{aligned}
v_\mu^0 &\equiv \frac{A_\mu^3 + \sqrt{3}A_\mu^8}{2} \sim \bar{U}_L \gamma_\mu U_L + \bar{U}_R \gamma_\mu U_R, \\
v_\mu^+ &\equiv \frac{A_\mu^1 - A_\mu^6}{2} - i \frac{A_\mu^2 - A_\mu^7}{2} \sim \bar{D}_L \gamma_\mu U_L + \bar{D}_L^c \gamma_\mu U_R, \\
v_\mu^- &\equiv \frac{A_\mu^1 - A_\mu^6}{2} + i \frac{A_\mu^2 - A_\mu^7}{2} \sim \bar{U}_L \gamma_\mu D_L + \bar{U}_R \gamma_\mu D_L^c, \\
a_\mu^0 &\equiv \frac{\sqrt{3}A_\mu^3 - A_\mu^8}{2} \sim \bar{U}_L \gamma_\mu U_L - \bar{U}_R \gamma_\mu U_R - 2\bar{D}_L \gamma_\mu D_L, \\
a_\mu^+ &\equiv \frac{A_\mu^1 + A_\mu^6}{2} - i \frac{A_\mu^2 + A_\mu^7}{2} \sim \bar{D}_L \gamma_\mu U_L - \bar{D}_L^c \gamma_\mu U_R, \\
a_\mu^- &\equiv \frac{A_\mu^1 + A_\mu^6}{2} + i \frac{A_\mu^2 + A_\mu^7}{2} \sim \bar{U}_L \gamma_\mu D_L - \bar{U}_R \gamma_\mu D_L^c, \\
\Omega_\mu^{++} &\equiv \frac{A_\mu^4 - iA_\mu^5}{\sqrt{2}} \sim \bar{U}_R^c \gamma_\mu U_L, \quad \Omega_\mu^{--} \equiv \frac{A_\mu^4 + iA_\mu^5}{\sqrt{2}} \sim \bar{U}_L \gamma_\mu U_R^c.
\end{aligned} \tag{B10}$$

The particle contents given above reproduce the corresponding results in [31] if one applies there the substitution $D_R \rightarrow D_L^c$.

Appendix C: Squared Mass Matrices

For the neutral scalar and pseudoscalar states, the charged and doubly charged states, the squared mass matrices are, respectively

$$\mathcal{M}_{h^0}^2 = \begin{pmatrix} 2v_\sigma^2(\lambda + 2\lambda') & 2v_\Delta v_\sigma(\lambda - 2\lambda'') \\ 2v_\Delta v_\sigma(\lambda - 2\lambda'') & 2(v_\sigma^2 \lambda'' + v_\Delta^2(\lambda + 4\lambda')) \end{pmatrix}, \tag{C1}$$

in the $\Re(\sigma^0), \Re(\Delta^0)$ basis,

$$\mathcal{M}_{\pi^0}^2 = \begin{pmatrix} 8v_\Delta^2 \lambda'' & 4v_\Delta v_\sigma \lambda'' \\ 4v_\Delta v_\sigma \lambda'' & 2v_\sigma^2 \lambda'' \end{pmatrix}, \tag{C2}$$

in the $\Im(\sigma^0), \Im(\Delta^0)$ basis,

$$\mathcal{M}_{h^\pm}^2 = \begin{pmatrix} 2v_\sigma^2(\lambda'' + \lambda') & 2\sqrt{2}v_\Delta v_\sigma(\lambda'' + \lambda') \\ 2\sqrt{2}v_\Delta v_\sigma(\lambda'' + \lambda') & 4v_\Delta^2(\lambda'' + \lambda') \end{pmatrix}, \tag{C3}$$

in the Δ^\pm, σ^\pm basis,

$$\mathcal{M}_{h^{\pm\pm}}^2 = \begin{pmatrix} 2v_\sigma^2 \lambda'' - 4(v_\Delta^2 - v_\sigma^2) \lambda' & 4v_\Delta^2 \lambda'' + 2v_\sigma^2 \lambda' \\ 4v_\Delta^2 \lambda'' + 2v_\sigma^2 \lambda' & 2v_\sigma^2 \lambda'' - 4(v_\Delta^2 - v_\sigma^2) \lambda' \end{pmatrix}, \tag{C4}$$

in the $\Delta^{\pm\pm}, \delta^{\pm\pm}$ basis.

We define, besides ϵ in Eq. (31), the following dimensionless parameters:

$$x = \frac{g_L v_w}{2m_A}, \quad t_\rho = \frac{\sqrt{2}v_\Delta}{v_\sigma}, \quad z_i = \left(\frac{g_{TC} v_w}{2m_A} \right)^2 r_i, \quad i = 1, 2, 3. \quad (\text{C5})$$

Then the non-zero terms of the charged vector boson squared mass matrix (which by definition is symmetric) are

$$\begin{aligned} (\mathcal{M}_{\tilde{W}}^2)_{1,1} &= m_A^2 \left[x^2 + \epsilon^2 \left(1 + z_1 - \frac{z_3}{2} (1 + c_\rho^2) \right) \right], \\ (\mathcal{M}_{\tilde{W}}^2)_{2,2} &= m_A^2 \left[1 + z_1 + z_2 s_{2\rho} - \frac{z_3}{2} (1 + c_\rho^2) \right], \quad (\mathcal{M}_{\tilde{W}}^2)_{1,2} = -\frac{\epsilon}{\sqrt{2}} (\mathcal{M}_{\tilde{W}}^2)_{2,2}, \\ (\mathcal{M}_{\tilde{W}}^2)_{3,3} &= m_A^2 \left[1 + z_1 - z_2 s_{2\rho} - \frac{z_3}{2} (1 + c_\rho^2) \right], \quad (\mathcal{M}_{\tilde{W}}^2)_{1,3} = -\frac{\epsilon}{\sqrt{2}} (\mathcal{M}_{\tilde{W}}^2)_{3,3}, \end{aligned} \quad (\text{C6})$$

in the $\tilde{W}_\mu^\pm, V_\mu^\pm, A_\mu^\pm$ basis, with furthermore the squared mass of the doubly charged vector boson given by

$$m_\Omega^2 = m_A^2 \left[1 + 2c_\rho^2 (z_1 + z_2) - \frac{z_3}{2} (1 + c_\rho^2) \right]. \quad (\text{C7})$$

Finally, the non-zero terms of the neutral vector boson squared mass matrix in the $\tilde{W}_\mu^3, B_\mu, V_\mu^3, A_\mu^3$ basis are

$$\begin{aligned} (\mathcal{M}_{\tilde{Z}}^2)_{1,1} &= m_A^2 \left[x^2 (1 + s_\rho^2) + \epsilon^2 \left(1 + z_1 - \frac{1}{2} (1 + c_\rho^2) z_3 + z_2 s_\rho^2 \right) \right], \\ (\mathcal{M}_{\tilde{Z}}^2)_{1,2} &= -m_A^2 \left[x^2 (1 + s_\rho^2) + \epsilon^2 (z_1 - c_\rho^2 (2z_1 - z_2) + z_2) \right] t_\xi, \\ (\mathcal{M}_{\tilde{Z}}^2)_{2,2} &= m_A^2 \left[x^2 (1 + s_\rho^2) + \epsilon^2 \left(3 + z_1 + c_\rho^2 (4z_1 - 5z_2) + z_2 - \frac{3}{2} (1 + c_\rho^2) z_3 \right) \right] t_\xi^2, \\ (\mathcal{M}_{\tilde{Z}}^2)_{3,3} &= m_A^2 \left[1 + 2c_\rho^2 (z_1 - z_2) - \frac{z_3}{2} (1 + c_\rho^2) \right], \\ (\mathcal{M}_{\tilde{Z}}^2)_{1,3} &= -\frac{\epsilon}{2} (\mathcal{M}_{\tilde{Z}}^2)_{3,3}, \quad (\mathcal{M}_{\tilde{Z}}^2)_{2,3} = -\frac{3}{2} \epsilon t_\xi (\mathcal{M}_{\tilde{Z}}^2)_{3,3}, \\ (\mathcal{M}_{\tilde{Z}}^2)_{4,4} &= m_A^2 \left[1 + z_1 - z_2 s_{2\rho} - \frac{z_3}{2} (1 + c_\rho^2) \right], \\ (\mathcal{M}_{\tilde{Z}}^2)_{1,4} &= -\frac{\sqrt{3}}{2} \epsilon (\mathcal{M}_{\tilde{Z}}^2)_{4,4}, \quad (\mathcal{M}_{\tilde{Z}}^2)_{2,4} = \frac{\sqrt{3}}{2} \epsilon t_\xi (\mathcal{M}_{\tilde{Z}}^2)_{4,4}. \end{aligned} \quad (\text{C8})$$

Appendix D: S and T Parameters for General Neutrino Mass Matrix

The most general mass terms for a pair of right- and left-handed neutrinos is defined by

$$\mathcal{L} \supset -m_E \bar{E}_R E_L - \frac{1}{2} n_L^T M n_L + \text{h.c.}, \quad M = \begin{pmatrix} M_L & m_D \\ m_D & M_R \end{pmatrix}, \quad n_L = (N_L, \bar{N}_R)^T \quad (\text{D1})$$

with eigenvalues

$$\lambda_{1,2} = \frac{1}{2} \left[(M_L + M_R) \pm \sqrt{(M_L - M_R)^2 + 4m_D^2} \right]. \quad (\text{D2})$$

The contributions of the corresponding heavy neutrinos mass eigenstates and of the heavy electron E to the S and T parameters have been derived in terms of integral functions in [43]. From those, we derived the corresponding explicit results:

$$\begin{aligned} S = & \frac{1}{12\pi} \left[1 + 2c_\zeta^4 (1 + \log \nu_1^2) - 2 \log \nu_E^2 + 2s_\zeta^4 (1 + \log \nu_2^2) \right] \\ & + \frac{s_{2\zeta}^2}{36\pi} \frac{9(1 - \log \nu_1^2) \nu_1^4 \nu_2^2 - 9(1 - \log \nu_2^2) \nu_1^2 \nu_2^4 - (1 - 3 \log \nu_1^2) \nu_1^6 + (1 - 3 \log \nu_2^2) \nu_2^6}{(\nu_1^2 - \nu_2^2)^3} \\ & - \frac{(-1)^\beta s_{2\zeta}^2 \nu_1 \nu_2 \left(\nu_1^4 - 2\nu_1^2 \nu_2^2 \log \frac{\nu_1^2}{\nu_2^2} - \nu_2^4 \right)}{8\pi (\nu_1^2 - \nu_2^2)^3}, \end{aligned} \quad (\text{D3})$$

$$\begin{aligned} T = & \frac{\Lambda_{NP}^2}{64\pi c_\xi^2 s_\xi^2 m_Z^2} \left[16c_\zeta^4 \nu_1^2 \log \nu_1^2 + 16s_\zeta^4 \nu_2^2 \log \nu_2^2 + 8\nu_E^2 \log \nu_E^2 \right. \\ & - s_{2\zeta}^2 \frac{(1 - 2 \log \nu_1^2) \nu_1^4 - (1 - 2 \log \nu_2^2) \nu_2^4}{\nu_1^2 - \nu_2^2} + 4(-1)^\beta s_{2\zeta}^2 \frac{(1 - \log \nu_1^2) \nu_1^3 \nu_2 - (1 - \log \nu_2^2) \nu_1 \nu_2^3}{\nu_1^2 - \nu_2^2} \\ & \left. + 4c_\zeta^2 \frac{(1 - 2 \log \nu_1^2) \nu_1^4 - (1 - 2 \log \nu_E^2) \nu_E^4}{\nu_1^2 - \nu_E^2} + 4s_\zeta^2 \frac{(1 - 2 \log \nu_2^2) \nu_2^4 - (1 - 2 \log \nu_E^2) \nu_E^4}{\nu_2^2 - \nu_E^2} \right] \end{aligned} \quad (\text{D4})$$

where Λ_{NP} is the given renormalization scale, ξ is the EW mixing angle, and

$$\nu_1 = \frac{\lambda_1}{\Lambda_{NP}}, \quad \nu_2 = \frac{\lambda_2}{\Lambda_{NP}}, \quad \nu_E = \frac{m_E}{\Lambda_{NP}}, \quad t_{2\zeta} = \frac{2m_D}{M_R - M_L}, \quad \beta = \frac{1}{2} \left[1 + \left(\sqrt{\frac{\lambda_1}{|\lambda_1|}} * \sqrt{\frac{\lambda_2}{|\lambda_2|}} \right)^2 \right]. \quad (\text{D5})$$

In the limit $M_R \rightarrow \infty$, and $M_L = m_E \equiv m_U$, one recovers the results in Eqs. (46).

-
- [1] G. Aad et al. (ATLAS), Phys.Lett. **B716**, 1 (2012), 1207.7214.
 - [2] S. Chatrchyan et al. (CMS), Phys.Lett. **B716**, 30 (2012), 1207.7235.
 - [3] The ATLAS collaboration, Tech. Rep. ATLAS-CONF-2013-034, CERN, Geneva (2013).
 - [4] The CMS collaboration, Tech. Rep. CMS-PAS-HIG-13-005, CERN, Geneva (2013).
 - [5] M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann, and K. Tuominen (2013), 1304.7006.
 - [6] M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann, and K. Tuominen (2013), 1305.4182.
 - [7] S. Weinberg, Phys.Rev. **D13**, 974 (1976).

- [8] L. Susskind, Phys.Rev. **D20**, 2619 (1979).
- [9] B. Holdom, Phys.Lett. **B150**, 301 (1985).
- [10] K. Yamawaki, M. Bando, and K.-i. Matumoto, Phys.Rev.Lett. **56**, 1335 (1986).
- [11] D. D. Dietrich, F. Sannino, and K. Tuominen, Phys.Rev. **D72**, 055001 (2005), hep-ph/0505059.
- [12] D. Elander, C. Nunez, and M. Piai, Phys.Lett. **B686**, 64 (2010), 0908.2808.
- [13] B. Grinstein and P. Uttayarat, JHEP **1107**, 038 (2011), 1105.2370.
- [14] O. Antipin, M. Mojaza, and F. Sannino, Phys.Lett. **B712**, 119 (2012), 1107.2932.
- [15] C. T. Hill and E. H. Simmons, Phys.Rept. **381**, 235 (2003), hep-ph/0203079.
- [16] R. Foadi, M. T. Frandsen, and F. Sannino (2012), 1211.1083.
- [17] S. Dimopoulos and L. Susskind, Nucl.Phys. **B155**, 237 (1979).
- [18] E. Eichten and K. D. Lane, Phys.Lett. **B90**, 125 (1980).
- [19] T. Appelquist and J. Terning, Phys.Rev. **D50**, 2116 (1994), hep-ph/9311320.
- [20] T. Appelquist, M. Piai, and R. Shrock, Phys.Rev. **D69**, 015002 (2004), hep-ph/0308061.
- [21] E. H. Simmons, Nucl.Phys. **B312**, 253 (1989).
- [22] A. Kagan and S. Samuel, Phys.Lett. **B270**, 37 (1991).
- [23] C. D. Carone and E. H. Simmons, Nucl.Phys. **B397**, 591 (1993), hep-ph/9207273.
- [24] C. D. Carone, E. H. Simmons, and Y. Su, Phys.Lett. **B344**, 287 (1995), hep-ph/9410242.
- [25] M. Antola, M. Heikinheimo, F. Sannino, and K. Tuominen, JHEP **1003**, 050 (2010), 0910.3681.
- [26] M. Dine, W. Fischler, and M. Srednicki, Nucl.Phys. **B189**, 575 (1981).
- [27] B. A. Dobrescu, Nucl.Phys. **B449**, 462 (1995), hep-ph/9504399.
- [28] M. Antola, S. Di Chiara, F. Sannino, and K. Tuominen, Eur.Phys.J. **C71**, 1784 (2011), 1001.2040.
- [29] M. Antola, S. Di Chiara, F. Sannino, and K. Tuominen, Nucl.Phys. **B864**, 664 (2012), 1111.1009.
- [30] F. Sannino and K. Tuominen, Phys.Rev. **D71**, 051901 (2005), hep-ph/0405209.
- [31] R. Foadi, M. T. Frandsen, T. A. Rytto, and F. Sannino, Phys.Rev. **D76**, 055005 (2007), 0706.1696.
- [32] M. E. Peskin and T. Takeuchi, Phys.Rev.Lett. **65**, 964 (1990).
- [33] M. E. Peskin and T. Takeuchi, Phys.Rev. **D46**, 381 (1992).

- [34] H. Georgi, Phys.Lett. **B298**, 187 (1993), hep-ph/9207278.
- [35] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, Phys.Lett. **B412**, 301 (1997), hep-ph/9706275.
- [36] M. Antola and K. Tuominen (2011), 1105.3178.
- [37] M. Bando, T. Kugo, S. Uehara, K. Yamawaki, and T. Yanagida, Phys.Rev.Lett. **54**, 1215 (1985).
- [38] M. Bando, T. Kugo, and K. Yamawaki, Phys.Rept. **164**, 217 (1988).
- [39] J. Beringer et al. (Particle Data Group), Phys.Rev. **D86**, 010001 (2012).
- [40] D. D. Dietrich and F. Sannino, Phys.Rev. **D75**, 085018 (2007), hep-ph/0611341.
- [41] R. Barbieri, A. Pomarol, R. Rattazzi, and A. Strumia, Nucl.Phys. **B703**, 127 (2004), hep-ph/0405040.
- [42] R. S. Chivukula, E. H. Simmons, H.-J. He, M. Kurachi, and M. Tanabashi, Phys.Rev. **D70**, 075008 (2004), hep-ph/0406077.
- [43] O. Antipin, M. Heikinheimo, and K. Tuominen, JHEP **0910**, 018 (2009), 0905.0622.
- [44] H.-J. He, N. Polonsky, and S.-f. Su, Phys.Rev. **D64**, 053004 (2001), hep-ph/0102144.
- [45] T. Alanne, S. Di Chiara, and K. Tuominen (2013), 1303.3615.
- [46] ATLAS-CONF-2012-086, ATLAS-COM-CONF-2012-132 (2012).
- [47] The ATLAS collaboration, Tech. Rep. ATLAS-CONF-2012-160, CERN, Geneva (2012).
- [48] The ATLAS collaboration, Tech. Rep. ATLAS-CONF-2012-161, CERN, Geneva (2012).
- [49] The ATLAS collaboration, Tech. Rep. ATLAS-CONF-2013-030, CERN, Geneva (2013).
- [50] The ATLAS collaboration, Tech. Rep. ATLAS-CONF-2013-012, CERN, Geneva (2013).
- [51] The ATLAS collaboration, Tech. Rep. ATLAS-CONF-2013-013, CERN, Geneva (2013).
- [52] The CMS collaboration, Tech. Rep. CMS-PAS-HIG-13-002, CERN, Geneva (2013).
- [53] The CMS collaboration, Tech. Rep. CMS-PAS-HIG-13-001, CERN, Geneva (2013).
- [54] The CMS collaboration, Tech. Rep. CMS-PAS-HIG-13-003, CERN, Geneva (2013).
- [55] The CMS collaboration, Tech. Rep. CMS-PAS-HIG-13-004, CERN, Geneva (2013).
- [56] T. Aaltonen et al. (CDF Collaboration, D0 Collaboration) (2013), 1303.6346.
- [57] J. F. Gunion, H. E. Haber, G. L. Kane, and S. Dawson, Front.Phys. **80**, 1 (2000).