

Using Quantum entanglement to study CP and CPT violations

Yu Shi

*Department of Physics, Fudan University
Shanghai, 200433, China*

** E-mail: yushi@fudan.edu.cn*

We report some general phenomenological results concerning CP and CPT violations in joint decays of entangled pseudoscalar neutral mesons.¹⁻⁴

1. Introduction

Quantum entanglement refers to the situation that the quantum state of a composite system is not a direct product of its subsystems. Its significance was discovered by Einstein, Podolsky and Rosen.⁵ Schödinger coined the term and regarded it as “the characteristic trait of quantum mechanics, the one that enforces its entire departure from classical lines of thought.”⁶ Pseudoscalar neutral mesons are copiously generated as pairs entangled in the flavor space.⁷⁻¹³ CP and CPT violating parameters can be measured by studying the joint decays of entangled meson pairs.^{1-4,7,12,14,15} This provides a venue of searching standard model extension.¹⁵

2. Single meson states

$|M^0\rangle$ and $|\bar{M}^0\rangle$ are eigenstates of parity P both with eigenvalue -1 , and of a characteristic flavor \mathcal{F} with eigenvalues ± 1 . $C|M^0\rangle = -|\bar{M}^0\rangle$, $C|\bar{M}^0\rangle = -|M^0\rangle$. Hence the eigenstates of CP are $|M_{\pm}\rangle = \frac{1}{\sqrt{2}}(|M^0\rangle \pm |\bar{M}^0\rangle)$, with eigenvalues ± 1 . In the Wigner-Weisskopf approximation, the weak decay of a meson can be described by $i\frac{\partial}{\partial t}|M(t)\rangle = H|M(t)\rangle$, where $H = \begin{pmatrix} H_{00} & H_{0\bar{0}} \\ H_{\bar{0}0} & H_{\bar{0}\bar{0}} \end{pmatrix}$. It is defined that $\frac{1-\epsilon_M}{1+\epsilon_M} \equiv \sqrt{\frac{H_{00}}{H_{0\bar{0}}}} \equiv \frac{q}{p}$, $\delta_M \equiv \frac{H_{0\bar{0}} - H_{\bar{0}0}}{\sqrt{H_{00}H_{\bar{0}\bar{0}}}}$.

- If CP or T is conserved indirectly, then $\epsilon_M = 0$.
- If CPT or CP is conserved indirectly, then $\delta_M = 0$.

The eigenstates $|M_S\rangle$ and $|M_L\rangle$ of H , corresponding to the eigenvalues λ_S and λ_L respectively, are found by diagonalizing H . Starting as $|M_S\rangle$, the state of a single meson evolves as $|M_S(t)\rangle = e^{-i\lambda_S t}|M_S\rangle$. Starting as $|M_L\rangle$, the state evolves as $|M_L(t)\rangle = e^{-i\lambda_L t}|M_L\rangle$. Based on this, one obtains $|M^0(t)\rangle$, which starts as $|M^0\rangle$; $|\bar{M}^0(t)\rangle$, which starts as $|\bar{M}^0\rangle$; $|M_+(t)\rangle$, which starts as $|M_+\rangle$; and $|M_-(t)\rangle$, which starts as $|M_-\rangle$.

To characterize direct violations, for decays into flavor eigenstates $|l^\pm\rangle$ with eigenvalue ± 1 , we define decay amplitudes $R^+ \equiv \langle l^+|\mathcal{H}|M^0\rangle$, $S^+ \equiv \langle l^+|\mathcal{H}|\bar{M}^0\rangle$, $S^- \equiv \langle l^-|\mathcal{H}|M^0\rangle$, $R^- \equiv \langle l^-|\mathcal{H}|\bar{M}^0\rangle$. They can be related to quantities a, b, c, d usually defined.^{2,7}

- If CP is conserved directly, then $R^+ = R^-$ and $S^+ = S^-$.
- If CPT is conserved directly, then $(R^+)^* = R^-$ and $(S^+)^* = S^-$.
- If $\Delta\mathcal{F} = \Delta Q$ rule is respected, then we have $S^\pm = 0$.

For decays into CP eigenstates $|h^\pm\rangle$ with eigenvalue ± 1 , we define decay amplitudes $Q^+ \equiv \langle h^+|\mathcal{H}|M_+\rangle$, $X^+ \equiv \langle h^+|\mathcal{H}|M_-\rangle$, $X^- \equiv \langle h^-|\mathcal{H}|M_+\rangle$, $Q^- \equiv \langle h^-|\mathcal{H}|M_-\rangle$. These newly defined quantities are convenient.

- If CP is conserved directly, then $X^\pm = 0$.
- If CPT is conserved directly, then X^\pm is purely imaginary.

3. Entangled states

The $C = -1$ entangled state of a pair of pseudoscalar mesons is $|\Psi_-\rangle = \frac{1}{\sqrt{2}}(|M^0\rangle_a|\bar{M}^0\rangle_b - |\bar{M}^0\rangle_a|M^0\rangle_b) = \frac{1}{\sqrt{2}}(|M_-\rangle_a|M_+\rangle_b - |M_+\rangle_a|M_-\rangle_b)$, which is produced for kaons at ϕ resonance,⁷ and for B mesons by $\Upsilon(4s)$ resonance^{8,9} and by $\Upsilon(5S)$ resonance with a large branch ratio.¹⁰⁻¹² The $C = +1$ entangled state is $|\Psi_+\rangle = \frac{1}{\sqrt{2}}(|M_0\rangle_a|\bar{M}_0\rangle_b + |\bar{M}_0\rangle_a|M_0\rangle_b) = \frac{1}{\sqrt{2}}(|M_+\rangle_a|M_+\rangle_b - |M_-\rangle_a|M_-\rangle_b)$, which is produced for B mesons by $\Upsilon(5S)$ resonance with some branch ratio¹⁰⁻¹² and above $\Upsilon(4s)$ resonance.¹³

Although physically a single meson cannot be in a CP eigenstate $|M_\pm\rangle$ because of CP violation, the entangled states can be *exactly* written in terms of $|M_\pm\rangle$, this makes the expression $|M_\pm(t)\rangle$ and the decay amplitudes Q^\pm and X^\pm meaningful and useful.

Starting from $|\Psi_-\rangle$, the entangled meson pair decay to certain products at t_a and t_b , hence the time-dependent state is $|\Psi_-(t_a, t_b)\rangle = \frac{1}{\sqrt{2}}(|M^0(t_a)\rangle_a|\bar{M}^0(t_b)\rangle_b - |\bar{M}^0(t_a)\rangle_a|M^0(t_b)\rangle_b) = \frac{1}{\sqrt{2}}(|M_-(t_a)\rangle_a|M_+(t_b)\rangle_b - |M_+(t_a)\rangle_a|M_-(t_b)\rangle_b)$. Similarly, starting as $|\Psi_+\rangle$, $|\Psi_+(t_a, t_b)\rangle = \frac{1}{\sqrt{2}}(|M^0(t_a)\rangle_a|\bar{M}^0(t_b)\rangle_b + |\bar{M}^0(t_a)\rangle_a|M^0(t_b)\rangle_b) = \frac{1}{\sqrt{2}}(|M_+(t_a)\rangle_a|M_+(t_b)\rangle_b - |M_-(t_a)\rangle_a|M_-(t_b)\rangle_b)$.

4. Joint decay rates

For state $|\Psi(t_a, t_b)\rangle$, the joint rate that particle a decays to f at t_a while particle b decays to g at t_b is $I(f, t_a; g, t_b) = |\langle f, g | \mathcal{H}_a \mathcal{H}_b | \Psi(t_a, t_b) \rangle|^2$, where \mathcal{H}_a and \mathcal{H}_b represent the Hamiltonians governing the weak decays of a and b , respectively. In experiments, it is more convenient to use the integrated rate $I(f, g, \Delta t) = \int_0^\infty I(f, t_a; g, t_a + \Delta t) dt_a$. Then one can find the asymmetry between the joint decays to f and g and the joint decays to f' and g' , $A(fg, f'g', \Delta t) \equiv \frac{I(f, t_a; g, t_a + \Delta t) - I(f', t_a; g', t_a + \Delta t)}{I(f, t_a; g, t_a + \Delta t) + I(f', t_a; g', t_a + \Delta t)} = \frac{I(f, g, \Delta t) - I(f', g', \Delta t)}{I(f, g, \Delta t) + I(f', g', \Delta t)}$.

We have considered the following cases of the final states: (1) the decay products are flavor eigenstates $|l^\pm\rangle$, with the equal-flavor asymmetry $A(l^+l^+, l^-l^-, \Delta t)$ and the unequal-flavor asymmetry $A(l^+l^-, l^-l^+, \Delta t)$; (2) the decay products are CP eigenstates $|h^\pm\rangle$, with the equal-CP asymmetry $A(h^+h^+, h^-h^-, \Delta t)$ and the unequal-CP asymmetry $A(h^+h^-, h^-h^+, \Delta t)$; (3) the decay products $|h_1\rangle$ and $|h_2\rangle$ are CP conjugates.

5. General results on joint decays of $|\Psi_-\rangle^2$

Theorem 1 If the equal-flavor asymmetry is nonzero, then there exists one or two of the following violations: (1) CP is violated indirectly, (2) both CP and CPT are violated directly.

Theorem 2 If the equal-flavor asymmetry is nonzero while CPT is assumed to be conserved both directly and indirectly, then in addition to indirect CP violation, we can draw the following conclusions: (1) $|q/p| \neq 1$, i.e. T must also be violated indirectly; (2) $|\langle l^+ | \mathcal{H} | \bar{M}_0 \rangle| \neq |\langle l^+ | \mathcal{H} | M^0 \rangle|$, $|\langle l^- | \mathcal{H} | M^0 \rangle| \neq |\langle l^- | \mathcal{H} | \bar{M}_0 \rangle|$, despite $\langle l^+ | \mathcal{H} | M^0 \rangle = \langle l^- | \mathcal{H} | \bar{M}_0 \rangle^*$ and $\langle l^+ | \mathcal{H} | \bar{M}_0 \rangle = \langle l^- | \mathcal{H} | M^0 \rangle^*$.

Theorem 3 If the unequal-flavor asymmetry is nonzero, then CP must be violated, directly or indirectly or both.

Theorem 4 If the unequal-flavor asymmetry is nonzero for $\Delta t \neq 0$ while CPT is assumed to be conserved both directly and indirectly, then we can draw the following conclusions: (1) $|\langle l^+ | \mathcal{H} | \bar{M}_0 \rangle| = |\langle l^- | \mathcal{H} | M^0 \rangle| \neq |\langle l^+ | \mathcal{H} | M^0 \rangle| = |\langle l^- | \mathcal{H} | \bar{M}_0 \rangle|$; (2) $\langle l^- | \mathcal{H} | M^0 \rangle = \langle l^+ | \mathcal{H} | \bar{M}_0 \rangle^* \neq 0$, which means $\Delta\mathcal{F} = \Delta Q$ rule must be violated.

Theorem 5 The equal-CP asymmetry is always a constant independent of Δt .

Theorem 6 For $\Delta t = 0$, the unequal-CP asymmetry vanishes, no matter whether CP or CPT is violated.

Theorem 7 If any equal-CP joint decay rate is nonzero, then CP must be violated, directly or indirectly or both.

6. General results on joint decays of $|\Psi_+\rangle^3$

Theorem 8: If the unequal-flavor asymmetry is nonzero, then CP must be violated indirectly.

Theorem 9: If the $I(l^+, l^-, \Delta t)$ and $I(l^-, l^+, \Delta t)$ depend on the first order of ϵ_M , then CP must also be violated directly.

Theorem 10: If the unequal-flavor asymmetry depends on the first or second order of ϵ_M , then $\Delta\mathcal{F} = \Delta Q$ rule is violated.

Theorem 11: The deviation of the unequal-CP joint decay rate $I(h^+, t_a; h^-, t_b)$ or $I(h^-, t_a; h^+, t_b)$ from zero implies direct CP violation.

Theorem 12: Suppose $|h_1\rangle$ and $|h_2\rangle$ are CP conjugates. If $I(h_1, h_2; \Delta t)$ and $I(h_2, h_1; \Delta t)$ depend on the first order of ϵ_M , then CP is violated directly.

In addition, we have derived various quantitative relations of the indirect violating parameters with the decay asymmetries of $|\Psi_-\rangle$,^{1,2} with those of $|\Psi_+\rangle$ ³ and with four asymmetries defined for some time-ordered integrated rates of $|\Psi_-\rangle$ and $|\Psi_+\rangle$.⁴

This work is supported by the National Science Foundation of China (Grant No. 10875028).

References

1. Y. Shi, Euro. Phys. J. C **72**, 1907(2012).
2. Y. Shi, arXiv:1306.2676.
3. Z. Huang and Y. Shi, arXiv:1307.4459.
4. Z. Huang and Y. Shi, Euro. Phys. J. C **72**, 1900(2012).
5. E. Einstein, B. Podolsky and N. Rosen, Phys. Rev. **47**, 777 (1935).
6. E. Schrödinger, Proc. Camb. Phi. Soc. **31**, 555 (1936).
7. A. Di Domenico (KLOE Collaboration), Found. Phys. **40**, 852 (2010).
8. B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **88**, 221802 (2002).
9. A. Go *et al.* (BELLE Collaboration), Phys. Rev. Lett. **99**, 131802 (2007).
10. O. Aquines *et al.* (CLEO Collaboration), Phys. Rev. Lett. **96**, 152001 (2006).
11. R. Louvot *et al.* (BELLE Collaboration), Phys. Rev. Lett. **102**, 021801 (2009).
12. D. Atwood and A. Soni, Phys. Rev. D **82**, 036003 (2010).
13. D. S. Skerib *et al.* (CLEO II Collaboration), Phys. Rev. Lett. **67**, 1692 (1991).
14. J. Bernabéu, F. J. Botella and J. Roldán, Phys. Lett. B **211**, 226 (1980); I. Dunietz, J. Hauser and J. L. Rosner, Phys. Rev. D **35**, 2166 (1987); C. D. Buchanan *et al.*, Phys. Rev. D **45**, 4088 (1992); G. D'Ambrosio, G. Isidori and A. Pugliese, in L. Maiani, G. Pancheri and N. Paver. (eds.), *The Second DAΦNE Physics Handbook*, (SIS-Pubblicazioni, Frascati, 1995).
15. V. A. Kostelecký, Phys. Rev. Lett. **80**, 1818 (1998); V. A. Kostelecký, Phys. Rev. D **61**, 016002 (1999); V. A. Kostelecký, Phys. Rev. D **64**, 076001 (2001).