

Mixing Patterns from the Groups $\Sigma(n\varphi)$

Claudia Hagedorn ^{a),b),c)}, Aurora Meroni ^{c),d)}
and Lorenzo Vitale ^{e)}

^{a)} Dipartimento di Fisica e Astronomia ‘G. Galilei’, Università di Padova,
Via Marzolo 8, I-35131 Padua, Italy

^{b)} INFN, Sezione di Padova, Via Marzolo 8, I-35131 Padua, Italy

^{c)} SISSA, Via Bonomea 265, I-34136 Trieste, Italy

^{d)} INFN, Sezione di Trieste, I-34126 Trieste, Italy

^{e)} Institut de Théorie des Phénomènes Physiques, EPFL,
Lausanne, Switzerland

E-mail: hagedorn@pd.infn.it, ameroni@sissa.it, lorenzo.vitale@epfl.ch

Abstract

We survey the mixing patterns which can be derived from the discrete groups $\Sigma(36 \times 3)$, $\Sigma(72 \times 3)$, $\Sigma(216 \times 3)$ and $\Sigma(360 \times 3)$, if these are broken to abelian subgroups G_e and G_ν in the charged lepton and neutrino sector, respectively. Since only $\Sigma(360 \times 3)$ possesses Klein subgroups, only this group allows neutrinos to be Majorana particles. We find a few patterns that can agree well with the experimental data on lepton mixing in scenarios with small corrections and that predict the reactor mixing angle θ_{13} to be $0.1 \lesssim \theta_{13} \lesssim 0.2$. All these patterns lead to a trivial Dirac phase. Patterns which instead reveal CP violation tend to accommodate the data not well. We also comment on the outer automorphisms of the discussed groups, since they can be useful for relating inequivalent representations of these groups.

1 Introduction

Finite discrete non-abelian flavor symmetries have been widely used in order to describe the three lepton mixing angles, for reviews see [1, 2], whose best fit values and 1σ errors are

$$\sin^2 \theta_{12} = 0.302_{-0.012}^{+0.013}, \quad \sin^2 \theta_{23} = 0.413_{-0.025}^{+0.037} \quad \text{and} \quad 0.594_{-0.022}^{+0.021}, \quad \sin^2 \theta_{13} = 0.0227_{-0.0024}^{+0.0023} \quad (1)$$

according to [3].¹ A particularly interesting approach is based on the assumption that a (lepton) flavor symmetry G_l is broken to two abelian subgroups G_e and G_ν in the charged lepton and neutrino sector, respectively [5, 6]. The mismatch of the embedding of these two subgroups into the group G_l determines the lepton mixing encoded in the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) mixing matrix. More precisely, the PMNS matrix is determined up to possible permutations of rows and columns (and phases) so that the results for the mixing angles belong to a small set of solutions. At the same time, the Jarlskog invariant [7] is determined up to a sign (and consequently the Dirac phase δ can be predicted). Left-handed leptons are assigned to an irreducible (faithful) three-dimensional representation of the group G_l in order to achieve at least two non-vanishing mixing angles. As regards the symmetries, G_e can be any abelian group which is capable of distinguishing among the three generations, while the nature of neutrinos decides on G_ν . For Dirac neutrinos G_ν is subject to the same constraints as G_e , while Majorana neutrinos require G_ν to be (a subgroup of) a Klein group. A prime example of this approach is tri-bimaximal (TB) mixing [8], $\sin^2 \theta_{12} = 1/3$, $\sin^2 \theta_{23} = 1/2$ and $\theta_{13} = 0$, which can be derived with the help of the permutation symmetry S_4 [9]. The vanishing of the reactor mixing angle $\theta_{13} = 0$, however, is not (well-)compatible with experimental data, since measurements have shown $\theta_{13} \approx 0.15$ [10]. This fact has triggered an intense search for discrete groups which can give rise to mixing patterns with non-vanishing θ_{13} , see e.g. [6, 11–15], as well as it has led to modifications of this approach, e.g. scenarios in which the symmetry of the neutrino sector is only partly contained in the group G_l [16] or setups in which the flavor symmetry is combined with a CP symmetry and the neutrino sector is invariant under a subgroup of G_l and under the CP symmetry [17, 18].²

We follow the approach to search for new discrete groups and focus in the present paper on the so-called “exceptional” finite groups $\Sigma(n\varphi)$: $\Sigma(36\varphi)$, $\Sigma(72\varphi)$, $\Sigma(216\varphi)$, $\Sigma(360\varphi)$ with $\varphi = 1, 3$, which are subgroups of $SU(3)$ ($\varphi = 3$) or of $SU(3)/C$ ($\varphi = 1$) with C being the center of $SU(3)$ [20]. Groups with $\varphi = 1$ are not suitable for our purposes,³ since they do not possess an irreducible faithful three-dimensional representation, while the groups with $\varphi = 3$ do have such representations. We compute all mixing patterns related to these groups, assuming neutrinos to be either Dirac or Majorana particles.⁴

¹Throughout this paper we use the values given in the left table of table 1 of [3] which are obtained by performing a global fit in which the reactor fluxes are left free and short baseline reactor data with $L \lesssim 100$ m are included, called “Free Fluxes + RSBL” in [3]. For other global fits which reach similar results see [4].

²For further literature and models with a flavor and a CP symmetry see [19].

³The group $\Sigma(36)$ is among the finite groups which can appear as symmetry group of the scalar sector of a three Higgs doublet model without leading to a potential with a continuous symmetry [21].

⁴This is in contrast to the recently performed scans of groups with an order smaller than 200 [15], an order smaller than 512 [12] or an order smaller than 1536 [13] that had the scope to only figure

Among these four groups only $\Sigma(360 \times 3)$ has Klein subgroups and is thus suitable as G_l , if neutrinos are assumed to be Majorana particles. $\Sigma(36 \times 3)$, $\Sigma(72 \times 3)$ and $\Sigma(216 \times 3)$ instead are only appropriate as G_l , if neutrinos are Dirac particles or only one of the Z_2 symmetries of the neutrino mass matrix is contained in G_l . In the latter case only one of the columns of the PMNS matrix is determined through the breaking of the flavor symmetry G_l to G_e and to G_ν . One can generalize this idea and consider as G_e or G_ν subgroups of G_l which are generated by elements that are represented by matrices with two degenerate eigenvalues. Then only one of the rows or columns of the PMNS matrix is fixed. This row/column is called “mixing vector” [12].

The authors of [22] have shown that the groups $\Sigma(72 \times 3)$, $\Sigma(216 \times 3)$ and $\Sigma(360 \times 3)$ are suitable for realizing a version of minimal flavor violation in the quark sector with a discrete group, since the products of (faithful) three-dimensional representations with their complex conjugates decompose in the same way as in $SU(3)$, i.e. $\mathbf{3} \times \mathbf{3}^* = \mathbf{1} + \mathbf{8}$ with $\mathbf{8}$ being irreducible in G_l . The group $\Sigma(168) \simeq PSL(2, Z_7)$ has also been recognized as interesting by the authors of [22]. However, the possible mixing patterns arising from this group have been already analyzed in [6], mainly under the assumption that neutrinos are Majorana particles.⁵

We find that the two smaller groups $\Sigma(36 \times 3)$ and $\Sigma(72 \times 3)$ only give rise to a few new mixing patterns, while the larger groups lead to several new patterns, since they have a richer structure of subgroups. Generically one sees (as already suggested by the scans of groups making use of the computer program GAP [12,13,15]) that only very few patterns are well-compatible with the experimental data of all three different mixing angles [3], if only small corrections are admitted, i.e. for each group one to two. Such small corrections, typically of order $\lambda^2 \approx 0.04$, are usually expected to arise in concrete models, see e.g. [1], and thus any pattern which can be brought into agreement with the experimental results including this type of corrections is considered as very interesting for model building. However, if one of the other patterns is realized in an explicit model one either has to invoke large corrections to the leading order results in order to reconcile them with the data or one needs to consider modifications of the breaking pattern, such as involving a CP symmetry [17,18]. It is noteworthy that patterns which fit the experimental data better usually do not lead to a non-trivial Dirac phase (although all mixing angles are different from 0 and $\pi/2$).⁶ On the other hand, among the patterns associated with the group $\Sigma(216 \times 3)$ we find several ones which give rise to non-vanishing CP violation and at the same time accommodate the solar and atmospheric mixing angles at the 3σ level or better. Only the reactor mixing angle turns out to be too large, see table 2. Furthermore, we apply outer automorphisms of the groups $\Sigma(n\varphi)$, $\varphi = 3$, in order to relate their irreducible faithful three-dimensional representations among each other. In this way, we see that it is sufficient to focus on a single three-dimensional representation in the discussion of the mixing patterns.

out the groups which, if broken in a non-trivial way, lead to mixing patterns, accommodating best the experimental data [3].

⁵For the sixth group of this type, called $\Sigma(60)$ and isomorphic to A_5 , the mixing patterns have also already been studied in detail for Majorana neutrinos with some comments on Dirac neutrinos [6,23]. This group however is not so interesting according to [22], since it permits several more invariants than the groups associated with discrete minimal flavor violation in the quark sector.

⁶Similar observations can also be made in the analyses found in [13–15].

The paper is structured as follows: in Section 2 we review the approach used in order to relate the mixing pattern with the group G_l and its breaking to the subgroups G_e and G_ν in the charged lepton and neutrino sectors. In Section 3 we discuss the mathematical properties of the different groups $\Sigma(n\varphi)$, the representations to which the three generations of left-handed leptons can be assigned, the relevant abelian subgroups and the resulting mixing patterns and predictions for mixing angles θ_{ij} and the Jarlskog invariant J_{CP} . We use as figure of merit χ^2 assuming the best fit values and 1σ errors of the mixing angles as given in eq.(1). We also comment on the mixing vectors for each group. We summarize in Section 4. Comments on outer automorphisms of the groups $\Sigma(36 \times 3)$, $\Sigma(72 \times 3)$ and $\Sigma(216 \times 3)$ can be found in Appendix A.

2 Approach

We briefly review how lepton mixing can be predicted with a non-abelian discrete flavor symmetry G_l which is broken, spontaneously or explicitly, to two different abelian subgroups G_e and G_ν in the charged lepton and neutrino sector [5, 6]. We assign the three generations of left-handed leptons to an irreducible three-dimensional representation, because we want to discuss patterns with at least two non-vanishing lepton mixing angles. We furthermore choose the representation to be faithful in the group G_l so that we (mostly) study mixing patterns which originate from this group itself and not from one of its proper subgroups.⁷ In order to fix the mixing angles we also have to assume that the three generations of left-handed leptons transform as three inequivalent one-dimensional representations under the residual symmetries G_e and G_ν . A possible set of generators of G_e and G_ν is called $g_{e,i}$, $i = 1, 2, \dots$ and $g_{\nu,j}$, $j = 1, 2, \dots$, respectively.⁸ Clearly, these are also elements of the group G_l . The requirement that a subgroup G_e is conserved entails that the combination $m_e m_e^\dagger$ (with the convention $\bar{\Psi}_L m_e \Psi_R$) has to be invariant under the symmetry G_e , i.e.

$$g_{e,i}^\dagger m_e m_e^\dagger g_{e,i} = m_e m_e^\dagger \quad , \quad i = 1, 2, \dots \quad , \quad (2)$$

while requiring that G_ν is a symmetry of the neutrino mass matrix m_ν entails that $m_\nu m_\nu^\dagger$ (m_ν itself) is invariant under G_ν for Dirac (Majorana) neutrinos

$$g_{\nu,j}^\dagger m_\nu m_\nu^\dagger g_{\nu,j} = m_\nu m_\nu^\dagger \quad , \quad j = 1, 2, \dots \quad \text{or} \quad g_{\nu,j}^T m_\nu g_{\nu,j} = m_\nu \quad , \quad j = 1, 2, \dots \quad (3)$$

For Majorana neutrinos G_ν is strongly constrained, since it has to be (a subgroup of) a Klein symmetry for three generations of neutrinos. If neutrinos are Dirac particles, G_ν is subject to the same constraint as G_e of the charged lepton sector, i.e. it can be any abelian group, capable of distinguishing the three generations. In our actual analysis we follow a minimality principle when choosing G_e and G_ν , since we consider all minimal

⁷The requirement of faithfulness of the three-dimensional representation also excludes the possibility to choose G_e or G_ν to be non-abelian, since a faithful representation of G_l usually decomposes into an irreducible representation of dimension larger than one in non-abelian G_e or G_ν so that a distinction among the three lepton generations becomes impossible.

⁸In abuse of notation we denote in the following the abstract elements of the groups G_e , G_ν and G_l with the same symbol as their matrix representatives.

(abelian) subgroups of the group G_l which allow the distinction of three generations.⁹ Furthermore, we wish to focus on mixing patterns which are associated with the group G_l and not only with one of its subgroups and thus we require that the generators of the subgroups G_e and G_ν give rise to the entire group G_l . We can change basis via the unitary transformations U_e and U_ν so that $g_{e,i}$ and $g_{\nu,j}$ are diagonalized ($g_{e(\nu),i(j)}^d$ are diagonal matrices), respectively,

$$U_e^\dagger g_{e,i} U_e = g_{e,i}^d \quad , \quad i = 1, 2, \dots \quad \text{and} \quad U_\nu^\dagger g_{\nu,j} U_\nu = g_{\nu,j}^d \quad , \quad j = 1, 2, \dots \quad (4)$$

Then, also $m_e m_e^\dagger$ and $m_\nu m_\nu^\dagger$ for Dirac (or m_ν for Majorana) neutrinos are diagonalized by U_e and U_ν , respectively, and the PMNS mixing matrix is derived as

$$U_{PMNS} = U_e^\dagger U_\nu \quad . \quad (5)$$

The matrices U_e and U_ν are uniquely determined up to permutations and phases of their column vectors, since we consider a scenario in which the three generations of left-handed leptons are distinguished, but no further assumption or prediction is made about the lepton masses. As a consequence, U_{PMNS} is fixed up to permutations of rows and columns and the multiplication with phase matrices from the left- and the right-hand side. In turn, the mixing angles θ_{ij} which only depend on the absolute values of the elements of the PMNS matrix $||U_{PMNS}||$, and the Jarlskog invariant J_{CP} which can be expressed as $J_{CP} = \text{Im}(U_{PMNS,11} U_{PMNS,13}^* U_{PMNS,31}^* U_{PMNS,33})$, take only values which belong to a small set (see [24] for our convention of mixing angles and CP phases). Note that an exchange of the roles of the subgroups G_e and G_ν transforms the PMNS matrix into its hermitian conjugate. Two pairs of subgroups, $\{G_e, G_\nu\}$ and $\{G'_e, G'_\nu\}$, lead to the same result for the PMNS matrix, if their generators $g_{e,i}$, $g_{\nu,j}$ and $g'_{e,i}$, $g'_{\nu,j}$ are related by a similarity transformation, i.e. if these pairs of groups are conjugate. Notice that two pairs $\{G_e, G_\nu\}$ and $\{G'_e, G'_\nu\}$ which are only isomorphic, but not conjugate, may or may not lead to different mixing patterns. In the case of the group $\Sigma(36 \times 3)$, for example, a combination $\{G_e, G_\nu\}$ of a certain type of subgroups gives, if admissible, always rise to the same mixing pattern, even if not all pairs $\{G_e, G_\nu\}$ are related by similarity transformations. In the case $G_l = \Sigma(216 \times 3)$, $G_e = Z_4$ and $G_\nu = Z_{18}$, see table 1 and 2, instead it is not sufficient to only specify the type of subgroups, since the combination $G_e = Z_4$ and $G_\nu = Z_{18}$ can give rise to pattern IV as well as to patterns IXa and IXb. If we exchange the roles of G_e and G_ν , i.e. $G_e = Z_{18}$ and $G_\nu = Z_4$, we see that also the patterns Ia and Ib in table 1 originate from the combination of these subgroups.

We also consider the case in which the generators of G_e or G_ν are represented by matrices with two degenerate eigenvalues. In this case only one generation can be distinguished from the other two ones and thus only one row or column can be fixed through the choice of G_e , G_ν and G_l . Such a row or column is called “mixing vector” [12].

Up to this point we have not fully specified the three-dimensional representation as which the three generations of left-handed leptons transform, apart from requesting that it should be irreducible and faithful in G_l . However, in general a group G_l possesses several representations which meet these requirements. If so, we have to analyze whether

⁹For example, if a subgroup $G_e = G_1$ is sufficient in order to distinguish among the generations, we do not discuss the case in which $G_e = G_1 \times G_2 \subset G_l$, since the presence of the group $G_2 \subset G_e$ would not add any information as far as lepton mixing is concerned.

the results for the mixing patterns do also depend on the choice of the three-dimensional representation and not only on the choice of the groups G_e , G_ν and G_l . We will show that in all groups $\Sigma(n\varphi)$ the mixing patterns are independent of the actual choice of the representation and that it is thus sufficient to study the mixing patterns for only one of them. In doing this, outer automorphisms are particularly useful, since they allow to relate inequivalent representations of a group.

3 Groups $\Sigma(n\varphi)$ and mixing patterns

The four groups we discuss in the following are $\Sigma(36 \times 3)$, $\Sigma(72 \times 3)$, $\Sigma(216 \times 3)$ and $\Sigma(360 \times 3)$. As we show, only the latter group has Klein subgroups, while the three other ones not. Thus, if G_ν is required to be a subgroup of G_l and neutrinos are Majorana particles, only $\Sigma(360 \times 3)$ can be chosen as G_l . In the other cases neutrinos are either Dirac particles or G_ν is only partly contained in G_l (i.e. one of the two Z_2 factors of the Klein group is accidental).

We first present a set of generators for each group together with the relations they have to satisfy. We then focus on the irreducible three-dimensional representations and discuss in some detail why it is sufficient for each of the four groups to only consider one of the representations when deriving the lepton mixing patterns. We detail the different abelian subgroups of the different $\Sigma(n\varphi)$ with $\varphi = 3$ and comment on those which are admissible as G_e and G_ν , because they are capable of distinguishing the three lepton generations (in our chosen three-dimensional representation), and on those which can only be employed in order to predict a mixing vector. As we will see, the groups $\Sigma(36 \times 3)$, $\Sigma(72 \times 3)$ and $\Sigma(360 \times 3)$ contain Z_2 and Z_6 generating elements which are represented by matrices with partly degenerate eigenvalues, while $\Sigma(216 \times 3)$ also comprises Z_9 generating elements with this property.

We show all patterns which can arise from a certain choice of G_l , G_e and G_ν , following our minimality principle for G_e and G_ν as well as imposing the requirement that the generators of these two groups give rise to the whole group G_l . As explained, the ordering of rows and columns is not fixed by the choice of the subgroups G_e and G_ν . We choose it in the presentation of our results in such a way that the value of χ^2 is minimized. The latter is computed in the standard way and with the experimental results for the mixing angles found in [3] in the left table of table 1. If all three mixing angles are in agreement with the experimental data at the 3σ level or better, the value of χ^2 is $\chi^2 \lesssim 27$ (the converse is in general not true and for this reason we always list the values of all three mixing angles). In several occasions we do not only show the pattern with rows and columns permuted in a way that χ^2 is minimized, but we also mention further patterns with a slightly larger χ^2 . The exact notion of “slightly larger” depends on the minimal value of χ^2 : for the latter being smaller than 100 we consider patterns with values of χ^2 up to 5% larger than the minimal one, while for a minimal value of χ^2 between 100 and 1000 we only mention patterns with values of χ^2 up to $1\% \div 2\%$ larger than the minimal one. In all cases we additionally require that at least one mixing angle agrees within 3σ with the experimental results [3]. If the minimal value of χ^2 is larger than 1000 we only show the pattern belonging to this value, unless there are two mixing angles which are in accordance with the experimental data within 3σ . If there are two permutations of

a pattern leading to a similar χ^2 and we display both, we denote them with the same Roman numeral, but a different letter, e.g. pattern Ia and Ib, as in table 1 for the group $\Sigma(216 \times 3)$. It turns out that in all cases the two permutations of a certain pattern are related through the exchange of the second and third rows of the PMNS matrix. Thus, they generally lead to the same reactor and solar mixing angles, while the atmospheric one changes from θ_{23} to $\pi/2 - \theta_{23}$, i.e. $\sin^2 \theta_{23}$ is replaced by $1 - \sin^2 \theta_{23}$. The tables 1, 2 and 4-6 summarize our results for the mixing patterns derived from the groups $\Sigma(216 \times 3)$ and $\Sigma(360 \times 3)$. We mark mixing angles that are in accordance with the experimental data at the 3σ level or better in these tables with a star (\star). For all patterns with $\chi^2 < 100$ we have checked that an agreement of the mixing angles with the experimental data at the 3σ level or better can be easily achieved in models with small corrections of order $\lambda^2 \approx 0.04$.¹⁰ Thus, we consider all these patterns as phenomenologically interesting.

3.1 Group $\Sigma(36 \times 3)$

The group $\Sigma(36 \times 3)$ (identification number [[108,15]] in the library SmallGroups [25] which is accessible via the computer algebra system GAP [26]) has 108 elements and can be represented in terms of three generators a , v and z which fulfill the following relations [27]

$$a^3 = 1, \quad v^4 = 1, \quad z^3 = 1, \quad av^{-1}zv = 1, \quad avz^{-1}v^{-1} = 1, \quad (az)^3 = 1. \quad (6)$$

Here and in the following, we denote the neutral element of the group with 1. For one of the irreducible faithful three-dimensional representations, identified with $\mathbf{3}^{(0)}$ of [27], the generators are given by the matrices

$$a = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad v = \frac{1}{\sqrt{3}i} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}, \quad z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \quad (7)$$

with $\omega = e^{2\pi i/3}$. Note that all generators have determinant +1. The group $\Sigma(36 \times 3)$ possesses in total eight faithful irreducible three-dimensional representations which form four complex conjugated pairs. The representation matrices of the generators a , v and z in the three other faithful three-dimensional representations $\mathbf{3}^{(p)}$, $p = 1, 2, 3$, are closely related to a , v and z of $\mathbf{3}^{(0)}$

$$a, \quad i^p v, \quad z \quad \text{with} \quad p = 1, 2, 3 \quad (8)$$

and those of (the complex conjugated representations) $(\mathbf{3}^{(p)})^*$, $p = 0, 1, 2, 3$, are obtained through complex conjugation. Apart from these, the irreducible representations of the group are: four one-dimensional representations, two real and two complex ones, and two real four-dimensional representations. Notice that they are all unfaithful in $\Sigma(36 \times 3)$. The character table of the group can be found, for example, in [27].

¹⁰We explicitly studied a scenario in which, due to an appropriate choice of basis, the lepton mixing arises at leading order only from the neutrino sector. Assuming that subleading corrections are present in the charged lepton sector which induce small mixing angles of order $\lambda^2 \approx 0.04$ there, all mixing angles can be accommodated within their 3σ ranges.

Clearly, the mixing patterns which can be derived by assuming that the three generations of leptons are assigned to any of the representations $\mathbf{3}^{(\mathbf{p})}$ have to coincide with those derived from assigning them to $\mathbf{3}^{(0)}$, since the additional factor i^p in the generator v cannot change the eigenvectors. Similarly, all mixing matrices originating from one of the complex conjugated representations $(\mathbf{3}^{(\mathbf{p})})^*$ are the same as those derived from $\mathbf{3}^{(\mathbf{p})}$, up to complex conjugation. In Appendix A.1 we comment on the outer automorphism group of $\Sigma(36 \times 3)$ and the possibility to understand relations among the three-dimensional representations using the latter. The products of two irreducible three-dimensional representations are of the following type

$$\mathbf{3}^{(\mathbf{p}_1)} \times \mathbf{3}^{(\mathbf{p}_2)} = (\mathbf{3}^{(\mathbf{p}_3)})^* + (\mathbf{3}^{(\mathbf{p}_4)})^* + (\mathbf{3}^{(\mathbf{p}_5)})^* \quad \text{and} \quad (\mathbf{3}^{(\mathbf{p}_1)}) \times (\mathbf{3}^{(\mathbf{p}_2)})^* = \mathbf{1}^{(\mathbf{q})} + \mathbf{4} + \mathbf{4}' , \quad (9)$$

with $p_i, q = 0, 1, 2, 3$, $i = 1, \dots, 5$, especially

$$\mathbf{3}^{(0)} \times \mathbf{3}^{(0)} = (\mathbf{3}^{(0)})^* + (\mathbf{3}^{(1)})^* + (\mathbf{3}^{(3)})^* \quad \text{and} \quad (\mathbf{3}^{(0)}) \times (\mathbf{3}^{(0)})^* = \mathbf{1}^{(0)} + \mathbf{4} + \mathbf{4}' . \quad (10)$$

Thus, this group does not belong to those singled out in the analysis of [22] as being a suitable candidate for the implementation of a version of minimal flavor violation with a discrete group. Nevertheless, we discuss the mixing patterns which can be derived from $\Sigma(36 \times 3)$, since its order is still rather small.

For analyzing all abelian subgroups of $\Sigma(36 \times 3)$ it is convenient to make use of the fact that all elements g of the group $\Sigma(36 \times 3)$ can be written, in the three-dimensional representations, in the form

$$g = \omega^o z^\zeta a^\alpha v^\chi \quad \text{with} \quad o, \zeta, \alpha = 0, 1, 2, \chi = 0, 1, 2, 3 \quad (11)$$

and that a representative (again in the three-dimensional representations) for each of the fourteen classes $c_1 \mathcal{C}_{c_2}$ (with c_1 denoting the number of elements of the class \mathcal{C}_{c_2} and c_2 their order) can be found [27]

$$\begin{aligned} 1 \mathcal{C}_1 : 1 , \quad 9 \mathcal{C}_2 : v^2 , \quad 1 \mathcal{C}_3 : aza^2z^2 = \omega 1 , \quad 1 \mathcal{C}_3 : a^2za^2z^2 = \omega^2 1 , \quad (12) \\ 12 \mathcal{C}_3 : z , \quad 12 \mathcal{C}_3 : za , \quad 9 \mathcal{C}_4 : v , \quad 9 \mathcal{C}_4 : v^3 , \quad 9 \mathcal{C}_6 : \omega v^2 , \\ 9 \mathcal{C}_6 : \omega^2 v^2 , \quad 9 \mathcal{C}_{12} : \omega v , \quad 9 \mathcal{C}_{12} : \omega^2 v , \quad 9 \mathcal{C}_{12} : \omega v^3 , \quad 9 \mathcal{C}_{12} : \omega^2 v^3 . \end{aligned}$$

The abelian subgroups of $\Sigma(36 \times 3)$ are nine Z_2 symmetries, 13 Z_3 symmetries (including C , the center of $SU(3)$), nine Z_4 and nine Z_6 symmetries, as well as nine Z_{12} groups. Furthermore, there are four groups $Z_3 \times Z_3$ where, however, one of the two Z_3 groups is the center of $SU(3)$. We have checked that all Z_2 symmetries, all Z_4 , all Z_6 and also all Z_{12} symmetries are conjugate to each other. The Z_3 symmetries fall instead into three categories: two with six such symmetries which are conjugate among each other and a third one which comprises only the center of $SU(3)$. Since $\Sigma(36 \times 3)$ does not contain a Klein group, neutrinos have to be either Dirac particles or the symmetry of the neutrino sector is only partly contained in G_l . Using the representation matrices in eq.(7) we see that 30 elements of $\Sigma(36 \times 3)$ are represented by matrices with (at least) two degenerate eigenvalues: the ones belonging to the center of $SU(3)$, the ones generating Z_2 groups as well as the ones generating Z_6 symmetries. Therefore these elements cannot be used as generators of the subgroups G_e and G_ν alone, if we wish to fix all three mixing angles through the breaking of G_l to G_e and to G_ν .

With this classification of subgroups of $\Sigma(36 \times 3)$ at hand we can distinguish the following cases in which the breaking of the group $\Sigma(36 \times 3)$ to G_e and to G_ν fixes the mixing pattern: $G_e = Z_3$ and $G_\nu = Z_3$, $G_e = Z_3$ and $G_\nu = Z_4$ or $G_\nu = Z_{12}$, $G_e = Z_4$ or $G_e = Z_{12}$ and $G_\nu = Z_4$ or $G_\nu = Z_{12}$. Out of these possibilities only two categories are interesting and can pass our constraint of generating the entire group $\Sigma(36 \times 3)$ with the generators of G_e and G_ν .¹¹ The first interesting case arises for $G_e = Z_3$ and $G_\nu = Z_4$ or $G_\nu = Z_{12}$. For both possibilities, $G_\nu = Z_4$ and $G_\nu = Z_{12}$, there are five independent choices of pairs $\{G_e, G_\nu\}$ whose sets of generators $\{g_e, g_\nu\}$ are not related among each other via similarity transformations. For one of them $G_e = Z_3$ is identified with the center of $SU(3)$ so that the three generations cannot be distinguished and thus this choice is not considered. The other four independent choices of generators instead fulfill all requirements, i.e. their generators allow for a distinction of the three generations and they generate together the original group $\Sigma(36 \times 3)$. All these admissible choices lead to the mixing pattern

$$||U_{PMNS}|| = \frac{1}{\sqrt{2(3+\sqrt{3})}} \begin{pmatrix} 1+\sqrt{3} & \sqrt{2} & 0 \\ 1 & \sqrt{2+\sqrt{3}} & \sqrt{3+\sqrt{3}} \\ 1 & \sqrt{2+\sqrt{3}} & \sqrt{3+\sqrt{3}} \end{pmatrix} \approx \begin{pmatrix} 0.888 & 0.460 & 0 \\ 0.325 & 0.628 & 0.707 \\ 0.325 & 0.628 & 0.707 \end{pmatrix}. \quad (13)$$

The mixing angles read $\sin^2 \theta_{12} \approx 0.211$, $\sin^2 \theta_{23} = 0.5$ and $\theta_{13} = 0$. Since the reactor mixing angle vanishes, also the Jarlskog invariant J_{CP} vanishes. The value of χ^2 is $\chi^2 \approx 151.5$ in this case and only the atmospheric mixing angle is in agreement with the experimental results within 3σ . Examples of generators of G_e and G_ν are $g_e = \omega^2 z a^2$ and $g_\nu = z a^2 v$ and $g_\nu = \omega z^2 a^2 v^3$, respectively. The pattern in eq.(13) has been recently found in a study in which the flavor group $\Delta(27)$ is combined with a CP symmetry [28].

If G_e and G_ν are instead both Z_4 or Z_{12} subgroups of $\Sigma(36 \times 3)$, we find that there exist three independent choices of pairs $\{G_e, G_\nu\}$ which are isomorphic, but not conjugate. One of these choices we have to exclude, because G_e and G_ν are the same or one is a subgroup of the other one, while the other two choices are admissible. We obtain for both of them as mixing pattern

$$||U_{PMNS}|| = \frac{1}{2\sqrt{2}} \begin{pmatrix} \sqrt{3+\sqrt{3}} & \sqrt{3} & \sqrt{2-\sqrt{3}} \\ \sqrt{2} & \sqrt{3-\sqrt{3}} & \sqrt{3+\sqrt{3}} \\ \sqrt{3-\sqrt{3}} & \sqrt{2+\sqrt{3}} & \sqrt{3} \end{pmatrix} \approx \begin{pmatrix} 0.769 & 0.612 & 0.183 \\ 0.500 & 0.398 & 0.769 \\ 0.398 & 0.683 & 0.612 \end{pmatrix} \quad (14)$$

leading to the mixing angles $\sin^2 \theta_{12} \approx 0.388$, $\sin^2 \theta_{23} \approx 0.612$ and $\sin^2 \theta_{13} \approx 0.033$. Again, J_{CP} vanishes. The value of χ^2 is 69.1. If the second and third rows of the PMNS matrix in eq.(14) are exchanged, the atmospheric mixing angle reads $\sin^2 \theta_{23} \approx 0.388$ like the solar one and the value of χ^2 slightly increases, $\chi^2 \approx 69.4$. In both cases only the atmospheric mixing angle agrees within 3σ with the experimental data. Examples for the generators of the groups G_e and G_ν are: if both are Z_4 subgroups, we can take $g_e = z a^2 v$ and $g_\nu = \omega^2 z^2 a v^3$; if G_e is a Z_4 and G_ν a Z_{12} subgroup, we can use $g_e = z a^2 v$ and $g_\nu = \omega z^2 a^2 v^3$ (we might also switch the roles of G_e and G_ν and still obtain the same mixing pattern); and for G_e and G_ν being both Z_{12} subgroups, we can choose for example

¹¹For the choice $G_e = Z_3$ and $G_\nu = Z_3$ we encounter only cases in which the mixing matrix is a permutation of the unit matrix or all its elements have the same absolute value or one of the subgroups G_e or G_ν is identified with the center of $SU(3)$ so that the three generations cannot be distinguished.

the generators $g_e = \omega z^2 a^2 v^3$ and $g_\nu = \omega^2 z v$. As one can see, all results which can be achieved with a Z_4 symmetry can also be obtained with an appropriate Z_{12} group. This happens, because all Z_{12} generating elements have a matrix representation which can be written as product of a matrix giving rise to a Z_4 generating element and a matrix representing a non-trivial element of the center of $SU(3)$.

Furthermore, we can consider the case in which one of the subgroups G_e or G_ν is generated by an element of $\Sigma(36 \times 3)$ which has two degenerate eigenvalues, i.e. G_e or G_ν is a Z_2 or Z_6 symmetry. The resulting mixing vectors can take only three possible forms (referring to the vectors with entries given in absolute values), namely a vector with only one non-vanishing entry, a vector with two equal entries and one vanishing one or a vector equal to the first column of the PMNS matrix shown in eq.(14). It is not by chance that the set of mixing vectors arising from Z_2 and Z_6 generating elements is the same, since one can check that all Z_6 generating elements are represented by matrices which are products of a matrix representing the generator of a Z_2 and a matrix which represents a non-trivial element of the center of $SU(3)$.

3.2 Group $\Sigma(72 \times 3)$

In order to generate the group $\Sigma(72 \times 3)$ which has the identification number [[216,88]] in the library SmallGroups we add one generator x to the set of generating elements a , v and z of the group $\Sigma(36 \times 3)$. This additional generator fulfills the relations [27]

$$x^2 = z^2 v^2 \quad \text{and} \quad x^4 = 1. \quad (15)$$

In the three-dimensional representation, called $\mathbf{3}^{(0,0)}$ in [27], the matrices of the generators can be chosen as a , v , z in eq.(7) together with

$$x = \frac{1}{\sqrt{3}i} \begin{pmatrix} 1 & 1 & \omega^2 \\ 1 & \omega & \omega \\ \omega & 1 & \omega \end{pmatrix}. \quad (16)$$

Notice also the matrix representative of the element x has determinant +1. This representation is one of the eight faithful irreducible three-dimensional representations of $\Sigma(72 \times 3)$ which form four complex conjugated pairs $\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)}$ and $(\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)})^*$ with $\mathbf{p}_{1,2} = 0, 1$. The representation matrices of the generators a , v , z and x in the three-dimensional representations $\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)}$ are given by

$$a, \quad (-1)^{\mathbf{p}_1} v, \quad (-1)^{\mathbf{p}_2} x, \quad z \quad \text{with} \quad \mathbf{p}_{1,2} = 0, 1 \quad (17)$$

and those of $(\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)})^*$ by the complex conjugated representation matrices. Like in the case of the three-dimensional representations of the group $\Sigma(36 \times 3)$ this shows that $\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)}$ and $(\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)})^*$, $\mathbf{p}_{1,2} = 0, 1$, all must lead to the same results for the mixing patterns. In Appendix A.2 we use the outer automorphisms of $\Sigma(72 \times 3)$ in order to relate the inequivalent three-dimensional representations. Apart from $\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)}$ and $(\mathbf{3}^{(\mathbf{p}_1, \mathbf{p}_2)})^*$ the group contains four one-dimensional, one two-dimensional, one eight-dimensional and two six-dimensional irreducible representations. The latter two are complex conjugated and faithful, while the former ones are all real and unfaithful. The character table of $\Sigma(72 \times 3)$ can be found in [27].

The generic form of the products among two three-dimensional representations is like in $SU(3)$, i.e.

$$\mathbf{3}^{(\mathbf{p}_{1,1}, \mathbf{p}_{2,1})} \times \mathbf{3}^{(\mathbf{p}_{1,2}, \mathbf{p}_{2,2})} = (\mathbf{3}^{(\mathbf{q}_1, \mathbf{q}_2)})^* + \mathbf{6}^* \quad \text{and} \quad \mathbf{3}^{(\mathbf{p}_{1,1}, \mathbf{p}_{2,1})} \times (\mathbf{3}^{(\mathbf{p}_{1,2}, \mathbf{p}_{2,2})})^* = \mathbf{1}^{(\mathbf{q}_1, \mathbf{q}_2)} + \mathbf{8}, \quad (18)$$

with $q_1 = p_{1,1} + p_{1,2} \bmod 2$ and $q_2 = p_{2,1} + p_{2,2} \bmod 2$, $p_{i,j} = 0, 1$, especially we have

$$\mathbf{3}^{(0,0)} \times \mathbf{3}^{(0,0)} = (\mathbf{3}^{(0,0)})^* + \mathbf{6}^* \quad \text{and} \quad \mathbf{3}^{(0,0)} \times (\mathbf{3}^{(0,0)})^* = \mathbf{1}^{(0,0)} + \mathbf{8}. \quad (19)$$

All elements g , in the three-dimensional representations, can be uniquely written in the form

$$g = \omega^o z^\zeta a^\alpha v^\chi x^\xi \quad \text{with} \quad o, \zeta, \alpha = 0, 1, 2, \quad \chi = 0, 1, 2, 3, \quad \xi = 0, 1. \quad (20)$$

Also here we display, for convenience, a representative (again in the three-dimensional representations) of each of the sixteen classes [27]

$$\begin{aligned} 1\mathcal{C}_1 : 1, \quad 9\mathcal{C}_2 : v^2, \quad 1\mathcal{C}_3 : aza^2z^2 = \omega 1, \quad 1\mathcal{C}_3 : a^2za^2z^2 = \omega^2 1, \quad (21) \\ 24\mathcal{C}_3 : z, \quad 18\mathcal{C}_4 : v, \quad 18\mathcal{C}_4 : x, \quad 18\mathcal{C}_4 : \omega^2 vx, \quad 9\mathcal{C}_6 : \omega v^2, \\ 9\mathcal{C}_6 : \omega^2 v^2, \quad 18\mathcal{C}_{12} : \omega v, \quad 18\mathcal{C}_{12} : \omega^2 v, \quad 18\mathcal{C}_{12} : \omega x, \quad 18\mathcal{C}_{12} : \omega^2 x, \\ 18\mathcal{C}_{12} : vx, \quad 18\mathcal{C}_{12} : \omega vx. \end{aligned}$$

Similarly to what we have seen for the representation $\mathbf{3}^{(0)}$ of the group $\Sigma(36 \times 3)$ also here 30 elements of the group are represented by matrices with at least two degenerate eigenvalues and clearly these coincide with those already found for $\Sigma(36 \times 3)$. The abelian subgroups of $\Sigma(72 \times 3)$ are: nine Z_2 symmetries, 13 Z_3 symmetries, 27 Z_4 groups, nine Z_6 symmetries, 27 Z_{12} groups as well as four $Z_3 \times Z_3$ groups. The Z_2 and Z_6 groups are all conjugate to each other, while the Z_3 symmetries form two categories (one with twelve members and the second one containing only the center of $SU(3)$). The Z_4 groups and the Z_{12} groups fall into three categories with nine members each which are conjugate to each other. In the following we will not use the four $Z_3 \times Z_3$ groups as G_e or G_ν , since one of the Z_3 factors is the center of $SU(3)$. As one can see, also $\Sigma(72 \times 3)$ does not possess Klein subgroups.

Considering all possible admissible choices for the groups G_e and G_ν we see that they have to be either both Z_4 subgroups or Z_{12} subgroups of $\Sigma(72 \times 3)$ or one is a Z_4 , while the other one is a Z_{12} subgroup. In all other cases the generators of G_e and G_ν cannot give rise to the entire group $\Sigma(72 \times 3)$. The mixing pattern arising from the admissible combinations is uniquely determined to be of the form

$$||U_{PMNS}|| = \frac{1}{2\sqrt{2}} \begin{pmatrix} \sqrt{3+\sqrt{3}} & \sqrt{4-\sqrt{3}} & 1 \\ \sqrt{2} & \sqrt{3+\sqrt{3}} & \sqrt{3-\sqrt{3}} \\ \sqrt{3-\sqrt{3}} & 1 & \sqrt{4+\sqrt{3}} \end{pmatrix} \approx \begin{pmatrix} 0.769 & 0.532 & 0.354 \\ 0.500 & 0.769 & 0.398 \\ 0.398 & 0.354 & 0.846 \end{pmatrix}. \quad (22)$$

For the mixing angles follows $\sin^2 \theta_{12} \approx 0.324$, $\sin^2 \theta_{23} \approx 0.181$ and $\sin^2 \theta_{13} = 0.125$. Consequently, the value of χ^2 is very large, $\chi^2 \approx 1984.1$, and only the solar mixing angle turns out to be within the 3σ range of the global fit [3]. Similar results are found, if the second and third rows of the PMNS matrix in eq.(22) are exchanged or the transpose (hermitian conjugate) of this matrix is considered as PMNS matrix. One interesting feature of this mixing matrix is that the Jarlskog invariant does not vanish:

$|J_{CP}| = \sqrt{3}/32 \approx 0.0541$. Examples of generators g_e and g_ν of the subgroups G_e and G_ν are: for both G_e and G_ν being Z_4 groups, $g_e = \omega^2 z^2 a^2 x$ and $g_\nu = z a^2 v$ can be chosen; for both G_e and G_ν being Z_{12} subgroups, we can take $g_e = z v x$ and $g_\nu = z^2 a v^2 x$, while for G_e being a Z_4 and G_ν a Z_{12} subgroup one possible combination of generators is $g_e = \omega^2 z^2 a^2 x$ and $g_\nu = z v x$. If G_e or G_ν is a Z_2 or a Z_6 subgroup, we can only fix one row or column of the PMNS matrix and the resulting mixing vectors coincide, as expected, with those found for the group $\Sigma(36 \times 3)$. The observations which have been made for the matrices representing Z_4 and Z_{12} and Z_2 and Z_6 generating elements, respectively, in $\Sigma(36 \times 3)$ are also valid in the case of the group $\Sigma(72 \times 3)$ and thus clearly the pattern in eq.(22) can be achieved with Z_4 or Z_{12} subgroups and the mixing vectors arising from matrices representing Z_2 or Z_6 generating elements have to be the same.

3.3 Group $\Sigma(216 \times 3)$

If we add the element w which fulfills the relations [27]

$$w^9 = 1, \quad w^{-1} a w = z a, \quad w^{-1} v^2 w = z v^2 \quad (23)$$

to the generators a , v and z of the group $\Sigma(36 \times 3)$, we arrive at the group $\Sigma(216 \times 3)$ (identification number [[648, 532]] in the library SmallGroups). Note that $x = w v w^{-1}$, see eqs.(15,16), is also an element of this group. For one of the faithful three-dimensional representations, denoted by $\mathbf{3}^{(0)}$ in [27], we choose the generators a , v , z and w to be represented by the matrices found in eq.(7) and

$$w = \begin{pmatrix} \epsilon & 0 & 0 \\ 0 & \epsilon & 0 \\ 0 & 0 & \epsilon \omega \end{pmatrix} \quad (24)$$

with $\epsilon = e^{4\pi i/9}$. Since $zw = wz$ holds, both generators z and w can be represented by diagonal matrices. Notice also that $w^3 = \omega^2 1$ in the three-dimensional representation $\mathbf{3}^{(0)}$. The group $\Sigma(216 \times 3)$ possesses in total seven irreducible three-dimensional representations: one of them, $\mathbf{3}^{(a)}$, is unfaithful and real, while the six other ones are faithful and form three pairs of complex conjugated representations $\mathbf{3}^{(p)}$ and $(\mathbf{3}^{(p)})^*$, $p = 0, 1, 2$. The representation matrices of the three faithful representations $\mathbf{3}^{(p)}$ can be chosen as

$$a, \quad v, \quad \omega^p w, \quad z \quad \text{with} \quad p = 0, 1, 2. \quad (25)$$

As is obvious from this, we can constrain ourselves to only consider the representation $\mathbf{3}^{(0)}$ in our analysis of the mixing patterns. Relations among the different irreducible and faithful three-dimensional representations can also be found by exploiting the outer automorphisms of the group $\Sigma(216 \times 3)$, as we show in Appendix A.3. Apart from the three-dimensional representations $\Sigma(216 \times 3)$ contains three one-, three two- and three eight-dimensional representations, which are organized as one real representation and one complex conjugated pair. All these representations are unfaithful, while the three complex conjugated pairs of six-dimensional and the pair of nine-dimensional representations are faithful in $\Sigma(216 \times 3)$.

The structure of the products of the faithful three-dimensional representations is similar to the one of the fundamental representation of $SU(3)$

$$\mathbf{3}^{(\mathbf{p}_1)} \times \mathbf{3}^{(\mathbf{p}_2)} = (\mathbf{3}^{(\mathbf{q}_1)})^* + (\mathbf{6}^{(\mathbf{q}_1)})^* \quad \text{and} \quad \mathbf{3}^{(\mathbf{p}_1)} \times (\mathbf{3}^{(\mathbf{p}_2)})^* = \mathbf{1}^{(\mathbf{q}_2)} + \mathbf{8}^{(\mathbf{q}_2)} \quad (26)$$

with $\mathbf{q}_1 = -\mathbf{p}_1 - \mathbf{p}_2 \bmod 3$, $\mathbf{q}_2 = \mathbf{p}_1 - \mathbf{p}_2 \bmod 3$ and $\mathbf{p}_{1,2} = 0, 1, 2$. Especially, we have

$$\mathbf{3}^{(0)} \times \mathbf{3}^{(0)} = (\mathbf{3}^{(0)})^* + (\mathbf{6}^{(0)})^* \quad \text{and} \quad \mathbf{3}^{(0)} \times (\mathbf{3}^{(0)})^* = \mathbf{1}^{(0)} + \mathbf{8}^{(0)}. \quad (27)$$

All elements g of the group $\Sigma(216 \times 3)$ can be written, at least in the faithful three-dimensional representations, as

$$g = \omega^o z^\zeta a^\alpha v^\chi x^\xi w^\kappa \quad \text{with} \quad o, \zeta, \alpha, \kappa = 0, 1, 2, \quad \chi = 0, 1, 2, 3, \quad \xi = 0, 1. \quad (28)$$

We mention one representative (in the faithful three-dimensional representations) for each of the 24 classes of this group [27]

$$\begin{aligned} 1\mathcal{C}_1 : 1, \quad 9\mathcal{C}_2 : v^2, \quad 1\mathcal{C}_3 : aza^2z^2 = \omega 1, \quad 1\mathcal{C}_3 : a^2zaz^2 = \omega^2 1, \quad (29) \\ 24\mathcal{C}_3 : z, \quad 72\mathcal{C}_3 : aw, \quad 72\mathcal{C}_3 : aw^2, \quad 54\mathcal{C}_4 : v, \quad 9\mathcal{C}_6 : \omega v^2, \\ 9\mathcal{C}_6 : \omega^2 v^2, \quad 12\mathcal{C}_9 : w, \quad 12\mathcal{C}_9 : \omega w, \quad 12\mathcal{C}_9 : \omega^2 w, \\ 12\mathcal{C}_9 : w^2, \quad 12\mathcal{C}_9 : \omega w^2, \quad 12\mathcal{C}_9 : \omega^2 w^2, \quad 54\mathcal{C}_{12} : \omega v, \\ 54\mathcal{C}_{12} : \omega^2 v, \quad 36\mathcal{C}_{18} : v^2 w, \quad 36\mathcal{C}_{18} : \omega v^2 w, \quad 36\mathcal{C}_{18} : \omega^2 v^2 w, \\ 36\mathcal{C}_{18} : vw^2, \quad 36\mathcal{C}_{18} : \omega vw^2, \quad 36\mathcal{C}_{18} : \omega^2 vw^2. \end{aligned}$$

Most of the abelian subgroups of $\Sigma(216 \times 3)$ form one category in which all members are conjugate to each other: the nine Z_2 subgroups, the 27 Z_4 groups, the nine Z_6 groups, the twelve Z_9 symmetries, the 27 Z_{12} groups as well as the 36 Z_{18} symmetries. Only the 85 Z_3 groups form instead three categories of conjugate groups: one which only contains the center of $SU(3)$, one with twelve members and the third one which comprises 72 Z_3 groups. Apart from these there are further abelian subgroups, $Z_3 \times Z_3$ and $Z_3 \times Z_9$, in which however one/the Z_3 factor is always given by the center of $SU(3)$ so that these groups are not relevant for our discussion. Notice also the group $\Sigma(216 \times 3)$ does not possess Klein subgroups and as consequence, neutrinos have to be either Dirac particles or G_ν can only be partly contained in $G_l = \Sigma(216 \times 3)$. 102 elements of this group are represented by matrices with at least two degenerate eigenvalues in the representation $\mathbf{3}^{(0)}$: these are the elements of the center of $SU(3)$, all Z_2 generating, all Z_6 as well as all Z_9 generating elements. All these cannot function as generators of the groups G_e and G_ν ; at least not alone, since the three generations of left-handed leptons cannot be distinguished in this way.

In table 1 and 2 on the next two pages we show the permutations of the mixing patterns with the smallest values of χ^2 for all combinations of subgroups G_e and G_ν which pass our constraints (i.e. their generators give rise to the entire group $\Sigma(216 \times 3)$ and they allow to distinguish the three generations).

Table 1: Mixing patterns associated with $\Sigma(216 \times 3)$. We mention the patterns, the subgroups G_e and G_ν , an example of their generators, the absolute value of J_{CP} , the sine squares of the mixing angles as well as the value of χ^2 . In all cases g_e and g_ν generate $\Sigma(216 \times 3)$. Note that we only give one example of g_e and g_ν . Analytic expressions of the patterns are given if sufficiently simple. We use as figure of merit χ^2 , computed using the results of the global fit in [3] in table 1 in which the reactor fluxes are left free and short baseline reactor data are included (called “Free Fluxes + RSBL”). We always display the permutation of the pattern with the minimum value of χ^2 . We show more than one permutation, if the χ^2 value of the further pattern(s) is very close to the minimum one. A star (\star) next to a result for a mixing angle $\sin^2 \theta_{ij}$ indicates that its contribution to the χ^2 value is less than 9 which is equivalent to being within the experimental 3σ interval. See the text for more details.

| # | Pattern | (G_e, G_ν) | Example of (g_e, g_ν) | $ J_{CP} $ | $(\sin^2 \theta_{12}, \sin^2 \theta_{23}, \sin^2 \theta_{13})$ | χ^2 |
|-----|--|---------------------------------------|---|------------|--|----------|
| Ia | $\frac{1}{2\sqrt{2(3+\sqrt{3})}} \begin{pmatrix} \sqrt{2}(2+\sqrt{3}) & \sqrt{2(3+\sqrt{3})} & -1+\sqrt{3} \\ \sqrt{6} & \sqrt{2(3+\sqrt{3})} & 3+\sqrt{3} \\ 2 & 2\sqrt{3+\sqrt{3}} & 2\sqrt{2+\sqrt{3}} \end{pmatrix} \approx \begin{pmatrix} 0.858 & 0.500 & 0.119 \\ 0.398 & 0.500 & 0.769 \\ 0.325 & 0.707 & 0.628 \end{pmatrix}$ | (Z_{18}, Z_4) (Z_{18}, Z_{12}) | $(\omega^2 zavxw, v)$ | 0 | $\sin^2 \theta_{12} \approx 0.254$ $\sin^2 \theta_{23} = 0.600 \star$ $\sin^2 \theta_{13} \approx 0.014$ | 28.3 |
| Ib | $\frac{1}{2\sqrt{2(3+\sqrt{3})}} \begin{pmatrix} \sqrt{2}(2+\sqrt{3}) & \sqrt{2(3+\sqrt{3})} & -1+\sqrt{3} \\ 2 & 2\sqrt{3+\sqrt{3}} & 2\sqrt{2+\sqrt{3}} \\ \sqrt{6} & \sqrt{2(3+\sqrt{3})} & 3+\sqrt{3} \end{pmatrix} \approx \begin{pmatrix} 0.858 & 0.500 & 0.119 \\ 0.325 & 0.707 & 0.628 \\ 0.398 & 0.500 & 0.769 \end{pmatrix}$ | (Z_{18}, Z_4) (Z_{18}, Z_{12}) | $(\omega^2 zavxw, v)$ | 0 | $\sin^2 \theta_{12} \approx 0.254$ $\sin^2 \theta_{23} = 0.400 \star$ $\sin^2 \theta_{13} \approx 0.014$ | 28.5 |
| II | $\frac{1}{2\sqrt{3}} \begin{pmatrix} 3 & \sqrt{3} & 0 \\ \sqrt{2} & \sqrt{6} & 2 \\ 1 & \sqrt{3} & 2\sqrt{2} \end{pmatrix} \approx \begin{pmatrix} 0.866 & 0.500 & 0 \\ 0.408 & 0.707 & 0.577 \\ 0.289 & 0.500 & 0.816 \end{pmatrix}$ | (Z_{18}, Z_{18}) | $(za^2 v^2 xw, zv^2 w)$ | 0 | $\sin^2 \theta_{12} = 0.250$ $\sin^2 \theta_{23} \approx 0.333$ $\sin^2 \theta_{13} = 0$ | 120.8 |
| III | $\begin{pmatrix} 0.844 & 0.525 & 0.110 \\ 0.449 & 0.804 & 0.390 \\ 0.293 & 0.279 & 0.914 \end{pmatrix}$ | (Z_3, Z_{18}) | $(\omega^2 z^2 av^3 xw, z^2 v^3 w)$ | 0 | $\sin^2 \theta_{12} \approx 0.279 \star$ $\sin^2 \theta_{23} \approx 0.154$ $\sin^2 \theta_{13} \approx 0.012$ | 131.2 |
| IV | $\frac{1}{2\sqrt{3+\sqrt{3}}} \begin{pmatrix} \sqrt{3(3+\sqrt{3})} & \sqrt{3+\sqrt{3}} & 0 \\ \sqrt{2+\sqrt{3}} & \sqrt{3(2+\sqrt{3})} & 2 \\ 1 & \sqrt{3} & 2\sqrt{2+\sqrt{3}} \end{pmatrix} \approx \begin{pmatrix} 0.866 & 0.500 & 0 \\ 0.444 & 0.769 & 0.460 \\ 0.230 & 0.398 & 0.888 \end{pmatrix}$ | (Z_4, Z_{18}) (Z_{12}, Z_{18}) | $(v, z^2 v^2 w)$ | 0 | $\sin^2 \theta_{12} = 0.250$ $\sin^2 \theta_{23} \approx 0.211$ $\sin^2 \theta_{13} = 0$ | 175.7 |
| V | $\begin{pmatrix} 0.804 & 0.578 & 0.139 \\ 0.525 & 0.800 & 0.290 \\ 0.279 & 0.160 & 0.947 \end{pmatrix}$ | (Z_3, Z_4) (Z_3, Z_{12}) | $(\omega^2 vw, v)$ | 0 | $\sin^2 \theta_{12} \approx 0.341$ $\sin^2 \theta_{23} \approx 0.086$ $\sin^2 \theta_{13} \approx 0.019 \star$ | 182.7 |
| VIa | $\begin{pmatrix} 0.874 & 0.429 & 0.228 \\ 0.429 & 0.621 & 0.657 \\ 0.228 & 0.657 & 0.719 \end{pmatrix}$ | (Z_3, Z_3) | $(\omega z^2 a^2 xw, \omega^2 z^2 av^3 xw)$ | 0.0321 | $\sin^2 \theta_{12} \approx 0.194$ $\sin^2 \theta_{23} \approx 0.455 \star$ $\sin^2 \theta_{13} \approx 0.052$ | 231.7 |
| VIb | $\begin{pmatrix} 0.874 & 0.429 & 0.228 \\ 0.228 & 0.657 & 0.719 \\ 0.429 & 0.621 & 0.657 \end{pmatrix}$ | (Z_3, Z_3) | $(\omega z^2 a^2 xw, \omega^2 z^2 av^3 xw)$ | 0.0321 | $\sin^2 \theta_{12} \approx 0.194$ $\sin^2 \theta_{23} \approx 0.545 \star$ $\sin^2 \theta_{13} \approx 0.052$ | 235.5 |

Table 2: Mixing patterns associated with $\Sigma(216 \times 3)$. For further details see the caption of table 1 and the text.

| # | Pattern | (G_e, G_ν) | Example of (g_e, g_ν) | $ J_{CP} $ | $(\sin^2 \theta_{12}, \sin^2 \theta_{23}, \sin^2 \theta_{13})$ | χ^2 |
|------|---|---------------------------------------|---------------------------------------|---------------------------------------|--|----------|
| VIIa | $\begin{pmatrix} 0.804 & 0.525 & 0.279 \\ 0.483 & 0.445 & 0.754 \\ 0.346 & 0.726 & 0.595 \end{pmatrix}$ | (Z_4, Z_3) (Z_{12}, Z_3) | $(v, \omega z^2 a^2 xw)$ | $\frac{1}{24} \approx 0.0417$ | $\sin^2 \theta_{12} \approx 0.299 \star$ $\sin^2 \theta_{23} \approx 0.616 \star$ $\sin^2 \theta_{13} \approx 0.078$ | 554.6 |
| VIIb | $\begin{pmatrix} 0.804 & 0.525 & 0.279 \\ 0.346 & 0.726 & 0.595 \\ 0.483 & 0.445 & 0.754 \end{pmatrix}$ | (Z_4, Z_3) (Z_{12}, Z_3) | $(v, \omega z^2 a^2 xw)$ | $\frac{1}{24} \approx 0.0417$ | $\sin^2 \theta_{12} \approx 0.299 \star$ $\sin^2 \theta_{23} \approx 0.384 \star$ $\sin^2 \theta_{13} \approx 0.078$ | 554.9 |
| VIII | $\begin{pmatrix} 0.804 & 0.525 & 0.279 \\ 0.517 & 0.723 & 0.458 \\ 0.293 & 0.449 & 0.844 \end{pmatrix}$ | (Z_{18}, Z_3) | $(za^2 v^2 xw, \omega^2 z^2 av^3 xw)$ | $\frac{1}{12\sqrt{3}} \approx 0.0481$ | $\sin^2 \theta_{12} \approx 0.299 \star$ $\sin^2 \theta_{23} \approx 0.227$ $\sin^2 \theta_{13} \approx 0.078$ | 608.7 |
| IXa | $\frac{1}{2\sqrt{6}} \begin{pmatrix} \sqrt{3(3+\sqrt{3})} & \sqrt{9-\sqrt{3}} & \sqrt{2(3-\sqrt{3})} \\ \sqrt{3(3-\sqrt{3})} & \sqrt{9+\sqrt{3}} & \sqrt{2(3+\sqrt{3})} \\ \sqrt{6} & \sqrt{6} & 2\sqrt{3} \end{pmatrix} \approx \begin{pmatrix} 0.769 & 0.550 & 0.325 \\ 0.398 & 0.669 & 0.628 \\ 0.500 & 0.500 & 0.707 \end{pmatrix}$ | (Z_4, Z_{18}) (Z_{12}, Z_{18}) | $(v, za^2 v^2 xw)$ | $\frac{1}{16} = 0.0625$ | $\sin^2 \theta_{12} \approx 0.339 \star$ $\sin^2 \theta_{23} \approx 0.441 \star$ $\sin^2 \theta_{13} \approx 0.106$ | 1255.5 |
| IXb | $\frac{1}{2\sqrt{6}} \begin{pmatrix} \sqrt{3(3+\sqrt{3})} & \sqrt{9-\sqrt{3}} & \sqrt{2(3-\sqrt{3})} \\ \sqrt{6} & \sqrt{6} & 2\sqrt{3} \\ \sqrt{3(3-\sqrt{3})} & \sqrt{9+\sqrt{3}} & \sqrt{2(3+\sqrt{3})} \end{pmatrix} \approx \begin{pmatrix} 0.769 & 0.550 & 0.325 \\ 0.500 & 0.500 & 0.707 \\ 0.398 & 0.669 & 0.628 \end{pmatrix}$ | (Z_4, Z_{18}) (Z_{12}, Z_{18}) | $(v, za^2 v^2 xw)$ | $\frac{1}{16} = 0.0625$ | $\sin^2 \theta_{12} \approx 0.339 \star$ $\sin^2 \theta_{23} \approx 0.559 \star$ $\sin^2 \theta_{13} \approx 0.106$ | 1257.5 |
| Xa | $\frac{1}{2\sqrt{3}} \begin{pmatrix} \sqrt{7} & \sqrt{3} & \sqrt{2} \\ \sqrt{3} & \sqrt{3} & \sqrt{6} \\ \sqrt{2} & \sqrt{6} & 2 \end{pmatrix} \approx \begin{pmatrix} 0.764 & 0.500 & 0.408 \\ 0.500 & 0.500 & 0.707 \\ 0.408 & 0.707 & 0.577 \end{pmatrix}$ | (Z_{18}, Z_{18}) | $(za^2 v^2 xw, za^2 v^3 w)$ | $\frac{1}{8\sqrt{3}} \approx 0.0722$ | $\sin^2 \theta_{12} = 0.300 \star$ $\sin^2 \theta_{23} = 0.600 \star$ $\sin^2 \theta_{13} \approx 0.167$ | 3753.2 |
| Xb | $\frac{1}{2\sqrt{3}} \begin{pmatrix} \sqrt{7} & \sqrt{3} & \sqrt{2} \\ \sqrt{2} & \sqrt{6} & 2 \\ \sqrt{3} & \sqrt{3} & \sqrt{6} \end{pmatrix} \approx \begin{pmatrix} 0.764 & 0.500 & 0.408 \\ 0.408 & 0.707 & 0.577 \\ 0.500 & 0.500 & 0.707 \end{pmatrix}$ | (Z_{18}, Z_{18}) | $(za^2 v^2 xw, za^2 v^3 w)$ | $\frac{1}{8\sqrt{3}} \approx 0.0722$ | $\sin^2 \theta_{12} = 0.300 \star$ $\sin^2 \theta_{23} = 0.400 \star$ $\sin^2 \theta_{13} \approx 0.167$ | 3753.4 |

As one can see, only the two patterns Ia and Ib agree well with the experimental results for all mixing angles, so that they can be considered as a good leading order form in a concrete model which only requires small corrections in order to achieve full accordance with the data. Three of the patterns lead to atmospheric and solar mixing angles which agree within 3σ with the experimental data (i.e. both these angles carry a star (\star)), while for the reactor mixing angle there is only one such pattern, pattern V. Patterns with smaller values of χ^2 , i.e. $\chi^2 \lesssim 200$, do not lead to a non-trivial Dirac phase, although all mixing angles are non-zero in the majority of the cases. On the other hand, patterns with larger χ^2 in general give rise to non-vanishing J_{CP} and thus CP violation. Interestingly, almost all patterns VIIa through Xb in table 2 accommodate both, solar and atmospheric, mixing angles well; however, the value of $\sin^2 \theta_{13}$ is too large (in all cases it is still at least a factor of two smaller than the other two $\sin^2 \theta_{ij}$). Notice that we find cases in which pairs of $\{G_e, G_\nu\}$ that are isomorphic, but not conjugate, do lead to different mixing patterns so that it is not sufficient to only specify the type of subgroups, e.g. the information $G_e = Z_4$ and $G_\nu = Z_{18}$ does not necessarily imply that we find the mixing pattern IV, because also the patterns IXa and IXb can arise from this combination of subgroups. If we furthermore consider the combinations in which the roles of G_e and G_ν are exchanged, the patterns Ia and Ib, see table 1, can result.

The pattern with the smallest value of χ^2 ($\chi^2 \approx 10.6$) is obtained for $G_e = Z_3$ and $G_\nu = Z_{18}$. We do not list this case in table 1 and 2, since the generators of G_e and G_ν do not give rise to the whole group $\Sigma(216 \times 3)$, but only to a group of order 162 with the mathematical structure $(Z_9 \times Z_3) \rtimes S_3$ [29]. It is denoted by $[[162, 14]]$ in the library SmallGroups. The mixing pattern reads

$$||U_{PMNS}|| = \frac{1}{\sqrt{6}} \begin{pmatrix} 2c_{18} & \sqrt{2} & 2s_{18} \\ c_{18} - \sqrt{3}s_{18} & \sqrt{2} & \sqrt{3}c_{18} + s_{18} \\ c_{18} + \sqrt{3}s_{18} & \sqrt{2} & \sqrt{3}c_{18} - s_{18} \end{pmatrix} \approx \begin{pmatrix} 0.804 & 0.577 & 0.142 \\ 0.279 & 0.577 & 0.767 \\ 0.525 & 0.577 & 0.625 \end{pmatrix} \quad (30)$$

where we have defined $s_{18} = \sin \pi/18 \approx 0.174$ and $c_{18} = \cos \pi/18 \approx 0.985$. The resulting mixing angles are: $\sin^2 \theta_{12} \approx 0.340$, $\sin^2 \theta_{23} \approx 0.601$ and $\sin^2 \theta_{13} \approx 0.020$ so that only the value of the solar mixing angle is marginally outside the 3σ range [3]. The Jarlskog invariant vanishes and thus the Dirac phase is trivial. A fit with a value of χ^2 only slightly larger than the optimal one, $\chi^2 \approx 10.8$, can be achieved, if the second and third rows are exchanged in the PMNS matrix in eq.(30). Then the atmospheric mixing angle is determined to be $\sin^2 \theta_{23} \approx 0.399$ which is also well within the 3σ range of [3]. Note that the PMNS matrix in eq.(30) can be written as the product of the TB mixing matrix and a rotation in the (13)-plane by the angle $\theta = -\pi/18$: $||U_{PMNS}|| = ||U_{TB}R_{13}(-\pi/18)||$. If we exchange the second and third rows in eq.(30), θ has to change sign, $\theta = \pi/18$.¹² Such type of pattern has already been found [6, 11, 13, 14], when studying groups belonging to the series $\Delta(6n^2)$, $n \in \mathbb{N}$. In [6, 11] it has been suggested that there might be a general rule, namely for $G_e = Z_3$, $G_\nu = Z_{2n}$ and $G_l = \Delta(6n^2)$ the angle θ reads $\theta = \pm\pi/(2n)$. In our case one would have to choose $n = 9$ and indeed the group $[[162, 14]]$ is a subgroup of $\Delta(6n^2)$ for $n = 9$. As example for the generators g_e and g_ν of G_e and G_ν , respectively, we can take $g_e = \omega^2 z^2 a v^3 x w$ and $g_\nu = a^2 v x w$. We note that the first column of the matrix in eq.(30) has been considered as mixing vector in [12], while the authors of [13]

¹²We use here the same conventions as used in [6, 11] for the signs in the TB mixing matrix U_{TB} and for the definition of the angle θ .

found the same mixing pattern in their scan for the group $(Z_{18} \times Z_6) \rtimes S_3$ (which has the identification number [[648,259]] in the library SmallGroups), assuming that neutrinos are Majorana particles and G_e is a Z_3 symmetry. Very recently, the same authors have performed a scan of smaller groups allowing neutrinos to be Dirac particles [15] and they also found the pattern in eq.(30) for the group with the identification number [[162,14]] in the library SmallGroups.

Assuming that one of the groups G_e or G_ν is generated by an element with order 2 or 6 which is represented by a matrix with two degenerate eigenvalues in the representation $\mathbf{3}^{(0)}$, we find as mixing vectors those which have already been found for the group $\Sigma(36 \times 3)$ and three additional ones that are: a vector which coincides (in absolute values) with the second column of pattern Ia found in table 1 and 2, a vector whose form is the one of the first row of pattern II and a vector with a form equal to the one of the second column of pattern III. The last mixing vector has also been considered in [12] and has been found for the groups with identification numbers [[162,14]] and [[486,61]] in the library SmallGroups, respectively. If we choose G_e or G_ν to be generated by an element of order 9, we find instead eight possible mixing vectors: a vector with only one non-vanishing entry and a vector with two equal entries (as regards the absolute value) and one vanishing one (these two are in common with the mixing vectors derived from $\Sigma(36 \times 3)$), a tri-maximal mixing vector, a vector with the same absolute values as the second and third rows of the TB mixing matrix, a vector which coincides (in absolute values) with the third row of pattern Ia, a vector equal to the third column of pattern II, a vector of the same form as the first column of pattern III and a vector equal to the third column of pattern IV. The vector coinciding with the first column of pattern III equals the columns, up to permutations, of a pattern mentioned in [28], in which the mixing matrices arising from the flavor group $\Delta(27)$ and a CP symmetry are analyzed.

As expected, we also find in our analysis of the patterns and mixing vectors of the group $\Sigma(216 \times 3)$ that Z_4 and Z_{12} subgroups lead to the same patterns, if chosen as G_e and/or G_ν , and that the representation matrices of Z_2 and Z_6 generating elements are related via the matrices representing the non-trivial elements of the center of $SU(3)$.

3.4 Group $\Sigma(360 \times 3)$

The last group we discuss is $\Sigma(360 \times 3)$ which has 1080 elements. Its identification number in the library SmallGroups is [[1080,260]]. We can define it in terms of four generators a , f , h and q . The relations they fulfill can be found in [20]

$$\begin{aligned} a^3 &= 1, \quad f^2 = 1, \quad h^2 = 1, \quad q^2 = 1, \\ (fq)^2 &= 1, \quad (ah)^2 = 1, \quad (af)^3 = 1, \quad (fh)^3 = 1, \quad (hq)^3 = 1, \quad (aq)^6 = 1. \end{aligned} \quad (31)$$

For an alternative set of generators (fulfilling a different set of relations) see also [20]. We mention its character table in table 3. This table is adapted from [30]. This group has four irreducible faithful three-dimensional representations which form two complex conjugated pairs. Apart from these representations $\Sigma(360 \times 3)$ has also a pair of complex conjugated six- and fifteen-dimensional representations which are faithful. Out of the three nine-dimensional representations only two are faithful which are complex, while the unfaithful one is real. The other six unfaithful representations have dimension one, five (two representations), eight (two representations) and ten and are all real.

Table 3: Character table of the group $\Sigma(360 \times 3)$, adapted from [30]. $c_1 \mathcal{C}_{c_2}$ denote the classes with c_1 elements which have order c_2 . G is a representative of the class $c_1 \mathcal{C}_{c_2}$ in terms of the generators a, f, h and q . Furthermore, $\rho = e^{2\pi i/5}$ and $\zeta = e^{2\pi i/15}$.

| | Classes | | | | | | | | | | | | | | | | |
|--------------------------------------|----------------------------|-----------------------------|----------------------------|----------------------------|------------------------------|------------------------------|-----------------------------|-----------------------------|-----------------------------|--------------------------------|--------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|
| | C_1 $1 \mathcal{C}_1$ | C_2 $45 \mathcal{C}_2$ | C_3 $1 \mathcal{C}_3$ | C_4 $1 \mathcal{C}_3$ | C_5 $120 \mathcal{C}_3$ | C_6 $120 \mathcal{C}_3$ | C_7 $90 \mathcal{C}_4$ | C_8 $72 \mathcal{C}_5$ | C_9 $72 \mathcal{C}_5$ | C_{10} $45 \mathcal{C}_6$ | C_{11} $45 \mathcal{C}_6$ | C_{12} $90 \mathcal{C}_{12}$ | C_{13} $90 \mathcal{C}_{12}$ | C_{14} $72 \mathcal{C}_{15}$ | C_{15} $72 \mathcal{C}_{15}$ | C_{16} $72 \mathcal{C}_{15}$ | C_{17} $72 \mathcal{C}_{15}$ |
| G | 1 | q | $(qa)^2$ | $(aqa)^2$ | a | hqa | fqh | $(hqa fq)^3$ | $(hqa fq)^6$ | qa | qa^2 | $a fq$ | $a^2 fq$ | $hqa fq$ | $(hqa fq)^2$ | $(hqa fq)^{13}$ | $(hqa fq)^{14}$ |
| 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 3⁽¹⁾ | 3 | -1 | 3ω | $3\omega^2$ | 0 | 0 | 1 | $-\rho^2 - \rho^3$ | $-\rho - \rho^4$ | $-\omega^2$ | $-\omega$ | ω^2 | ω | $-\zeta^2 - \zeta^8$ | $-\zeta - \zeta^4$ | $-\zeta^{11} - \zeta^{14}$ | $-\zeta^7 - \zeta^{13}$ |
| 3⁽²⁾ | 3 | -1 | 3ω | $3\omega^2$ | 0 | 0 | 1 | $-\rho - \rho^4$ | $-\rho^2 - \rho^3$ | $-\omega^2$ | $-\omega$ | ω^2 | ω | $-\zeta^{11} - \zeta^{14}$ | $-\zeta^7 - \zeta^{13}$ | $-\zeta^2 - \zeta^8$ | $-\zeta - \zeta^4$ |
| (3⁽¹⁾)[*] | 3 | -1 | $3\omega^2$ | 3ω | 0 | 0 | 1 | $-\rho^2 - \rho^3$ | $-\rho - \rho^4$ | $-\omega$ | $-\omega^2$ | ω | ω^2 | $-\zeta^7 - \zeta^{13}$ | $-\zeta^{11} - \zeta^{14}$ | $-\zeta - \zeta^4$ | $-\zeta^2 - \zeta^8$ |
| (3⁽²⁾)[*] | 3 | -1 | $3\omega^2$ | 3ω | 0 | 0 | 1 | $-\rho - \rho^4$ | $-\rho^2 - \rho^3$ | $-\omega$ | $-\omega^2$ | ω | ω^2 | $-\zeta - \zeta^4$ | $-\zeta^2 - \zeta^8$ | $-\zeta^7 - \zeta^{13}$ | $-\zeta^{11} - \zeta^{14}$ |
| 5⁽¹⁾ | 5 | 1 | 5 | 5 | 2 | -1 | -1 | 0 | 0 | 1 | 1 | -1 | -1 | 0 | 0 | 0 | 0 |
| 5⁽²⁾ | 5 | 1 | 5 | 5 | -1 | 2 | -1 | 0 | 0 | 1 | 1 | -1 | -1 | 0 | 0 | 0 | 0 |
| 6 | 6 | 2 | 6ω | $6\omega^2$ | 0 | 0 | 0 | 1 | 1 | $2\omega^2$ | 2ω | 0 | 0 | ω | ω^2 | ω | ω^2 |
| 6[*] | 6 | 2 | $6\omega^2$ | 6ω | 0 | 0 | 0 | 1 | 1 | 2ω | $2\omega^2$ | 0 | 0 | ω^2 | ω | ω^2 | ω |
| 8⁽¹⁾ | 8 | 0 | 8 | 8 | -1 | -1 | 0 | $-\rho^2 - \rho^3$ | $-\rho - \rho^4$ | 0 | 0 | 0 | 0 | $-\rho - \rho^4$ | $-\rho^2 - \rho^3$ | $-\rho^2 - \rho^3$ | $-\rho - \rho^4$ |
| 8⁽²⁾ | 8 | 0 | 8 | 8 | -1 | -1 | 0 | $-\rho - \rho^4$ | $-\rho^2 - \rho^3$ | 0 | 0 | 0 | 0 | $-\rho^2 - \rho^3$ | $-\rho - \rho^4$ | $-\rho - \rho^4$ | $-\rho^2 - \rho^3$ |
| 9⁽¹⁾ | 9 | 1 | 9 | 9 | 0 | 0 | 1 | -1 | -1 | 1 | 1 | 1 | 1 | -1 | -1 | -1 | -1 |
| 9⁽²⁾ | 9 | 1 | 9ω | $9\omega^2$ | 0 | 0 | 1 | -1 | -1 | ω^2 | ω | ω^2 | ω | $-\omega$ | $-\omega^2$ | $-\omega$ | $-\omega^2$ |
| (9⁽²⁾)[*] | 9 | 1 | $9\omega^2$ | 9ω | 0 | 0 | 1 | -1 | -1 | ω | ω^2 | ω | ω^2 | $-\omega^2$ | $-\omega$ | $-\omega^2$ | $-\omega$ |
| 10 | 10 | -2 | 10 | 10 | 1 | 1 | 0 | 0 | 0 | -2 | -2 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 | 15 | -1 | 15ω | $15\omega^2$ | 0 | 0 | -1 | 0 | 0 | $-\omega^2$ | $-\omega$ | $-\omega^2$ | $-\omega$ | 0 | 0 | 0 | 0 |
| 15[*] | 15 | -1 | $15\omega^2$ | 15ω | 0 | 0 | -1 | 0 | 0 | $-\omega$ | $-\omega^2$ | $-\omega$ | $-\omega^2$ | 0 | 0 | 0 | 0 |

The form of the product of $\mathbf{3}^{(1)}$ with itself and with its complex conjugated representation is like in $SU(3)$ ¹³

$$\mathbf{3}^{(1)} \times \mathbf{3}^{(1)} = (\mathbf{3}^{(1)})^* + \mathbf{6}^* \quad \text{and} \quad \mathbf{3}^{(1)} \times (\mathbf{3}^{(1)})^* = \mathbf{1} + \mathbf{8}^{(1)}. \quad (32)$$

The generators of the group can be represented by the following matrices in the representation $\mathbf{3}^{(1)}$: a can be chosen as in eq.(7) and

$$f = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad h = \frac{1}{2} \begin{pmatrix} -1 & \mu_- & \mu_+ \\ \mu_- & \mu_+ & -1 \\ \mu_+ & -1 & \mu_- \end{pmatrix}, \quad q = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -\omega \\ 0 & -\omega^2 & 0 \end{pmatrix} \quad (33)$$

with $\mu_{\pm} = \frac{1}{2}(-1 \pm \sqrt{5})$. All these matrices have determinant +1. A set of representation matrices for the other three-dimensional representations $\mathbf{3}^{(2)}$, $(\mathbf{3}^{(1)})^*$ and $(\mathbf{3}^{(2)})^*$ can be constructed by using outer automorphisms of the group $\Sigma(360 \times 3)$ which relate the other representations and their matrix realizations to $\mathbf{3}^{(1)}$ and a, f, h and q , see eqs. (7,33),

$$\begin{aligned} \mathbf{3}^{(1)} \rightarrow \mathbf{3}^{(2)} &: \quad a \rightarrow ahq, \quad f \rightarrow fahf, \quad h \rightarrow q, \quad q \rightarrow a^2h, \\ \mathbf{3}^{(1)} \rightarrow (\mathbf{3}^{(1)})^* &: \quad a \rightarrow a^2hq, \quad f \rightarrow fa^2hf, \quad h \rightarrow q, \quad q \rightarrow ah, \\ \mathbf{3}^{(1)} \rightarrow (\mathbf{3}^{(2)})^* &: \quad a \rightarrow a^2, \quad f \rightarrow f, \quad h \rightarrow h, \quad q \rightarrow q. \end{aligned} \quad (34)$$

These outer automorphisms can be nicely seen as symmetries of the character table: the first one exchanges the representations $\mathbf{3}^{(1)}$ and $\mathbf{3}^{(2)}$ as well as their complex conjugates and $\mathbf{8}^{(1)}$ with $\mathbf{8}^{(2)}$ and at the same time the classes: $C_8 \leftrightarrow C_9$, $C_{14} \leftrightarrow C_{16}$ and $C_{15} \leftrightarrow C_{17}$, while the second one exchanges all representations with their complex conjugates, i.e. $\mathbf{3}^{(1)}$ with $(\mathbf{3}^{(1)})^*$, $\mathbf{3}^{(2)}$ with $(\mathbf{3}^{(2)})^*$, $\mathbf{6}$ and $\mathbf{6}^*$, $\mathbf{9}^{(2)}$ with $(\mathbf{9}^{(2)})^*$ and $\mathbf{15}$ with $\mathbf{15}^*$ and it also exchanges the classes: $C_3 \leftrightarrow C_4$, $C_{10} \leftrightarrow C_{11}$, $C_{12} \leftrightarrow C_{13}$, $C_{14} \leftrightarrow C_{17}$ and $C_{15} \leftrightarrow C_{16}$. The third automorphism exchanges the representations $\mathbf{3}^{(1)}$ and $(\mathbf{3}^{(2)})^*$ as well as their complex conjugates, $\mathbf{6}$ with $\mathbf{6}^*$, the two real eight-dimensional representations $\mathbf{8}^{(1)}$ and $\mathbf{8}^{(2)}$, the complex conjugated pair of nine-dimensional representations as well as the fifteen-dimensional representations; at the same time the following classes have to be exchanged: $C_3 \leftrightarrow C_4$, $C_8 \leftrightarrow C_9$, $C_{10} \leftrightarrow C_{11}$, $C_{12} \leftrightarrow C_{13}$, $C_{14} \leftrightarrow C_{15}$ and $C_{16} \leftrightarrow C_{17}$. The group of outer automorphisms is a Klein group $Z_2 \times Z_2$. Thus, the mentioned mappings have order two and commute among each other, as can be checked.

The abelian subgroups of $\Sigma(360 \times 3)$ which are generated by a single element are 45 Z_2 groups, 121 Z_3 symmetries, 45 Z_4 symmetries, 36 Z_5 groups, 45 Z_6 symmetries, 45 Z_{12} groups as well as 36 Z_{15} symmetries. All these Z_n subgroups are conjugate among each other apart from the Z_3 symmetries which form three categories (one of which contains only the center of $SU(3)$, a second one with 60 conjugate members as well as a third one with also 60 members). There are 30 Klein groups which form two categories comprising 15 conjugate groups each. There are also abelian subgroups of the form $Z_3 \times Z_3$ and $Z_2 \times Z_6$ which however contain the center of $SU(3)$ and thus are not relevant for our discussion.

The patterns we find for $\Sigma(360 \times 3)$ are listed in table 4-6. Patterns Ia through XVIII all arise from subgroups G_e and G_ν which are generated by a single element each and

¹³The form of the products is very similar for $\mathbf{3}^{(2)}$. The combinations $\mathbf{3}^{(1)} \times \mathbf{3}^{(2)}$ and $\mathbf{3}^{(1)} \times (\mathbf{3}^{(2)})^*$ and alike are instead irreducible in $\Sigma(360 \times 3)$.

thus require neutrinos to be Dirac particles. The only exception are the patterns XIa and XIb which can also be achieved, if both groups G_e and G_ν are Klein symmetries. Furthermore, there are several patterns, called KI through KVII, that are generated, if one of the two residual symmetries, G_e or G_ν , is a Klein group. Indeed for all patterns apart from KI the permutation with the minimum value of χ^2 is compatible with G_ν being a Klein group. Thus, neutrinos can naturally be Majorana particles. Overall there are only a few patterns with a value of χ^2 smaller than 100: patterns Ia through IIIb. Unfortunately, all these predict vanishing CP violation. This is very similar to what we have already observed in the case of the group $\Sigma(216 \times 3)$, see table 1 and 2. The pattern with the smallest χ^2 ($\chi^2 \approx 338.0$) and with non-vanishing J_{CP} is pattern VII which, however, does not fit any of the three mixing angles at the 3σ level or better. Pattern XVIII turns out to be the only one which leads to maximal CP violation, $|\sin \delta| = 1$, as we indicate with a circle (o) in table 6. Although CP violation is not maximal in the other cases, we frequently find patterns with $|\sin \delta| \gtrsim 0.9$. Almost half of the patterns Ia through XVIII fits the atmospheric mixing angle well, while none leads to a reactor mixing angle which is compatible at the 3σ level with the results found in [3]. On top of that only pattern Ia is able to accommodate two of the three mixing angles well. Among the patterns Ki, $i=I, \dots, VII$, the smallest value of χ^2 is $\chi^2 \approx 148.1$ which is larger than that of the patterns Ia to IIIb. Although pattern KI has $\chi^2 \approx 148.1$ and only fits the solar mixing angle well, it might be interesting, since J_{CP} is non-vanishing in this case. This is at variance to what we have found for the patterns which are produced in setups with G_e and G_ν being both generated by one element each. It is likely to be a coincidence, but indeed all patterns Ki, $i=I, \dots, VII$ lead to non-zero J_{CP} . Two of these, pattern KV and KVI, have also been mentioned in [31]. Generally speaking, all patterns would require additional ingredients to be present which allow them to be compatible with the experimental data [3]. These might be suitable corrections coming from contributions to the mass matrices of charged leptons and neutrinos which are not invariant under G_e and G_ν , respectively, in a concrete model or modifications of the breaking pattern, e.g. by involving CP [17, 18]. As one can see from tables 4-6, the choice of Z_4 or Z_{12} subgroups and of Z_5 or Z_{15} subgroups, respectively, leads to the same mixing patterns. This is, like in the other groups $\Sigma(n\varphi)$ with $\varphi = 3$, due to the relation of these subgroups via the center of $SU(3)$.

As one can check there are 138 matrices among the representation matrices in $\mathbf{3}^{(1)}$ which have at least two degenerate eigenvalues: these are all Z_2 and all Z_6 generating elements as well as the elements which comprise the center of $SU(3)$. The former can be identified with the generators of G_e or G_ν and then give rise to one of the fourteen different mixing vectors: the three vectors which have been found in the analysis of the group $\Sigma(36 \times 3)$, a vector of the same form (in absolute values) as the first column of the TB mixing matrix, a vector which is tri-maximal, a vector which equals the first row of pattern KI (see table 6), another one of the same form of the third row of pattern KI, a vector equal to the second column of pattern KII, another one which is of the same form as the third column of pattern KII, a vector which coincides with the third column of pattern KIII, vectors like the second and the third columns of pattern KIVa, a vector with the same form as the third column of pattern KV as well as a vector which coincides with the second column of pattern KVI.

Table 4: Mixing patterns associated with $\Sigma(360 \times 3)$. For more details see the caption of table 1 and the text.

| # | Pattern | (G_e, G_ν) | Example of (g_e, g_ν) | $ J_{CP} $ | $(\sin^2 \theta_{12}, \sin^2 \theta_{23}, \sin^2 \theta_{13})$ | χ^2 |
|------|--|--|--------------------------------|---|--|----------|
| Ia | $\frac{1}{4} \begin{pmatrix} \frac{1+\sqrt{5}}{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}} & \frac{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}}{\sqrt{8+\sqrt{3}-\sqrt{15}}} & \frac{\sqrt{5+\sqrt{3}-\sqrt{5}-\sqrt{15}}}{\sqrt{3+\sqrt{5}}} \\ \frac{\sqrt{5+\sqrt{3}-\sqrt{5}-\sqrt{15}}}{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}} & \frac{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}}{\sqrt{3+\sqrt{5}}} & \frac{\sqrt{5+\sqrt{3}-\sqrt{5}-\sqrt{15}}}{\sqrt{8-\sqrt{3}+\sqrt{15}}} \end{pmatrix} \approx \begin{pmatrix} 0.809 & 0.554 & 0.197 \\ 0.554 & 0.605 & 0.572 \\ 0.197 & 0.572 & 0.796 \end{pmatrix}$ | (Z_4, Z_4) (Z_4, Z_{12}) (Z_{12}, Z_{12}) | $(f q h, a f h q a^2)$ | 0 | $\sin^2 \theta_{12} \approx 0.319 \star$ $\sin^2 \theta_{23} \approx 0.341 \star$ $\sin^2 \theta_{13} \approx 0.039$ | 58.0 |
| Ib | $\frac{1}{4} \begin{pmatrix} \frac{1+\sqrt{5}}{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}} & \frac{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}}{\sqrt{8+\sqrt{3}-\sqrt{15}}} & \frac{\sqrt{5+\sqrt{3}-\sqrt{5}-\sqrt{15}}}{\sqrt{3+\sqrt{5}}} \\ \frac{\sqrt{5+\sqrt{3}-\sqrt{5}-\sqrt{15}}}{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}} & \frac{\sqrt{5-\sqrt{3}-\sqrt{5}+\sqrt{15}}}{\sqrt{8-\sqrt{3}+\sqrt{15}}} & \frac{\sqrt{5+\sqrt{3}-\sqrt{5}-\sqrt{15}}}{\sqrt{3+\sqrt{5}}} \end{pmatrix} \approx \begin{pmatrix} 0.809 & 0.554 & 0.197 \\ 0.197 & 0.572 & 0.796 \\ 0.554 & 0.605 & 0.572 \end{pmatrix}$ | (Z_4, Z_4) (Z_4, Z_{12}) (Z_{12}, Z_{12}) | $(f q h, a f h q a^2)$ | 0 | $\sin^2 \theta_{12} \approx 0.319 \star$ $\sin^2 \theta_{23} \approx 0.659$ $\sin^2 \theta_{13} \approx 0.039$ | 58.9 |
| IIa | $\frac{1}{20} \begin{pmatrix} 5-\sqrt{5}+\sqrt{30(5+\sqrt{5})} & 4\sqrt{5} & 2\sqrt{5(7-\sqrt{5}-\sqrt{6(5-\sqrt{5})})} \\ 4\sqrt{5} & -5+\sqrt{5}+\sqrt{30(5+\sqrt{5})} & 2\sqrt{5(7-\sqrt{5}+\sqrt{6(5-\sqrt{5})})} \\ 2\sqrt{5(7-\sqrt{5}-\sqrt{6(5-\sqrt{5})})} & 2\sqrt{5(7-\sqrt{5}+\sqrt{6(5-\sqrt{5})})} & 2(5+\sqrt{5}) \end{pmatrix} \approx \begin{pmatrix} 0.875 & 0.447 & 0.186 \\ 0.447 & 0.598 & 0.665 \\ 0.186 & 0.665 & 0.724 \end{pmatrix}$ | (Z_5, Z_5) (Z_5, Z_{15}) (Z_{15}, Z_{15}) | $((h q a f q)^3, a^2 f q h)$ | 0 | $\sin^2 \theta_{12} \approx 0.207$ $\sin^2 \theta_{23} \approx 0.458 \star$ $\sin^2 \theta_{13} \approx 0.035$ | 84.6 |
| IIb | $\frac{1}{20} \begin{pmatrix} 5-\sqrt{5}+\sqrt{30(5+\sqrt{5})} & 4\sqrt{5} & 2\sqrt{5(7-\sqrt{5}-\sqrt{6(5-\sqrt{5})})} \\ 2\sqrt{5(7-\sqrt{5}-\sqrt{6(5-\sqrt{5})})} & 2\sqrt{5(7-\sqrt{5}+\sqrt{6(5-\sqrt{5})})} & 2(5+\sqrt{5}) \\ 4\sqrt{5} & -5+\sqrt{5}+\sqrt{30(5+\sqrt{5})} & 2\sqrt{5(7-\sqrt{5}+\sqrt{6(5-\sqrt{5})})} \end{pmatrix} \approx \begin{pmatrix} 0.875 & 0.447 & 0.186 \\ 0.186 & 0.665 & 0.724 \\ 0.447 & 0.598 & 0.665 \end{pmatrix}$ | (Z_5, Z_5) (Z_5, Z_{15}) (Z_{15}, Z_{15}) | $((h q a f q)^3, a^2 f q h)$ | 0 | $\sin^2 \theta_{12} \approx 0.207$ $\sin^2 \theta_{23} \approx 0.542 \star$ $\sin^2 \theta_{13} \approx 0.035$ | 88.9 |
| IIIa | $\begin{pmatrix} 0.894 & 0.430 & 0.124 \\ 0.250 & 0.710 & 0.659 \\ 0.372 & 0.558 & 0.742 \end{pmatrix}$ | (Z_5, Z_4) (Z_{15}, Z_4) (Z_5, Z_{12}) (Z_{15}, Z_{12}) | $((h q a f q)^3, a^2 f h q a)$ | 0 | $\sin^2 \theta_{12} \approx 0.188$ $\sin^2 \theta_{23} \approx 0.441 \star$ $\sin^2 \theta_{13} \approx 0.015$ | 93.0 |
| IIIb | $\begin{pmatrix} 0.894 & 0.430 & 0.124 \\ 0.372 & 0.558 & 0.742 \\ 0.250 & 0.710 & 0.659 \end{pmatrix}$ | (Z_5, Z_4) (Z_{15}, Z_4) (Z_5, Z_{12}) (Z_{15}, Z_{12}) | $((h q a f q)^3, a^2 f h q a)$ | 0 | $\sin^2 \theta_{12} \approx 0.188$ $\sin^2 \theta_{23} \approx 0.559 \star$ $\sin^2 \theta_{13} \approx 0.015$ | 95.0 |
| IV | $\frac{1}{20} \begin{pmatrix} 2\sqrt{5(7+\sqrt{5}+\sqrt{6(5+\sqrt{5})})} & 4\sqrt{5} & -5-\sqrt{5}+\sqrt{30(5-\sqrt{5})} \\ 2\sqrt{5(7+\sqrt{5}-\sqrt{6(5+\sqrt{5})})} & 5+\sqrt{5}+\sqrt{30(5-\sqrt{5})} & 4\sqrt{5} \\ 2(5-\sqrt{5}) & 2\sqrt{5(7+\sqrt{5}-\sqrt{6(5+\sqrt{5})})} & 2\sqrt{5(7+\sqrt{5}+\sqrt{6(5+\sqrt{5})})} \end{pmatrix} \approx \begin{pmatrix} 0.890 & 0.447 & 0.093 \\ 0.364 & 0.817 & 0.447 \\ 0.276 & 0.364 & 0.890 \end{pmatrix}$ | (Z_5, Z_5) (Z_5, Z_{15}) (Z_{15}, Z_{15}) | $((h q a f q)^3, q h q a f h)$ | 0 | $\sin^2 \theta_{12} \approx 0.202$ $\sin^2 \theta_{23} \approx 0.202$ $\sin^2 \theta_{13} \approx 0.009$ | 171.0 |
| V | $\begin{pmatrix} 0.795 & 0.602 & 0.076 \\ 0.590 & 0.739 & 0.324 \\ 0.139 & 0.302 & 0.943 \end{pmatrix}$ | (Z_5, Z_4) (Z_{15}, Z_4) (Z_5, Z_{12}) (Z_{15}, Z_{12}) | $((h q a f q)^3, f q h)$ | 0 | $\sin^2 \theta_{12} \approx 0.364$ $\sin^2 \theta_{23} \approx 0.106$ $\sin^2 \theta_{13} \approx 0.006$ | 227.7 |
| VI | $\frac{1}{2\sqrt{5}} \begin{pmatrix} \sqrt{2(5+\sqrt{15})} & \sqrt{2(5-\sqrt{15})} & 0 \\ \sqrt{5-\sqrt{15}} & \sqrt{5+\sqrt{15}} & \sqrt{10} \\ \sqrt{5-\sqrt{15}} & \sqrt{5+\sqrt{15}} & \sqrt{10} \end{pmatrix} \approx \begin{pmatrix} 0.942 & 0.336 & 0 \\ 0.237 & 0.666 & 0.707 \\ 0.237 & 0.666 & 0.707 \end{pmatrix}$ | (Z_5, Z_4) (Z_{15}, Z_4) (Z_5, Z_{12}) (Z_{15}, Z_{12}) | $((h q a f q)^3, q a f a^2)$ | 0 | $\sin^2 \theta_{12} \approx 0.113$ $\sin^2 \theta_{23} \approx 0.500 \star$ $\sin^2 \theta_{13} = 0$ | 328.2 |
| VII | $\frac{1}{6} \begin{pmatrix} 2\sqrt{3+\sqrt{5}} & \sqrt{2(9-\sqrt{5})} & -1+\sqrt{5} \\ -1+\sqrt{5} & -1+\sqrt{5} & 2\sqrt{6+\sqrt{5}} \\ \sqrt{2(9-\sqrt{5})} & 2\sqrt{3+\sqrt{5}} & -1+\sqrt{5} \end{pmatrix} \approx \begin{pmatrix} 0.763 & 0.613 & 0.206 \\ 0.206 & 0.206 & 0.957 \\ 0.613 & 0.763 & 0.206 \end{pmatrix}$ | (Z_3, Z_3) | $(h q a, h f)$ | $\frac{1}{108} \sqrt{6(3-\sqrt{5})} \approx 0.0198$ | $\sin^2 \theta_{12} \approx 0.392$ $\sin^2 \theta_{23} \approx 0.956$ $\sin^2 \theta_{13} \approx 0.042$ | 338.0 |

Table 5: Mixing patterns associated with $\Sigma(360 \times 3)$. Note that patterns XIa and XIb can also be achieved, if both G_e and G_ν are Klein subgroups of $\Sigma(360 \times 3)$. For further explanation see the caption of table 4.

| # | Pattern | (G_e, G_ν) | Example of (g_e, g_ν) | $ J_{CP} $ | $(\sin^2 \theta_{12}, \sin^2 \theta_{23}, \sin^2 \theta_{13})$ | χ^2 |
|--------------------------------------|---|---|--|---|--|----------|
| VIII $\frac{1}{4}$ | $\begin{pmatrix} \frac{\sqrt{5+\sqrt{3}+\sqrt{5}+\sqrt{15}}}{-1+\sqrt{5}} & \frac{\sqrt{8-\sqrt{6(3+\sqrt{5})}}}{\sqrt{5+\sqrt{3}+\sqrt{5}+\sqrt{15}}} & \frac{\sqrt{3-\sqrt{5}}}{\sqrt{5-\sqrt{3}+\sqrt{5}-\sqrt{15}}} \\ \frac{\sqrt{5-\sqrt{3}+\sqrt{5}-\sqrt{15}}}{\sqrt{3-\sqrt{5}}} & \frac{\sqrt{8+\sqrt{6(3+\sqrt{5})}}}{\sqrt{3-\sqrt{5}}} \end{pmatrix} \approx \begin{pmatrix} 0.896 & 0.387 & 0.219 \\ 0.309 & 0.896 & 0.319 \\ 0.319 & 0.219 & 0.922 \end{pmatrix}$ | (Z_4, Z_4) (Z_4, Z_{12}) (Z_{12}, Z_{12}) | $(f q h, a^2 f h q a)$ | 0 | $\sin^2 \theta_{12} \approx 0.157$ $\sin^2 \theta_{23} \approx 0.107$ $\sin^2 \theta_{13} \approx 0.048$ | 397.6 |
| IX $\frac{1}{2\sqrt{3(3-\sqrt{5})}}$ | $\begin{pmatrix} \frac{\sqrt{11-\sqrt{3}-\sqrt{5}(3-\sqrt{3})}}{\sqrt{11+\sqrt{3}-\sqrt{5}(3+\sqrt{3})}} & \frac{\sqrt{11+\sqrt{3}-\sqrt{5}(3+\sqrt{3})}}{\sqrt{11-\sqrt{3}-\sqrt{5}(3-\sqrt{3})}} & \frac{3-\sqrt{5}}{3-\sqrt{5}} \\ \frac{3-\sqrt{5}}{3-\sqrt{5}} & \frac{3-\sqrt{5}}{2\sqrt{2}} \end{pmatrix} \approx \begin{pmatrix} 0.838 & 0.484 & 0.252 \\ 0.484 & 0.838 & 0.252 \\ 0.252 & 0.252 & 0.934 \end{pmatrix}$ | (Z_4, Z_3) (Z_{12}, Z_3) | $(f q h, q h a)$ | $\frac{1}{48}(\sqrt{5}-1) \approx 0.0258$ | $\sin^2 \theta_{12} \approx 0.251$ $\sin^2 \theta_{23} \approx 0.068$ $\sin^2 \theta_{13} \approx 0.064$ | 511.2 |
| Xa $\frac{1}{6}$ | $\begin{pmatrix} \frac{\sqrt{2(9+\sqrt{5})}}{1+\sqrt{5}} & \frac{1+\sqrt{5}}{2\sqrt{3-\sqrt{5}}} & \frac{2\sqrt{3-\sqrt{5}}}{1+\sqrt{5}} \\ \frac{2\sqrt{3-\sqrt{5}}}{1+\sqrt{5}} & \frac{1+\sqrt{5}}{\sqrt{2(9+\sqrt{5})}} \end{pmatrix} \approx \begin{pmatrix} 0.790 & 0.539 & 0.291 \\ 0.539 & 0.647 & 0.539 \\ 0.291 & 0.539 & 0.790 \end{pmatrix}$ | (Z_3, Z_3) | $(h f, h a q)$ | $\frac{1}{36\sqrt{3}}(1+\sqrt{5}) \approx 0.0519$ | $\sin^2 \theta_{12} \approx 0.318 \star$ $\sin^2 \theta_{23} \approx 0.318$ $\sin^2 \theta_{13} \approx 0.085$ | 716.2 |
| Xb $\frac{1}{6}$ | $\begin{pmatrix} \frac{\sqrt{2(9+\sqrt{5})}}{2\sqrt{3-\sqrt{5}}} & \frac{1+\sqrt{5}}{1+\sqrt{5}} & \frac{2\sqrt{3-\sqrt{5}}}{\sqrt{2(9+\sqrt{5})}} \\ \frac{2\sqrt{3-\sqrt{5}}}{1+\sqrt{5}} & \frac{1+\sqrt{5}}{2\sqrt{6-\sqrt{5}}} & \frac{1+\sqrt{5}}{1+\sqrt{5}} \end{pmatrix} \approx \begin{pmatrix} 0.790 & 0.539 & 0.291 \\ 0.291 & 0.539 & 0.790 \\ 0.539 & 0.647 & 0.539 \end{pmatrix}$ | (Z_3, Z_3) | $(h f, h a q)$ | $\frac{1}{36\sqrt{3}}(1+\sqrt{5}) \approx 0.0519$ | $\sin^2 \theta_{12} \approx 0.318 \star$ $\sin^2 \theta_{23} \approx 0.682$ $\sin^2 \theta_{13} \approx 0.085$ | 718.6 |
| XIa $\frac{1}{4}$ | $\begin{pmatrix} \frac{1+\sqrt{5}}{2} & \frac{2}{2\sqrt{2}} & \frac{-1+\sqrt{5}}{2} \\ \frac{2}{-1+\sqrt{5}} & \frac{2}{2} & \frac{1+\sqrt{5}}{1+\sqrt{5}} \end{pmatrix} \approx \begin{pmatrix} 0.809 & 0.500 & 0.309 \\ 0.500 & 0.707 & 0.500 \\ 0.309 & 0.500 & 0.809 \end{pmatrix}$ | (Z_4, Z_4) (Z_4, Z_{12}) (Z_{12}, Z_{12}) $(Z_2 \times Z_2, Z_2 \times Z_2)$ | $(f q h, q a f a^2)$ $(\{f q, h f q h\}, \{a f a^2, a q a^2\})$ | $\frac{\sqrt{3}}{32} \approx 0.0541$ | $\sin^2 \theta_{12} \approx 0.276 \star$ $\sin^2 \theta_{23} \approx 0.276$ $\sin^2 \theta_{13} \approx 0.095$ | 993.5 |
| XIb $\frac{1}{4}$ | $\begin{pmatrix} \frac{1+\sqrt{5}}{-1+\sqrt{5}} & \frac{2}{2\sqrt{2}} & \frac{-1+\sqrt{5}}{2} \\ \frac{2}{-1+\sqrt{5}} & \frac{2}{2\sqrt{2}} & \frac{1+\sqrt{5}}{2} \end{pmatrix} \approx \begin{pmatrix} 0.809 & 0.500 & 0.309 \\ 0.309 & 0.500 & 0.809 \\ 0.500 & 0.707 & 0.500 \end{pmatrix}$ | (Z_4, Z_4) (Z_4, Z_{12}) (Z_{12}, Z_{12}) $(Z_2 \times Z_2, Z_2 \times Z_2)$ | $(f q h, q a f a^2)$ $(\{f q, h f q h\}, \{a f a^2, a q a^2\})$ | $\frac{\sqrt{3}}{32} \approx 0.0541$ | $\sin^2 \theta_{12} \approx 0.276 \star$ $\sin^2 \theta_{23} \approx 0.724$ $\sin^2 \theta_{13} \approx 0.095$ | 1000.0 |
| XII $\frac{1}{2\sqrt{15}}$ | $\begin{pmatrix} \frac{\sqrt{25+\sqrt{5}+\sqrt{30(5-\sqrt{5})}}}{\sqrt{25+\sqrt{5}-\sqrt{30(5-\sqrt{5})}}} & \frac{\sqrt{25+\sqrt{5}-\sqrt{30(5-\sqrt{5})}}}{\sqrt{25+\sqrt{5}+\sqrt{30(5-\sqrt{5})}}} & \frac{\sqrt{2(5-\sqrt{5})}}{\sqrt{2(5-\sqrt{5})}} \\ \frac{\sqrt{2(5-\sqrt{5})}}{\sqrt{2(5-\sqrt{5})}} & \frac{\sqrt{2(5-\sqrt{5})}}{\sqrt{2(5-\sqrt{5})}} & \frac{\sqrt{2(5-\sqrt{5})}}{2\sqrt{10+\sqrt{5}}} \end{pmatrix} \approx \begin{pmatrix} 0.778 & 0.550 & 0.304 \\ 0.550 & 0.778 & 0.304 \\ 0.304 & 0.304 & 0.903 \end{pmatrix}$ | (Z_5, Z_3) (Z_{15}, Z_3) | $((h q a f q)^3, q h)$ | $\frac{1}{60}\sqrt{2(5-\sqrt{5})} \approx 0.0392$ | $\sin^2 \theta_{12} \approx 0.333 \star$ $\sin^2 \theta_{23} \approx 0.101$ $\sin^2 \theta_{13} \approx 0.092$ | 1034.3 |
| XIII $\frac{1}{\sqrt{12}}$ | $\begin{pmatrix} \frac{\sqrt{5+\sqrt{15}}}{\sqrt{5-\sqrt{15}}} & \frac{\sqrt{2}}{2\sqrt{2}} & \frac{\sqrt{5-\sqrt{15}}}{\sqrt{5+\sqrt{15}}} \\ \frac{\sqrt{2}}{\sqrt{5-\sqrt{15}}} & \frac{2\sqrt{2}}{\sqrt{2}} & \frac{\sqrt{2}}{\sqrt{5+\sqrt{15}}} \end{pmatrix} \approx \begin{pmatrix} 0.860 & 0.408 & 0.306 \\ 0.408 & 0.816 & 0.408 \\ 0.306 & 0.408 & 0.860 \end{pmatrix}$ | (Z_4, Z_3) (Z_{12}, Z_3) | $(f q h, a f h f a)$ | $\frac{1}{24} \approx 0.0417$ | $\sin^2 \theta_{12} \approx 0.184$ $\sin^2 \theta_{23} \approx 0.184$ $\sin^2 \theta_{13} \approx 0.094$ | 1091.6 |
| XIV $\frac{1}{\sqrt{6(3-\sqrt{5})}}$ | $\begin{pmatrix} \frac{\sqrt{2}}{\sqrt{8-3\sqrt{5}+\sqrt{3(9-4\sqrt{5})}}} & \frac{\sqrt{2}}{\sqrt{8-3\sqrt{5}-\sqrt{3(9-4\sqrt{5})}}} & \frac{3-\sqrt{5}}{\sqrt{2}} \\ \frac{\sqrt{2}}{\sqrt{8-3\sqrt{5}-\sqrt{3(9-4\sqrt{5})}}} & \frac{\sqrt{2}}{\sqrt{8-3\sqrt{5}+\sqrt{3(9-4\sqrt{5})}}} & \frac{\sqrt{2}}{\sqrt{2}} \end{pmatrix} \approx \begin{pmatrix} 0.661 & 0.661 & 0.357 \\ 0.609 & 0.439 & 0.661 \\ 0.439 & 0.609 & 0.661 \end{pmatrix}$ | (Z_4, Z_3) (Z_{12}, Z_3) | $(f q h, h q a)$ | $\frac{1}{48}(1+\sqrt{5}) \approx 0.0674$ | $\sin^2 \theta_{12} = 0.500$ $\sin^2 \theta_{23} \approx 0.500 \star$ $\sin^2 \theta_{13} \approx 0.127$ | 2238.5 |
| XV $\frac{1}{\sqrt{6(5+\sqrt{5})}}$ | $\begin{pmatrix} \frac{\sqrt{2(6+\sqrt{5}+\sqrt{15+6\sqrt{5}})}}{1+\sqrt{5}} & \frac{1+\sqrt{5}}{\sqrt{2(9+\sqrt{5})}} & \frac{\sqrt{2(6+\sqrt{5}-\sqrt{15+6\sqrt{5}})}}{1+\sqrt{5}} \\ \frac{1+\sqrt{5}}{\sqrt{2(6+\sqrt{5}-\sqrt{15+6\sqrt{5}})}} & \frac{1+\sqrt{5}}{1+\sqrt{5}} & \frac{\sqrt{2(6+\sqrt{5}+\sqrt{15+6\sqrt{5}})}}{1+\sqrt{5}} \end{pmatrix} \approx \begin{pmatrix} 0.791 & 0.491 & 0.366 \\ 0.491 & 0.719 & 0.491 \\ 0.366 & 0.491 & 0.791 \end{pmatrix}$ | (Z_5, Z_3) (Z_{15}, Z_3) | $((h q a f q)^3, h a q)$ | $\frac{1}{60}\sqrt{2(5+\sqrt{5})} \approx 0.0634$ | $\sin^2 \theta_{12} \approx 0.278 \star$ $\sin^2 \theta_{23} \approx 0.278$ $\sin^2 \theta_{13} \approx 0.134$ | 2269.1 |
| XVI $\frac{1}{\sqrt{2(5+\sqrt{5})}}$ | $\begin{pmatrix} \frac{\sqrt{4+\sqrt{5}+\sqrt{5+2\sqrt{5}}}}{\sqrt{4+\sqrt{5}-\sqrt{5+2\sqrt{5}}}} & \frac{\sqrt{4+\sqrt{5}-\sqrt{5+2\sqrt{5}}}}{\sqrt{4+\sqrt{5}+\sqrt{5+2\sqrt{5}}}} & \frac{\sqrt{2}}{\sqrt{2}} \\ \frac{\sqrt{4+\sqrt{5}-\sqrt{5+2\sqrt{5}}}}{\sqrt{2}} & \frac{\sqrt{4+\sqrt{5}+\sqrt{5+2\sqrt{5}}}}{\sqrt{2}} & \frac{\sqrt{2}}{1+\sqrt{5}} \end{pmatrix} \approx \begin{pmatrix} 0.802 & 0.467 & 0.372 \\ 0.467 & 0.802 & 0.372 \\ 0.372 & 0.372 & 0.851 \end{pmatrix}$ | (Z_5, Z_4) (Z_{15}, Z_4) (Z_5, Z_{12}) (Z_{15}, Z_{12}) | $((h q a f q)^3, a^2 f q h a)$ | $\frac{1}{80}\sqrt{6(5-\sqrt{5})} \approx 0.0509$ | $\sin^2 \theta_{12} \approx 0.253$ $\sin^2 \theta_{23} \approx 0.160$ $\sin^2 \theta_{13} \approx 0.138$ | 2532.8 |

Table 6: Mixing patterns associated with $\Sigma(360 \times 3)$. Patterns called Ki require that one of the groups G_e or G_ν is a Klein group. Notice that only pattern KI requires G_e to be a Klein group instead of G_ν . The circle (o) next to the result of $|J_{CP}|$ for pattern XVIII indicates that this value corresponds to maximal CP violation $|\sin \delta| = 1$. For further explanation see table 4.

| # | Pattern | (G_e, G_ν) | Example of (g_e, g_ν) | $ J_{CP} $ | $(\sin^2 \theta_{12}, \sin^2 \theta_{23}, \sin^2 \theta_{13})$ | χ^2 |
|-------|--|--|------------------------------------|---|--|----------|
| XVII | $\frac{1}{2\sqrt{5+\sqrt{5}}} \begin{pmatrix} \sqrt{7+\sqrt{5}+\sqrt{2(5+\sqrt{5})}} & 1+\sqrt{5} & \sqrt{7+\sqrt{5}-\sqrt{2(5+\sqrt{5})}} \\ 1+\sqrt{5} & 2\sqrt{2} & 1+\sqrt{5} \\ \sqrt{7+\sqrt{5}-\sqrt{2(5+\sqrt{5})}} & 1+\sqrt{5} & \sqrt{7+\sqrt{5}+\sqrt{2(5+\sqrt{5})}} \end{pmatrix} \approx \begin{pmatrix} 0.671 & 0.602 & 0.433 \\ 0.602 & 0.526 & 0.602 \\ 0.433 & 0.602 & 0.671 \end{pmatrix}$ | (Z_5, Z_4) (Z_{15}, Z_4) (Z_5, Z_{12}) (Z_{15}, Z_{12}) | $((hqa fq)^3, af hqa^2)$ | $\frac{1}{80} \sqrt{6(5+\sqrt{5})} \approx 0.0824$ | $\sin^2 \theta_{12} \approx 0.445$ $\sin^2 \theta_{23} \approx 0.445 \star$ $\sin^2 \theta_{13} \approx 0.188$ | 5060.1 |
| XVIII | $\frac{1}{\sqrt{6}} \begin{pmatrix} \sqrt{3} & 1 & 1 \\ 1 & \sqrt{2} & \sqrt{2} \\ 1 & \sqrt{2} & \sqrt{2} \end{pmatrix} \approx \begin{pmatrix} 0.775 & 0.447 & 0.447 \\ 0.447 & 0.632 & 0.632 \\ 0.447 & 0.632 & 0.632 \end{pmatrix}$ | (Z_5, Z_5) (Z_5, Z_{15}) (Z_{15}, Z_{15}) | $((hqa fq)^3, qha fq)$ | $\frac{\sqrt{15}}{80} \approx 0.0775$ o | $\sin^2 \theta_{12} = 0.250$ $\sin^2 \theta_{23} = 0.500 \star$ $\sin^2 \theta_{13} = 0.200$ | 5715.1 |
| KI | $\frac{1}{4} \begin{pmatrix} 1+\sqrt{5} & \sqrt{5+\sqrt{15}-\sqrt{2(4+\sqrt{15})}} & \sqrt{5+\sqrt{3}-\sqrt{5(1+\sqrt{3})}} \\ 2 & \sqrt{2(3+\sqrt{3})} & \sqrt{2(3-\sqrt{3})} \\ -1+\sqrt{5} & \sqrt{5-\sqrt{3}+\sqrt{5(1-\sqrt{3})}} & \sqrt{5+\sqrt{15}+\sqrt{2(4+\sqrt{15})}} \end{pmatrix} \approx \begin{pmatrix} 0.809 & 0.554 & 0.197 \\ 0.500 & 0.769 & 0.398 \\ 0.309 & 0.319 & 0.896 \end{pmatrix}$ | $(Z_2 \times Z_2, Z_4)$ $(Z_2 \times Z_2, Z_{12})$ | $(\{f, q\}, af qha^2)$ | $\frac{1}{32} \approx 0.0313$ | $\sin^2 \theta_{12} \approx 0.319 \star$ $\sin^2 \theta_{23} \approx 0.165$ $\sin^2 \theta_{13} \approx 0.039$ | 148.1 |
| KII | $\frac{1}{\sqrt{2(5+\sqrt{5})}} \begin{pmatrix} \sqrt{4+\sqrt{5}+\sqrt{15+6\sqrt{5}}} & \sqrt{2} & \sqrt{4+\sqrt{5}-\sqrt{15+6\sqrt{5}}} \\ \sqrt{2} & 1+\sqrt{5} & \sqrt{2} \\ \sqrt{4+\sqrt{5}-\sqrt{15+6\sqrt{5}}} & \sqrt{2} & \sqrt{4+\sqrt{5}+\sqrt{15+6\sqrt{5}}} \end{pmatrix} \approx \begin{pmatrix} 0.894 & 0.372 & 0.250 \\ 0.372 & 0.851 & 0.372 \\ 0.250 & 0.372 & 0.894 \end{pmatrix}$ | $(Z_5, Z_2 \times Z_2)$ $(Z_{15}, Z_2 \times Z_2)$ | $((hqa fq)^3, \{a^2 fa, a^2 qa\})$ | $\frac{1}{80} \sqrt{2(5-\sqrt{5})} \approx 0.0294$ | $\sin^2 \theta_{12} \approx 0.147$ $\sin^2 \theta_{23} \approx 0.147$ $\sin^2 \theta_{13} \approx 0.063$ | 553.5 |
| KIII | $\frac{\sqrt{2}}{\sqrt{3(1+\sqrt{5})}} \begin{pmatrix} \sqrt{7+3\sqrt{5}} & 1 & 1 \\ 1 & \sqrt{7+3\sqrt{5}} & 1 \\ 1 & 1 & \sqrt{7+3\sqrt{5}} \end{pmatrix} \approx \begin{pmatrix} 0.934 & 0.252 & 0.252 \\ 0.252 & 0.934 & 0.252 \\ 0.252 & 0.252 & 0.934 \end{pmatrix}$ | $(Z_3, Z_2 \times Z_2)$ | $(hafa, \{f, q\})$ | $\frac{1}{144} \sqrt{6(3-\sqrt{5})} \approx 0.0149$ | $\sin^2 \theta_{12} \approx 0.068$ $\sin^2 \theta_{23} \approx 0.068$ $\sin^2 \theta_{13} \approx 0.064$ | 844.7 |
| KIVa | $\frac{1}{2\sqrt{5+\sqrt{5}}} \begin{pmatrix} \sqrt{7+\sqrt{5}+\sqrt{6(5+\sqrt{5})}} & 1+\sqrt{5} & \sqrt{7+\sqrt{5}-\sqrt{6(5+\sqrt{5})}} \\ \sqrt{7+\sqrt{5}-\sqrt{6(5+\sqrt{5})}} & 1+\sqrt{5} & \sqrt{7+\sqrt{5}+\sqrt{6(5+\sqrt{5})}} \\ 1+\sqrt{5} & 2\sqrt{2} & 1+\sqrt{5} \end{pmatrix} \approx \begin{pmatrix} 0.739 & 0.602 & 0.302 \\ 0.302 & 0.602 & 0.739 \\ 0.602 & 0.526 & 0.602 \end{pmatrix}$ | $(Z_5, Z_2 \times Z_2)$ $(Z_{15}, Z_2 \times Z_2)$ | $((hqa fq)^3, \{afa^2, aqa^2\})$ | $\frac{1}{80} \sqrt{2(5+\sqrt{5})} \approx 0.0476$ | $\sin^2 \theta_{12} \approx 0.398$ $\sin^2 \theta_{23} \approx 0.602 \star$ $\sin^2 \theta_{13} \approx 0.091$ | 915.3 |
| KIVb | $\frac{1}{2\sqrt{5+\sqrt{5}}} \begin{pmatrix} \sqrt{7+\sqrt{5}+\sqrt{6(5+\sqrt{5})}} & 1+\sqrt{5} & \sqrt{7+\sqrt{5}-\sqrt{6(5+\sqrt{5})}} \\ 1+\sqrt{5} & 2\sqrt{2} & 1+\sqrt{5} \\ \sqrt{7+\sqrt{5}-\sqrt{6(5+\sqrt{5})}} & 1+\sqrt{5} & \sqrt{7+\sqrt{5}+\sqrt{6(5+\sqrt{5})}} \end{pmatrix} \approx \begin{pmatrix} 0.739 & 0.602 & 0.302 \\ 0.602 & 0.526 & 0.602 \\ 0.302 & 0.602 & 0.739 \end{pmatrix}$ | $(Z_5, Z_2 \times Z_2)$ $(Z_{15}, Z_2 \times Z_2)$ | $((hqa fq)^3, \{afa^2, aqa^2\})$ | $\frac{1}{80} \sqrt{2(5+\sqrt{5})} \approx 0.0476$ | $\sin^2 \theta_{12} \approx 0.398$ $\sin^2 \theta_{23} \approx 0.398 \star$ $\sin^2 \theta_{13} \approx 0.091$ | 915.5 |
| KV | $\frac{1}{\sqrt{3(3-\sqrt{5})}} \begin{pmatrix} 1 & 1 & \sqrt{7-3\sqrt{5}} \\ 1 & \sqrt{7-3\sqrt{5}} & 1 \\ \sqrt{7-3\sqrt{5}} & 1 & 1 \end{pmatrix} \approx \begin{pmatrix} 0.661 & 0.661 & 0.357 \\ 0.661 & 0.357 & 0.661 \\ 0.357 & 0.661 & 0.661 \end{pmatrix}$ | $(Z_3, Z_2 \times Z_2)$ | $(af ha^2, \{f, q\})$ | $\frac{1}{144} \sqrt{6(3+\sqrt{5})} \approx 0.0389$ | $\sin^2 \theta_{12} = 0.500$ $\sin^2 \theta_{23} = 0.500 \star$ $\sin^2 \theta_{13} \approx 0.127$ | 2238.5 |
| KVI | $\frac{1}{2\sqrt{2}} \begin{pmatrix} \sqrt{3+\sqrt{3}} & \sqrt{2} & \sqrt{3-\sqrt{3}} \\ \sqrt{2} & 2 & \sqrt{2} \\ \sqrt{3-\sqrt{3}} & \sqrt{2} & \sqrt{3+\sqrt{3}} \end{pmatrix} \approx \begin{pmatrix} 0.769 & 0.500 & 0.398 \\ 0.500 & 0.707 & 0.500 \\ 0.398 & 0.500 & 0.769 \end{pmatrix}$ | $(Z_4, Z_2 \times Z_2)$ $(Z_{12}, Z_2 \times Z_2)$ | $(fqh, \{f, afa^2\})$ | $\frac{\sqrt{5}}{32} \approx 0.0699$ | $\sin^2 \theta_{12} \approx 0.297 \star$ $\sin^2 \theta_{23} \approx 0.297$ $\sin^2 \theta_{13} \approx 0.158$ | 3360.7 |
| KVII | $\frac{1}{\sqrt{6}} \begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{pmatrix} \approx \begin{pmatrix} 0.816 & 0.408 & 0.408 \\ 0.408 & 0.816 & 0.408 \\ 0.408 & 0.408 & 0.816 \end{pmatrix}$ | $(Z_3, Z_2 \times Z_2)$ | $(a, \{fq, hfqh\})$ | $\frac{\sqrt{15}}{72} \approx 0.0538$ | $\sin^2 \theta_{12} = 0.200$ $\sin^2 \theta_{23} = 0.200$ $\sin^2 \theta_{13} \approx 0.167$ | 3892.3 |

As regards the vector which is of the same form (in absolute values) as the second column of the pattern KIVa we notice that it has been discussed as mixing vector in [12]. There it has been associated with the group $\Sigma(60)$ which is isomorphic to A_5 . Due to the relation between the representation matrices of Z_2 and Z_6 generating elements (via the center of $SU(3)$) the sets of mixing vectors have to coincide.

4 Summary

Non-abelian discrete flavor symmetries G_l which are broken to subgroups G_e and G_ν in the charged lepton and neutrino sector, respectively, are still considered to be an interesting possibility to explain the peculiar lepton mixing pattern. The discovery of a large reactor mixing angle $\theta_{13} \approx 0.15$, however, strongly disfavors many mixing patterns such as TB mixing, which is related to the flavor group S_4 . Thus, efforts have been made to find new symmetries, usually larger than the group S_4 , which could predict the experimentally observed mixing parameters. We have pursued such an attempt in the present paper and have discussed the “exceptional” finite groups $\Sigma(n\varphi)$, $\varphi = 3$, which are subgroups of $SU(3)$. These four groups are interesting, since they all contain irreducible faithful three-dimensional representations and have several types of abelian subgroups, suitable as G_e and G_ν . Since one of them, the group $\Sigma(360 \times 3)$, also comprises Klein subgroups, neutrinos could be Majorana particles as well.

Having in mind a realization of the mixing patterns in a scenario with only small corrections to the leading order results, we find that only a few of our patterns can give a good fit to the experimental data: one pattern associated with the group $\Sigma(36 \times 3)$, see eq.(14), which leads to $\theta_{13} \approx 0.18$, one associated with $\Sigma(216 \times 3)$ with $\theta_{13} \approx 0.12$, see table 1,¹⁴ as well as two patterns associated with $\Sigma(360 \times 3)$ which give rise to $\theta_{13} \approx 0.20$ and $\theta_{13} \approx 0.19$, respectively, see table 4. Although they are associated with $\Sigma(360 \times 3)$, also the latter two patterns require neutrinos to be Dirac particles. Interestingly enough, all these patterns lead to vanishing J_{CP} . A non-trivial CP phase only arises from patterns that do not accommodate the data well, i.e. the minimum value of χ^2 is larger than 100, see table 1, 2 and 4-6. Notice that among these are all patterns which can be derived for Majorana neutrinos from $G_l = \Sigma(360 \times 3)$. If they are implemented in a concrete model, large corrections to the pattern, derived from the breaking of G_l to G_e and to G_ν , have to be present so that the mixing angles agree reasonably well with the data. Another possibility could be to modify the breaking pattern, for example by reducing one of the residual symmetries G_e or G_ν or by involving CP symmetry in the breaking. In this way, all the presented patterns could function as starting point for the search for new patterns which lead to mixing parameters agreeing well with the data. For the group $\Sigma(216 \times 3)$ we find several patterns with $|J_{CP}| \neq 0$ which allow the solar and atmospheric mixing angles to be within the experimental 3σ ranges, but lead to a too large value of θ_{13} . It would be interesting to analyze whether a class of models can be constructed in which θ_{13} can be corrected appropriately, while the other two mixing angles (and the prediction of the Jarlskog invariant) only undergo small corrections.

¹⁴In our analysis of $\Sigma(216 \times 3)$ the pattern which fits the data best, see eq.(30), is not associated with $\Sigma(216 \times 3)$, but only with one of its subgroups of order 162. Also this pattern predicts a trivial Dirac phase.

Eventually, we have studied the outer automorphisms of the groups $\Sigma(n\varphi)$, $\varphi = 3$, since we can relate inequivalent irreducible representations with their help. In particular, we could show that all irreducible faithful three-dimensional representations of a group $\Sigma(n\varphi)$ lead to the same set of mixing patterns, and thus that it is sufficient to only consider one of these representations in our discussion.

Acknowledgements

We thank Ferruccio Feruglio for encouragement. CH would like to thank Martin Holthausen and Michael A. Schmidt for help with the computer program GAP as well as Roman Zwicky for email correspondence. CH is supported by the ERC Advanced Grant no. 267985, “Electroweak Symmetry Breaking, Flavour and Dark Matter: One Solution for Three Mysteries” (DaMeSyFla). CH thanks the Galileo Galilei Institute for Theoretical Physics in Arcetri, Florence, for the hospitality and the INFN for partial support during the final stages of this work. AM acknowledges partial support by the INFN program on “Astroparticle Physics” and the Italian MIUR program on “Neutrinos, Dark Matter and Dark Energy in the Era of LHC” and thanks the University of Padua for hospitality and support during her visits. The work of LV is supported by the Swiss National Science Foundation under the contract 200021-125237.

A Comments on outer automorphisms of $\Sigma(n\varphi)$

We would like to consider outer automorphisms of the groups $\Sigma(n\varphi)$, since these allow to relate inequivalent irreducible representations of a group with each other. The group of outer automorphisms is defined as the group of all automorphisms divided by the inner ones. The latter are mappings represented by conjugation with an element of the group. The connection between (outer) automorphisms of a group and symmetries of its character table is also discussed in [18].

A.1 Outer automorphisms of $\Sigma(36 \times 3)$

The outer automorphisms of $\Sigma(36 \times 3)$ form a Klein group and thus three non-trivial mappings exist which act in general non-trivially on the inequivalent irreducible representations of the group. The first one can be defined on the generators of the group as

$$a \rightarrow z^2 a, \quad z \rightarrow za, \quad v \rightarrow azv. \quad (35)$$

It exchanges the representations $\mathbf{3}^{(0)}$ and $\mathbf{3}^{(2)}$ with their complex conjugates and $\mathbf{3}^{(1)}$ with $(\mathbf{3}^{(3)})^*$ as well as $\mathbf{3}^{(3)}$ with $(\mathbf{3}^{(1)})^*$. This transformation is a symmetry of the character table, if we perform at the same time the following exchanges among the classes: $C_2 \leftrightarrow C_3$, $C_7 \leftrightarrow C_8$, $C_{10} \leftrightarrow C_{11}$ and $C_{13} \leftrightarrow C_{14}$ (we refer here to the notation of the classes as found in [27]). The second mapping acts on the generators as follows

$$a \rightarrow za^2z, \quad z \rightarrow za^2, \quad v \rightarrow z^2a^2zv^3. \quad (36)$$

In terms of representations it interchanges the one-dimensional representations $\mathbf{1}^{(1)}$ and $\mathbf{1}^{(3)}$ as well as $\mathbf{3}^{(1)}$ with $\mathbf{3}^{(3)}$ (and consequently also $(\mathbf{3}^{(1)})^*$ with $(\mathbf{3}^{(3)})^*$). It is a symmetry of the character table, if we exchange $C_9 \leftrightarrow C_{12}$, $C_{10} \leftrightarrow C_{13}$ and $C_{11} \leftrightarrow C_{14}$. The third mapping is found if the two presented ones are performed one after another (the ordering is irrelevant, because they form a Klein group). It corresponds to an interchange of all representations with their complex conjugates.

A.2 Outer automorphisms of $\Sigma(72 \times 3)$

The group of outer automorphisms of $\Sigma(72 \times 3)$ is the permutation group S_3 . Thus, we can define two mappings as generators. The first one acts as

$$a \rightarrow z^2, \quad z \rightarrow a^2, \quad v \rightarrow v^3, \quad x \rightarrow z^2 v^3 x \quad (37)$$

and thus exchanges the following representations: $\mathbf{1}^{(1,0)}$ and $\mathbf{1}^{(1,1)}$, $\mathbf{3}^{(1,0)}$ with $(\mathbf{3}^{(1,1)})^*$ and their complex conjugates as well as $\mathbf{3}^{(0,0)}$, $\mathbf{3}^{(0,1)}$ and $\mathbf{6}$ with their complex conjugates. If we exchange the classes C_2 and C_3 , C_6 and C_7 , C_9 and C_{10} , C_{11} and C_{16} , C_{12} with C_{15} and C_{13} with C_{14} (referring to the character table given in [27]), the character table remains invariant. Note this mapping generates a Z_2 symmetry. The second outer automorphism we can define sends the generators a , z , v and x into

$$a \rightarrow za, \quad z \rightarrow aza^2, \quad v \rightarrow z^2 az v^3 x, \quad x \rightarrow zaza^2 v. \quad (38)$$

Its effect on the representations is: $\mathbf{1}^{(1,0)} \rightarrow \mathbf{1}^{(1,1)} \rightarrow \mathbf{1}^{(0,1)} \rightarrow \mathbf{1}^{(1,0)}$ and similarly $\mathbf{3}^{(1,0)} \rightarrow \mathbf{3}^{(1,1)} \rightarrow \mathbf{3}^{(0,1)} \rightarrow \mathbf{3}^{(1,0)}$ and the same holds for their complex conjugated representations. If we cyclicly permute the classes $C_8 \rightarrow C_{16} \rightarrow C_{11} \rightarrow C_8$, $C_9 \rightarrow C_{14} \rightarrow C_{12} \rightarrow C_9$ and $C_{10} \rightarrow C_{15} \rightarrow C_{13} \rightarrow C_{10}$ together with the representations, this mapping is a symmetry of the character table. Notice that it generates a Z_3 symmetry and that it does not commute with the action of the mapping defined in eq.(37). The latter is expected since the two mappings should give rise to the group S_3 . Thus, there are in addition three non-trivial mappings: the map in eq.(38) applied twice, the application of the first map followed by the second one as well as vice versa.

A.3 Outer automorphisms of $\Sigma(216 \times 3)$

The outer automorphism group is a Z_6 symmetry. We can define a mapping of the generators as¹⁵

$$a \rightarrow a^2, \quad z \rightarrow z, \quad v \rightarrow v^3, \quad w \rightarrow (zw)^2. \quad (39)$$

Applying this mapping we see that the three-dimensional representations and their complex conjugates transform as

$$\mathbf{3}^{(0)} \rightarrow (\mathbf{3}^{(1)})^*, \quad \mathbf{3}^{(1)} \rightarrow (\mathbf{3}^{(2)})^*, \quad \mathbf{3}^{(2)} \rightarrow (\mathbf{3}^{(0)})^*. \quad (40)$$

¹⁵Note that if applied only to the generators a , v and z which give rise to the group $\Sigma(36 \times 3)$ this mapping only generates a Z_2 symmetry instead of a Z_6 .

The other representations are mapped as follows: $\mathbf{1}^{(1)} \leftrightarrow \mathbf{1}^{(2)}$, $\mathbf{2}^{(1)} \leftrightarrow \mathbf{2}^{(2)}$, $\mathbf{6}^{(0)} \rightarrow (\mathbf{6}^{(1)})^*$, $\mathbf{6}^{(1)} \rightarrow (\mathbf{6}^{(2)})^*$, $\mathbf{6}^{(2)} \rightarrow (\mathbf{6}^{(0)})^*$ (and similarly their complex conjugates), as well as $\mathbf{8}^{(1)} \leftrightarrow \mathbf{8}^{(2)}$ and $\mathbf{9} \leftrightarrow \mathbf{9}^*$. Using the convention of [27] for the character table, we see that we have to exchange the classes $C_2 \leftrightarrow C_3$, $C_6 \leftrightarrow C_7$, $C_9 \leftrightarrow C_{10}$, $C_{11} \leftrightarrow C_{12}$ as well as apply the transformations $C_{13} \rightarrow C_{16} \rightarrow C_{15} \rightarrow C_{17} \rightarrow C_{14} \rightarrow C_{18} \rightarrow C_{13}$ and $C_{19} \rightarrow C_{24} \rightarrow C_{21} \rightarrow C_{22} \rightarrow C_{20} \rightarrow C_{23} \rightarrow C_{19}$, if we want to promote this mapping to a symmetry of the character table. As one can check, applying this mapping repeatedly we can relate all faithful irreducible three-dimensional representations to $\mathbf{3}^{(0)}$.

References

- [1] G. Altarelli, F. Feruglio, Rev. Mod. Phys. **82** (2010) 2701-2729 [arXiv:1002.0211 [hep-ph]]; H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, M. Tanimoto, Prog. Theor. Phys. Suppl. **183** (2010) 1-163 [arXiv:1003.3552 [hep-th]]; S. F. King and C. Luhn, Rept. Prog. Phys. **76** (2013) 056201 [arXiv:1301.1340 [hep-ph]].
- [2] W. Grimus and P. O. Ludl, J. Phys. A **45** (2012) 233001 [arXiv:1110.6376 [hep-ph]].
- [3] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, JHEP **1212** (2012) 123 [arXiv:1209.3023 [hep-ph]].
- [4] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo, A. M. Rotunno, Phys. Rev. **D84** (2011) 053007 [arXiv:1106.6028 [hep-ph]]; D. V. Forero, M. Tortola and J. W. F. Valle, Phys. Rev. D **86** (2012) 073012 [arXiv:1205.4018 [hep-ph]].
- [5] C. S. Lam, Phys. Lett. B **656** (2007) 193 [arXiv:0708.3665 [hep-ph]]; A. Blum, C. Hagedorn and M. Lindner, Phys. Rev. D **77** (2008) 076004 [arXiv:0709.3450 [hep-ph]].
- [6] R. de Adelhart Toorop, F. Feruglio and C. Hagedorn, Nucl. Phys. B **858** (2012) 437 [arXiv:1112.1340 [hep-ph]].
- [7] C. Jarlskog, Phys. Rev. Lett. **55** (1985) 1039.
- [8] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B **530** (2002) 167 [hep-ph/0202074]; Z. -z. Xing, Phys. Lett. B **533** (2002) 85 [hep-ph/0204049].
- [9] C. S. Lam, Phys. Lett. B **656** (2007) 193 [arXiv:0708.3665 [hep-ph]]; C. S. Lam, Phys. Rev. Lett. **101** (2008) 121602 [arXiv:0804.2622 [hep-ph]]; C. S. Lam, Phys. Rev. D **78** (2008) 073015 [arXiv:0809.1185 [hep-ph]].
- [10] K. Abe *et al.* [T2K Collaboration], Phys. Rev. Lett. **107** (2011) 041801 [arXiv:1106.2822 [hep-ex]]; P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. Lett. **107** (2011) 181802 [arXiv:1108.0015 [hep-ex]]; Y. Abe *et al.* [Double-Chooz Collaboration], Phys. Rev. Lett. **108** (2012) 131801 [arXiv:1112.6353 [hep-ex]]; F. P. An *et al.* [Daya-Bay Collaboration], Phys. Rev. Lett. **108** (2012) 171803 [arXiv:1203.1669 [hep-ex]]; J. K. Ahn *et al.* [Reno Collaboration], Phys. Rev. Lett. **108** (2012) 191802 [arXiv:1204.0626 [hep-ex]]; F. P. An *et al.* [Daya-Bay Collaboration], Chin. Phys. C **37** (2013) 011001 [arXiv:1210.6327 [hep-ex]].

- [11] R. de Adelhart Toorop, F. Feruglio and C. Hagedorn, Phys. Lett. B **703** (2011) 447 [arXiv:1107.3486 [hep-ph]].
- [12] C. S. Lam, Phys. Rev. D **87** (2013) 013001 [arXiv:1208.5527 [hep-ph]].
- [13] M. Holthausen, K. S. Lim and M. Lindner, Phys. Lett. B **721** (2013) 61 [arXiv:1212.2411 [hep-ph]].
- [14] S. F. King, T. Neder and A. J. Stuart, Phys. Lett. B **726** (2013) 312 [arXiv:1305.3200 [hep-ph]].
- [15] M. Holthausen and K. S. Lim, Phys. Rev. D **88** (2013) 033018 [arXiv:1306.4356 [hep-ph]].
- [16] S. -F. Ge, D. A. Dicus and W. W. Repko, Phys. Lett. B **702** (2011) 220 [arXiv:1104.0602 [hep-ph]]; S. -F. Ge, D. A. Dicus and W. W. Repko, Phys. Rev. Lett. **108** (2012) 041801 [arXiv:1108.0964 [hep-ph]]; D. Hernandez and A. Y. Smirnov, Phys. Rev. D **86** (2012) 053014 [arXiv:1204.0445 [hep-ph]]; D. Hernandez and A. Y. Smirnov, Phys. Rev. D **87** (2013) 053005 [arXiv:1212.2149 [hep-ph]].
- [17] F. Feruglio, C. Hagedorn and R. Ziegler, JHEP **1307** (2013) 027 [arXiv:1211.5560 [hep-ph]].
- [18] M. Holthausen, M. Lindner and M. A. Schmidt, JHEP **1304** (2013) 122 [arXiv:1211.6953 [hep-ph]].
- [19] P. F. Harrison and W. G. Scott, Phys. Lett. B **535** (2002) 163 [hep-ph/0203209]; P. F. Harrison and W. G. Scott, Phys. Lett. B **547** (2002) 219 [hep-ph/0210197]; P. F. Harrison and W. G. Scott, Phys. Lett. B **594** (2004) 324 [hep-ph/0403278]; W. Grimus and L. Lavoura, Phys. Lett. B **579** (2004) 113 [hep-ph/0305309]; S. Gupta, A. S. Joshipura and K. M. Patel, Phys. Rev. D **85** (2012) 031903 [arXiv:1112.6113 [hep-ph]]; R. N. Mohapatra and C. C. Nishi, Phys. Rev. D **86** (2012) 073007 [arXiv:1208.2875 [hep-ph]]; R. Krishnan, P. F. Harrison and W. G. Scott, JHEP **1304** (2013) 087 [arXiv:1211.2000 [hep-ph]]; G. -J. Ding, S. F. King, C. Luhn and A. J. Stuart, JHEP **1305** (2013) 084 [arXiv:1303.6180 [hep-ph]]; F. Feruglio, C. Hagedorn and R. Ziegler, arXiv:1303.7178 [hep-ph].
- [20] G. A. Miller, H. F. Blichfeldt, L. E. Dickson, "Theory and Applications of Finite Groups", John Wiley & Sons, New York, 1916, and Dover Edition, 1961; W. M. Fairbairn, T. Fulton, and W. H. Klink, J. Math. Phys. **5** (1964) 1038.
- [21] I. P. Ivanov and E. Vdovin, Eur. Phys. J. C **73** (2013) 2309 [arXiv:1210.6553 [hep-ph]].
- [22] R. Zwicky and T. Fischbacher, Phys. Rev. D **80** (2009) 076009 [arXiv:0908.4182 [hep-ph]].
- [23] C. S. Lam, Phys. Rev. D **83** (2011) 113002 [arXiv:1104.0055 [hep-ph]].
- [24] J. Beringer *et al.* [Particle Data Group], Phys. Rev. D **86** (2012) 010001.

- [25] H. U. Besche, B. Eick, E. O'Brien, "SmallGroups- library of all 'small' groups, GAP package, Version included in GAP 4.5.5", (2002), <http://www.gap-system.org/Packages/sgl.html>.
- [26] GAP, "GAP – Groups, Algorithms, and Programming, Version 4.5.5", (2012), <http://www.gap-system.org>.
- [27] W. Grimus and P. O. Ludl, J. Phys. A **43** (2010) 445209 [arXiv:1006.0098 [hep-ph]]; A. Merle and R. Zwicky, JHEP **1202** (2012) 128 [arXiv:1110.4891 [hep-ph]].
- [28] C. C. Nishi, Phys. Rev. D **88** (2013) 033010 [arXiv:1306.0877 [hep-ph]].
- [29] P. O. Ludl, J. Phys. A **44** (2011) 255204 [Erratum-ibid. A **45** (2012) 069502] [arXiv:1101.2308 [math-ph]].
- [30] A. Hanany and Y. -H. He, JHEP **9902** (1999) 013 [hep-th/9811183].
- [31] B. Hu, Phys. Rev. D **87** (2013) 033002 [arXiv:1212.2819 [hep-ph]].