

PREPARED FOR SUBMISSION TO JHEP

R_2 vertices for the effective ggH theory

Ben Page and Roberto Pittau

*Departamento de Física Teórica y del Cosmos and CAFPE,
Campus Fuentenueva s. n., Universidad de Granada E-18071 Granada, Spain*

E-mail: page@ugr.es, pittau@ugr.es

ABSTRACT: We list all possible R_2 Feynman rules needed in NLO computations involving couplings of Higgs and gluons mediated by an infinitely heavy top loop. They provide the rational contribution generated by the $(d - 4)$ -dimensional part of the amplitude, paving the way for four-dimensional automatic NLO methods in Higgs phenomenology.

Contents

1	Introduction	1
2	R₂ in a nutshell	2
3	The R₂ vertices generated by the <i>GGH</i> operator	2
4	Conclusions	5

1 Introduction

Automation of NLO calculations is an important issue in modern high energy particle physics. The complexity of the processes one has to deal with forbids a case by case approach, forcing the use of automatic tools [1–9].

In the last few years, a lot of progress has been achieved and, among the various available techniques [10–14], the Ossola-Papadopoulos-Pittau (OPP) method [15] appears to be a rather convenient one. OPP is a procedure which allows one to numerically extract, from dimensionally regulated one-loop amplitudes, the Cut-Constructible part plus a rational piece - called R₁ - strictly linked to the four-dimensional numerator of the Feynman diagrams. A second contribution to the rational part, dubbed R₂, cannot be obtained numerically in four-dimensions, and must be provided externally. A particularly convenient way to account for R₂ is via effective Feynman rules. This is because the R₂ contribution disappears when a sufficiently large number of particles undergo the scattering and, in renormalizable theories, only up to four-particle-vertices have to be considered.

The R₂ vertices for QCD [16], the Electroweak Model [17–19] and MSSM [20] have already been presented in the literature. In this paper, we list the R₂ Feynman rules generated by the interaction of one Higgs field H with two, three and four gluons - mediated by an infinitely heavy top loop - encoded in the effective Lagrangian [21, 22]

$$\mathcal{L}_{\text{eff}} = -\frac{1}{4}AHG_{\mu\nu}^a G^{a,\mu\nu}, \quad (1.1)$$

where

$$A = \frac{\alpha_S}{3\pi v} \left(1 + \frac{11}{4} \frac{\alpha_S}{\pi} \right) \quad (1.2)$$

and v is the vacuum expectation value, $v^2 = (G_F\sqrt{2})^{-1}$.

There is a very important physical motivation that led us to study this problem: the main Higgs production mechanism at the LHC is via gluon fusion [23, 24], and considering NLO corrections to processes generated by the effective operator in eq. (1.1) provides an

easy way to model, at two-loop precision, Higgs + n jets final states for jets with p_T not much larger than the top mass [25–27]. The Hjj and $Hjjj$ signatures have been recently considered [28, 29], in the context of a d -dimensional integrand reduction technique, within the GoSam [8]/Sherpa [30] framework. In this paper we provide the only missing ingredient needed by 4-dimensional NLO approaches, such as OPP, Open Loops [31] or FDR [32], to attack the problem in a fully automatic fashion.

In the next section we review the basics of R_2 . In section 3 we list the contributing R_2 vertices and section 4 contains our conclusions.

2 R_2 in a nutshell

The full theory of R_2 can be found in [33, 34]. For the purposes of this paper, it is enough to recall that, starting from a one-loop amplitude computed in d dimensions, R_2 is generated by the explicit splits

$$\begin{aligned} d &= 4 + \epsilon, \\ \bar{q}^2 &= q^2 + \tilde{q}^2 \end{aligned} \tag{2.1}$$

in the numerators of the contributing Feynman diagrams, where \bar{q}^2 is the d -dimensional integration momentum squared, q^2 its four-dimensional part and \tilde{q}^2 the difference between the two. In renormalizable theories, in the limit $\epsilon \rightarrow 0$, the ϵ and \tilde{q}^2 pieces generate constant terms in one particle irreducible Green’s functions up to four legs. The effective operator in eq. (1.1) has dimension five and so a non-vanishing contribution can survive in up to five-leg vertices.

In Dimensional Reduction like schemes [35], one is free to set $d = 4$ in eq. (2.1) right from the beginning, while the \tilde{q}^2 part is always necessary in order to keep gauge invariance. The transition rules between Dimensional Reduction and the usual ’t Hooft-Veltman Dimensional Regularization [36], in which $d = 4 + \epsilon$, are well known [34, 37]. To allow the reading of our rules in both schemes we introduced a parameter λ_{HV} in our formulae, such that

$$d = 4 + \lambda_{\text{HV}} \epsilon. \tag{2.2}$$

Thus, Dimensional Reduction corresponds to $\lambda_{\text{HV}} = 0$ and Dimensional Regularization to $\lambda_{\text{HV}} = 1$.

3 The R_2 vertices generated by the GGH operator

By power counting, a non-zero R_2 contribution is present only in the following five interactions

$$Hgg, \quad Hggg, \quad Hgggg, \quad Hq\bar{q}, \quad Hq\bar{q}g. \tag{3.1}$$

We generated and computed, with the help of QGRAF [38] and FORM [39], all possible contributing diagrams¹ from which we extracted R_2 via eq. (2.1). Two independent calculations were performed, with slightly different strategies. In the first computation a Feynman parametrization was applied before eq. (2.1); in the second, eq. (2.1) was used to classify independent integrals, which were computed at a later stage of the calculation. Both procedures gave the same expressions, listed in figure 1, which represent the main result of this paper.

The Hgg vertex is fully proportional to the QCD three gluon vertex

$$V_{m_1 m_2 m_3}^{\mu_1 \mu_2 \mu_3}(p_1, p_2, p_3) = g_s f_{m_1 m_2 m_3} \left[g^{\mu_1 \mu_2} (p_2 - p_1)^{\mu_3} + g^{\mu_2 \mu_3} (p_3 - p_2)^{\mu_1} + g^{\mu_3 \mu_1} (p_1 - p_3)^{\mu_2} \right], \quad (3.2)$$


while the tensor involved in the Higgs/4-Gluons vertex reads

$$X_{m_1 m_2 m_3 m_4}^{\mu_1 \mu_2 \mu_3 \mu_4} = \begin{aligned} & + \text{Tr}(T^{m_1} T^{m_2} T^{m_3} T^{m_4}) \left[+ 21 g^{\mu_1 \mu_2} g^{\mu_3 \mu_4} - 41 g^{\mu_1 \mu_3} g^{\mu_2 \mu_4} + 21 g^{\mu_1 \mu_4} g^{\mu_2 \mu_3} \right] \\ & + \text{Tr}(T^{m_1} T^{m_2} T^{m_4} T^{m_3}) \left[+ 21 g^{\mu_1 \mu_2} g^{\mu_3 \mu_4} + 21 g^{\mu_1 \mu_3} g^{\mu_2 \mu_4} - 41 g^{\mu_1 \mu_4} g^{\mu_2 \mu_3} \right] \\ & + \text{Tr}(T^{m_1} T^{m_3} T^{m_2} T^{m_4}) \left[- 41 g^{\mu_1 \mu_2} g^{\mu_3 \mu_4} + 21 g^{\mu_1 \mu_3} g^{\mu_2 \mu_4} + 21 g^{\mu_1 \mu_4} g^{\mu_2 \mu_3} \right], \end{aligned} \quad (3.3)$$

where the color structure is compactly expressed in terms of traces of color matrices T^a in the *adjoint* representation. The translation to traces of Gell-Mann matrices t^a in the fundamental representation of $SU(N_c)$ is given by the formula

$$\text{Tr}(T^a T^b T^c T^d) = N_c \left[\text{Tr}(t^a t^b t^c t^d) + \text{Tr}(t^d t^c t^b t^a) \right] + \frac{\delta_{ab} \delta_{cd} + \delta_{ac} \delta_{bd} + \delta_{ad} \delta_{bc}}{2}. \quad (3.4)$$

Any λ_{HV} contribution is entirely contained in the R_2 part. However, in physical combinations no dependence on λ_{HV} remains. Naturally this means that the vertices involving only gluons are devoid of λ_{HV} . Furthermore one can show that in the physical sum of diagrams, given by




there is no dependence on λ_{HV} .


Finally, it is important to realize that, being one particle irreducible, the vertices in figure 1 are the building blocks from which the R_2 contribution to *any* amplitude involving one coupling of order A ² and *any* number of additional jets can be derived.

¹We used the QCD Feynman rules in appendix B of [16], together with the effective Higgs-gluons couplings listed in [40].


²See eq. (1.2).




$$= \frac{iAg_s^2 N_c \delta^{m_1 m_2}}{384\pi^2} \left\{ p_1^{\mu_1} p_2^{\mu_2} + 89 p_1^{\mu_2} p_2^{\mu_1} + 14 (p_1^{\mu_1} p_1^{\mu_2} + p_2^{\mu_1} p_2^{\mu_2}) - [17 (p_1^2 + p_2^2) + 93 (p_1 \cdot p_2)] g^{\mu_1 \mu_2} \right\}$$




$$= -\frac{15Ag_s^2 N_c}{128\pi^2} V_{m_1 m_2 m_3}^{\mu_1 \mu_2 \mu_3} (p_1, p_2, p_3)$$



$$= \frac{iAg_s^4}{128\pi^2} X_{m_1 m_2 m_3 m_4}^{\mu_1 \mu_2 \mu_3 \mu_4}$$



$$= \frac{iAg_s^2}{32\pi^2} \left(\frac{N_c^2 - 1}{2N_c} \right) \delta^{j_1 j_2} \lambda_{\text{HV}} (\not{p}_1 - \not{p}_2)$$



$$= \frac{iAg_s^3}{64\pi^2} \gamma_\mu t_{j_2 j_1}^m \left[\frac{2\lambda_{\text{HV}} + 1}{N_c} - (2\lambda_{\text{HV}} + 3)N_c \right]$$

Figure 1. The R_2 vertices generated by the effective Lagrangian in eq. (1.1). All momenta are incoming, N_c is the number of colors and $\lambda_{\text{HV}} = 1$ ($\lambda_{\text{HV}} = 0$) in Dimensional Regularization (Reduction). Quarks are massless and the dashed line represents the Higgs field. The tensors $V_{m_1 m_2 m_3}^{\mu_1 \mu_2 \mu_3} (p_1, p_2, p_3)$ and $X_{m_1 m_2 m_3 m_4}^{\mu_1 \mu_2 \mu_3 \mu_4}$ are defined in the text.

4 Conclusions

We have presented the complete set of effective R_2 Feynman rules generated by the dimension five operator GGH . They encode the missing analytical information needed to apply four-dimensional automatic integrand reduction techniques to Higgs physics.

Our result can be used to study both production and decay processes involving one Higgs particle and any number of jets. This is particularly useful for Higgs phenomenology at the LHC, where, due to the large amount of available energy, the Higgs particle is very often produced in association with a large number of jets.

Acknowledgments

This work was performed in the framework of the ERC grant 291377, “LHCtheory: Theoretical predictions and analyses of LHC physics: advancing the precision frontier”. We also thank the support of the MICINN project FPA2011-22398 (LHC@NLO) and the Junta de Andalucía project P10-FQM-6552.

References

- [1] G. Ossola, C. G. Papadopoulos, and R. Pittau, *CutTools: A Program implementing the OPP reduction method to compute one-loop amplitudes*, *JHEP* **0803** (2008) 042, [[arXiv:0711.3596](#)].
- [2] W. Giele and G. Zanderighi, *On the Numerical Evaluation of One-Loop Amplitudes: The Gluonic Case*, *JHEP* **0806** (2008) 038, [[arXiv:0805.2152](#)].
- [3] A. van Hameren, C. Papadopoulos, and R. Pittau, *Automated one-loop calculations: A Proof of concept*, *JHEP* **0909** (2009) 106, [[arXiv:0903.4665](#)].
- [4] V. Hirschi, R. Frederix, S. Frixione, M. V. Garzelli, F. Maltoni, et al., *Automation of one-loop QCD corrections*, *JHEP* **1105** (2011) 044, [[arXiv:1103.0621](#)].
- [5] R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, R. Pittau, et al., *Four-lepton production at hadron colliders: aMC@NLO predictions with theoretical uncertainties*, *JHEP* **1202** (2012) 099, [[arXiv:1110.4738](#)].
- [6] G. Bevilacqua, M. Czakon, M. Garzelli, A. van Hameren, A. Kardos, et al., *HELAC-NLO*, *Comput.Phys.Comm.* **184** (2013) 986–997, [[arXiv:1110.1499](#)].
- [7] Z. Bern, L. Dixon, F. Febres Cordero, S. Hoeche, H. Ita, et al., *Next-to-Leading Order $W + 5$ -Jet Production at the LHC*, [[arXiv:1304.1253](#)].
- [8] G. Cullen, N. Greiner, G. Heinrich, G. Luisoni, P. Mastrolia, et al., *Automated One-Loop Calculations with GoSam*, *Eur.Phys.J.* **C72** (2012) 1889, [[arXiv:1111.2034](#)].
- [9] P. Mastrolia, G. Ossola, T. Reiter, and F. Tramontano, *Scattering Amplitudes from Unitarity-based Reduction Algorithm at the Integrand-level*, *JHEP* **1008** (2010) 080, [[arXiv:1006.0710](#)].
- [10] Z. Bern, L. J. Dixon, D. C. Dunbar, and D. A. Kosower, *Fusing gauge theory tree amplitudes into loop amplitudes*, *Nucl.Phys.* **B435** (1995) 59–101, [[hep-ph/9409265](#)].

- [11] R. Britto, F. Cachazo, and B. Feng, *Generalized unitarity and one-loop amplitudes in $N=4$ super-Yang-Mills*, *Nucl.Phys.* **B725** (2005) 275–305, [[hep-th/0412103](#)].
- [12] D. Forde, *Direct extraction of one-loop integral coefficients*, *Phys.Rev.* **D75** (2007) 125019, [[arXiv:0704.1835](#)].
- [13] C. Berger, Z. Bern, L. Dixon, F. Febres Cordero, D. Forde, et al., *An Automated Implementation of On-Shell Methods for One-Loop Amplitudes*, *Phys.Rev.* **D78** (2008) 036003, [[arXiv:0803.4180](#)].
- [14] R. K. Ellis, K. Melnikov, and G. Zanderighi, *Generalized unitarity at work: first NLO QCD results for hadronic $W + 3$ jet production*, *JHEP* **0904** (2009) 077, [[arXiv:0901.4101](#)].
- [15] G. Ossola, C. G. Papadopoulos, and R. Pittau, *Reducing full one-loop amplitudes to scalar integrals at the integrand level*, *Nucl.Phys.* **B763** (2007) 147–169, [[hep-ph/0609007](#)].
- [16] P. Draggiotis, M. Garzelli, C. Papadopoulos, and R. Pittau, *Feynman Rules for the Rational Part of the QCD 1-loop amplitudes*, *JHEP* **0904** (2009) 072, [[arXiv:0903.0356](#)].
- [17] M. Garzelli, I. Malamos, and R. Pittau, *Feynman rules for the rational part of the Electroweak 1-loop amplitudes*, *JHEP* **1001** (2010) 040, [[arXiv:0910.3130](#)].
- [18] M. Garzelli, I. Malamos, and R. Pittau, *Feynman rules for the rational part of the Electroweak 1-loop amplitudes in the R_ξ gauge and in the Unitary gauge*, *JHEP* **1101** (2011) 029, [[arXiv:1009.4302](#)].
- [19] H.-S. Shao, Y.-J. Zhang, and K.-T. Chao, *Feynman Rules for the Rational Part of the Standard Model One-loop Amplitudes in the 't Hooft-Veltman γ_5 Scheme*, *JHEP* **1109** (2011) 048, [[arXiv:1106.5030](#)].
- [20] H.-S. Shao and Y.-J. Zhang, *Feynman Rules for the Rational Part of One-loop QCD Corrections in the MSSM*, *JHEP* **1206** (2012) 112, [[arXiv:1205.1273](#)].
- [21] M. A. Shifman, A. Vainshtein, M. Voloshin, and V. I. Zakharov, *Low-Energy Theorems for Higgs Boson Couplings to Photons*, *Sov.J.Nucl.Phys.* **30** (1979) 711–716.
- [22] S. Dawson and R. Kauffman, *Higgs boson plus multi-jet rates at the SSC*, *Phys.Rev.Lett.* **68** (1992) 2273–2276.
- [23] **ATLAS Collaboration** Collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys.Lett.* **B716** (2012) 1–29, [[arXiv:1207.7214](#)].
- [24] **CMS Collaboration** Collaboration, S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys.Lett.* **B716** (2012) 30–61, [[arXiv:1207.7235](#)].
- [25] V. Del Duca, W. Kilgore, C. Oleari, C. Schmidt, and D. Zeppenfeld, *Higgs + 2 jets via gluon fusion*, *Phys.Rev.Lett.* **87** (2001) 122001, [[hep-ph/0105129](#)].
- [26] V. Del Duca, W. Kilgore, C. Oleari, C. Schmidt, and D. Zeppenfeld, *Gluon fusion contributions to $H + 2$ jet production*, *Nucl.Phys.* **B616** (2001) 367–399, [[hep-ph/0108030](#)].
- [27] F. Campanario and M. Kubocz, *Higgs boson production in association with three jets via gluon fusion at the LHC: Gluonic contributions*, [[arXiv:1306.1830](#)].
- [28] H. van Deurzen, N. Greiner, G. Luisoni, P. Mastrolia, E. Mirabella, et al., *NLO QCD corrections to the production of Higgs plus two jets at the LHC*, *Phys.Lett.* **B721** (2013) 74–81, [[arXiv:1301.0493](#)].

- [29] G. Cullen, H. van Deurzen, N. Greiner, G. Luisoni, P. Mastrolia, et al., *NLO QCD corrections to Higgs boson production plus three jets in gluon fusion*, [arXiv:1307.4737](#).
- [30] T. Gleisberg, S. Hoeche, F. Krauss, M. Schonherr, S. Schumann, et al., *Event generation with SHERPA 1.1*, *JHEP* **0902** (2009) 007, [[arXiv:0811.4622](#)].
- [31] F. Cascioli, P. Maierhofer, and S. Pozzorini, *Scattering Amplitudes with Open Loops*, *Phys.Rev.Lett.* **108** (2012) 111601, [[arXiv:1111.5206](#)].
- [32] R. Pittau, *A four-dimensional approach to quantum field theories*, *JHEP* **1211** (2012) 151, [[arXiv:1208.5457](#)].
- [33] G. Ossola, C. G. Papadopoulos, and R. Pittau, *On the Rational Terms of the one-loop amplitudes*, *JHEP* **0805** (2008) 004, [[arXiv:0802.1876](#)].
- [34] R. Pittau, *Primary Feynman rules to calculate the epsilon-dimensional integrand of any 1-loop amplitude*, *JHEP* **1202** (2012) 029, [[arXiv:1111.4965](#)].
- [35] W. Siegel, *Supersymmetric Dimensional Regularization via Dimensional Reduction*, *Phys.Lett.* **B84** (1979) 193.
- [36] G. 't Hooft and M. Veltman, *Regularization and Renormalization of Gauge Fields*, *Nucl.Phys.* **B44** (1972) 189–213.
- [37] A. Signer, *Helicity method for next-to-leading order corrections in QCD*, *Ph.D. Thesis* (1995).
- [38] P. Nogueira, *Automatic Feynman graph generation*, *J.Comput.Phys.* **105** (1993) 279–289.
- [39] J. Vermaseren, *New features of FORM*, [math-ph/0010025](#).
- [40] R. P. Kauffman, S. V. Desai, and D. Risal, *Production of a Higgs boson plus two jets in hadronic collisions*, *Phys.Rev.* **D55** (1997) 4005–4015, [[hep-ph/9610541](#)].