Compact Stars in the QCD Phase Diagram III (CSQCD III)
December 12-15, 2012, Guarujá, SP, Brazil
http://www.astro.iag.usp.br/~foton/CSQCD3

Strong magnetic fields, nuclear matter, and anomalous magnetic moments

[1]*Eduardo L. Coelho,* [2]*Marcelo Chiapparini*
Instituto de Física, Universidade do Estado do Rio de Janeiro, 20559-900, Rio de Janeiro, RJ, Brazil
Email: [1]eduardo.lenho@gmail.com

Mirian E. Bracco
Faculdade de Tecnologia, Universidade do Estado do Rio de Janeiro, 27537-000, Resende, RJ, Brazil

arXiv:1307.7298v1 [hep-ph] 27 Jul 2013

1 Introduction

Strong magnetics fields of magnitudes about 10^{15} to 10^{19} G are suppose to exists on the surface of pulsars. So, it is of interest to study the properties of nuclear matter in the presence of magnetic fields. In this work we study the influence of a strong magnetic field on the composition of nuclear matter at $T = 0$ including the anomalous magnetic moment (AMM) of baryons. We use a relativistic mean field (MF) theory of nuclear matter which describes correctly the nuclear ground state properties and elastic scattering of nucleons. The Lagrangian that describes this model, with an uniform magnetic field B along the z axis, is given by [1]

$$\begin{aligned}\mathcal{L} &= \sum_b \overline{\psi}_b \left[i\gamma_\mu D^\mu - m_b - g_{\sigma b}\sigma - g_{\omega b}\gamma_\mu \omega^\mu - \frac{1}{2} g_{\rho b}\gamma_\mu \boldsymbol{\tau}\cdot\boldsymbol{\rho}^\mu - \frac{1}{2}k_b\,\sigma_{\mu\nu}F^{\mu\nu} \right]\psi_b \\ &+\frac{1}{2}\partial_\mu\sigma\partial^\mu\sigma - \frac{1}{2}m_\sigma^2\sigma^2 - U(\sigma) - \frac{1}{4}\omega_{\mu\nu}\omega^{\mu\nu} + \frac{1}{2}m_\omega^2\omega_\mu\omega^\mu - \frac{1}{4}\rho_{\mu\nu}\rho^{\mu\nu} \\ &-\frac{1}{2}m_\rho^2\boldsymbol{\rho_\mu}\cdot\boldsymbol{\rho^\mu} + \sum_{l=e,\mu}\overline{\psi}_l(i\gamma_\mu\partial^\mu - m_l)\psi_l - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}, \end{aligned} \quad (1)$$

where $D = \partial + ieA^\mu$, were $A^0 = 0$, $\vec{A} = (0, xB, 0)$, ψ_b is the Dirac spinor for the baryon octet $\{n, p, \Lambda, \Sigma, \Xi\}$ with masses m_b, $\{\sigma, \omega, \rho\}$ are the meson fields and $\{m_\sigma, m_\omega, m_\rho\}$ their masses. The summation includes the free Lagrangian of baryons, mesons and leptons and the interaction terms between baryons and mesons. The scalar self-interaction potential $U(\sigma)$ is

$$U(\sigma) = \frac{1}{3}bm_N(g_{\sigma N}\sigma)^3 + \frac{1}{4}c(g_{\sigma N}\sigma)^4, \quad (2)$$

$F_{\mu\nu}$ is the electromagnetic tensor, $\kappa_b = (\mu_b/\mu_N - q_b m_p/m_b)\ \mu_N$ is the anomalous magnetic moment of baryon b, $\mu_N = e\hbar/(2m_p) = 3.15 \times 10^{-18}$ MeV G^{-1} is the nuclear magneton and $\sigma_{\mu\nu} = \frac{i}{2}[\gamma_\mu, \gamma_\nu]$.

The Euler-Lagrange equations for the meson fields, written in the mean field approximations for infinite nuclear matter reads [2]

$$\omega_0 = \left(\frac{g_{\omega N}}{m_\omega}\right)^2 \sum_b \chi_{\omega b}\rho_b, \tag{3}$$

$$\rho_{03} = \left(\frac{g_{\rho N}}{m_\rho}\right)^2 \sum_b \chi_{\rho b} I_{3b}\rho_b, \tag{4}$$

$$m_N^* = m_N + bm_N\left(\frac{g_{\sigma N}}{m_\sigma}\right)^2 (m_N - m_N^*)^2 + c(m_N - m_N^*)^3 - n_s, \tag{5}$$

where ρ_b is the number density of baryon b (defined below), I_{3b} is the $3th$-component of the isospin, $\chi_{\sigma b} = \frac{g_{\sigma b}}{g_{\sigma N}}$, $\chi_{\omega b} = \frac{g_{\omega b}}{g_{\omega N}}$, $\chi_{\rho b} = \frac{g_{\rho b}}{g_{\rho N}}$, $m_N^* = m_N - g_{\sigma N}\sigma$, $n_s = n_s^{q=0} + n_s^{q\neq 0}$ is the scalar density, with

$$n_s^{q\neq 0} = \frac{B}{2\pi^2}\sum_{\substack{b\\ q\neq 0}} |q_b| m_b^* \sum_{s=-1}^{1}\sum_{\nu=0}^{\nu_{b,s}^m} \frac{\sqrt{m_b^{*2} + 2\nu|q_b|B} + s\kappa_b B}{\sqrt{m_b^{*2} + 2\nu|q_b|B}} \times \ln\left|\frac{\mu_b^* + k_{b,\nu,s}^F}{\sqrt{m_b^{*2} + 2\nu|q_b|B} + s\kappa_b B}\right|, \tag{6}$$

$$n_s^{q=0} = \frac{1}{4\pi^2}\sum_{\substack{b\\ q=0}} m_b^* \sum_{s=-1,1}\left[k_{b,s}^F\mu_b^* - (m_b^* + s\kappa_b B)^2 \ln\left|\frac{\mu_b^* + k_{b,s}^F}{m_b^* + s\kappa_b B}\right|\right], \tag{7}$$

and

$$\left(k_{b,\nu,s}^F\right)^2 = \mu_b^{*2} - \left(\sqrt{m_b^{*2} + 2\nu|q_b|B} + s\kappa_b B\right)^2 \quad (q \neq 0), \tag{8}$$

$$\left(k_{b,s}^F\right)^2 = \mu_b^{*2} - (m_b^* + s\kappa_b B)^2 \quad (q = 0), \tag{9}$$

$$\left(k_{l,\nu}^F\right)^2 = \mu_l^2 - m_l^2 + 2\nu|q_l|B, \tag{10}$$

where $m_b^* = m_b - \chi_{\sigma b} g_{\sigma N}\sigma$. $|q_b|$ and μ_b^* are the electrical charge and effective chemical potential for barion b, k^F is the Fermi momentum and ν is the Landau principal quantum number, which can take all possible positive integer values including zero. Analogous definitions holds for leptons. Baryon chemical potentials are given by $\mu_b = \chi_{\omega b} g_{\omega N}\omega_0 + \chi_{\rho b} g_{\rho N} I_{3b}\rho_{03} + \mu_b^*$.

The baryon and lepton densities are given by the expressions

$$\rho_b^{q\neq 0} = \frac{|q_b|B}{2\pi^2}\sum_{s=-1}^{1}\sum_{\nu=0}^{\nu_{b,s}^m} k_{b,\nu,s}^F, \tag{11}$$

$$\rho_b^{q=0} = \frac{1}{2\pi^2}\sum_{s=-1}^{1}\left\{\frac{1}{3}\left(k_{b,s}^F\right)^3 + \frac{1}{2}s\kappa_b B\left[\left(m_b^* + s\kappa_b B\right)k_{b,s}^F + \mu_b^{*2}\left(\arcsin\left(\frac{m_b^* + s\kappa_b B}{\mu_b^*}\right) - \frac{\pi}{2}\right)\right]\right\}, \tag{12}$$

$$\rho_l = \frac{|q_l|\,B}{2\pi^2}\sum_{\nu=0}^{\nu_l^m} k_{l,\nu}^F. \tag{13}$$

The maximum value for the Landau quantum numbers, $\nu_{b,s}^m$ and ν_l^m, are defined by the conditions $\left(k^F_{b,\nu^m_{b,s},s}\right)^2 = 0$ and $\left(k^F_{l,\nu_l^m}\right)^2 = 0$, thus we have

$$\nu_{b,s}^m = \text{int}\left[\frac{(\mu_b^* - sk_bB)^2 - m_b^{*2}}{2|q_b|B}\right] \quad \text{and} \quad \nu_l^m = \text{int}\left[\frac{\mu_l^2 - m_l^2}{2|q_l|B}\right]. \tag{14}$$

In the β-equilibrium, it is possible to write any chemical potential as a linear combination of two of them, for example μ_n and μ_e, as $\mu_i = q_b^{(i)}\mu_n + q^{(i)}\mu_e$, with $q_b^{(i)}$ and $q^{(i)}$ being the baryonic and electric charge of the ith particle. These two independent chemical potentials are the Lagrange multipliers associated with the conservation of baryon number and electric charge, respectively

$$\sum_b \rho_b = \rho, \quad \sum_b q_b\rho_b - \sum_l \rho_l = 0 \tag{15}$$

2 Results

We obtained the relative population of each specie of particle as a function of the baryon density. We used the coupling constants given by Set 2 of Ref.[3] and a strong magnetic field of 3.3×10^{19} G for illustration. The upper panel of Fig 1 shows the results for the relative population of particles without taking into account the anomalous magnetic moments. The relative population of particles with anomalous magnetic moments are showed in lower panel of the same Figure. It can be seen that the incorporation of the anomalous magnetic moment increases the strangeness content of matter at medium and high densities. Besides, the proton and electron fraction increases. This effect is important for neutrino emissivity by direct Urca process, having an impact on the cooling of neutron stars.




Figure 1: Particle fractions in cold neutron star matter in β-equilibrium without and with anomalous magnetic moment of baryons included (upper and lower panels respectively.)

References

[1] N.K. Glendennig, Compact Star, Springer, New York, 2000, pp. 189.

[2] S. Chakrabarty, D. Bandyopadhyay, and S. Pal, Phys. Rev. Lett., **78**, 2898 (1997).

[3] M. Chiapparini, M.E. Bracco, A. Delfino, M. Malheiro, D.P. Menezes, C. Providência, Nucl. Phys. A **826**, 178 (2009).