# OPTIMAL SWITCHING ON AND OFF THE ENTIRE SERVICE CAPACITY OF A PARALLEL QUEUE

EUGENE FEINBERG

Department of Applied Mathematics and Statistics, Stony Brook University, Stony Brook, NY 11794, efeinberg@notes.cc.sunysb.edu

#### XIAOXUAN ZHANG

IBM T.J. Watson Research Center, Yorktown Heights, NY 10598, zhangxiaoxuan@live.com

This paper studies optimal switching on and off of the entire service capacity of an  $M/M/\infty$  queue with holding, running and switching costs where the running costs depend only on whether the system is running or not. The goal is to minimize average costs per unit time. The main result is that an average-optimal policy either always runs the system or is an (M, N)-policy defined by two thresholds M and N, such that the system is switched on upon an arrival epoch when the system size accumulates to N and is switched off upon a departure epoch when the system size decreases to M. It is shown that this optimization problem can be reduced to a problem with a finite number of states and actions, and an averageoptimal policy can be computed via linear programming. An example, in which the optimal (M, N)-policy outperforms the best (0, N)-policy, is provided. Thus, unlike the case of single-server queues studied in the literature, (0, N)-policies may not be average-optimal.

Key words: Parallel Queue, Optimal Policy, Markov Decision Process

#### **1. INTRODUCTION**

This paper studies optimal control of a parallel M/M/ $\infty$  queue with Poisson arrivals and an unlimited number of independent identical servers with exponentially distributed service times. The cost to switch the system on is  $s_1$  and the cost to switch the system off is  $s_0$ . The other costs include the linear holding cost h for each unit of time that a customer spends in the system, the running cost  $c_1$  per unit time when the system is on and the idling cost  $c_0$  per unit time when the system is off. It is assumed that  $s_0, s_1 \ge 0, s_0 + s_1 > 0, h > 0$ , and  $c_1 > c_0$ . Denote  $c = c_1 - c_0$ . Without loss of generality, let  $c_0 = 0$  and  $c_1 = c > 0$ . The goal is to minimize average costs per unit time. The main result of this paper is that either the policy that always keeps the system on is averageoptimal or, for some integers M and N, where  $N > M \ge 0$ , the so-called (M, N)-policy is averageoptimal. The (M, N)-policy switches the running system off when the number of customers in the system is not greater than M and it switches the idling system on when the number of customers in the queue reaches or exceeds N. It is shown in this paper that this optimization problem can be reduced to a problem with finite number of states and actions and an average-optimal policy can be computed via linear programming. An example when the best (0, N)-policy is not optimal is provided.

Studies on control problems for queues started around fifty years ago, and one of the first papers on this topic, Yadin & Naor [41], dealt with switching on and off a server of a single-server queue. Heyman [17] showed the optimality of a (0, N)-policy, which is usually called an *N*-policy, for M/G/1 queues. The early results on switching servers in single-server queues led to two relevant research directions:

(i) optimality of (0, *N*)-policies or their ramifications under very general assumptions such as batch arrivals, start-up and shut-down times and costs, nonlinear holding costs, known workload, and so on; see Lee & Srinivasan [29], Federgruen & So [9], Altman & Nain [1], Denardo et al. [8], and Feinberg & Kella [12];

(ii) decomposition results for queues with vacations; see Fuhrmann & Cooper [13], Hofri [18], Shanthikumar [38], Kella [23], and Kella & Whitt [24]. Sobel [39] studied (M, N)-policies for GI/G/1 queues.

As for general multi-server parallel queues, switching on and off individual servers for a parallel queue is a more difficult problem. Even for an M/M/n queue, there's no known description of an optimal switching policy for individual servers when n > 2; see Bell [3, 4], Rhee & Sivazlian [36], and Igaki [21]. Studies of stationary distributions and performance evaluations for parallel queues

with vacations (Levy & Yechiali [30], Huang et al. [19], Kao & Narayanan [22], Browne & Kella [5], Chao & Zhao [6] and Li & Alfa [31]) usually assume that vacations begin when the system is empty. Observe that, if vacations start when the system becomes empty and end simultaneously

for all the servers, the model describes a particular case of switching the entire service capacity of the system on and off. Browne & Kella [5] studied  $M/G/\infty$  queues with vacations and described how to compute the best (0, N)-policy for switching on and off the entire service capacity.

This research is motivated by two observations: (i) the problem of switching on and off the entire service capacity of the facility has an explicit solution described in this paper while there is no known explicit solution for problems with servers that can be switched on and off individually, and (ii) with the development of internet and high performance computing, many applications behave in the way described in this paper. For example, consider a service provider that uses cloud computing and pays for the time the cloud is used. When there are many service requests, it is worth paying for using the cloud, and when there is a small number of service requests, it may be too expensive to use the cloud. This paper analyzes such a situation and finds an optimal solution. In fact, the performance analysis literature treats cloud computing as a parallel queue with thousands of servers (see IBM SmartCloud Provisioning [20]), and the number of servers in the models will increase with the development of technologies. Many papers and research works on cloud computing model cloud computing facilities as multi-server queues; see Mazzucco et al. [32] and Khazaei et al. [25]. Mazzucco et al. [32] studies the revenue management problem from the perspective of a cloud computing provider and investigates the resource allocation via dynamically powering the servers on or off. However, there can be a huge number of servers in a cloud computing center, typically of the order of hundreds or thousands; see Greenberg et al. [14], Windows Azure [40], and Amazon EC2 [2]. Given that the number of servers is large and tends to increase over time with the development of new technologies, it is natural to model controlling of the facility as an  $M/M/\infty$  queue rather than an M/M/n queue if this leads to analytical advantages. Here we study a model based on an  $M/M/\infty$  queue and find a solution.

In addition to cloud computing, another example comes from the application to IT software maintenance. Kulkarni et al. [28] studied the software maintenance problem as a control problem for a queue formed by software maintenance requests generated by software bugs experienced by customers. Once a customer is served and the appropriate bug is fixed in the new software release or patch, it also provides solutions to some other customers in the queue and these customers are served simultaneously. In Kulkarni et al. [28], it was assumed that the number of customers leaving the queue at a service completion time has a binomial distribution. This problem was modeled in Kulkarni et al. [28] as an optimal switching problem for an M/G/1 queue in which a binomially distributed number of customers depending on the queue size are served each time, and the best policy was found among the policies that switch the system off when it is empty and switch it on when there are N or more customers in the system. Here we observe that after an appropriate scaling, the software maintenance problem with exponential service times and the optimal switching problem for an  $M/M/\infty$  queue have the same fluid approximations. So, the result on average-optimality of (M, N)-policies described here provides certain insights to the software maintenance problem studied in Kulkarni et al. [28].

As a conclusion to the introduction, we describe the structure of the paper and some of the main ideas. There are two main obstacles in the analysis of the  $M/M/\infty$  switching problem compared to a single-server one. First, the service intensities are unbounded, and therefore the standard reduction of continuous-time problems to discrete time via uniformization can not be applied. Second, there are significantly more known decomposition and performance analysis results for single-server queues than for parallel queues and, in particular, we are not aware of such results for  $M/M/\infty$  queues with vacations that can start when the queue is not empty. The first obstacle is resolved by

5

reducing the discounted version of the problem to negative dynamic programming instead of to discounted dynamic programming. The second obstacle is resolved by solving a discounted problem for the system that cannot be switched off. This problem is solved by using optimal stopping, where the stopping decision corresponds to starting the servers, and its solution is used to derive useful inequalities and to reduce the problem for the original  $M/M/\infty$  queue to a control problem of a semi-Markov process with finite state and action sets representing the system being always on when the number of customers exceeds a certain level.

The optimal switching problem for an M/M/∞ queue is modeled in Section 2 as a Continuous-Time Markov Decision Process (CTMDP) with unbounded transition rates. Such a CTMDP cannot be reduced to discrete time via uniformization; see, e.g., Piunovskiy & Zhang [34]. The available results for average costs require that any stationary policy defines an ergodic continuous-time Markov chain; see Guo & Zhu [16] and Guo & Hernández-Lerma [15, Assumptions 7.4. 7.5 on p. 107]. These assumptions do not hold for the problem we consider because the policy that always keeps the system off defines a transient Markov chain. Therefore, in this paper we provide a direct analysis of the problem.

Section 3 studies expected total discounted costs. Such a CTMDP can be reduced to a discretetime problem with the expected total costs; see Feinberg [11], Piunovskiy & Zhang [34]. Since transition rates are unbounded, expected total costs for the discrete-time problem cannot be presented as expected total discounted costs with the discount factor smaller than 1. However, since all the costs are nonnegative, the resulting discrete-time problem belongs to the class of negative MDPs that deal with minimizing expected total nonnegative costs, which is equivalent to maximizing expected total nonpositive rewards. For this negative MDP we derive the optimality equation, show that the value function is finite, and establish the existence of stationary discount-optimal policies; see Theorem 1.

Subsection 3.2 investigates the discounted total-cost problem limited to the policies that never switch the running system off. Such policies are called full-service policies. By using the fact that the number of customers in an  $M/G/\infty$  queue at each time has a Poisson distribution (see Ross [37, p. 70]), we compute value functions for full-service policies and then compute the discountoptimal full-service policy in Theorem 3. This is done by analyzing the optimality equation for an optimal stopping problem when stopping, in-fact, corresponds to the decision to start the system. The optimal full-service policy is defined by a number  $n_{\alpha}$  such that the system should be switched on as soon as the number of customers is greater than or equal to  $n_{\alpha}$ , where  $\alpha > 0$  is the discount rate. The important feature of the function  $n_{\alpha}$  is that it is increasing in  $\alpha$  and therefore bounded when  $\alpha \in (0, \alpha^*]$  for any  $\alpha^* \in (0, \infty)$ . In Section 3.3, the problem with the expected discounted total costs is reduced to a problem with finite state and action sets by showing in Lemma 7 that the system should always be on, if the number of customers is greater or equal than  $n_{\alpha}$ . In Section 4, by using the vanishing discount rate arguments, we prove the existence of stationary average-optimal policies and describe the (M, N)-policy in Theorem 5. A linear program (LP) for their computation is provided in Section 5. An example in which the best (0, N)-policy is not average-optimal is given in Section 6.

#### 2. PROBLEM FORMULATION

We model the above described control problem for an  $M/M/\infty$  queue as a CTMDP with a countable state space and a finite number of actions; see Kitaev & Rykov [26] and Guo & Hernández-Lerma [15]. In general, such a CTMDP is defined by the tuple {Z, A, A(z), q, c}, where Z is a countable state space, A is a finite action set, A(z) are sets of actions available in states  $z \in Z$ , and q and c are transition and cost rates, respectively. A general policy can be time-dependent, historydependent, and at a jump epoch the action that controls the process is the action selected at the previous state; see Kitaev & Rykov [26, p. 138]. An initial state  $z \in Z$  and a policy  $\pi$  define a stochastic process  $z_t$  and the expectations for this stochastic process are denoted by  $E_z^{\pi}$ . Let C(t) be the cumulative costs incurred during the time interval [0, t]. For  $\alpha > 0$ , the expected total discounted cost is

$$V_{\alpha}^{\pi}(z) = E_{z}^{\pi} \int_{0}^{\infty} e^{-\alpha t} dC(t), \qquad (1)$$

and the average cost per unit time is

$$v^{\pi}(z) = \limsup_{t \to \infty} t^{-1} E_z^{\pi} C(t).$$
<sup>(2)</sup>

Let

$$V_{\alpha}(z) = \inf_{\pi} V_{\alpha}^{\pi}(z), \tag{3}$$

$$v(z) = \inf_{\pi} v^{\pi}(z).$$
 (4)

A policy  $\pi$  is called discount-optimal if  $V^{\pi}_{\alpha}(z) = V_{\alpha}(z)$  for all initial states  $z \in Z$ . A policy  $\pi$  is called average-optimal if  $v^{\pi}(z) = v(z)$  for all initial states  $z \in Z$ .

For our problem, the states of the system change only at arrival and departure epochs, which we call jump epochs. The state of the system at time  $t \ge 0$  is  $z_t = (x_t, \delta_t)$ , where  $x_t$  is the number of customers in the system at time t, and  $\delta_t$  is the status of the servers that an arrival or departure saw at the last jump epoch. If  $\delta_t = 0$ , the severs at the last jump epoch during the interval [0, t] were off, and, if  $\delta_t = 1$ , they were on. In particular, if the last jump epoch was departure,  $\delta_t = 1$ . If the last jump epoch was an arrival, then  $\delta_t = 1$ , if the servers were on at that epoch, and  $\delta_t = 0$  otherwise. The initial state  $z_0 = (x_0, \delta_0)$  is given.

The state space is  $Z = \mathbb{N} \times \{0, 1\}$ , where  $\mathbb{N} = \{0, 1, ...\}$ , and the action set is  $A = \{0, 1\}$ , with 0 means that the system should be off and 1 meaning that the system should be on. If at time *t* the state is  $z_t = (x_t, \delta_t)$ , this means that  $x_t$  is the number of customers in the system at time *t*, and  $\delta_t \in \{0, 1\}$  is the control used at the last jump epoch during the interval [0, t]. The action sets A(z) = A for all

 $z \in Z$ . A stationary policy chooses actions deterministically at jump epochs and follows them until the next jump. In addition, the choice of an action depends only on the state of the system  $z = (x, \delta)$ , where x is the number of customers in the system and  $\delta \in \{0, 1\}$  is the status of the system prior to the last jump.

The transition rate from a state  $z = (i, \delta)$  with an action  $a \in A$  to a state z' = (j, a), where  $j \neq i$ , is q(z'|z, a) = q(j|i, a), with

$$q(j|i,a) = \begin{cases} \lambda, & \text{if } j = i+1, \\ i\mu, & \text{if } i > 0, \ a = 1, \ j = i-1, \\ 0, & \text{otherwise}, \end{cases}$$
(5)

where  $\lambda$  is the intensity of the arrival process and  $\mu$  is the service rate of individual servers. At state  $z = (i, \delta)$ , define  $q(z, a) = q(i, a) = \sum_{j \in \mathbb{N} \setminus \{i\}} q(j|i, a)$  and q(z|z, a) = q(i|i, a) = -q(i, a).

The costs include the linear holding cost *h* for a unit time that a customer spends in the system, the running cost *c* per unit time when the system is on, the start-up cost  $s_1$ , and the shut-down cost  $s_0$ , where h, c > 0,  $s_0, s_1 \ge 0$ , and  $s_0 + s_1 > 0$ . At state  $z = (i, \delta)$ , if action 1 is taken, the cost rate is hi + c; if action 0 is taken, the cost rate is hi. At state  $z = (i, \delta)$ , if action 1 is taken, the instantaneous cost  $(1 - \delta)s_1$  is incurred; if action 0 is taken, the instantaneous cost  $\delta s_0$  is incurred. The presence of instantaneous switching costs  $s_0$  and  $s_1$  complicates the situation, because standard models of CTMDPs deal only with cost rates. To resolve this complication, observe that, since  $s_0 + s_1 > 0$ , the number of times the system's status (on or off) changes up to any time N(t), when  $t < \infty$ , should be finite with probability 1 for a policy  $\pi$  and an initial state *z*. Otherwise,  $V_a^{\pi}(z) = v^{\pi}(z) = \infty$  for all  $\alpha > 0$ . Let  $0 \le t_1 < t_2 < ...$  be the times when the system is switched on or off. Let a(t) be the action 0 or 1, selected at time t. If this function has a finite number of jumps on each finite interval, we consider the function *a* being left continuous. This is consistent with the definition of a general policy as a predictable function; see Kitaev & Rykov [26, p. 138]. For an initial state *z* and a policy  $\pi$ , if the value of *N*(*t*) is finite with probability 1 for all finite *t*, we define

$$C(t) = \int_0^t (hx_t + ca(t)) dt + \sum_{n=1}^{N(t)} s_{a(t_n+)} |a(t_n+) - a(t_n)|,$$

and consider the expected total discounted costs  $V_{\alpha}^{\pi}(z)$  and the expected average costs per unit time  $v^{\pi}(z)$  defined in (1) and (2), respectively. For some  $t < \infty$ , if  $N(t) = \infty$  with positive probability, we set  $V_{\alpha}^{\pi}(z) = v^{\pi}(z) = \infty$ .

### 3. DISCOUNTED COST CRITERION

In this section we study the expected total discounted cost criterion. We first reduce the CTMDP to the discrete time MDP and then study the so called full-service policies, which are used to reduce the original problem to an equivalent problem with a finite state space.

## 3.1. REDUCTION TO DISCRETE TIME AND EXISTENCE OF STATIONARY

#### **DISCOUNT-OPTIMAL POLICIES**

In this subsection, we formulate the optimality equation and prove the existence of stationary discount-optimal policy. This is done by reducing the problem to discrete time.

When the system is on and there are i customers, the time until the next jump has an exponential distribution with intensity  $q(i, 1) = \lambda + i\mu \rightarrow \infty$  as  $i \rightarrow \infty$ . Since the jump rates are unbounded, it is impossible to present the problem as a discounted MDP in discrete-time with a discount factor smaller than 1. Thus, we shall present our problem as minimization of the expected total costs. If the decisions are chosen only at jump times, the expected total discounted sojourn time until the next jump epoch is  $\tau_{\alpha}(z,a) = \int_{0}^{\infty} (\int_{0}^{t} e^{-\alpha s} ds)q(z,a)e^{-q(z,a)t} dt = \int_{0}^{\infty} e^{-\alpha t}e^{-q(z,a)t} dt = \frac{1}{\alpha + q(z,a)}$ , and the one-step cost is  $C_{\alpha}(z,a) = |a - \delta|s_{a} + (hi + ac)\tau_{\alpha}(z,a)$  with  $z = (i, \delta)$ . For  $\alpha = 0$ , we denote  $\tau_{0}(z,a)$  and  $C_{0}(z,a)$  as  $\tau(z,a)$  and C(z,a) respectively.

By Feinberg [11, Theorem 5.6], there exists a stationary discount-optimal policy, the value function  $V_{\alpha}(z)$  satisfies the discount-optimality equation,

$$V_{\alpha}(z) = \min_{a \in A(z)} \{ C_{\alpha}(z, a) + \sum_{z' \in Z} \frac{q(z'|z, a)}{\alpha + q(z, a)} V_{\alpha}(z') \}, \qquad z \in Z,$$
(6)

and a stationary policy  $\phi$  is discount-optimal if and only if

$$V_{\alpha}(z) = C_{\alpha}(z,\phi(z)) + \sum_{z'\in Z} \frac{q(z'|z,\phi(z))}{\alpha + q(z,\phi(z))} V_{\alpha}(z), \quad z \in Z.$$

$$\tag{7}$$

Formulae (6) and (7) imply that the discounted version of the problem is equivalent to finding a policy that minimizes the expected total costs for the discrete-time MDP { $Z, A, A(z), p_{\alpha}, C_{\alpha}$ } with sub-stochastic transition probabilities  $p_{\alpha}(z'|z, a) = q(z'|z, a)/(\alpha + q(z, a))$  and with one-step cost  $C_{\alpha}(z, a)$ , where  $\alpha > 0$ .

As mentioned above, classic CTMDPs do not deal with instantaneous costs described in the previous section. However, if we replace the instantaneous costs  $s_a$ ,  $a \in \{0, 1\}$ , with the cost rates  $s(z, a) = s_a |a - \delta| (\alpha + q(z, a))$ , where  $z = (i, \delta)$ , then the expected total discounted cost until the next jump does not change for policies that change actions only at jump epochs. For an arbitrary policy, the expected total discounted cost until the next jump can either decrease or remain unchanged, if instantaneous costs are replaced with the described cost rates. This follows from Feinberg [10, Theorem 1], which implies that the defined cost rates s(z, a) correspond to the situation when only the first nonzero switching cost after the last jump is charged and the remaining switchings are free (in particular, if  $s_0, s_1 > 0$ , only the first switching is charged). Thus, a discount-optimal policy for the problem with the switching costs, and the optimality equation (6) is also the optimality equation for the original problem with the goal to minimize the expected total discounted costs.

The following lemma computes the value function under the policy that always runs the system. This lemma produces an upper bound for the value function  $V_{\alpha}$  and, in addition, it shows that the value function takes finite values. LEMMA 1. Let  $\phi$  be a policy that always runs the system. For all i = 0, 1, ...,

$$V_{\alpha}^{\phi}(i,\delta) = (1-\delta)s_1 + \frac{hi}{\mu+\alpha} + \frac{h\lambda}{\alpha(\mu+\alpha)} + \frac{c}{\alpha} < \infty.$$
(8)

*Proof.*  $V^{\phi}_{\alpha}(i,0) = s_1 + V^{\phi}_{\alpha}(i,1)$ , or equivalently,  $V^{\phi}_{\alpha}(i,\delta) = (1-\delta)s_1 + V^{\phi}_{\alpha}(i,1)$ . Observe that

$$V_{\alpha}^{\phi}(0,1) = E\left[\int_{0}^{\infty} e^{-\alpha t} \left(hX_{0}(t) + c\right) dt\right] = hE\left[\int_{0}^{\infty} e^{-\alpha t}X_{0}(t)dt\right] + \frac{c}{\alpha} = \frac{h\lambda}{\alpha(\mu+\alpha)} + \frac{c}{\alpha},\tag{9}$$

where  $X_0(t)$  is the number of busy servers at time *t* if at time 0 the system is empty. The last equality in (9) holds because, according to Page 70 in Ross [37],  $X_0(t)$  has a Poisson distribution with the mean  $\lambda \int_0^t e^{-\mu t} dt = \frac{\lambda}{\mu} (1 - e^{-\mu t})$ . Thus,

$$E\left[\int_0^\infty e^{-\alpha t} X_0(t) dt\right] = \int_0^\infty e^{-\alpha t} \frac{\lambda}{\mu} (1 - e^{-\mu t}) dt = \frac{\lambda}{\alpha(\mu + \alpha)}.$$

Also observe that

$$V^{\phi}_{\alpha}(i,1) = G_{\alpha}(i) + V^{\phi}_{\alpha}(0,1) = iG_{\alpha}(1) + V^{\phi}_{\alpha}(0,1),$$

where  $G_{\alpha}(i)$  is the expected total discounted holding cost to serve *i* customers that are in the system at time 0. Since the service times are exponential,  $G_{\alpha}(1) = E\left[\int_{0}^{\xi} e^{-\alpha t} h dt\right] = \frac{h}{\mu + \alpha}$ , where  $\xi \sim \exp(\mu)$ . In addition,  $V_{\alpha}^{\phi}(i, 0) = s_1 + V_{\alpha}^{\phi}(i, 1)$ .

We follow the conventions that  $V_{\alpha}(-1, \delta) = 0$ ,  $\sum_{\emptyset} = 0$ , and  $\prod_{\emptyset} = 1$ . The following theorem is the main result of this subsection.

THEOREM 1. For any  $\alpha > 0$  the following statements hold:

(*i*) for all i = 0, 1, ...,

$$V_{\alpha}(i,\delta) \le (1-\delta)s_1 + \frac{hi}{\mu+\alpha} + \frac{h\lambda}{\alpha(\mu+\alpha)} + \frac{c}{\alpha};$$
(10)

(ii) for all i = 0, 1, ... and for all  $\delta = 0, 1$ , the value function  $V_{\alpha}(i, \delta)$  satisfies the discount-optimality equation

$$V_{\alpha}(i,\delta) = \min_{a \in \{0,1\}} \left\{ C_{\alpha}((i,\delta),a) + \frac{q(i-1|i,a)}{\alpha + q(i,a)} V_{\alpha}(i-1,a) + \frac{q(i+1|i,a)}{\alpha + q(i,a)} V_{\alpha}(i+1,a) \right\}$$

E. Feinberg and X. Zhang

$$= \min\left\{ (1-\delta)s_1 + \frac{hi+c}{\alpha+\lambda+i\mu} + \frac{\lambda}{\alpha+\lambda+i\mu}V_{\alpha}(i+1,1) + \frac{i\mu}{\alpha+\lambda+i\mu}V_{\alpha}(i-1,1), \\ \delta s_0 + \frac{hi}{\alpha+\lambda} + \frac{\lambda}{\alpha+\lambda}V_{\alpha}(i+1,0)\right\};$$
(11)

(iii) there exists a stationary discount-optimal policy, and a stationary policy  $\phi$  is discount-optimal if and only if for all i = 0, 1, ... and for all  $\delta = 0, 1$ ,

$$\begin{aligned} V_{\alpha}(i,\delta) &= \min_{\phi(i,\delta) \in \{0,1\}} \left\{ C_{\alpha}((i,\delta),\phi(i,\delta)) + \frac{q(i-1|i,a)}{\alpha + q(i,\phi(i,\delta))} V_{\alpha}(i-1,\phi(i,\delta)) \right. \\ &+ \frac{q(i+1|i,a)}{\alpha + q(i,\phi(i,\delta))} V_{\alpha}(i+1,\phi(i,\delta)) \right\}. \end{aligned}$$

*Proof.* (i) Consider the policy  $\phi$  that always runs the system. Then  $V_{\alpha}(i, \delta) \leq V_{\alpha}^{\phi}(i, \delta)$ , and (10) follows from Lemma 1.

Statements (ii) and (iii) are corollaries from Feinberg [11, Theorem 5.6].

By Theorem 1(iii), we consider only stationary policies in the remaining part of the paper. Define  $V^1_{\alpha}(i,\delta)$  and  $V^0_{\alpha}(i,\delta)$  as

$$V_{\alpha}^{1}(i,\delta) = (1-\delta)s_{1} + \frac{hi+c}{\alpha+\lambda+i\mu} + \frac{\lambda}{\alpha+\lambda+i\mu}V_{\alpha}(i+1,1) + \frac{i\mu}{\alpha+\lambda+i\mu}V_{\alpha}(i-1,1),$$
  

$$V_{\alpha}^{0}(i,\delta) = \delta s_{0} + \frac{hi}{\alpha+\lambda} + \frac{\lambda}{\alpha+\lambda}V_{\alpha}(i+1,0).$$
(12)

#### **3.2. FULL-SERVICE POLICIES**

The class of policies that never switch the running system off is the class of all policies for the case when all the action sets A(i, 1) are reduced to the set {1}. This is a sub-model of our original model defined by (3) with the action sets A(i, 1) reduced to {1} for all i = 0, 1, ... Let  $U_{\alpha}(i, \delta)$ , i = 0, 1, ..., be the optimal expected total discounted cost under the policies that never switch the system off.

THEOREM 2. For any  $\alpha > 0$  the following statements hold:

(*i*) for all i = 0, 1, ...,

$$U_{\alpha}(i,1) = \frac{hi}{\mu + \alpha} + \frac{h\lambda}{\alpha(\mu + \alpha)} + \frac{c}{\alpha};$$

(ii) for all i = 0, 1, ..., the value function  $U_{\alpha}(i, 0)$  satisfies the optimality equation

$$U_{\alpha}(i,0) = \min\left\{s_{1} + \frac{hi + c}{\alpha + \lambda + i\mu} + \frac{\lambda}{\alpha + \lambda + i\mu}U_{\alpha}(i+1,1) + \frac{i\mu}{\alpha + \lambda + i\mu}U_{\alpha}(i-1,1), \frac{hi}{\alpha + \lambda} + \frac{\lambda}{\alpha + \lambda}U_{\alpha}(i+1,0)\right\}.$$
(13)

*Proof.* (i) For a policy  $\phi$  that never switches the running system off,  $U_{\alpha}(i, 1) = V_{\alpha}^{\phi}(i, 1)$ , and the rest follows from Lemma 1.

(ii) Since  $U_{\alpha}(i, 0)$  is the optimal discounted cost for the sub-model of the original MDP, it satisfies the discount-optimality equation. Thus, (13) follows from (11).

DEFINITION 1. For an integer  $n \ge 0$ , a policy is called an *n*-full-service policy if it never switches the running system off and switches the inactive system on if and only if there are *n* or more customers in the system. In particular, the 0-full-service policy switches the system on at time 0, if it is off, and always keeps it on. A policy is called a full-service policy if and only if it is an *n*-full-service policy for some  $n \ge 0$ .

The following theorem implies that an *n*-full-service policy is discount-optimal within the class of policies that never switch the running system off.

THEOREM 3. A policy  $\phi$  is discount-optimal within the class of the policies that never switch off the running system if and only if for all i = 0, 1, ...,

$$\phi(i,0) = \begin{cases} 1, & \text{if } i > A(\alpha), \\ 0, & \text{if } i < A(\alpha), \end{cases}$$

where

$$A(\alpha) = \frac{(\mu + \alpha)(c + \alpha s_1)}{h\mu}.$$
(14)

Before proving Theorem 3, we introduce the definition of passive policies and some lemmas. In particular, the passive policy never changes the status of the system.

DEFINITION 2. The policy  $\varphi$ , with  $\varphi(i, \delta) = \delta$  for all i = 0, 1, ... and for all  $\delta = 0, 1$ , is called passive.

LEMMA 2. For any  $\alpha > 0$ , the passive policy  $\varphi$  is not discount-optimal within the class of policies that never switch off the running system. Furthermore,  $V^{\varphi}_{\alpha}(i,0) > U_{\alpha}(i,0)$  for all i = 0, 1, ...

*Proof.* For the passive policy  $\varphi$ ,

$$V^{\varphi}_{\alpha}(i,0) = \sum_{k=0}^{\infty} \left(\frac{\lambda}{\lambda+\alpha}\right)^{k} \frac{h(i+k)}{\lambda+\alpha} = \frac{hi}{\alpha} + \frac{h\lambda}{\alpha^{2}}$$

For the policy  $\phi$  that always runs the system,

$$V^{\phi}_{\alpha}(i,0) = s_1 + \frac{hi}{\mu + \alpha} + \frac{h\lambda}{\alpha(\mu + \alpha)} + \frac{c}{\alpha}.$$
(15)

Thus

$$V_{\alpha}^{\varphi}(i,0) - V_{\alpha}^{\phi}(i,0) = \left(\frac{hi}{\alpha} + \frac{h\lambda}{\alpha^2}\right) - \left(s_1 + \frac{hi}{\mu + \alpha} + \frac{h\lambda}{\alpha(\mu + \alpha)} + \frac{c}{\alpha}\right) = \frac{hi\mu}{\alpha(\mu + \alpha)} + \frac{h\lambda\mu}{\alpha^2(\mu + \alpha)} - s_1 - \frac{c}{\alpha} > 0,$$

when *i* is large enough. Let  $i^*$  be the smallest natural integer such that the last inequality holds with  $i = i^*$ . Let the initial state be (i, 0) with  $i < i^*$ . Consider a policy  $\pi$  that keeps the system off in states  $(j, 0), j < i^*$ , and switches to a discount-optimal policy, when the number of customers in the system reaches  $i^*$ . Then  $V^{\varphi}_{\alpha}(i, 0) > V^{\pi}_{\alpha}(i, 0) \ge U_{\alpha}(i, 0)$ , where the first inequality holds because, before the process hits the state  $(i^*, 0)$ , the policies  $\varphi$  and  $\pi$  coincide, and, after the process hits the state  $(i^*, 0)$ , the policy  $\pi$ , which starting from that event coincides with  $\phi$ , incurs lower total discounting costs than the passive policy  $\varphi$ .

LEMMA 3. Let  $\psi$  be the policy that switches the system on at time 0 and keeps it on forever, and  $\pi$  be the policy that waits for one arrival and then switches the system on and keeps it on forever. Then

$$\begin{cases} V_{\alpha}^{\pi}(i,0) > V_{\alpha}^{\psi}(i,0), & \text{if } i > A(\alpha), \\ V_{\alpha}^{\pi}(i,0) < V_{\alpha}^{\psi}(i,0), & \text{if } i < A(\alpha), \\ V_{\alpha}^{\pi}(i,0) = V_{\alpha}^{\psi}(i,0), & \text{if } i = A(\alpha), \end{cases}$$

where  $A(\alpha)$  is as in (14).

Proof.

$$\begin{split} V_{\alpha}^{\pi}(i,0) - V_{\alpha}^{\psi}(i,0) &= \left(\frac{hi}{\lambda+\alpha} + \frac{\lambda}{\lambda+\alpha}\left(s_{1} + U_{\alpha}(i+1,1)\right)\right) - \left(s_{1} + U_{\alpha}(i,1)\right) \\ &= \left[\frac{hi}{\lambda+\alpha} + \frac{\lambda}{\lambda+\alpha}\left(s_{1} + \frac{h(i+1)}{\mu+\alpha} + \frac{h\lambda}{\alpha(\mu+\alpha)} + \frac{c}{\alpha}\right)\right] - \left[s_{1} + \frac{hi}{\mu+\alpha} + \frac{h\lambda}{\alpha(\mu+\alpha)} + \frac{c}{\alpha}\right] \\ &= \frac{hi}{\lambda+\alpha}\frac{\mu}{\mu+\alpha} - \frac{\alpha}{\lambda+\alpha}\left(s_{1} + \frac{c}{\alpha}\right) = \frac{h\mu}{(\lambda+\alpha)(\mu+\alpha)}\left(i - A(\alpha)\right), \end{split}$$

where the second equality holds in view of Theorem 2(i) and the rest is straightforward.

*Proof of Theorem 3.* Let  $\phi$  be a stationary discount-optimal policy within the class of the policies that never switch off the running system. Let  $\psi$  be the policy that switches the system on at time 0 and keeps it on forever, and  $\pi$  be the policy that waits for one arrival and then switches the system on and keeps it on forever. By (13),

$$V_{\alpha}^{\phi}(i,0) = \min\left\{s_1 + U_{\alpha}(i,1), \ \frac{hi}{\lambda + \alpha} + \frac{\lambda}{\lambda + \alpha}U_{\alpha}(i+1,0)\right\}.$$
(16)

We show that if  $i > A(\alpha)$ , then  $\phi(i, 0) = 1$ . Assume  $\phi(i, 0) = 0$  for some  $i > A(\alpha)$ . By Lemma 2,  $\phi(j, 0) = 1$  for some j > i. Thus, there exists an  $i^* \ge i$  such that  $\phi(i^*, 0) = 0$  and  $\phi(i^* + 1, 0) = 1$ . This implies that  $V^{\psi}_{\alpha}(i^*, 0) \ge V^{\pi}_{\alpha}(i^*, 0)$ , where  $i^* > A(\alpha)$ . By Lemma 3, this is a contradiction. Thus  $\phi(i, 0) = 1$  for all  $i > A(\alpha)$ .

If  $i < A(\alpha)$ , then Lemma 3 implies  $V_{\alpha}^{\pi}(i, 0) < V_{\alpha}^{\psi}(i, 0)$ . Thus  $\phi(i, 0) = 0$  for all  $i < A(\alpha)$ .

Let  $A(\alpha)$  be an integer and  $i = A(\alpha)$ . In this case, Lemma 3 implies  $V_{\alpha}^{\psi}(i, 0) = V_{\alpha}^{\pi}(i, 0)$ . From (13),  $V_{\alpha}^{\psi}(i, 0) = V_{\alpha}^{\pi}(i, 0) = U_{\alpha}(i, 0) = \min \left\{ V_{\alpha}^{\psi}(i, 0), V_{\alpha}^{\pi}(i, 1) \right\}$ . Thus  $\phi(i, 0) = 1$  or  $\phi(i, 0) = 0$ .

COROLLARY 1. Let

$$n_{\alpha} = \lceil A(\alpha) \rceil, \tag{17}$$

where  $A(\alpha)$  is as in (14). Then the following statements hold:

(i) if  $A(\alpha)$  is not an integer then the  $n_{\alpha}$ -full service policy is the unique stationary discountoptimal policy within the class of policies that never switch the running system off;

(ii) if  $A(\alpha)$  is an integer then there are exactly two stationary discount-optimal policies within the class of policies that never switch the running system off, and these policies are  $n_{\alpha}$ - and  $(n_{\alpha} + 1)$ -full-service policies;

(iii)

$$U_{\alpha}(i,0) = \begin{cases} \sum_{k=0}^{n_{\alpha}-i-1} \left(\frac{\lambda}{\lambda+\alpha}\right)^{k} \frac{h(i+k)}{\lambda+\alpha} + \left(\frac{\lambda}{\lambda+\alpha}\right)^{n_{\alpha}-i} \left(s_{1} + \frac{hn_{\alpha}}{\mu+\alpha} + \frac{h\lambda}{\alpha(\mu+\alpha)} + \frac{c}{\alpha}\right), & \text{if } i < n_{\alpha}, \\ s_{1} + \frac{hi}{\mu+\alpha} + \frac{h\lambda}{\alpha(\mu+\alpha)} + \frac{c}{\alpha}, & \text{if } i \ge n_{\alpha}. \end{cases}$$
(18)

*Proof.* Statements (i) and (ii) follow directly from Theorem 3 and Definition 1. Statements (i) and (ii) imply that  $V_{\alpha}^{\phi} = U_{\alpha}$ , where  $\phi$  is the  $n_{\alpha}$ -full service policy. The first line of (18) is the discounted cost to move from state (*i*, 0) to state ( $n_{\alpha}$ , 0), when the system is off, plus the discounted cost  $U_{\alpha}(n_{\alpha}, 0)$ . The second line of (18) follows from (15).

COROLLARY 2. Let  $n = \lfloor \frac{c}{h} + 1 \rfloor$ . Then there exists  $\alpha^* > 0$  such that the n-full-service policy is discount-optimal within the class of the policies that never switch the running system off for all discount rates  $\alpha \in (0, \alpha^*]$ .

*Proof.* In view of (14), the function  $A(\alpha)$  is strictly monotone when  $\alpha > 0$ . In addition,  $A(\alpha) \searrow \frac{c}{h}$  when  $\alpha \searrow 0$ . This implies that  $n_{\alpha} = n$  for all  $\alpha \in (0, \alpha^*]$ , where  $\alpha^*$  can be found by solving the quadratic inequality  $A(\alpha) \le n$ . The rest follows from Corollary 1 (i) and (ii).

## 3.3. PROPERTIES OF DISCOUNT-OPTIMAL POLICIES AND REDUCTION TO A PROBLEM WITH A FINITE STATE SPACE

This subsection introduces the properties of the discount-optimal policies formulated in Lemma 4 and Lemma 5, describes the inequalities between the major thresholds in Lemma 7 that lead to the reduction of the original infinite-state problem to a finite state problem. This reduction essentially follows from Corollary 4. Certain structural properties of discount-optimal policies are described in Theorem 4.

Define

1

$$M_{\alpha}^{*} = \begin{cases} \max\{i \ge 0 : V_{\alpha}^{0}(i,1) \le V_{\alpha}^{1}(i,1)\}, & \text{if } \{i \ge 0 : V_{\alpha}^{0}(i,1) \le V_{\alpha}^{1}(i,1)\} \neq \emptyset, \\ -1, & \text{otherwise.} \end{cases}$$
(19)

LEMMA 4. Let  $\phi$  be a stationary discount-optimal policy. Then  $\phi(i, 1) = 1$  for  $i \ge \frac{h\lambda + c(\mu + \alpha) - s_0\alpha(\mu + \alpha)}{h\mu}$ .

*Proof.* Let  $\phi(i, 1) = 0$ . Then  $V_{\alpha}^{\phi}(i, 1) > s_0 + hi/\alpha$ , since the number of customers in the system is always greater or equal than *i* and after the first arrival it is greater than *i*. Observe that  $V_{\alpha}^{\phi}(i, 1) = V_{\alpha}(i, 1) \leq U_{\alpha}(i, 1)$ . From (8),

$$s_0 + \frac{hi}{\alpha} < \frac{hi}{\mu + \alpha} + \frac{h\lambda}{\alpha(\mu + \alpha)} + \frac{c}{\alpha}$$

This inequality implies  $i < \frac{h\lambda + c(\mu + \alpha) - s_0\alpha(\mu + \alpha)}{h\mu}$ . Thus, the opposite inequality implies  $\phi(i, 1) = 1$ .

Corollary 3. For all  $\alpha > 0$ 

$$M_{\alpha}^* \le \frac{\lambda}{\mu} + \frac{(c+s_0\mu)^2}{4s_0h\mu} < \infty.$$
<sup>(20)</sup>

*Proof.* According to Lemma 4,  $M_{\alpha}^* \leq f(\alpha)$ , where  $f(\alpha) = \frac{h\lambda + c(\mu + \alpha) - s_0\alpha(\mu + \alpha)}{h\mu}$ . For  $\alpha > 0$ , the maximum of  $f(\alpha)$  equals to the expression on the right-hand side of (20).

LEMMA 5. Let  $\phi$  be a stationary discount-optimal policy. Then for any integer  $j \ge 0$  there exists an integer  $i \ge j$  such that  $\phi(i, 0) = 1$ .

*Proof.* If  $\phi(i, 0) = 0$  for all  $i \ge j$  then by Lemma 2,  $V_{\alpha}^{\phi}(j, 0) > U_{\alpha}(j, 0) \ge V_{\alpha}(j, 0)$ . This contradicts the optimality of  $\phi$ .

Define

$$N_{\alpha}^{*} = \min\{i > M_{\alpha}^{*} : V_{\alpha}^{1}(i,0) \le V_{\alpha}^{0}(i,0)\}.$$
(21)

Lemma 5 implies that  $N_{\alpha}^*$  is well defined and  $N_{\alpha}^* < \infty$  for all  $\alpha > 0$ . Before proving the relationship between  $M_{\alpha}^*$  and  $N_{\alpha}^*$ , we introduce the following lemma.

LEMMA 6. The following properties hold for the function  $V_{\alpha}(i, \delta)$ :

- (*i*) if  $V_{\alpha}(i, 0) = V_{\alpha}^{1}(i, 0)$ , then  $V_{\alpha}^{1}(i, 1) < V_{\alpha}^{0}(i, 1)$ ;
- (*ii*) if  $V_{\alpha}(i, 1) = V_{\alpha}^{0}(i, 1)$ , then  $V_{\alpha}^{0}(i, 0) < V_{\alpha}^{1}(i, 0)$ ;
- (*iii*)  $-s_1 \le V_{\alpha}(i, 1) V_{\alpha}(i, 0) \le s_0$ .

*Proof.* (i) If  $V_{\alpha}(i,0) = V_{\alpha}^{1}(i,0)$ , then  $V_{\alpha}^{1}(i,0) \le V_{\alpha}^{0}(i,0)$ . Hence  $V_{\alpha}(i,1) = V_{\alpha}(i,0) - s_1 < V_{\alpha}(i,0) + s_0 = V_{\alpha}^{0}(i,1)$ , where the inequality follows from the assumption that  $s_0 + s_1 > 0$ . This implies  $V_{\alpha}^{1}(i,1) < V_{\alpha}^{0}(i,1)$ .

(ii) If  $V_{\alpha}(i,1) = V_{\alpha}^{0}(i,1)$ , then  $V_{\alpha}^{0}(i,1) \le V_{\alpha}^{1}(i,1)$ . Hence  $V_{\alpha}(i,0) = V_{\alpha}(i,1) - s_{0} < V_{\alpha}(i,1) + s_{1} = V_{\alpha}^{1}(i,0)$ .

(iii) 
$$V_{\alpha}(i,0) \le s_1 + V_{\alpha}(i,1)$$
 because  $V_{\alpha}(i,0) = \min\{s_1 + V_{\alpha}(i,1), V_{\alpha}^0(i,0)\} \le s_1 + V_{\alpha}(i,1)$ , and  $V_{\alpha}(i,1) \le s_0 + V_{\alpha}(i,0)$  because  $V_{\alpha}(i,1) = \min\{V_{\alpha}^1(i,1), s_0 + V_{\alpha}(i,0)\} \le s_0 + V_{\alpha}(i,0)$ .

The next Lemma shows the orders among  $M_{\alpha}^*$ ,  $N_{\alpha}^*$  and  $n_{\alpha}$ . This leads to the description of the properties of discount-optimal policies in Corollary 4 that essentially reduces the problem to a finite state space problem.

LEMMA 7.  $M^*_{\alpha} < N^*_{\alpha} \le n_{\alpha}$  for all  $\alpha > 0$ .

*Proof.* The definition (21) of  $N_{\alpha}^*$  implies that  $M_{\alpha}^* < N_{\alpha}^*$ . Thus, we need only to prove that  $N_{\alpha}^* \le n_{\alpha}$ .

If  $M_{\alpha}^* = -1$ , according to (19), a discount-optimal policy should never switch the running system system off and therefore  $V_{\alpha} = U_{\alpha}$ . In view of Corollary 1,  $V_{\alpha}^{0}(i,0) < V_{\alpha}^{1}(i,0)$ , when  $i = 0, ..., n_{\alpha} - 1$ , and  $V_{\alpha}^{0}(n_{\alpha}, 0) = V_{\alpha}^{1}(n_{\alpha}, 0)$ . Thus, in this case,  $N_{\alpha}^* = n_{\alpha}$ .

Let  $M_{\alpha}^* \ge 0$ . Consider a stationary discount-optimal policy  $\varphi$  that switches the system on at state  $(N_{\alpha}^*, 0)$ . Such a policy exists in view of the definition of  $N_{\alpha}^*$ . It follows from the definition of  $M_{\alpha}^*$  that  $V_{\alpha}^1(i, 1) < V_{\alpha}^0(i, 1)$  for  $i > M_{\alpha}^*$ . Thus, the discount-optimal policy  $\varphi$  always keeps running the active system at states (i, 1) when  $i > M_{\alpha}^*$ . Observe that

$$V^{0}_{\alpha}(N^{*}_{\alpha}-1,0) < V^{1}_{\alpha}(N^{*}_{\alpha}-1,0).$$
(22)

If  $M_{\alpha}^* < N_{\alpha}^* - 1$ , (22) follows from the definition of  $N_{\alpha}^*$ . If  $M_{\alpha}^* = N_{\alpha}^* - 1$ , (22) follows from  $V_{\alpha}^0(M_{\alpha}^*, 1) \le V_{\alpha}^1(M_{\alpha}^*, 1)$  and from Lemma 6 (ii). Thus, starting from the state  $(N_{\alpha}^* - 1, 0)$ , the discount-optimal policy  $\varphi$  waits until the next arrival, then switches the system on and runs it until the number of customers in queue becomes  $M_{\alpha}^* \le N_{\alpha}^* - 1$ . For  $i = 0, 1, ..., \text{ let } F_{\alpha}^1(i)$  be the expected total discounted cost incurred until the first time  $\theta(i)$  when the number of customers in the system is *i* and the system is running, if at time 0 the system is off, there are *i* customers in queue, and the system is switched on after the first arrival and is kept on as long as the number of customers in system is greater than *i*. Let  $\theta = \theta(N_{\alpha}^* - 1)$ . Since  $\varphi$  is the discount-optimal policy,  $V_{\alpha}(N_{\alpha}^* - 1, 0) = F_{\alpha}^1(N_{\alpha}^* - 1) + [Ee^{-\alpha\theta}]V_{\alpha}(N_{\alpha}^* - 1, 1)$ .

Let  $\pi$  be a policy that switches the system on in state  $(N_{\alpha}^* - 1, 0)$  and then follows a discountoptimal policy. Then, in view of (22), the policy  $\pi$  is not discount-optimal at the initial state  $(N_{\alpha}^* - 1, 0)$ . Thus,  $V_{\alpha}^{\pi}(N_{\alpha}^* - 1, 0) > V_{\alpha}(N_{\alpha}^* - 1, 0)$ . Since  $V_{\alpha}^{\pi}(N_{\alpha}^* - 1, 0) = s_1 + V_{\alpha}(N_{\alpha}^* - 1, 1)$ ,

$$F_{\alpha}^{1}(N_{\alpha}^{*}-1) + [Ee^{-\alpha\theta}]V_{\alpha}(N_{\alpha}^{*}-1,1) < s_{1} + V_{\alpha}(N_{\alpha}^{*}-1,1),$$

and this is equivalent to

$$\left(1 - [Ee^{-\alpha\theta}]\right) V_{\alpha}(N_{\alpha}^* - 1, 1) > F_{\alpha}^1(N_{\alpha}^* - 1) - s_1.$$
(23)

Assume that  $n_{\alpha} < N_{\alpha}^*$ . Then  $n_{\alpha} \le N_{\alpha}^* - 1$  and, in view of Theorem 3,  $\psi(N_{\alpha}^* - 1, 0) = 1$  for a stationary discount-optimal policy  $\psi$  within the class of policies that never switches the system off. Thus,  $U_{\alpha}(N_{\alpha}^* - 1, 0) = V_{\alpha}^{\psi}(N_{\alpha}^* - 1, 0) = s_1 + U_{\alpha}(N_{\alpha}^* - 1, 1)$ . In addition,  $U_{\alpha}(N_{\alpha}^* - 1, 0) \le V_{\alpha}^{\varphi}(N_{\alpha}^* - 1, 0) = F_{\alpha}^1(N_{\alpha}^* - 1) + [Ee^{-\alpha\theta}]V_{\alpha}(N_{\alpha}^* - 1, 1)$ . Thus,

$$(1 - [Ee^{-\alpha\theta}])U_{\alpha}(N_{\alpha}^* - 1, 1) \le F_{\alpha}^1(i) - s_1.$$
(24)

Observe that  $\theta$  is greater than the time until the first arrival, which has the positive expectation  $\lambda^{-1}$ . Thus,  $[Ee^{-\alpha\theta}] < 1$  and  $U_{\alpha}(N_{\alpha}^* - 1, 1) \ge V_{\alpha}(N_{\alpha}^* - 1, 1)$ . (24) contradicts (23). Thus  $N_{\alpha}^* \le n_{\alpha}$ . LEMMA 8. For each  $\alpha > 0$ , the inequality  $V^1_{\alpha}(i,0) \le V^0_{\alpha}(i,0)$  holds when  $i \ge n_{\alpha}$ .

*Proof.* Fix any  $\alpha > 0$ . Consider two cases: case (i) the best full-service policy is discount-optimal, and case (ii) the best full-service policy is not discount-optimal.

Case (i). According to Corollary 1, the  $n_{\alpha}$ -full-service policy is discount-optimal. This implies that  $V_{\alpha}^{1}(i,0) \leq V_{\alpha}^{0}(i.0)$  for all  $i \geq n_{\alpha}$ .

Case (ii). Let  $\phi$  be a stationary discount-optimal policy. Assume that there exists an integer  $j \ge n_{\alpha}$ such that  $\phi(j, 0) = 0$ . Then, in view of Lemma 5, there is  $i \ge j$  such that  $\phi(i, 0) = 0$  and  $\phi(i+1, 0) = 1$ . As shown in Lemma 7,  $n_{\alpha} > M_{\alpha}^*$  and therefore  $\phi(\ell, 1) = 1$  for all  $\ell > M_{\alpha}^*$ . Thus,  $\phi(\ell, 1) = 1$  for all  $\ell > i$ . We have

$$V_{\alpha}^{\phi}(i,0) = F_{\alpha}^{1}(i) + [Ee^{-\alpha\theta(i)}]V_{\alpha}(i,1) \le s_{1} + V_{\alpha}(i,1) \Rightarrow F_{\alpha}^{1}(i) - s_{1} \le (1 - [Ee^{-\alpha\theta(i)}])V_{\alpha}(i,1), \quad (25)$$

where the stopping time  $\theta(i)$  and the expected total discounted cost  $F_{\alpha}^{1}(i)$  are defined in the proof of Lemma 7. On the other hand, since  $i \ge n_{\alpha}$ , under  $n_{\alpha}$ -full-service policy  $\pi$  we have

$$V_{\alpha}^{\pi}(i,0) = s_1 + U_{\alpha}(i,1) \le F_{\alpha}^1(i) + [Ee^{-\alpha\theta(i)}]U_{\alpha}(i,1) \Rightarrow (1 - [Ee^{-\alpha\theta(i)}])U_{\alpha}(i,1) \le F_{\alpha}^1(i) - s_1.$$
(26)

By (25) and (26), we have  $U_{\alpha}(i, 1) \leq V_{\alpha}(i, 1)$ . Since the best full-service policy is not discountoptimal,  $U_{\alpha}(i, 1) > V_{\alpha}(i, 1)$ . This contradiction implies the correctness of the lemma.

COROLLARY 4. Let  $\alpha > 0$  and  $\alpha' \in (0, \alpha]$ . For a stationary discount-optimal policy  $\phi$  for the discount rate  $\alpha'$ , consider the stationary policy  $\phi'$ ,

$$\phi'(i,\delta) = \begin{cases} \phi(i,\delta), & \text{if } i < n_{\alpha}, \\ 1, & \text{if } i \ge n_{\alpha}. \end{cases}$$

$$(27)$$

Then the policy  $\phi'$  is also discount-optimal for the discount rate  $\alpha'$ .

*Proof.* Let  $\alpha' = \alpha$ . By the definition (19) of  $M_{\alpha}^*$ , the inequality  $V_{\alpha}^1(i, 1) \leq V_{\alpha}^0(i, 1)$  holds for all  $i > M_{\alpha}^*$ . By Lemma 8 and by Corollary 1,  $V_{\alpha}^1(i, 0) \leq V_{\alpha}^0(i, 0)$  for all  $i \geq n_{\alpha}$ . In view of Lemma 7,  $M_{\alpha}^* < n_{\alpha}$ . Thus,  $V_{\alpha}^1(i, \delta) \leq V_{\alpha}^0(i, \delta)$  for all  $i \geq n_{\alpha}$  and for all  $\delta = 0, 1$ . This implies the discount-optimality of  $\phi'$  for the discount rate  $\alpha = \alpha'$ . Now let  $\alpha' \in (0, \alpha)$ . Since  $\alpha > \alpha' > 0$ , then  $n_{\alpha'} \leq n_{\alpha}$ , and 1 is an optimal decision for the discount rate  $\alpha'$  at each state  $(i, \delta)$  with  $i \geq n_{\alpha}$ .

Corollary 4 means that the system should be always run, if there are  $n_{\alpha}$  or more customers and the discount rate is not greater than  $\alpha$ . This essentially means that, in order to find a discountoptimal policy for discount rates  $\alpha' \in (0, \alpha]$ , the decision maker should find such a policy only for a finite set of states  $(i, \delta)$  with  $i = 0, 1, ..., n_{\alpha} - 1$  and  $\delta = 0, 1$ . Thus, Lemma 4 reduces the original problem of optimization of the total discounted costs to a finite-state problem, and for every  $\alpha > 0$ this finite state set is the same for all discount factors between 0 and  $\alpha$ . The following theorem describes structural properties of a discount-optimal policy for a fixed discount factor.

THEOREM 4. For each  $\alpha > 0$ , either the  $n_{\alpha}$ -full-service policy is discount-optimal, or there exists a stationary discount-optimal policy  $\phi_{\alpha}$  with the following properties:

$$\phi_{\alpha}(i,\delta) = \begin{cases} 1, & \text{if } i > M_{\alpha}^{*} \text{ and } \delta = 1, \\ 1, & \text{if } i = N_{\alpha}^{*} \text{ and } \delta = 0, \\ 1, & \text{if } i \ge n_{\alpha} \text{ and } \delta = 0, \\ 0, & \text{if } i = M_{\alpha}^{*} \text{ and } \delta = 1, \\ 0, & \text{if } M_{\alpha}^{*} \le i < N_{\alpha}^{*} \text{ and } \delta = 0. \end{cases}$$
(28)

*Proof.* Consider a stationary discount-optimal policy  $\psi$  for the discount rate  $\alpha > 0$ , and change it to  $\phi_{\alpha}$  according to (28) on the set of states specified on the right-hand side of (28). The optimality of the new policy, denoted by  $\phi_{\alpha}$ , follows from the definitions of  $M_{\alpha}^{*}$  and  $N_{\alpha}^{*}$ , and from Corollary 4.  $\Box$ 

## 4. THE EXISTENCE AND STRUCTURE OF AVERAGE-OPTIMAL POLICIES

In this section we study the average cost criteria, prove the existence of average-optimal policies and describe their properties.

DEFINITION 3. For two nonnegative integers *M* and *N* with N > M, a stationary policy is called an (M, N)-policy if

$$\phi(i,\delta) = \begin{cases} 1, & \text{if } i > M \text{ and } \delta = 1, \\ 1, & \text{if } i \ge N \text{ and } \delta = 0, \\ 0, & \text{if } i \le M \text{ and } \delta = 1, \\ 0, & \text{if } i < N \text{ and } \delta = 0. \end{cases}$$

THEOREM 5. There exists a stationary average-optimal policy and, depending on the model parameters, either the n-full service policy is average-optimal for n = 0, 1, ..., or an (M, N)-policy is average-optimal for some  $N > M \ge 0$  and  $N \le n^*$ , where

$$n^* = \lfloor \frac{c}{h} + 1 \rfloor. \tag{29}$$

In addition, the optimal average-cost value  $v(i, \delta)$  is the same for all initial states  $(i, \delta)$ ; that is,  $v(i, \delta) = v$ .

*Proof.* We first prove that either the  $n^*$ -full-service policy is average-optimal or an (M, N)policy is average-optimal for some  $N > M \ge 0$  and  $N \le n^*$ . For the initial CTMDP, consider a
sequence  $\alpha_k \downarrow 0$  as  $k \to \infty$ . Let  $\phi^k$  be a stationary discount-optimal policy for the discount rate  $\alpha_k$ .
According to Theorem 4, for each k this policy can be selected either as an  $n_{\alpha_k}$ -full-service policy
or as a  $\phi_{\alpha_k}$  policy satisfying (28). Since  $n_{\alpha_k} \le n_{\alpha_1} < (\mu + \alpha_1)(c + \alpha_1 s_1)/h\mu + 1 < \infty$  for all k = 1, 2, ...,there exists a subsequence  $\{\alpha_{k_\ell}\}$ ,  $\ell = 1, 2, ...,$  of the sequence  $\{\alpha_k\}$ , k = 1, 2, ... such that all the
policies  $\phi^{k_\ell} = \phi$ , where  $\phi$  is a stationary policy such that either (i) the policy  $\phi$  is an n-full-service

policy for some integer *n* or (ii) the policy  $\phi$  satisfies the conditions on the right hand side of (28) with the same  $M_{\alpha}^* = M$  and  $N_{\alpha}^* = N$  for  $\alpha = \alpha_{k_{\ell}}$ .

Observe that the values of  $v^{\phi}(i, \delta)$  do not depend on the initial state  $(i, \delta)$ . Indeed, in case (i), when the policy  $\phi$  is an  $n^*$ -full-service policy, the stationary policy  $\phi$  defines a Markov chain with a single positive recurrent class  $\{(i, 1) \in Z : i = 0, 1, ...\}$ , and all the states in its complement  $\{(i, 0) \in Z : i = 0, 1, ...\}$  are transient. The same is true for case (ii) with the positive recurrent class  $Z^* = \{(i, 1) \in Z : i = M, M + 1, ...\} \cup \{(i, 0) \in Z : i = M, M + 1, ..., N\}$  and with the set of transient states  $Z \setminus Z^*$ . In each case, the Markov chain leaves the set of transient states in a finite expected amount of time incurring a finite expected cost until the time the chain enters the single positive recurrent class. Thus, in each case  $v^{\phi}(i, \delta) = v^{\phi}$  does not depend on  $(i, \delta)$ .

For all initial states  $(i, \delta)$  and for an arbitrary policy  $\pi$ , we have

$$v^{\phi} = \lim_{t \to \infty} t^{-1} E^{\phi}_{(i,\delta)} C(t) \le \lim_{\alpha \downarrow 0} \alpha V^{\phi}_{\alpha}(i,\delta) \le \limsup_{\alpha \downarrow 0} \alpha V^{\pi}_{\alpha}(i,\delta) \le \limsup_{t \to \infty} t^{-1} E^{\pi}_{(i,\delta)} C(t) = v^{\pi}(i,\delta),$$

where the first equality holds because of the definition of average costs per unit time and the limit exists because both  $n^*$ -full-service policy and (M, N)-policy define regenerative processes, the second and the last inequalities follow from the Tauberian theorem (see, e.g., Korevaar [27]), and the last equality is the definition of the average cost per unit time. Since  $\pi$  is an arbitrary policy, the policy  $\phi$  is average-optimal. In addition, if  $\alpha > 0$  is sufficiently close to 0 then  $n_{\alpha} = \lceil c/h \rceil$  if c/h is not integer, and  $n_{\alpha} = c/h + 1$ , if c/h is integer. This explains why  $n^* = \lfloor \frac{c}{h} + 1 \rfloor$  in Theorem 5. In conclusion,  $v(i, \delta) = v$ , since  $v^{\phi}(i, \delta) = v^{\phi}$ . In addition, if  $n^*$ -full-service policy  $\phi$  is average-optimal, and  $\psi$  is an *n*-full-service policy for n = 0, 1, ..., then  $v^{\psi} = v^{\phi}$ .

## 5. COMPUTATION OF AN AVERAGE-OPTIMAL POLICY

In this section, we show how an optimal policy can be computed via Linear Programming. According to Theorem 5, there is an optimal policy  $\phi$  with  $\phi(i, \delta) = 1$  when  $i \ge n^* = \lfloor \frac{c}{h} + 1 \rfloor$ . Thus, the goal is to find the values of  $\phi(i, \delta)$  when  $i = 0, 1, ..., n^* - 1$  and  $\delta = 0, 1$ . To do this, we truncate the state space Z to  $Z' = \{0, 1, ..., n^* - 1\} \times \{0, 1\}$ . If the action 1 is selected at state  $(n^* - 1, 1)$ , the system moves to the state  $(n^* - 2, 1)$ , if the next change of the number of the customers in the system is a departure and the system remains in  $(n^* - 1, 1)$ , if an arrival takes place. In the latter case, the number of customers increases by one at the arrival time and then it moves according to the random work until it hits the state  $(n^* - 1, 1)$  again. Thus the system can jump from the state  $(n^* - 1, 1)$  to itself and therefore it cannot be described as a CTMDP. However, it can be described as a Semi-Markov Decision Process (SMDP); see Mine & Osaki [33, Chapter 5] and Puterman [35, Chapter 11].

We formulate our problem as an SMDP with the state set Z' and the action set  $A(z) = A = \{0, 1\}$ . If an action a is selected at state  $z \in Z'$ , the system spends an average time  $\tau'$  in this state until it moves to the next state  $z' \in Z'$  with the probability p(z'|z, a). During this time the expected cost C'(z, a) is incurred. For  $z = (i, \delta)$  with  $i = 0, 1, ..., n^* - 2$  and  $\delta = 0, 1$ , these characteristics are the same as for the original CTMDP and are given by

$$p(z'|z,a) = \begin{cases} 1, & \text{if } a = 0, \ z' = (i+1,0), \\ \frac{\lambda}{\lambda + i\mu}, & \text{if } a = 1, \ z' = (i+1,1), \\ \frac{i\mu}{\lambda + i\mu}, & \text{if } a = 1, \ z' = (i-1,1), \\ 0, & \text{otherwise}, \end{cases}$$
(30)

$$\tau'((i,\delta),a) = \begin{cases} \frac{1}{\lambda}, & \text{if } a = 0, \\ \frac{1}{\lambda + i\mu} & \text{if } a = 1, \end{cases}$$
(31)

and  $C'((i, \delta), a) = |a - \delta|s_a + (hi + ac)\tau'((i, \delta), a)$ . The transition probabilities in states  $(n^* - 1, \delta)$  with  $\delta = 0, 1$  are defined by  $p((n^* - 2, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1, \delta), 1) = (n^* - 1)\mu/(\lambda + (n^* - 1)\mu), p((n^* - 1, 1)|(n^* - 1)\mu)$ 

 $(1, \delta), 1) = \lambda/(\lambda + (n^* - 1)\mu)$ , and  $p((n^* - 1, 1)|(n^* - 1, \delta), 0) = 1$ . In the last case, the number of customers increases by 1 to  $n^*$ , the system switches on, and eventually the number of customers becomes  $n^* - 1$ .

Let  $T_i$  be the expected time between an arrival seeing *i* customers in an  $M/M/\infty$  queue and the next time when a departure leaves *i* customers behind, i = 0, 1, .... Applying the memoryless property of the exponential distribution,  $T_i = B_{i+1} - B_i$ , where  $B_i$  is the expected busy period for  $M/M/\infty$  starting with *i* customers in the system and  $B_0 = 0$ . By formula (34b) in Browne & Kella [5],

$$B_{i} = \frac{1}{\lambda} \left( e^{\rho} - 1 + \sum_{k=1}^{i-1} \frac{k!}{\rho^{k}} \left( e^{\rho} - \sum_{j=0}^{k} \frac{\rho^{j}}{j!} \right) \right),$$
(32)

where  $\rho = \frac{\lambda}{\mu}$ . Thus  $T_{n^*-1} = B_{n^*} - B_{n^*-1} = \frac{1}{\lambda} \sum_{k=0}^{\infty} \frac{\rho^{k+1}}{n^*(n^*+1)\dots(n^*+k)}$ .

The expected time  $\tau'((n^* - 1, \delta), 1)$ , where  $\delta = 0, 1$ , is the expected time until the next arrival plus  $T_{n^*-1}$ , if the next event is an arrival. Thus,  $\tau'((n^* - 1, \delta), 1) = \frac{\lambda}{\lambda + (n^* - 1)\mu} \left(\frac{1}{\lambda} + T_{n^*-1}\right)$ ,  $\delta = 0, 1$ . In addition  $\tau'((n^* - 1, \delta), 0) = \frac{1}{\lambda} + T_{n^*-1}$ ,  $\delta = 0, 1$ .

To compute the one-step cost  $C'((n^* - 1, 1), 1)$ , we define  $m_i$  as the average number of visits to state (i, 1) starting from state  $(n^* - 1, 1)$  and before revisiting state  $(n^* - 1, 1), i = n^* - 1, n^*, \ldots$  And define  $m_{i,i+1}$  as the expected number of jumps from (i, 1) to  $(i + 1, 1), i = n^* - 1, n^*, \ldots$ , and  $m_{i,i-1}$ as the expected number of jumps from (i, 1) to  $(i - 1, 1), i = n^*, n^* + 1, \ldots$ . Then  $m_{i,i+1} = \frac{\lambda}{\lambda + i\mu}m_i$ ,  $m_{i,i-1} = \frac{i\mu}{\lambda + i\mu}m_i$  and  $m_{i,i+1} = m_{i+1,i}$ . Since  $m_{n^*-1} = 1$ ,  $m_i = \prod_{i=0}^{i-n^*} \frac{\lambda}{\lambda + (n^* - 1 + j)\mu} \frac{\lambda + (n^* + j)\mu}{(n^* + j)\mu}$ ,  $i = n^*, n^* + 1, \ldots$ . (33)

Thus,

$$C'((n^*-1,1),1) = \sum_{i=n^*-1}^{\infty} m_i C((i,1),1) = \sum_{i=n^*-1}^{\infty} m_i \frac{hi+c}{\lambda+i\mu}$$

where  $C((i, 1), 1) = \frac{hi + c}{\lambda + i\mu}$ ,  $i = n^* - 1, n^*, ...$  is the cost incurred in state (i, 1) under action 1 for the original state space model; see Section 3.1. The one-step cost  $C'((n^* - 1, 0), 1) = s_1 + C'((n^* - 1, 1), 1)$ .

Let  $C_{n^*}$  be the total cost incurred in  $M/M/\infty$  until the number of customers becomes  $(n^* - 1)$  if at time 0 there are  $n^*$  customers in the system and the system is running. Then

$$C'((n^*-1,1),1) = \frac{h(n^*-1)+c}{\lambda+(n^*-1)\mu} + \frac{\lambda}{\lambda+(n^*-1)\mu}C_{n^*}$$

and this implies

$$C_{n^*} = \left(1 + \frac{(n^* - 1)\mu}{\lambda}\right) C'((n^* - 1, 1), 1) - \frac{h(n^* - 1) + c}{\lambda}.$$

We also have  $C'((n^* - 1, 0), 0) = \frac{h(n^* - 1)}{\lambda} + s_1 + C_{n^*}, C'((n^* - 1, 0), 1) = s_1 + C'((n^* - 1, 1), 1), \text{ and}$  $C'((n^* - 1, 1), 0) = s_0 + C'((n^* - 1, 0), 0).$ 

With the definitions of the transition mechanisms, sojourn times, and one-step costs for the SMDP, now we formulate the LP according to Section 5.5 in Mine & Osaki [33] or Theorem 11.4.2 and formula 11.4.17 in Puterman [35] as

$$\begin{aligned} \text{Minimize} & \sum_{z \in Z'} \sum_{a \in A} C'(z, a) x_{z,a} \\ \text{s.t.} & \sum_{a \in A(z)} x_{z,a} - \sum_{z' \in Z'} \sum_{a \in A(z)} p(z|z', a) x_{z,a} = 0, \ z \in Z', \\ & \sum_{z \in Z'} \sum_{a \in A(z)} \tau'(z, a) x_{z,a} = 1, \\ & x_{z,a} \ge 0, \ z \in Z', \ a \in A. \end{aligned}$$

$$(34)$$

Let  $x^*$  be the optimal basic solution of (34). According to general results on SMDPs in Denardo [7, Section III], for each  $z \in Z'$ , there exists at most one  $a \in \{0, 1\}$  such that  $x^*_{z,a} > 0$ . If  $x^*_{z,a} > 0$ , then for the average-optimal policy  $\phi$ ,  $\phi(z) = a$ , for a = 0, 1. If  $x^*_{z,0} = x^*_{z,1} = 0$ , then  $\phi(z)$  can be either 0 or 1. For our problem, Theorem 6 explains how  $x^* := \{x^*_{z,a} : z \in Z', a \in A\}$  can be used to construct a stationary average-optimal policy  $\phi$  with the properties stated in Theorem 5. THEOREM 6. For an optimal basic solution  $x^*$  of (34), the following statements hold:

- (i) if  $x_{(0,1),1}^* > 0$ , then any n-full-service policy is average-optimal, n = 0, 1, ...;
- (ii) If  $x_{(0,1),0}^* > 0$ , then the (0, N)-policy is average-optimal with

$$N = \begin{cases} n^*, & \text{if } \min\{i = 1, \dots, n^* - 1 : x^*_{(i,0),1} > 0\} = \emptyset; \\ \min\{i = 1, \dots, n^* - 1 : x^*_{(i,0),1} > 0\}, & \text{if } \min\{i = 1, \dots, n^* - 1 : x^*_{(i,0),1} > 0\} \neq \emptyset; \end{cases}$$
(35)

(*iii*) if  $x_{(0,1),0}^* = x_{(0,1),1}^* = 0$ , then the (M, N)-policy is average-optimal with  $M = \min\{i = 1, ..., n^* - 1 : x_{(i,1),0}^* > 0\} > 0$  and N being the same as in (35).

*Proof.* Let  $\phi^*$  be a stationary average-optimal policy defined by the optimal basic solution  $x^*$  of LP (34). Since at most one of the values  $\{x^*_{(0,1),0}, x^*_{(0,1),1}\}$  is positive and they both are nonnegative, cases (i)–(iii) are mutually exclusive and cover all the possibilities.

(i) If  $x_{(0,1),1}^* > 0$ , then the state (0, 1) is recurrent under the policy  $\phi^*$  and  $\phi^*(0, 1) = 1$ . Since the state (0, 1) is recurrent and the system should be kept on in this state, the policy  $\phi^*$  always keeps the running system on. The states corresponding to the inactive system are transient. Thus, for any *n*-full-service policy  $\phi$ , n = 0, 1, ..., we have that  $v^{\phi}(j, 0) = v^{\phi^*}(i, 0) = v$  for all i, j = 1, 2, ... Thus, any *n*-full-service policy is average-optimal.

(ii) If  $x_{(0,1),0}^* > 0$  then the state (0, 1) is recurrent under the policy  $\phi^*$  and  $\phi^*(0, 1) = 0$ . Since the state (0, 1) is recurrent, the policy  $\phi^*$  always keeps the running system on as long as the system is nonempty. By Lemma 6 (ii),  $\phi^*(0,0) = 0$ . The first constraint in LP (34) implies that  $x_{(1,0),0}^* + x_{(1,0),1}^* > 0$ . In general, if  $x_{(i,0),0}^* + x_{(i,0),1}^* > 0$  for some  $i = 1, ..., n^* - 1$ , then  $\phi^*(j,0) = 0$  if  $x_{(j,0),1}^* = 0$  for j = 0, ..., i - 1, and  $\phi^*(i,0) = 1$  if  $x_{(i,0),1}^* > 0$ . Otherwise, if  $x_{(i,0),0}^* + x_{(i,0),1}^* = 0$  for all  $i = 1, ..., n^* - 1$ ,  $\phi^*(i,0)$  can be arbitrary and we define  $\phi^*(i,0) = 0$  for  $i = 0, 1, ..., n^* - 1$ . Thus, formula (35) defines the minimal number *N* of customers in the system, at which the inactive system should be switched on by the average-optimal policy  $\phi^*$ . We recall that the SMDP is defined for the LP in the way

that the system always starts on in state  $(n^*, 0)$ . Thus, the policy  $\phi^*$  always keep running the active system if the system is not empty, switches it off when the system becomes empty, and switches on the inactive system when the number of customers becomes *N*. If there are more than *N* customers when the system is inactive, the corresponding states are transient. The defined (0, N)-policy starts the system in all these states, and therefore it is average-optimal.

(iii) If  $x_{(0,1),0}^* = x_{(0,1),1}^* = 0$  then the state (0, 1) is transient under the policy  $\phi^*$ . In transient states the average-optimal policy  $\phi^*$  can be defined arbitrary. First observe that  $x_{(i,1),0}^* > 0$  for some  $i = 1, \ldots, n^* - 1$  and therefore M is well-defined in the theorem. Indeed, if  $x_{(i,1),0}^* = 0$  for all states  $i = 0, \ldots, n^* - 1$ , we can set  $\phi^*(i, 1) = 1$  for all these values of i. This means that in the original Markov chain, where the running system is always kept on when the number of customers in the system is greater or equal than  $n^*$ , the system is always on. Since the birth-and-death for an  $M/M/\infty$  system is positive recurrent, we have a contradiction. Since the state (M, 1) is recurrent for the Markov chain defined by the policy  $\phi^*$ , this policy always keeps the running system on when the number of the customers in the system is M or more. Since  $x_{(i,\delta),a}^* = 0$  for i < M and for all  $\delta, a = 0, 1$ , we can define  $\phi^*(i, \delta)$  arbitrarily when i < M. Let  $\phi(i, \delta) = 0$ , when i < M and  $\delta = 0, 1$ . Similar to case (i), the policy  $\phi^*$  prescribes to keep inactive system off as long as the number of customers in the system is less than N, switches it on when this number becomes N, and it can be prescribed to switch the inactive system on when the number of customers is greater than N, because all such states are transient. Thus, the defined (M, N)-policy is optimal.

Similar to (34), the LP can be formulated to find the discount-optimal policy. However, this paper focuses on average-optimality criteria, so we do not elaborate the LP for discount-optimality here.

## 6. FINDING THE BEST (0, N)-POLICY AND ITS NON-OPTIMALITY

In this section we discuss how to compute the best (0, N)-policy and show that it may not be average-optimal. To do the latter, we consider an example.

Before providing the example, we show how to find the best (0, N)-policy. This problem was studied by Browne & Kella [5] for the M/G/1 queue without the running cost. Here we extend their solution to the case with running cost. Let  $\psi_N$  be a (0, N)-policy. The average cost under  $\psi_N$ can be found by formula (26) in Browne & Kella [5] by replacing the set up cost there with the sum of switching costs and running costs  $s_0 + s_1 + cB_N$ , where  $B_N$  is as in (32) or formula (34b) in Browne & Kella [5]. This implies

$$v^{\psi_N} = hl_N + \frac{s_0 + s_1 + cB_N}{N/\lambda + B_N},$$
(36)

where  $l_N$  is the expected number of customers in the system under (0, *N*)-policy. By formulae (22), (23) in Browne & Kella [5],

$$l_N = \rho + \frac{N-1}{2} \frac{N}{N+\lambda B_N}.$$
(37)

The optimal  $N^*$  for the best (0, N)-policy is found by

$$N^* = \underset{N}{\arg\min} v^{\psi_N}.$$
 (38)

The following theorem extends Theorem 6 in Browne & Kella [5] to non-negative running cost  $c \ge 0$ .

THEOREM 7. Let

$$\tilde{N} = \min\left\{N \ge \frac{c}{h} : \frac{N(N+1)}{2\lambda} \ge \frac{s_0 + s_1}{h}\right\},\tag{39}$$

then for every  $N \ge \tilde{N}$  we have that  $v^{\psi_N} < v^{\psi_{N+1}}$ , hence  $\inf_{N\ge 1} v^{\psi_N} = \min_{1\le N\le \tilde{N}} v^{\psi_N}$ .

*Proof.* To avoid notation conflict, let  $b_N$  be  $a_N$  defined as in formula (29) in Browne & Kella [5]. Note that  $\frac{1}{\lambda} + T_N = b_{N-1}, N \ge 1$  and  $B_N + \frac{N}{\lambda} = \sum_{i=0}^{N-1} b_i$ . By (36) and (37), we have for (0, *N*)-policy

 $\psi_N$  that

$$v^{\psi_N} = h\left(\rho + \frac{N-1}{2} \frac{N/\lambda}{\sum_{i=0}^{N-1} b_i}\right) + \frac{s_0 + s_1 + c\left(\sum_{i=0}^{N-1} b_i - N/\lambda\right)}{\sum_{i=0}^{N-1} b_i}.$$

Thus  $v^{\psi_N} < v^{\psi_{N+1}}$ , if hN - c > 0 and

$$\left(\frac{h(N-1)}{2\lambda} + \frac{s_0 + s_1}{N} - \frac{c}{\lambda}\right) / \left(\frac{hN}{\lambda} - \frac{c}{\lambda}\right) < \frac{\sum_{i=0}^{N-1} b_i}{Nb_N}$$

Straightforward calculations show that, if  $N \ge \tilde{N}$  in (39), the left hand side  $\le 1$  in the above inequality, and since the right hand side is always greater than 1 since  $\{b_i : i \ge 0\}$  is decreasing, thus the result follows.

Theorem 7 implies that an average-optimal (0, N)-policy can also be found by solving the LP (34) with the state space  $Z'' = \{(i, \delta) : i = 0, 1, ..., \tilde{N} - 1, \delta = 0, 1\}$  and with the new action set  $A''(\cdot)$  defined as  $A''(0, 1) = \{0\}$ ,  $A''(i, 1) = \{1\}$  for  $i = 1, ..., \tilde{N} - 1$ , and  $A''(i, 0) = \{0, 1\}$  for  $i = 1, ..., \tilde{N} - 1$ .

EXAMPLE 1. We consider our model with arrival rate  $\lambda = 2$ , service rate  $\mu = 1$  for each server, holding cost rate h = 1, service cost rate c = 100, and switching cost  $s_0 = s_1 = 100$ . We implement it in (34) and run the LP with CPLEX in MatLab. We compute  $n^*$  as  $n^* = \lfloor \frac{c}{h} + 1 \rfloor = 101$ . Thus  $Z' = \{(i, \delta)\}$ , with i = 0, 1, ..., 100 and  $\delta = 0, 1$ . For the found solutions of (34),  $x_{(i,0),1}^* > 0$ , for i = $38; x_{(i,0),0}^* > 0$ , for  $i = 4, ..., 38; x_{(i,1),1}^* > 0$ , for  $i = 5, ..., 40; x_{(i,1),0}^* > 0$ , for i = 4; and  $x_{z,a}^* = 0$  for all the other  $z \in Z', a \in A'$ . By Theorem 6, the average-optimal policy  $\phi$  is (M, N)-policy with M = 4and N = 39. The average cost of the (4, 39)-policy is  $v^{\phi} \approx 43.39$ . The best (0, N)-policy can be found by solving (38). Substituting (32) and (37) in (36), we have  $N^* = 47$  and the corresponding average cost is  $v^{\psi_{N^*}} \approx 51.03 > v^{\phi}$ .

#### Acknowledgment

This research was partially supported by NSF grants CMMI-0928490 and CMMI-1335296. The authors thank Manasa Mandava for useful comments.

#### References

- 1. Altman, E. & Nain, P. (1993). Optimal control of the M/G/1 queue with repeated vacations of the server. *Automatic Control, IEEE Transactions on*, 38(12), 1766–1775.
- Amazon EC2 (2012). Amazon Elastic Compute Cloud, User Guide. http://aiweb.techfak.uni-bielefeld. de/content/bworld-robot-control-software/. Online; accessed 03/01/2013.
- 3. Bell, C. E. (1975). Turning off a server with customers present: Is this any way to run an M/M/c queue with removable servers? *Operations Research*, 23(3), 571–574.
- 4. Bell, C. E. (1980). Optimal operation of an M/M/2 queue with removable servers. *Operations Research*, 28(5), 1189–1204.
- 5. Browne, S. & Kella, O. (1995). Parallel service with vacations. *Operations Research*, 43(5), 870–878.
- 6. Chao, X. & Zhao, Y. Q. (1998). Analysis of multi-server queues with station and server vacations. *European Journal of Operational Research*, 110(2), 392–406.
- 7. Denardo, E. (1970). On linear programming in a Markov Decision Problem. *Management Science*, 16(5), 281–288.
- 8. Denardo, E., Feinberg, E., & Kella, O. (1997). Stochastic monotonicity for stationary recurrence times of first passage heights. *The Annals of Applied Probability*, 7(2), 326–339.
- 9. Federgruen, A. & So, K. (1991). Optimality of threshold policies in single-server queueing systems with server vacations. *Advances in Applied Probability*, 23(2), 388–405.
- Feinberg, E. (1994). A generalization of 'expectation equals reciprocal of intensity' to nonstationary distributions. *Journal of Applied Probability*, 31(1), 262–267.
- 11. Feinberg, E. (2012). Reduction of discounted continuous-time MDPs with unbounded jump and reward rates to discrete-time total-reward MDPs. In *In Optimization, Control, and Applications of Stochastic Systems, D. Hernaández-Hernaández and J.A. Minjárez-Sosa Birkhäuser (Eds.)* (pp. 77–97). Springer, New York. http://www.ams.sunysb.edu/~feinberg/public/Feinberg\_HLFest\_Revision.pdf.
- 12. Feinberg, E. & Kella, O. (2002). Optimality of D-policies for an M/G/1 queue with a removable server. *Queueing Systems*, 42(4), 355–376.
- Fuhrmann, S. & Cooper, R. B. (1985). Stochastic decompositions in the M/G/1 queue with generalized vacations. *Operations Research*, 33(5), 1117–1129.

- 14. Greenberg, A., Hamilton, J., Maltz, D., & Patel, P. (2008). The cost of a cloud: research problems in data center networks. *ACM SIGCOMM Computer Communication Review*, 39(1), 68–73.
- 15. Guo, X. & Hernández-Lerma, O. (2005). Continuous-Time Markov Decision Processes. Springer-Verlag, Berlin.
- 16. Guo, X. & Zhu, W. (2002). Denumerable-state continuous-time Markov Decision Processes with unbounded transition and reward rates under the discounted criterion. *Journal of Applied Probability*, 39(2), 233–250.
- 17. Heyman, D. (1968). Optimal operating policies for M/G/1 queuing systems. *Operations Research*, 16(2), 362–382.
- 18. Hofri, M. (1986). Queueing systems with a procrastinating server. *ACM SIGMETRICS Performance Evaluation Review*, 14(1), 245–253.
- 19. Huang, C., Brumelle, S., Sawaki, K., & Vertinsky, I. (1977). Optimal control for multi-servers queueing systems under periodic review. *Naval Research Logistics Quarterly*, 24(1), 127–135.
- 20. IBM SmartCloud Provisioning (2009). Ibm smartcloud provisioning, version 2.1. http://pic.dhe.ibm.com/ infocenter/tivihelp/v48r1/topic/com.ibm.scp.doc\_2.1.0/Raintoc.pdf.
- Igaki, N. (1992). Exponential two server queue with n-policy and general vacations. *Queueing Systems*, 10(4), 279–294.
- 22. Kao, E. P. & Narayanan, K. S. (1991). Analyses of an M/M/N/queue with server's vacations. *European Journal of Operational Research*, 54(2), 256–266.
- 23. Kella, O. (1989). The threshold policy in the M/G/1 queue with server vacations. *Naval Research Logistics*, 36(1), 111–123.
- 24. Kella, O. & Whitt, W. (1991). Queues with server vacations and lévy processes with secondary jump input. *The Annals of Applied Probability*, 1(1), 104–117.
- 25. Khazaei, H., Misic, J., & Misic, V. (2012). Performance analysis of cloud computing centers using M/G/m/m+ r queuing systems. *Parallel and Distributed Systems, IEEE Transactions on*, 23(5), 936–943.
- 26. Kitaev, M. & Rykov, V. (1995). Controlled Queueing Systems. CRC Press, Boca Raton.
- 27. Korevaar, J. (2004). Tauberian Theory: A Century of Developments, volume 329. Springer, New York.

- 28. Kulkarni, V., Kumar, S., Mookerjee, V., & Sethi, S. (2009). Optimal allocation of effort to software maintenance: A queuing theory approach. *Production and Operations Management*, 18(5), 506–515.
- Lee, H. & Srinivasan, M. (1989). Control policies for the M<sup>x</sup>/G/1 queueing system. Management Science, 35(6), 708–721.
- 30. Levy, Y. & Yechiali, U. (1976). An M/M/s queue with servers' vacations. *Information Systems and Operational Research*, 14(2), 153–163.
- 31. Li, W. & Alfa, A. S. (2000). Optimal policies for M/M/m queue with two different kinds of (n, t)-policies. *Naval Research Logistics (NRL)*, 47(3), 240–258.
- 32. Mazzucco, M., Dyachuk, D., & Deters, R. (2010). Maximizing cloud providers' revenues via energy aware allocation policies. In *Cloud Computing (CLOUD), 2010 IEEE 3rd International Conference on* (pp. 131–138).
- Mine, H. & Osaki, S. (1970). *Markovian Decision Processes*, volume 25. Elsevier Publishing Company, New York.
- 34. Piunovskiy, A. & Zhang, Y. (2012). The transformation method for continuous-time Markov Decision Processes. *Journal of Optimization Theory and Applications*, 154(2), 691–712.
- 35. Puterman, M. (1994). *Markov Decision Processes: Discrete Stochastic Dynamic Programming*. John Wiley & Sons, Inc., New York.
- 36. Rhee, H.-K. & Sivazlian, B. (1990). Distribution of the busy period in a controllable M/M/2 queue operating under the triadic (0, k, n, m) policy. *Journal of Applied Probability*, (pp. 425–432).
- 37. Ross, S. (1996). Stochastic Processes. John Wiley & Sons, Inc., New York.
- 38. Shanthikumar, J. (1988). On stochastic decomposition in m/g/1 type queues with generalized server vacations. *Operations Research*, 36(4), 566–569.
- 39. Sobel, M. (1969). Optimal average-cost policy for a queue with start-up and shut-down costs. *Operations Research*, 17(1), 145–162.
- 40. Windows Azure (2010). Introducing the windows azure platform. David Chappell & Associates White Paper.
- 41. Yadin, M. & Naor, P. (1963). Queueing systems with a removable service station. *Operations Research*, 14(4), 393–405.