

# The network-level reproduction number and extinction threshold for vector-borne diseases

Ling Xue \*, Caterina Scoglio

Kansas State Epicenter, Department of Electrical & Computer Engineering,  
Kansas State University, U.S. 66506

\* E-mail: lxue@ksu.edu

## Abstract

The reproduction number of deterministic models is an essential quantity to predict whether an epidemic will spread or die out. Thresholds for disease extinction contribute crucial knowledge on disease control, elimination, and mitigation of infectious diseases. Relationships between the basic reproduction numbers of two network-based ordinary differential equation vector-host models, and extinction thresholds of corresponding continuous-time Markov chain models are derived under some assumptions. Numerical simulation results for malaria and Rift Valley fever transmission on heterogeneous networks are in agreement with analytical results without any assumptions, reinforcing the relationships may always exist and proposing a mathematical problem of proving their existences in general. Moreover, numerical simulations show that the reproduction number is not monotonically increasing or decreasing with the extinction threshold. Key parameters in predicting uncertainty of extinction thresholds are identified using Latin Hypercube Sampling/Partial Rank Correlation Coefficient. Consistent trends of extinction probability observed through numerical simulations provide novel insights into mitigation strategies to increase the disease extinction probability. Research findings may improve understandings of thresholds for disease persistence in order to control vector-borne diseases.

Keywords: Extinction thresholds; Reproduction number; Network; Branching process; Vector-borne diseases

## 1 Introduction

Vector-borne diseases greatly impact health of humans and animals and are among the leading causes of worldwide death every year [16]; almost half of the world's population is infected with at least one type of vector-borne diseases and millions of people die of vector-borne diseases every year [9]. These diseases also cause significant economic losses in areas of animal trade, agriculture, health care, tourism, as well as destroy ecosystems of throughout the world. Therefore, control and prevention of vector-borne diseases are both economical and humane. Efficient interventions require a good understanding of disease transmission and persistence, and dynamic modeling of vector-borne diseases may contribute greatly to this end [14]. A model may be used to learn many characteristics of an outbreak such as: whether or not an outbreak may occur, the size of the outbreak, the duration time of the outbreak, or the probability for the epidemic to die out [6]. Efficient mitigation strategies deduced from model results may stop an outbreak at early stages by reducing spreading parameters [6].

Globalization of trade and travel is one of the key factors driving the emergence of vector-borne diseases; heterogeneous structure also plays an important role on dynamics of infectious diseases [20]. Modeling the spatial spread of vector-borne diseases is a challenging task [3], but one possible approach is to consider a meta-population as a directed graph, or a network, with each vertex representing a subpopulation in a location, and links placed between two locations

if there is a possibility of transmission, such as movement or proximity [5]. Network models are more widely used in epidemiology to understand the spread of infectious diseases through connected populations [27, 39].

The basic reproduction number,  $R_0$  defined as the number of secondary cases produced by an infected individual in a naive population [12] is an important threshold on epidemiology among many others, such as type reproduction number [32], target reproduction [34], and threshold index for epidemicity [18]. The basic reproduction number is an important metric, predicting whether a disease will spread or die out in deterministic population and communicable disease theory [2]. If  $R_0 > 1$ , one infectious individual generally produces more than one infection, leading to spread of an epidemic, whereas if  $R_0 < 1$ , one infectious individual generates less than one infection on average [10], and epidemic may die out [12]. The same trajectory can always be observed with deterministic models given the same initial conditions [21]. If it is possible for an epidemic to occur again, a real world epidemic does not allow us to observe that the same infection happens to the same person at the same time [21]. Moreover, deterministic models have the shortcoming that the number of infected individuals may go to less than one [23].

In comparison, Markov chain models are more realistic in the sense of only taking integer values instead of continuously varying quantities [23] and taking into account chances by approximating or mimicking the random or probabilistic factors. The last infectious individual may recover before the infection is transmitted to other susceptible individuals so that the disease may become extinct [23]. Consequently, an infection introduced to a completely susceptible population may not invade the system even if  $R_0 > 1$  [23]. Threshold for the extinction of an infectious disease to occur and probability of disease extinction are of interests. Bienaymé-Galton-Watson branching processes are widely used to study extinction of diseases involving multi-type infections.

Lloyd [23] reviewed theory of branching processes and computed extinction probability using branching processes for Ross malaria model [33] taking into account stochasticity and heterogeneity. Pénişon [28] presented several statistical tools to study extinction of populations consisted of different types of individuals, and their behaviors before extinction and in the case of a very late extinction. Allen and Lahodny Jr [1] computed reproduction numbers for deterministic models, and extinction thresholds for corresponding continuous-time Markov chain (CTMC) models using continuous-time branching process, and derived their relationships. A CTMC model is a stochastic counterpart of a deterministic ordinary differential equation (ODE) model [1]. Lahodny Jr and Allen [22] estimated probability of disease extinction for a Susceptible-Infected-Susceptible (SIS) multipatch model and illustrated some differences between thresholds for deterministic models and stochastic models numerically. Allen and van den Driessche [2] established connections between extinction thresholds for continuous-time models and discrete-time models and illustrated the relations through numerical simulations. Although probability of disease extinction is defined as the probability for the number of infections to become zero when time goes to infinity, various numerical approximations for many types of models within finite time showed good agreement with predicted extinction probability using branching processes [1, 2, 22].

Deriving relationships between reproduction numbers and extinction thresholds is a complex task for vector-borne diseases transmitted on heterogeneous networks due to too many parameters and large size matrices. According to current knowledge, very little research has studied it. The objectives of this research are to relate the extinction threshold,  $E_0$  in a stochastic setting and the reproduction number,  $R_0$  in a deterministic setting for vector-host meta-population models, as well as gain understandings in how to increase extinction probability.

The contribution of our work is summarized as follows.

1. Relationships between extinction thresholds and the reproduction numbers are derived for network-based vector-host models under some assumptions.
2. Numerical simulations show that the relationships still exist after removing above assumptions.
3. Consistent trends of extinction probability varying with disease parameters are observed through extensive numerical simulations.
4. The key parameters in predicting uncertainty of the extinction threshold are identified using Latin Hypercube Sampling/Partial Rank Correlation Coefficient (LHS/PRCC).
5. The relationship between varying disease parameters and potential mitigation strategies is biologically interpreted.

This paper is organized as follows. Section 2 reviews the next generation matrix approach for computing  $R_0$  and the branching process for deriving  $E_0$ . Section 3 calculates  $R_0$  for a deterministic vector-host model in which transmission dynamics of vectors are described by a Susceptible-Infected (SI) model and transmission dynamics of hosts are described by an SIS model. We relate  $E_0$  of corresponding CTMC model and  $R_0$  analytically. In Section 4, an analogue of results in Section 3 is obtained for a model in which transmission dynamics of vectors are described by a Susceptible-Exposed-Infected (SEI) model and transmission dynamics of hosts are described by a Susceptible-Exposed-Infected-Recovered (SEIR) model. Local transmission and trans-location transmission due to proximity for vector-borne diseases are both considered in Sections 3 and 4. In Section 5, the relationships derived in Sections 3 and 4 are numerically shown to hold without any assumptions for simplified malaria and Rift Valley fever meta-population models. The sensitivity test sorted out the key parameters in predicting uncertainty of extinction probability. Relationships between varying parameters and extinction probabilities are explored through extensive simulations for homogeneous populations and a two-node network. Section 6 provides a summary and discussion of mathematical derivations and simulation results.

## 2 Preliminary

The next generation matrix approach used to compute  $R_0$  for compartmental models is reviewed here, followed by a review of the multitype branching process approximation used to derive  $E_0$  for corresponding CTMC models.

### 2.1 Computation of $R_0$ using the next generation matrix approach

The next generation matrix approach is frequently used to compute  $R_0$ . In this section, we quickly review this approach. For more details, we refer to [11, Chapter 5], [38]. For simplicity, let  $Y_i, i = 1, \dots, m$  stand for compartments that are only related to infected and asymptotically infected individuals. The original nonlinear system of ODEs including these compartments can be written as  $\frac{\partial Y_i}{\partial t} = \mathcal{F} - \mathcal{V}$ , where  $\mathcal{F} = (\mathcal{F}_i)$  and  $\mathcal{V} = (\mathcal{V}_i)$  represent new infections and transfer between compartments, respectively. Moreover,  $\mathcal{F}_i$  represents the rate at which new infections appear in compartment  $i$ , and  $\mathcal{V}_i = \mathcal{V}_i^- - \mathcal{V}_i^+$ , where  $\mathcal{V}_i^-$  (resp.  $\mathcal{V}_i^+$ ) represents the rate at which individuals transfer from (resp. into) compartment  $i$ . The Jacobian matrices  $F$

representing transmission, and  $V$  representing transition are defined as:

$$F = [\frac{\partial \mathcal{F}_i(x^0)}{\partial x_j}], \quad V = [\frac{\partial \mathcal{V}_i(x^0)}{\partial x_j}], \quad (1)$$

where  $x^0$  denotes disease free equilibrium (DFE), and  $x_j$  is the number or proportion of infected individuals in compartment  $j$ , where  $j = 1, \dots, m$ . Matrix  $F$  is nonnegative and  $V$  is a nonsingular M-matrix.

Matrix  $FV^{-1}$  is called the next generation matrix. The  $(i, k)$  entry of  $FV^{-1}$  indicates the expected number of new infections in compartment  $i$  produced by the infected individual originally introduced into compartment  $k$ , where  $i, k = 1, \dots, m$ .

The *reproduction number*,  $R_0$ , is defined as the spectral radius of  $FV^{-1}$ , denoted by  $\rho(FV^{-1})$ .

## 2.2 Deriving $E_0$ using branching process approximation

Calculating the probability of disease extinction is one of the most interesting applications of branching process. The branching process may lead to disease extinction or persistence. We are interested in the conditions under which a disease may become extinct and the probability for this event to occur. First, we review the approach of using branching process to compute extinction threshold and extinction probability for multi-type infections.

We refer to [1, 28] for the rest of this section. Let  $\vec{X}(t) = (X_1(t), \dots, X_n(t))^T : t \in (0, \infty)$  be a set of discrete-valued vector random variables. Assume that individuals of type  $i$  produce individuals of type  $j$  and that the number of infected individuals produced by type  $i$  are independent of the number of infected individuals produced by other individuals of type  $i$  or type  $j$  for  $i, j = 1, \dots, n, i \neq j$ . Additionally, individuals of type  $i$  have identical probability generating function (pgf). Let  $\{X_{ji}\}_{j=1}^n$  be the offspring random variables for type  $i$ , where  $X_{ji}$  is the number of infected individuals of type  $j$  produced by individuals of type  $i$ . The probability that one individual of type  $i$  produces  $x_j$  infected individuals of type  $j$  is given as

$$P_i(x_1, \dots, x_n) = \text{Prob}\{X_{1i} = x_1, \dots, X_{ni} = x_n\}.$$

The offspring pgf array  $(g_1, \dots, g_n) : [0, 1]^n \rightarrow [0, 1]^n$ , is defined as

$$g_i(w_1, \dots, w_n) = \sum_{x_n=0}^{\infty} \dots \sum_{x_1=0}^{\infty} P_i(x_1, \dots, x_n) w_1^{x_1} \dots w_n^{x_n}. \quad (2)$$

Note that a trivial fixed point of  $(g_1, \dots, g_n)$  always exists at  $\mathbf{1} = (1, \dots, 1)$ .

We denote by  $M = [m_{ij}]_{n \times n}$  the expectation matrix of offspring distribution which is non-negative, where  $m_{ij} := \frac{\partial g_i}{\partial w_j}|_{x=1} < \infty$  represents the expected number of new infected individuals of type  $j$  produced by an individual of type  $i$ .

The *extinction threshold*,  $E_0$  is defined as the spectral radius of the expectation matrix, denoted by  $\rho(M)$ .

Recall that  $(B_0)$  and  $(B_1)$  assumptions in [28] are as follows.

$(B_0)$   $g_i$  is not simple. Here, a function is called simple if it is linear with no constant term.

$(B_1)$  Matrix  $M$  is irreducible.

If  $E_0 > 1$ , under assumptions  $(B_0)$  and  $(B_1)$ , the pgf has at most one fixed point in  $(0, 1)^n$ , denoted by  $w^* = (w_1^*, \dots, w_n^*)$ , if extinction array  $w^*$  in  $(0, 1)^n$  exists. In the following, extinction array only refer to  $w^* \in (0, 1)^n$ . If  $I_j(0) = i_j$ , then disease extinction probability, denoted by  $P_E$ , is

$$P_E = \lim_{t \rightarrow \infty} \text{Prob}\{\vec{I}(t) = 0\} = w_1^{*i_1} \dots w_n^{*i_n} < 1. \quad (3)$$

If  $E_0 \leq 1$ , then

$$P_E = \lim_{t \rightarrow \infty} \text{Prob}\{\vec{I}(t) = 0\} = 1.$$

### 3 SI vector model and SIS host metapopulation model

In this section, a deterministic vector-host model in which disease transmission dynamics of vectors are described by an SI model, while transmission dynamics of hosts are described by an SIS model. The reproduction number and extinction threshold for corresponding CTMC model are analytically related.

#### 3.1 The reproduction number

The model for vectors consists of compartment  $G$  representing susceptible vectors, and compartment  $J$  representing infected vectors. Disease dynamics of hosts are modeled by an SIS model.

$$\begin{aligned} \frac{dG_i}{dt} &= \eta_i - \beta_i G_i I_i / N_i - \sum_{j=1, j \neq i}^n \omega_{ji} G_i I_j / N_j - \mu_i G_i \\ \frac{dJ_i}{dt} &= \beta_i G_i I_i / N_i + \sum_{j=1, j \neq i}^n \omega_{ji} G_i I_j / N_j - \mu_i J_i \\ \frac{dS_i}{dt} &= \psi_i + \gamma_i I_i - \alpha_i S_i J_i / N_i - \sum_{j=1, j \neq i}^n \sigma_{ji} S_i J_j / N_i - d_i S_i \\ \frac{dI_i}{dt} &= \alpha_i S_i J_i / N_i + \sum_{j=1, j \neq i}^n \sigma_{ji} S_i J_j / N_i - \gamma_i I_i - d_i I_i \end{aligned} \quad (4)$$

The recruitment rate of vectors (resp. hosts) in node  $i$  is  $\eta_i$  (resp.  $\psi_i$ ) for all  $i = 1, \dots, n$ . The rate of new infections in vectors in node  $i$  produced by local hosts, and hosts in other nodes are  $\beta_i G_i I_i / N_i$  and  $\sum_{j=1, j \neq i}^n \omega_{ji} G_i I_j / N_j$ , respectively. The death rate of susceptible and infected vectors in node  $i$  are  $\mu_i G_i$  and  $\mu_i J_i$ , respectively. The rate of host infection in node  $i$  produced by local vectors, and vectors in other nodes are  $\alpha_i S_i J_i / N_i$  and  $\sum_{j=1, j \neq i}^n \sigma_{ji} S_i J_j / N_i$ , respectively. The death rates of susceptible and infected hosts in node  $i$  are  $d_i S_i$  and  $d_i I_i$ , respectively. The rate of recovery for hosts in node  $i$  is  $\gamma_i I_i$ .

Since  $J_i$  and  $I_i$ ,  $i = 1, \dots, n$  are only compartments related to infected and asymptotically infected, system of ODEs (4) can be rewritten as follows.

$$\frac{d}{dt} \begin{bmatrix} J_1 & \dots & J_n & I_1 & \dots & I_n \end{bmatrix}^T = \mathcal{F} - \mathcal{V}.$$

A unique solution at DFE, represented by  $(G_i^0, 0, N_i^0, 0)$  exists, where  $G_i^0 = \frac{\eta_i}{\mu_i}$  and  $N_i^0 = \frac{\psi_i}{d_i}$ . The Jacobian matrices  $F$  and  $V$  defined in (1) for this model are

$$F = \begin{bmatrix} 0 & \mathcal{A} \\ \mathcal{B} & 0 \end{bmatrix}, \quad V = \begin{bmatrix} \Lambda_1 & 0 \\ 0 & \Lambda_2 \end{bmatrix},$$

where

$$\mathcal{A} = \begin{bmatrix} \hat{\beta}_1 & \hat{\omega}_{21} & \cdots & \hat{\omega}_{n1} \\ \hat{\omega}_{12} & \hat{\beta}_2 & \cdots & \hat{\omega}_{n2} \\ \cdots & \cdots & \ddots & \cdots \\ \hat{\omega}_{1n} & \hat{\omega}_{2n} & \cdots & \hat{\beta}_n \end{bmatrix}, \quad \mathcal{B} = \begin{bmatrix} \alpha_1 & \sigma_{21} & \cdots & \sigma_{n1} \\ \sigma_{12} & \alpha_2 & \cdots & \sigma_{n2} \\ \cdots & \cdots & \ddots & \cdots \\ \sigma_{1n} & \alpha_2 & \cdots & \alpha_n \end{bmatrix}, \quad (5)$$

$$\Lambda_1 = \text{diag}(\mu_1, \cdots, \mu_n), \quad \Lambda_2 = \text{diag}(d_1 + \gamma_1, \cdots, d_n + \gamma_n). \quad (6)$$

Here

$$\hat{\beta}_i = \frac{\beta_i G_i^0}{N_i^0} \quad \text{and} \quad \hat{\omega}_{ij} = \frac{\omega_{ij} G_j^0}{N_i^0}.$$

The notation  $\text{diag}(\mu_1, \mu_2, \cdots, \mu_n)$  represents the diagonal matrix with diagonal entries  $\mu_1, \cdots, \mu_n$ . To calculate  $R_0$ , we first prove the following lemma.

**Lemma 1.** *Let  $A_1, A_2$  be square matrices of the same size and  $A = \begin{bmatrix} 0 & A_1 \\ A_2 & 0 \end{bmatrix}$ , then  $\rho(A) = \sqrt{\rho(A_2 A_1)}$ .*

*Proof.* For any  $\lambda \neq 0$ ,

$$|\lambda I - A| = \begin{vmatrix} \lambda I & -A_1 \\ -A_2 & \lambda I \end{vmatrix} = \begin{vmatrix} \lambda I & -A_1 \\ 0 & \lambda I - \frac{A_2 A_1}{\lambda} \end{vmatrix} = |\lambda^2 I - A_2 A_1|. \quad (7)$$

Therefore,  $\rho(A) = \sqrt{\rho(A_2 A_1)}$  if  $\rho(A_2 A_1) \neq 0$ .

If  $\rho(A_2 A_1) = 0$ , we assume that  $\rho(A) \neq 0$ . Then there exists a  $\lambda' \neq 0$  such that  $|\lambda' I - A| = 0$ . By (7),  $|\lambda'^2 I - A_2 A_1| = 0$  for a nonzero  $\lambda'$ , contradicting the assumption that  $\rho(A_2 A_1) = 0$ . Therefore,  $\rho(A) = \sqrt{\rho(A_2 A_1)}$ .  $\square$

A direct calculation gives  $FV^{-1} = \begin{bmatrix} 0 & \mathcal{A}\Lambda_2^{-1} \\ \mathcal{B}\Lambda_1^{-1} & 0 \end{bmatrix}$ . By Lemma 1, we have the following proposition.

**Proposition 1.** *The reproduction number of the model (4) is*

$$R_0 = \sqrt{\rho(\mathcal{B}\Lambda_1^{-1}\mathcal{A}\Lambda_2^{-1})}. \quad (8)$$

### 3.2 The threshold for extinction probability

In this section, we compute  $E_0$  for corresponding CTMC of model (4). See Table 1 for state transitions and rates.

The pgfs are:

$$g_i(w_1, \cdots, w_n, u_1, \cdots, u_n) = \begin{cases} \frac{\alpha_i w_i u_i + \sum_{j=1, j \neq i}^n \sigma_{ij} w_i u_j + \mu_i}{\alpha_i + \sum_{j=1, j \neq i}^n \sigma_{ij} + \mu_i}, & \text{if } 1 \leq i \leq n, \\ \frac{\hat{\beta}_k u_k w_k + \sum_{j=1, j \neq k}^n \hat{\omega}_{kj} u_k w_j + d_k + \gamma_k}{\hat{\beta}_k + \sum_{j=1, j \neq k}^n \hat{\omega}_{kj} + d_k + \gamma_k}, & \text{if } n+1 \leq i \leq 2n, \end{cases}$$

| Description               | State transition $a \rightarrow b$              | Rate $P(a, b)$              |
|---------------------------|-------------------------------------------------|-----------------------------|
| Host birth                | $(S, I, G, J) \rightarrow (S + 1, I, G, J)$     | $\psi$                      |
| Death of $S$              | $(S, I, G, J) \rightarrow (S - 1, I, G, J)$     | $dS$                        |
| Host local infection      | $(S, I, G, J) \rightarrow (S - 1, I + 1, G, J)$ | $\alpha SJ/N$               |
| Host infection by $J_j$   | $(S, I, G, J) \rightarrow (S - 1, I + 1, G, J)$ | $\sigma_{ji} S_i J_j / N_i$ |
| Host recovery             | $(S, I, G, J) \rightarrow (S + 1, I - 1, G, J)$ | $\gamma I$                  |
| Death of $I$              | $(S, I, G, J) \rightarrow (S, I - 1, G, J)$     | $dI$                        |
| Vector birth              | $(S, I, G, J) \rightarrow (S, I, G + 1, J)$     | $\eta$                      |
| Death of $G$              | $(S, I, G, J) \rightarrow (S, I, G - 1, J)$     | $\mu G$                     |
| Vector local infection    | $(S, I, G, J) \rightarrow (S, I, G - 1, J + 1)$ | $\beta GI/N$                |
| Vector infection by $I_j$ | $(S, I, G, J) \rightarrow (S, I, G - 1, J + 1)$ | $\omega_{ji} G_i I_j / N_j$ |
| Death of $J$              | $(S, I, G, J) \rightarrow (S, I, G, J - 1)$     | $\mu J$                     |

Table 1: State transitions and rates for corresponding continuous-time Markov chain for deterministic model (4) omitting node index  $i$ .

where  $j = 1, \dots, n$ , the index  $k = i - n$  for  $n + 1 \leq i \leq 2n$ ,  $w_i$  represents  $I_{V_i} = 1, I_{H_i} = 0$ , and  $u_i$  represents  $I_{H_i} = 1, I_{V_i} = 0$  for  $i = 1, \dots, n$ .

The expectation matrix  $M$  is:

$$M = \begin{bmatrix} \Lambda_3 \Lambda_4 & \mathcal{A} \Lambda_5 \\ \mathcal{B} \Lambda_4 & \Lambda_6 \Lambda_5 \end{bmatrix}, \quad (9)$$

where  $\mathcal{A}, \mathcal{B}$  are the same as those in (5), and

$$\begin{aligned} \Lambda_3 &= \text{diag}(\alpha_1 + \sum_{i \neq 1} \sigma_{1i}, \dots, \alpha_n + \sum_{i \neq n} \sigma_{ni}), & \Lambda_4 &= \text{diag}(\frac{1}{C_1}, \dots, \frac{1}{C_n}), \\ \Lambda_6 &= \text{diag}(\hat{\beta}_1 + \sum_{i \neq 1} \hat{w}_{1i}, \dots, \hat{\beta}_n + \sum_{i \neq n} \hat{w}_{ni}), & \Lambda_5 &= \text{diag}(\frac{1}{D_1}, \dots, \frac{1}{D_n}), \\ C_i &= \alpha_i + \sum_{j \neq i} \sigma_{ij} + \mu_i, & D_i &= \hat{\beta}_i + \sum_{j \neq i} \hat{w}_{ij} + d_i + \gamma_i, \quad \text{for } i = 1, \dots, n. \end{aligned}$$

Note that if both  $\mathcal{A}$  and  $\mathcal{B}$  are positive matrices, then the assumptions  $(B_0)$  and  $(B_1)$  in [28] hold for this model.

**Lemma 2.** Let  $A_1, A_2$  be nonnegative square matrices with the same size such that  $\rho(A_2 A_1)$  is an eigenvalue of  $A_2 A_1$  and  $\Lambda, \Lambda'$  be nonnegative diagonal matrices such that  $0 \leq k_1 I \leq \begin{bmatrix} \Lambda & 0 \\ 0 & \Lambda' \end{bmatrix} \leq k_2 I$  for some real numbers  $k_1, k_2$ . Then the spectral radius of  $B = \begin{bmatrix} \Lambda & A_1 \\ A_2 & \Lambda' \end{bmatrix}$  satisfies that

$$\sqrt{\rho(A_2 A_1)} + k_1 \leq \rho(B) \leq \sqrt{\rho(A_2 A_1)} + k_2.$$

*Proof.* Since  $0 \leq \begin{bmatrix} k_1 I & A_1 \\ A_2 & k_1 I \end{bmatrix} \leq B \leq \begin{bmatrix} k_2 I & A_1 \\ A_2 & k_2 I \end{bmatrix}$ , by Theorem 4 in [40],

$$\rho\left(\begin{bmatrix} k_1 I & A_1 \\ A_2 & k_1 I \end{bmatrix}\right) \leq \rho(B) \leq \rho\left(\begin{bmatrix} k_2 I & A_1 \\ A_2 & k_2 I \end{bmatrix}\right). \quad (10)$$

By hypothesis and (7),  $\rho\left(\begin{bmatrix} 0 & A_1 \\ A_2 & 0 \end{bmatrix}\right)$  is an eigenvalue of  $\begin{bmatrix} 0 & A_1 \\ A_2 & 0 \end{bmatrix}$ . Following the fact that  $|\lambda' + k| < \lambda + k$  for any  $k > 0$  if  $|\lambda'| < \lambda$ ,

$$\rho\left(\begin{bmatrix} k_1 I & A_1 \\ A_2 & k_1 I \end{bmatrix}\right) = \rho\left(\begin{bmatrix} 0 & A_1 \\ A_2 & 0 \end{bmatrix}\right) + k_1 = \sqrt{\rho(A_2 A_1)} + k_1.$$

Similarly,  $\rho\left(\begin{bmatrix} k_2 I & A_1 \\ A_2 & k_2 I \end{bmatrix}\right) = \sqrt{\rho(A_2 A_1)} + k_2$ . Lemma follows (10) and Lemma 1.  $\square$

**Remark 1.** If both  $A_1$  and  $A_2$  are positive matrices, then  $\rho(A_2 A_1)$  is an eigenvalue of  $A_2 A_1$  by Perron-Frobenius theorem.

By Lemma 2, we have the following proposition.

**Proposition 2.** The extinction threshold of model (4) satisfies that

$$\begin{aligned} & \min_{1 \leq i \leq n} \left( \frac{\alpha_i + \sum_{j=1, j \neq i}^n \sigma_{ij}}{C_i}, \frac{\hat{\beta}_i + \sum_{j=1, j \neq i}^n \hat{\omega}_{ij}}{D_i} \right) + \sqrt{\rho(\mathcal{B}\Lambda_5 \mathcal{A}\Lambda_4)} \leq E_0 \\ & \leq \max_{1 \leq i \leq n} \left( \frac{\alpha_i + \sum_{j=1, j \neq i}^n \sigma_{ij}}{C_i}, \frac{\hat{\beta}_i + \sum_{j=1, j \neq i}^n \hat{\omega}_{ij}}{D_i} \right) + \sqrt{\rho(\mathcal{B}\Lambda_5 \mathcal{A}\Lambda_4)}. \end{aligned}$$

### 3.3 The relationship between $R_0$ and $E_0$

To obtain a theoretical relationship between  $R_0$  in (8) and  $E_0$ , we assume that

$$\frac{\mu_i}{C_i} = k_1 \quad \text{and} \quad \frac{d_i + \gamma_i}{D_i} = k_2, \quad \forall i = 1, \dots, n \quad (11)$$

for constant numbers  $k_1, k_2 \in [0, 1]$  throughout this section. The assumption can be interpreted biologically as: the probability of natural death is identical for vectors from each node, and the probability of natural death is identical for hosts from each node. The assumption shall be removed for numerical simulations in the next section.

**Theorem 1.** Under the assumption (11),

- (1) If  $R_0 \leq \frac{1-k_2}{1-\sqrt{k_1 k_2}} \leq 1$  or  $E_0 \leq \frac{1-k_2}{1-\sqrt{k_1 k_2}} \leq 1$ , then  $R_0 \leq E_0$ ;
- (2) If  $R_0 \geq \frac{1-k_1}{1-\sqrt{k_1 k_2}} \geq 1$  or  $E_0 \geq \frac{1-k_1}{1-\sqrt{k_1 k_2}} \geq 1$ , then  $R_0 \geq E_0$ .

*Proof.* Under the assumption (11),  $\Lambda_1 \Lambda_4 = k_1 I$ ,  $\Lambda_3 \Lambda_4 = (1 - k_1)I$ ,  $\Lambda_2 \Lambda_5 = k_2 I$  and  $\Lambda_6 \Lambda_5 = (1 - k_2)I$ , where  $I$  is the identity matrix. Therefore,  $M$  in (9) can be rewritten as follows,

$$M = \begin{bmatrix} 0 & k_2 \mathcal{A} \Lambda_2^{-1} \\ k_1 \mathcal{B} \Lambda_1^{-1} & 0 \end{bmatrix} + \begin{bmatrix} (1 - k_1)I & 0 \\ 0 & (1 - k_2)I \end{bmatrix}.$$

Without loss of generality, we assume that  $k_1 < k_2$ . By Lemma 2 and (8),

$$R_0 \sqrt{k_1 k_2} + 1 - k_2 \leq E_0 \leq R_0 \sqrt{k_1 k_2} + 1 - k_1. \quad (12)$$

Following (12),

$$R_0(1 - \sqrt{k_1 k_2}) - (1 - k_1) \leq R_0 - E_0 \leq R_0(1 - \sqrt{k_1 k_2}) - (1 - k_2),$$

$$\frac{1}{\sqrt{k_1 k_2}}(E_0(1 - \sqrt{k_1 k_2}) - (1 - k_1)) \leq R_0 - E_0 \leq \frac{1}{\sqrt{k_1 k_2}}(E_0(1 - \sqrt{k_1 k_2}) - (1 - k_2)).$$

Theorem follows the above two inequalities.  $\square$

**Corollary 1.** *If the further assumption is made that  $k_1 = k_2$  except assumption (11), then  $R_0 \leq 1$  if and only if  $E_0 \leq 1$ . Moreover,  $|R_0 - 1| \geq |E_0 - 1|$ .*

*Proof.* By Theorem 1 (1), if  $R_0 \leq 1$ , then  $R_0 \leq E_0$ . Assuming that  $E_0 > 1$ , by Theorem 1 (2),  $R_0 \geq E_0$ , which is a contradiction. Conversely, if  $E_0 > 1$ , then  $R_0 \leq 1$  following a similar argument. Hence,  $R_0 \leq 1$  if and only if  $E_0 \leq 1$ . This proves the first part. The second part directly follows Theorem 1.  $\square$

## 4 SEI vector model and SEIR host metapopulation model

A deterministic model in which vectors are divided into compartments  $S, E$ , and  $I$ , and hosts are classified into compartments  $S, E, I$ , and  $R$  is presented. The reproduction number for this model and the extinction threshold for corresponding CTMC model are connected.

### 4.1 The reproduction number

The following model extends the model in Section 3.1 by adding compartment  $Z$  for exposed vectors, and compartment  $E$  for exposed hosts. Other terms have identical meanings as corresponding ones in model (4). The rate at which exposed vectors and exposed hosts in node  $i$  transfer to infected compartments are  $\varphi_i Z_i$  and  $\varepsilon_i E_i$ , respectively.

$$\begin{aligned} \frac{dG_i}{dt} &= \eta_i - \beta_i G_i I_i / N_i - \sum_{j=1, j \neq i}^n \omega_{ji} G_i I_j / N_j - \mu_i G_i \\ \frac{dZ_i}{dt} &= \beta_i G_i I_i / N_i + \sum_{j=1, j \neq i}^n \omega_{ji} G_i I_j / N_j - \varphi_i Z_i - \mu_i Z_i \\ \frac{dJ_i}{dt} &= \varphi_i Z_i - \mu_i J_i \\ \frac{dS_i}{dt} &= \psi_i - \alpha_i S_i J_i / N_i - \sum_{j=1, j \neq i}^n \sigma_{ji} S_i J_j / N_i - d_i S_i \\ \frac{dE_i}{dt} &= \alpha_i S_i J_i / N_i + \sum_{j=1, j \neq i}^n \sigma_{ji} S_i J_j / N_i - \varepsilon_i E_i - d_i E_i \\ \frac{dI_i}{dt} &= \varepsilon_i E_i - \gamma_i I_i - d_i I_i \\ \frac{dR_i}{dt} &= \gamma_i I_i - d_i R_i \end{aligned} \tag{13}$$

Compartments related to infected and asymptotically infected are  $Z_i, E_i, J_i$  and  $I_i$ ,  $i = 1, \dots, n$ . The unique solution at DFE is  $(G_i^0, 0, 0, N_i^0, 0, 0, 0)$ , where  $G_i^0$  and  $N_i^0$  are the same as

those in Section 3.1. The above system of ODEs including these compartments can be rewritten as follows.

$$\frac{d}{dt} \begin{bmatrix} Z_1 & \cdots & Z_n & E_1 & \cdots & E_n & J_1 & \cdots & J_n & I_1 & \cdots & I_n \end{bmatrix}^T = \mathcal{F} - \mathcal{V}.$$

The Jacobian matrices  $F$  and  $V$  at DFE are

$$F = \begin{bmatrix} 0 & 0 & 0 & \mathcal{A} \\ 0 & 0 & \mathcal{B} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad V = \begin{bmatrix} \Lambda_7 & 0 & 0 & 0 \\ 0 & \Lambda_8 & 0 & 0 \\ -\Lambda_9 & 0 & \Lambda_1 & 0 \\ 0 & -\Lambda_{10} & 0 & \Lambda_2 \end{bmatrix},$$

where  $\Lambda_1$  and  $\Lambda_2$  are given in (6); matrices  $\mathcal{A}$  and  $\mathcal{B}$  are in Equation (5); and

$$\begin{aligned} \Lambda_7 &= \text{diag}(\varphi_1 + \mu_1, \dots, \varphi_n + \mu_n), & \Lambda_8 &= \text{diag}(\varepsilon_1 + d_1, \dots, \varepsilon_n + d_n), \\ \Lambda_9 &= \text{diag}(\varphi_1, \dots, \varphi_n), & \Lambda_{10} &= \text{diag}(\varepsilon_1, \dots, \varepsilon_n). \end{aligned}$$

By a direct calculation,

$$FV^{-1} = \begin{bmatrix} 0 & \mathcal{A}\Lambda_2^{-1}\Lambda_{10}\Lambda_8^{-1} & 0 & \mathcal{A}\Lambda_2^{-1} \\ \mathcal{B}\Lambda_1^{-1}\Lambda_9\Lambda_7^{-1} & 0 & \mathcal{B}\Lambda_1^{-1} & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Following Lemma 1,

**Proposition 3.** *The reproduction number of the model (13) is*

$$R_0 = \sqrt{\rho(\mathcal{B}\Lambda_1^{-1}\Lambda_9\Lambda_7^{-1}\mathcal{A}\Lambda_2^{-1}\Lambda_{10}\Lambda_8^{-1})}. \quad (14)$$

## 4.2 The threshold for extinction probability

State transitions and rates for corresponding CTMC of model (13) are listed in Table 2. The pgfs are:

$$g_i(w_1, \dots, w_{2n}, u_1, \dots, u_{2n}) = \begin{cases} \frac{\varphi_i u_i + \mu_i}{\varphi_i + \mu_i}, & \text{if } 1 \leq i \leq n, \\ \frac{\varepsilon_k u_i + d_k}{\varepsilon_k + d_k}, & \text{if } n+1 \leq i \leq 2n, \\ \frac{\alpha_p u_p w_{p+n} + \sum_{j=1, j \neq p}^n \sigma_{pj} u_p w_{j+n} + \mu_p}{\alpha_p + \sum_{j=1, j \neq p}^n \sigma_{pj} + \mu_p}, & \text{if } 2n+1 \leq i \leq 3n, \\ \frac{\hat{\beta}_q u_{q+n} w_q + \sum_{j=1, j \neq q}^n \hat{\omega}_{qj} u_{q+n} w_j + d_q + \gamma_q}{\hat{\beta}_q + \sum_{j=1, j \neq q}^n \hat{\omega}_{qj} + d_q + \gamma_q}, & \text{if } 3n+1 \leq i \leq 4n, \end{cases}$$

where  $w_i$  represents only  $Z_i = 1$ ,  $w_{i+n}$  represents  $E_i = 1$ ,  $u_i$  represents  $J_i = 1$ , and  $u_{i+n}$  represents  $I_i = 1$  for  $i = 1, \dots, n$ . The indexes  $k = i - n$  for  $1 \leq i \leq n$ ,  $p = i - 2n$  for  $n+1 \leq i \leq 2n$ , and  $q = i - 3n$  for  $3n+1 \leq i \leq 4n$ .

The expectation matrix  $M$  is:

$$M = \begin{bmatrix} 0 & 0 & 0 & \mathcal{A}\Lambda_5 \\ 0 & 0 & \mathcal{B}\Lambda_4 & 0 \\ \Lambda_9\Lambda_7^{-1} & 0 & I - \Lambda_1\Lambda_4 & 0 \\ 0 & \Lambda_{10}\Lambda_8^{-1} & 0 & I - \Lambda_2\Lambda_5 \end{bmatrix}.$$

Similarly, the assumptions  $(B_0)$  and  $(B_1)$  in [28] hold for this model if both  $\mathcal{A}$  and  $\mathcal{B}$  are positive matrices. By Lemmas 1 and 2, as well as Remark 1, we have the following proposition.

| Description                 | State transition $a \rightarrow b$                                | Rate<br>$P(a, b)$      |
|-----------------------------|-------------------------------------------------------------------|------------------------|
| Host birth                  | $(S, E, I, R, G, Z, J) \rightarrow (S + 1, E, I, R, G, Z, J)$     | $\psi$                 |
| Death of $S$                | $(S, E, I, R, G, Z, J) \rightarrow (S - 1, E, I, R, G, Z, J)$     | $dS$                   |
| Death of $E$                | $(S, E, I, R, G, Z, J) \rightarrow (S, E - 1, I, R, G, Z, J)$     | $dE$                   |
| Death of $I$                | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I - 1, R, G, Z, J)$     | $dI$                   |
| Death of $R$                | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R - 1, G, Z, J)$     | $dR$                   |
| Host local infection        | $(S, E, I, R, G, Z, J) \rightarrow (S - 1, E + 1, I, R, G, Z, J)$ | $\alpha SJ/N$          |
| Host infection by $J_j$     | $(S, E, I, R, G, Z, J) \rightarrow (S - 1, E + 1, I, R, G, Z, J)$ | $\sigma_{ji} SJ_j/N$   |
| Host recovery               | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I - 1, R + 1, G, Z, J)$ | $\gamma I$             |
| Host Latent to infectious   | $(S, E, I, R, G, Z, J) \rightarrow (S, E - 1, I + 1, R, G, Z, J)$ | $\varepsilon E$        |
| Vector birth                | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R, G + 1, Z, J)$     | $\eta$                 |
| Death of $G$                | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R, G - 1, Z, J)$     | $\mu G$                |
| Death of $Z$                | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R, G, Z - 1, J)$     | $\mu Z$                |
| Death of $J$                | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R, G, Z, J - 1)$     | $\mu J$                |
| Vector local infection      | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R, G - 1, Z + 1, J)$ | $\beta GI/N$           |
| Vector infection by $I_j$   | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R, G - 1, Z + 1, J)$ | $\omega_{ji} GI_j/N_j$ |
| Vector Latent to infectious | $(S, E, I, R, G, Z, J) \rightarrow (S, E, I, R, G, Z - 1, J + 1)$ | $\varphi Z$            |

Table 2: State transitions and rates for corresponding continuous-time Markov chain for deterministic model (13) omitting node index  $i$ .

**Proposition 4.** *The extinction threshold of the model (13) satisfies that*

$$\begin{aligned}
& \sqrt[4]{\rho(\Lambda_{10}\Lambda_8^{-1}\mathcal{B}\Lambda_4\Lambda_9\Lambda_7^{-1}\mathcal{A}\Lambda_5)} + \min_{1 \leq i \leq n} \left( \frac{\alpha_i + \sum_{j \neq i} \sigma_{ij}}{C_i}, \frac{\hat{\beta}_i + \sum_{j \neq i} \hat{\omega}_{ij}}{D_i} \right) \leq E_0 \\
& \leq \sqrt[4]{\rho(\Lambda_{10}\Lambda_8^{-1}\mathcal{B}\Lambda_4\Lambda_9\Lambda_7^{-1}\mathcal{A}\Lambda_5)} + \max_{1 \leq i \leq n} \left( \frac{\alpha_i + \sum_{j \neq i} \sigma_{ij}}{C_i}, \frac{\hat{\beta}_i + \sum_{j \neq i} \hat{\omega}_{ij}}{D_i} \right).
\end{aligned}$$

### 4.3 The relationship between $R_0$ and $E_0$

In this section, the assumption (11) holds and  $k_1 < k_2$ . Under the assumption (11), by Lemma 2,

$$\begin{aligned}
& \sqrt[4]{k_1 k_2 \rho(\Lambda_{10}\Lambda_8^{-1}\mathcal{B}\Lambda_1^{-1}\Lambda_9\Lambda_7^{-1}\mathcal{A}\Lambda_2^{-1})} + 1 - k_2 \leq E_0 \\
& \leq \sqrt[4]{k_1 k_2 \rho(\Lambda_{10}\Lambda_8^{-1}\mathcal{B}\Lambda_1^{-1}\Lambda_9\Lambda_7^{-1}\mathcal{A}\Lambda_2^{-1})} + 1 - k_1.
\end{aligned} \tag{15}$$

Recall that, for any square matrices  $A, B$  with the same size,  $\rho(AB) = \rho(BA)$ . By this property,

$$\rho(\Lambda_{10}\Lambda_8^{-1}\mathcal{B}\Lambda_4\Lambda_9\Lambda_7^{-1}\mathcal{A}\Lambda_2^{-1}) = \rho(\mathcal{B}\Lambda_4\Lambda_9\Lambda_7^{-1}\mathcal{A}\Lambda_2^{-1}\Lambda_{10}\Lambda_8^{-1}). \tag{16}$$

By (14), (15) and (16),

$$\sqrt{R_0} \sqrt[4]{k_1 k_2} + 1 - k_2 \leq E_0 \leq \sqrt{R_0} \sqrt[4]{k_1 k_2} + 1 - k_1.$$

Hence,

$$\begin{aligned} \sqrt{R_0}(1 - \sqrt[4]{k_1 k_2}) - (1 - k_1) &\leq \sqrt{R_0} - E_0 \leq \sqrt{R_0}(1 - \sqrt[4]{k_1 k_2}) - (1 - k_2), \\ \frac{1}{\sqrt[4]{k_1 k_2}}(E_0(1 - \sqrt[4]{k_1 k_2}) - (1 - k_1)) &\leq \sqrt{R_0} - E_0 \leq \frac{1}{\sqrt[4]{k_1 k_2}}(E_0(1 - \sqrt[4]{k_1 k_2}) - (1 - k_2)). \end{aligned}$$

Similarly, we have the following theorem.

**Theorem 2.** *Under assumption (11),*

- (1) *If  $\sqrt{R_0} \leq \frac{1-k_2}{1-\sqrt[4]{k_1 k_2}} \leq 1$  or  $E_0 \leq \frac{1-k_2}{1-\sqrt[4]{k_1 k_2}} \leq 1$ , then  $\sqrt{R_0} \leq E_0$ ;*
- (2) *If  $\sqrt{R_0} \geq \frac{1-k_1}{1-\sqrt[4]{k_1 k_2}} \geq 1$  or  $E_0 \geq \frac{1-k_1}{1-\sqrt[4]{k_1 k_2}} \geq 1$ , then  $\sqrt{R_0} \geq E_0$ .*

**Corollary 2.** *If a further assumption is made that  $k_1 = k_2$  besides assumption (11), then  $\sqrt{R_0} \leq 1$  if and only if  $E_0 \leq 1$ . Furthermore,  $|\sqrt{R_0} - 1| \geq |E_0 - 1|$ .*

*Proof.* The proof is similar to that of Corollary 1. □

## 5 Numerical results

We show numerically the general relations between  $R_0$  and  $E_0$  for two models on heterogeneous networks. Significant parameters in predicting the uncertainty in  $E_0$  are found by Latin Hypercube Sampling/Partial Rank Correlation Coefficient (LHS/PRCC) analysis. Finally, the trends of parameters varying with extinction array is summarized.


### 5.1 Numerical results on relations between $R_0$ and $E_0$

Model (4) is applied to study thresholds for malaria transmission through numerical simulations. Five thousand realizations with parameters uniformly distributed within the ranges listed in Table 3 on a four-node network give rise to  $R_0$  ranging from 0.7668 to 63.8111 and  $E_0$  from 0.8965 to 1.9140. The ranges of  $R_0$  and  $E_0$  vary with the number of nodes of a network and the assumed ranges of vector (host) recruitment rates with ranges of other parameters fixed. The values of  $R_0$  are sorted from small values to large valued in Figure 1(a) and 1(b), and  $E_0$  are ranked from small values to large values in Figure 1(c) and 1(d). The largest value of  $E_0$  is 0.9980 when all values of  $R_0$  are smaller than 1 and  $R_0 \leq E_0$ , as shown in Figure 1(a). The smallest value of  $E_0$  is 1.003 when all values of  $R_0$  are greater than 1 and  $R_0 \geq E_0$ , as shown in Figure 1(b). The largest value of  $R_0$  is 0.9947 when all values of  $E_0$  are smaller than 1, as shown in Figure 1(c). The smallest value of  $R_0$  is 1.006 when all values of  $E_0$  are greater than 1, as shown in Figure 1(d). The value of  $E_0$  is not monotonically increasing with the increase of  $R_0$ , as shown Figure 1(a) and 1(b). Similarly,  $R_0$  fluctuates as  $E_0$  increases, as shown in Figure 1(c) and 1(d).


Model (13) is applied to numerical examine the relationship between  $R_0$  and  $E_0$  for Rift Valley fever. See Table 4 for descriptions and ranges of parameters. Five thousand realizations produce  $R_0$  ranging between 0.2289 and 54.5086 and  $E_0$  from 0.6757 to 1.9763. The values of  $R_0$  are ordered from small to large magnitudes in Figure 2(a) and 2(b), and the values of  $E_0$  are from small to large values in Figure 2(c) and 2(d). The largest value of  $E_0$  is 1 when all values of  $R_0$  are smaller than 1, and  $\sqrt{R_0} \leq E_0$ , as shown in Figure 2(a). The smallest value of  $E_0$  is 1.005 when all values of  $R_0$  are greater than 1, and  $\sqrt{R_0} \geq E_0$ , as shown in Figure 2(b). The

| Parameter | Description                          | Range              | Dimension | Source |
|-----------|--------------------------------------|--------------------|-----------|--------|
| $\alpha$  | Contact rate: mosquitoes to humans   | 0.010 – 0.27       | 1/day     | [8]    |
| $\beta$   | Contact rate: humans to mosquitoes   | 0.072 – 0.64       | 1/day     | [8]    |
| $\mu$     | Per capita death rate for mosquitoes | 0.020 – 0.27       | 1/day     | [8]    |
| $d$       | Per capita death rate for humans     | 0.000027 – 0.00014 | 1/day     | [8]    |
| $\gamma$  | Per capita recovery rate for humans  | 0.0014 – 0.0017    | 1/day     | [8]    |
| $\eta$    | Mosquito recruitment rate            | 1 – 5              | 1/day     | Assume |
| $\psi$    | Human recruitment rate               | 1 – 60             | 1/day     | Assume |


Table 3: Parameters of the malaria metapopulation model.




(a) When  $R_0 \leq 1$ ,  $R_0 \leq E_0$ .



(b) When  $R_0 \geq 1$ ,  $R_0 \geq E_0$ .



(c) When  $E_0 \leq 1$ ,  $R_0 \leq E_0$ .



(d) When  $E_0 \geq 1$ ,  $R_0 \geq E_0$ .

Figure 1: Relationships between  $R_0$  and  $E_0$  for malaria model.

| Parameter    | Description                           | Range           | Dimension | Source                      |
|--------------|---------------------------------------|-----------------|-----------|-----------------------------|
| $\alpha$     | Contact rate: mosquito to livestock   | 0.0021 – 0.2762 | 1/day     | [7, 17, 19, 24, 30, 36, 37] |
| $\beta$      | Contact rate: livestock to mosquitoes | 0 – 0.32        | 1/day     | [7, 17, 19, 24, 30, 35]     |
| $1/\mu$      | Longevity of mosquitoes               | 3 – 60          | 1/day     | [4, 26, 30]                 |
| $1/d$        | Longevity of livestock                | 360 – 3600      | 1/day     | [31]                        |
| $1/\gamma$   | Recover rate in livestock             | 2 – 5           | 1/day     | [13]                        |
| $1/\varphi$  | Incubation period in mosquitoes       | 4 – 8           | days      | [36]                        |
| $1/\epsilon$ | Incubation period in livestock        | 2 – 6           | days      | [29]                        |
| $\eta$       | Mosquito recruitment rate             | 1 – 500         | 1/day     | Assume                      |
| $\psi$       | Livestock recruitment rate            | 1 – 10          | 1/day     | Assume                      |

Table 4: Parameters of the Rift Valley fever metapopulation model.


| Parameter   | PRCC    | p-value   |
|-------------|---------|-----------|
| $\alpha$    | 0.5649  | $< 0.001$ |
| $\beta$     | 0.6039  | $< 0.001$ |
| $1/\mu$     | 0.8061  | $< 0.001$ |
| $1/d$       | −0.4660 | $< 0.001$ |
| $1/\gamma$  | 0.5524  | $< 0.001$ |
| $1/\varphi$ | −0.0284 | $< 0.05$  |
| $\eta$      | 0.5785  | $< 0.001$ |
| $\psi$      | −0.5036 | $< 0.001$ |

Table 5: Sensitivity testing results using Partial Rank Correlation Coefficients for model (13) for homogeneous populations. Only significant parameters are shown.


largest value of  $R_0$  is 0.9998 when all values of  $E_0$  are smaller than 1, and  $\sqrt{R_0} \geq E_0$ , as shown in Figure 2(c). The smallest value of  $R_0$  is 1.008 when all values of  $E_0$  are greater than 1, and  $\sqrt{R_0} \geq E_0$ , as shown in Figure 2(d). When  $R_0$  increases,  $E_0$  does not always increase, as shown in Figure 2(a) and 2(b). Similarly,  $R_0$  fluctuates as  $E_0$  increases, as shown in Figure 2(c) and 2(d).

## 5.2 Sensitivity analysis


We employ Latin Hypercube Sampling/Partial Rank Correlation Coefficient (LHS/PRCC) analysis [25] to identify key parameters whose uncertainty contribute to predict uncertainty of  $E_0$  for model (13) and rank the parameters by their significances. The parameters shown to be significant with large PRCC values ( $> 0.5$ ) or small  $p$ -values ( $< 0.05$ ) [15] by the sensitivity test with 5000 sets of parameter values, are listed in Table 5. The magnitude of PRCC value represent the contribution to the prediction for the imprecision of  $E_0$ , and a negative sign indicates that the parameter is inversely proportional to the magnitude of  $E_0$ . The closer PRCC value is to +1 or −1, the more greatly the parameter impacts the outcome of  $E_0$ .




(a) When  $\sqrt{R_0} \leq 1$ ,  $\sqrt{R_0} \leq E_0$ .



(b) When  $\sqrt{R_0} \geq 1$ ,  $\sqrt{R_0} \geq E_0$ .



(c) When  $E_0 \leq 1$ ,  $\sqrt{R_0} \leq E_0$ .



(d) When  $E_0 \geq 1$ ,  $\sqrt{R_0} \geq E_0$ .

Figure 2: Relationships between  $R_0$  and  $E_0$  for Rift Valley fever model.

### 5.3 Trends of extinction array with varying parameters

Consistent trends of  $w^*$  are observed by numerical simulations for homogeneous populations and a two-node network for Model (13). Table 6 lists three different values for each parameter and corresponding extinction array for homogeneous populations as an example. Table 7 shows the trends of extinction array by varying one parameter at a time, keeping other parameters fixed and  $E_0 > 1$  for homogeneous populations and a two-node network. If at least one entry of extinction array increases and others remain constant, then we say that the array increases. The extinction array  $w^*$  decreases with the increase of contact rates from local vectors and vectors in other nodes to local hosts, contact rates from local hosts and hosts in other nodes to local vectors, death rates of hosts, recruitment rates of vectors, and incubation rates of vectors and hosts, whereas increases with the increase of vector death rates, vector recovery rates, host recruitment rates.

| Changing parameter | $(w_1^*, w_2^*, u_1^*, u_2^*)$   |
|--------------------|----------------------------------|
| $\alpha = 0.0601$  | (0.9965, 0.9978, 0.9961, 0.9978) |
| $\alpha = 0.0766$  | (0.8648, 0.9212, 0.8467, 0.9212) |
| $\alpha = 0.0781$  | (0.8546, 0.9158, 0.8352, 0.9158) |
| $\beta = 0.0639$   | (0.9158, 0.9824, 0.9046, 0.9824) |
| $\beta = 0.1026$   | (0.6623, 0.8967, 0.6173, 0.8966) |
| $\beta = 0.1426$   | (0.5448, 0.8224, 0.4841, 0.8223) |
| $\mu = 1/60$       | (0.1955, 0.4961, 0.1419, 0.4956) |
| $\mu = 1/59$       | (0.1996, 0.5016, 0.1453, 0.5110) |
| $\mu = 1/56$       | (0.2127, 0.5188, 0.1565, 0.5182) |
| $d = 1/3477$       | (0.4621, 0.7398, 0.3904, 0.7395) |
| $d = 1/3370$       | (0.4554, 0.7312, 0.3828, 0.7310) |
| $d = 1/3311$       | (0.4518, 0.7265, 0.3787, 0.7262) |
| $\gamma = 1/5$     | (0.4247, 0.6877, 0.3480, 0.6874) |
| $\gamma = 1/4$     | (0.4698, 0.7491, 0.3992, 0.7488) |
| $\gamma = 1/3$     | (0.5451, 0.8226, 0.4845, 0.8224) |
| $\epsilon = 1/6$   | (0.4700, 0.7493, 0.3994, 0.7489) |
| $\epsilon = 1/4$   | (0.4698, 0.7491, 0.3992, 0.7488) |
| $\epsilon = 1/2$   | (0.4697, 0.7489, 0.3990, 0.7488) |
| $\varphi = 1/8$    | (0.5494, 0.7784, 0.4293, 0.7782) |
| $\varphi = 1/7$    | (0.5312, 0.7715, 0.4218, 0.7712) |
| $\varphi = 1/6$    | (0.5119, 0.7643, 0.4142, 0.7641) |
| $\eta = 19$        | (0.5412, 0.8195, 0.4801, 0.8193) |
| $\eta = 76$        | (0.5264, 0.8069, 0.4632, 0.8066) |
| $\eta = 482$       | (0.2907, 0.3169, 0.1961, 0.3162) |
| $\psi = 1$         | (0.4698, 0.7491, 0.3992, 0.7488) |
| $\psi = 2$         | (0.6859, 0.9123, 0.6553, 0.9122) |
| $\psi = 3$         | (0.9219, 0.9838, 0.9115, 0.9838) |

Table 6: The extinction array changes with one parameter within the range at a time for homogeneous populations, while keeping other parameters fixed and  $E_0 > 1$  for model (13). Fixed parameters are:  $\alpha = 0.2$ ,  $\beta = 0.19$ ,  $\mu = 1/30$ ,  $d = 1/3600$ ,  $\gamma = 1/4$ ,  $\epsilon = 1/2$ ,  $\varphi = 1/4$ ,  $\eta = 100$ ,  $\psi = 1$  in this example. Same trends are obtained with various sets of fixed parameters.

| Increasing parameter                                                                                               | $(w_1^*, \dots, w_n^*, u_1^*, \dots, u_n^*)$ |
|--------------------------------------------------------------------------------------------------------------------|----------------------------------------------|
| $\alpha_i, \beta_i, d_i, \epsilon_i, \varphi_i, \eta_i, \sigma_{ij}, \omega_{ij} \ (i, j = 1, \dots, n, i \neq j)$ | decreases                                    |
| $\mu_i, \gamma_i, \psi_i \ (i = 1, \dots, n)$                                                                      | increases                                    |

Table 7: Summary of trends for extinction array changing with one parameter at a time, while keeping other parameters fixed and  $E_0 > 1$  for model (13) for homogeneous populations and a two-node network throughout various simulations.

## 6 Discussions

The reproduction number,  $R_0$  for deterministic vector-host models and thresholds for extinction probabilities,  $E_0$  for corresponding CTMC models are analytically and numerically connected. For model (4), our analysis show that  $R_0 \leq 1$ , if and only if  $E_0 \leq 1$ , and  $|R_0 - 1| \geq |E_0 - 1|$  under certain assumptions. Numerical simulations for a malaria model on heterogeneous networks with different number of nodes show that Corollary 1 holds without any assumptions. For model (13), analytical results show that  $\sqrt{R_0} < 1$  if and only if  $E_0 < 1$ , and  $|\sqrt{R_0} - 1| \geq |E_0 - 1|$  by the same assumption in (11). Extensive numerical simulation results for a Rift Valley fever model on networks with various number of nodes show that Corollary 2 holds without any assumptions.

**Conjecture 1.** *Theorems 1, 2 and Corollary 1, 2 hold without assumption (11), i.e.,  $R_0 \leq 1$  if and only if  $E_0 \leq 1$  for both models (4) and (13), besides,  $|R_0 - 1| \geq |E_0 - 1|$  for model (4), and  $|\sqrt{R_0} - 1| \geq |E_0 - 1|$  for model (13) without assumption (11).*

The first part was proven by Allen and van den Driessche under the assumption (16) in [2], i.e.,  $(F - V)^T = W(M - I)$ , where  $F$  and  $V$  are Jacobian matrices defined in (1),  $M$  is a mean matrix of offspring distribution defined in Section 2.2,  $I$  is the identity matrix, and  $W$  is a positive diagonal matrix with each entry  $w_i$  representing the rate parameter at which lifespan of group  $i$  are exponentially distributed for  $i = 1, \dots, n$  [28]. This assumption does not hold for both model (4) and model (13).

Consistent trends in the extinction array  $w^*$  while changing one parameter through numerical simulations is helpful in deducing trends of extinction probability and possible interventions for vector-borne diseases. According to Equation (3), the probability of disease extinction is monotonically increasing (decreasing) with the increase (decrease) of the extinction array when the initial number of infection is fixed. The following biological interpretations on disease extinction or persistence are in terms of fixed initial number of infections. If contact rates from vectors to hosts ( $\alpha, \sigma$ ), or those from hosts to vectors ( $\beta, \omega$ ) increase, the probability for the disease to persist is higher. If death rates of hosts ( $d$ ) increase, the number of vectors is relatively dominant. Consequently, the disease is more likely to persist. Similarly, growing vector recruitment rates ( $\eta$ ) increase the probability for disease persistence. The higher incubation rates in vectors ( $\varphi$ ) or that in hosts ( $\epsilon$ ) lead to faster vector or host infections, such that the disease is prone to persist. On the contrary, increasing death rates of vectors ( $\mu$ ) may reduce rates of host infection, and ultimately, may increase the likelihood of disease extinction. Increasing recovery rates of hosts ( $\gamma$ ) may reduce the number of infections, such that the probability of disease extinction increases. Increasing recruitment rate of hosts ( $\psi$ ) may reduce vector infection rates and increase probability of disease extinction.

The findings show that we may increase extinction probability of vector-borne diseases by properly controlling vector and host population size, and promptly detecting and applying treatment for hosts. Analytical and numerical results shed light on deriving relationships between  $R_0$  and  $E_0$ , as well as connections between varying parameters and increasing extinction probabilities for many other vector-borne diseases transmitted among heterogeneous works. In summary, the resulting mathematical derivations and numerical simulations facilitate understanding thresholds for the spread of vector-borne diseases, as well as providing novel insights into disease prevention, mitigation and control strategies.

## Acknowledgments

This material is based upon work supported by the U. S. Department of Homeland Security under Grant Award Number 2010-ST061-AG0001, a grant through the Kansas Biosciences Authority, and is funded by U. S. Department of Agriculture - the Agricultural Research Service with Specific Cooperative Agreement 58-5430-1-0356. The views and conclusions contained in this publication are those of the authors and should not be interpreted as necessarily representing the official policies, either explicit or implicit, of the U. S. Department of Homeland Security and Kansas Biosciences Authority. We are grateful to Dr. Lee Cohnstaedt and Dr. H. Morgan Scott, and Dr. Linda J. S. Allen for helpful conversations.

## References

- [1] L. J. Allen and G.E. Lahodny Jr. Extinction thresholds in deterministic and stochastic epidemic models. *J Biol Dyn*, 6(2):590–611, 2012.
- [2] L. J. Allen and P. van den Driessche. Relations between deterministic and stochastic thresholds for disease extinction in continuous- and discrete-time infectious disease models. *Math Biosci*, 243:99–108, 2013.
- [3] J. Arino and P. van den Driessche. The basic reproduction number in a multi-city compartmental epidemic model. *Posit Sys, Proc*, 294:135–142, 2003.
- [4] M. Bates. The natural history of mosquitoes. *Am J Public Health*, 39(12):1592, 1949.
- [5] D. Bisanzio, L. Bertolotti, L. Tomassone, G. Amore, C. Ragagli, A. Mannelli, M. Giacobini, and P. Provero. Modeling the spread of vector-borne diseases on bipartite networks. *PloS One*, 5(11):e13796, 2010.
- [6] T. Britton and D. Lindenstrand. Epidemic modelling: Aspects where stochasticity matters. *Math Biosci*, 222:109–116, 2009.
- [7] D. V. Canyon, J. L. K. Hii, and R. Muller. The frequency of host biting and its effect on oviposition and survival in *Aedes aegypti* (Diptera: Culicidae). *Bull Entomol Res*, 89(1):35–39, 1999.
- [8] Cushing J.M. Chitnis N., Hyman J.M. Determining important parameters in the spread of malaria through the sensitivity analysis of a mathematical model. *Bull of Math Biol*, 70(5):1272–1296, 2008.
- [9] CIESIN. Changes in the incidence of vector-borne diseases attributable to climate change. <http://www.ciesin.columbia.edu/TG/HH/veclev2.html>, 2007. (Center for International Earth Science Information Network). Accessed June 20, 2013.
- [10] A. Cintrn-Arias, C. Castillo-Chvezl, L. M. Bettencourt, and H. T. Banks A. L. Lloyd. The estimation of the effective reproductive number from disease outbreak data. *Math Biosci Eng*, 255:261–282, 2009.
- [11] O. Diekmann and J. A. P. Heesterbeek. *Mathematical Epidemiology of Infectious Diseases*. Wiley, Chichester, 2000.

- [12] O. Diekmann, J.A.P. Heesterbeek, and J.A.J. Metz. On the definition and the computation of the basic reproduction ratio  $R_0$  in models for infectious diseases in heterogeneous populations. *J Math Biol*, 28(4):365–382, 1990.
- [13] B. J. Erasmus and J. A. W. Coetzer. The symptomatology and pathology of Rift Valley fever in domestic animals. *Contrib. Epidemiol. Biostat.*, 3:77–82, 1981.
- [14] B. F. Finkenstadt, O. N. Bjornstad, and B. T. Grenfell. A stochastic model for extinction and recurrence of epidemics: estimation and inference for measles outbreaks. *Biostat*, 3(4):493–510, 2002.
- [15] B. Gomero. Latin Hypercube Sampling and Partial Rank Correlation Coefficient analysis applied to an optimal control problem. Master’s thesis, the University of Tennessee, the United States, 2012.
- [16] N. G. Gratz. Emerging and resurging vector-borne diseases. *Annu Rev Entomol*, 44:51–75, 1999.
- [17] R. O. Hayes, C. H. Tempelis, A. D. Hess, and W. C. Reeves. Mosquito host preference studies in Hale County, Texas. *Am J Trop Med Hyg*, 22(2):270–277, 1973.
- [18] G. R. Hosack, P. A. Rossignol, and P. Van den Driessche. The control of vector-borne disease epidemics. *J Theor Biol*, 255:16–25, 2008.
- [19] C. J. Jones and J. E. Lloyd. Mosquitos feeding on sheep in southeastern Wyoming. *J Am Mosq Control Assoc*, 1(4):530–532, 1985.
- [20] R.R. Kao. Networks and models with heterogeneous population structure in epidemiology. *Netw Sci*, pages 51–84, 2010.
- [21] M.J. Keeling and P. Rohani. *Modeling infectious diseases in humans and animals*. Princeton University Press, 2008.
- [22] G. E. Lahodny Jr and L. J. Allen. Probability of a disease outbreak in stochastic multipatch epidemic models. *Bull Math Biol*, 75(7):1157–1180, 2013.
- [23] A. L. Lloyd, J. Zhang, and A. M. Root. Stochasticity and heterogeneity in host-vector models. *J R Soc Interface*, 4(16):851–863, 2007.
- [24] L. A. Magnarelli. Host feeding patterns of Connecticut mosquitos (Diptera: Culicidae). *Am J Trop Med Hyg*, 26(3):547–552, 1977.
- [25] S. Marino, I. B. Hogueand, C. J. Ray, and D. E. Kirschner. A methodology for performing global uncertainty and sensitivity analysis in systems biology. *J Theor Biol*, 254:178–196, 2008.
- [26] C. G. Moore, R. G. McLean, C. J. Mitchell, R. S. Nasci, T. F. Tsai, C. H. Caslisher, A. A. Marfin, P. S. Moorsee, and D. J. Gubler. *Guidelines for Arbovirus Surveillance Programs in the United States*. Centers for Disease Control and Prevention, 1993.
- [27] F. Natale, A. Giovannini, L. Savini, D. Palma, L. Possenti, G. Fiore, and P. Calistri. Network analysis of italian cattle trade patterns and evaluation of risks for potential disease spread. *Prev Vet Med*, 92(4):341–350, 2009.

- [28] S. Pénisson. *Conditional limit theorems for multitype branching processes and illustration in epidemiological risk analysis*. PhD thesis, Institut für Mathematik der Universität Potsdam, Germany, 2010.
- [29] C. J. Peters and K. J. Linthicum. Rift Valley fever. In G.B. Beran, editor, *Handbook of Zoonoses*, Section B: Viral, pages 125–138. CRC Press, Inc., Boca Raton, Fl, second edition, 1994.
- [30] H. D. Pratt and C. G. Moore. *Vector-borne disease control: mosquitoes of public health importance and their control*. U.S. Department of Health and Human Services, Atlanta, GA, 1993.
- [31] O. M. Radostits. *Herd healthy: food animal production medicine*. Saunders, 2001.
- [32] M. G. Roberts and J. A. P. Heesterbeek. A new method for estimating the effort required to control an infectious disease. *Proc Biol Sci*, 270(1522):1359–1364, 2003.
- [33] R. Ross. Some quantitative studies in epidemiology. *Nature*, 4:466–467, 1911.
- [34] Z. Shuai, J. A. P. Heesterbeek, and P. van den Driessche. Extending the type reproduction number to infectious disease control targeting contacts between types. *J Mathl Biol*, doi: 10.1007/s00285-012-0579-9, 2012.
- [35] M. J. Turell and C. L. Bailey. Transmission studies in mosquitoes (Diptera: Culicidae) with disseminated Rift Valley fever virus infections. *J Med Entomol*, 24(1):11–18, 1987.
- [36] M. J. Turell, C. L. Bailey, and J. R. Beaman. Vector competence of a Houston, Texas strain of *Aedes Albopictus* for Rift Valley fever virus. *J Am Mosq Control Assoc*, 4(1):94–96, 1988.
- [37] M. J. Turell, M. E. Faran, M. Cornet, and C. L. Bailey. Vector competence of senegalese *Aedes fowleri* (Diptera: Culicidae) for Rift Valley fever virus. *J Med Entomol*, 25(4):262–266, 1988.
- [38] P. van den Driessche and J. Watmough. Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Math Biosci*, 180:29–48, 2002.
- [39] M. C. Vernon and M. J. Keeling. Representing the uk’s cattle herd as static and dynamic networks. *Proc Biol Sci*, 276(1656):469–476, 2009.
- [40] L. Xue and C. Scoglio. The network level reproduction number for infectious diseases with both vertical and horizontal transmission. *Math Biosci*, 243(1):67–80, 2013.