

On the relative importance of second-order terms in relativistic dissipative fluid dynamics

E. Molnár,^{1,2} H. Niemi,^{3,4} G.S. Denicol,⁵ and D.H. Rischke⁶

¹MTA-DE Particle Physics Research Group, H-4010 Debrecen, P.O. Box 105, Hungary

²Frankfurt Institute for Advanced Studies, Ruth-Moufang-Str. 1, D-60438 Frankfurt am Main, Germany

³Department of Physics, P.O. Box 35 (YFL), FI-40014 University of Jyväskylä, Finland

⁴Helsinki Institute of Physics, P.O.Box 64, FI-00014 University of Helsinki, Finland

⁵Department of Physics, McGill University, 3600 rue University, Montréal, Canada

⁶Institut für Theoretische Physik, Johann Wolfgang Goethe-Universität,
Max-von-Laue-Str. 1, D-60438 Frankfurt am Main, Germany

In Denicol *et al.*, Phys. Rev. D **85**, 114047 (2012), the equations of motion of relativistic dissipative fluid dynamics were derived from the relativistic Boltzmann equation. These equations contain a multitude of terms of second order in Knudsen number, in inverse Reynolds number, or their product. Terms of second order in Knudsen number give rise to non-hyperbolic (and thus acausal) behavior and must be neglected in (numerical) solutions of relativistic dissipative fluid dynamics. The coefficients of the terms which are of the order of the product of Knudsen and inverse Reynolds numbers have been explicitly computed in the above reference, in the limit of a massless Boltzmann gas. Terms of second order in inverse Reynolds number arise from the collision term in the Boltzmann equation, upon expansion to second order in deviations from the single-particle distribution function in local thermodynamical equilibrium. In this work, we compute these second-order terms for a massless Boltzmann gas with constant scattering cross section. Consequently, we assess their relative importance in comparison to the terms which are of the order of the product of Knudsen and inverse Reynolds numbers.

PACS numbers: 12.38.Mh, 24.10.Nz, 47.75.+f, 51.10+y

I. INTRODUCTION AND CONCLUSIONS

Relativistic fluid dynamics has found widespread applications in heavy-ion physics, in modelling nuclear collisions at ultrarelativistic bombarding energies [1, 2], in astrophysics, for instance in modelling binary mergers of compact stellar objects, see e.g. Ref. [3], as well as in cosmology [4–8]. In the past, in order to solve the equations of motion of relativistic fluid dynamics, one has often made the assumption that the fluid is ideal, i.e., one demands instantaneous local thermodynamical equilibrium, which in turn allows to neglect all dissipative effects. However, there are no ideal fluids in nature, as can be seen for instance from the fact that the shear viscosity coefficient may attain a lower limit, but never vanishes [9–11].

A more realistic modelling of the dynamics of relativistic fluids thus demands that one uses the equations of relativistic *dissipative* fluid dynamics. First attempts to formulate such equations were made by Eckart [12] and Landau and Lifshitz [13] based on a relativistic generalization of the non-relativistic Navier-Stokes equations. However, their equations suffer from instabilities and acausal signal propagation [14]. The reason for this is the (erroneous) assumption that the dissipative quantities, like bulk viscous pressure Π , particle diffusion current n^μ , and shear-stress tensor $\pi^{\mu\nu}$, react *instantaneously* to the thermodynamic forces, like gradients of the fluid velocity or temperature and chemical potential. If one relaxes this assumption by introducing certain time scales τ_Π , τ_n , and τ_π , on which the dissipative quantities are allowed to approach the values determined by the corresponding thermodynamic forces, these problems can be cured [provided the relaxation times fulfill certain conditions [15]]. Similarly to earlier works by Grad [16] and Müller [17–19] in the non-relativistic context, Israel and Stewart (IS) were among the first to suggest equations of motion for relativistic dissipative fluids that were stable and causal [20–22].

In recent years, relativistic dissipative fluid dynamics based on the IS formulation was extensively applied to describe the dynamics of nuclear collisions. At the same time, the theoretical foundations of this theory were further explored both from kinetic theory [23–40] and from irreversible thermodynamics [41–51]. In particular, in Refs. [23–25] it has been investigated how to derive the equations of motion for the dissipative quantities using the Boltzmann equation as underlying microscopic theory.

In Ref. [24] a derivation of the equations of motion of relativistic dissipative fluid dynamics was presented, which is based on a systematic power-counting scheme in Knudsen and inverse Reynolds number. The Knudsen number, $\text{Kn}=\lambda/L$, is the ratio between a characteristic microscopic time/length scale, λ , e.g., the mean-free path between collisions, and a characteristic macroscopic scale of the fluid L . In this context, the inverse Reynolds numbers are the ratios of dissipative quantities and (local) equilibrium values of macroscopic fields, e.g. like $R_\Pi^{-1} = |\Pi|/P_0$, $R_n^{-1} = |n^\mu|/n_0$, or $R_\pi^{-1} = |\pi^{\mu\nu}|/P_0$, where P_0 is the thermodynamic pressure and n_0 is the particle density in

equilibrium. The time scales τ_Π , τ_n , and τ_π are identified with the slowest *microscopic* time scales of the Boltzmann equation.

The physical picture that emerges is that microscopic processes (i.e., in the case of the Boltzmann equation, binary collisions) occur on time scales smaller than (or at most as large as) τ_Π , τ_n , and τ_π . These processes affect that the dissipative quantities Π , n^μ , and $\pi^{\mu\nu}$ approach the values given by the (relativistic generalization of the) Navier-Stokes equations on the time scales τ_Π , τ_n , and τ_π . Since microscopic physics influences the motion of the fluid only on short time scales, the term “transient fluid dynamics” was coined for such theories of relativistic dissipative fluid dynamics. The fact that the microscopic dynamics of the Boltzmann equation gives rise to relaxation-type equations of motion for the dissipative quantities, i.e., where these quantities exponentially decay towards the values given by the Navier-Stokes equations, was confirmed in Ref. [52]. It was also shown in that paper that approaches based on the AdS/CFT correspondence lead to equations of motion which are of the type encountered for an underdamped harmonic oscillator. The relaxation towards the Navier-Stokes values is then not exponential but oscillatory.

Let us recall the relaxation-type equations for the dissipative quantities derived in Ref. [24],

$$\tau_\Pi \dot{\Pi} + \Pi = -\zeta\theta + \mathcal{J} + \mathcal{K} + \mathcal{R}, \quad (1)$$

$$\tau_n \dot{n}^{(\mu)} + n^\mu = \kappa I^\mu + \mathcal{J}^\mu + \mathcal{K}^\mu + \mathcal{R}^\mu, \quad (2)$$

$$\tau_\pi \dot{\pi}^{(\mu\nu)} + \pi^{\mu\nu} = 2\eta\sigma^{\mu\nu} + \mathcal{J}^{\mu\nu} + \mathcal{K}^{\mu\nu} + \mathcal{R}^{\mu\nu}, \quad (3)$$

where the over-dot denotes the proper time derivative, $\dot{A} \equiv DA = u^\mu \partial_\mu A$. Here, ζ is the coefficient of the bulk viscosity, κ the coefficient of particle diffusion (which is related to that of heat conduction) and η the coefficient of shear viscosity. Furthermore, with the fluid 4-velocity u^μ (chosen in the Landau frame), where $u^\mu u_\mu = 1$, with $\Delta^{\mu\nu} = g^{\mu\nu} - u^\mu u^\nu$ being the 3-projector onto the subspace orthogonal to u^μ , and with $\nabla^\mu = \Delta^{\mu\nu} \partial_\nu$ being the 3-gradient, $\theta = \nabla_\mu u^\mu$ is the expansion scalar, $\sigma^{\mu\nu} \equiv \nabla^{(\mu} u^{\nu)} = \frac{1}{2}(\nabla^\mu u^\nu + \nabla^\nu u^\mu) - \frac{1}{3}\theta\Delta^{\mu\nu}$ is the shear tensor, while $I^\mu = \nabla^\mu \alpha_0$ is the gradient of $\alpha_0 = \mu/T$, the ratio of chemical potential μ and temperature T .

In the above equations, the tensors \mathcal{J} , \mathcal{J}^μ , and $\mathcal{J}^{\mu\nu}$ contain all terms of first order in the product of Knudsen and inverse Reynolds number,

$$\begin{aligned} \mathcal{J} &= -\ell_{\Pi n} \nabla \cdot n - \tau_{\Pi n} n \cdot F - \delta_{\Pi\Pi} \Pi\theta - \lambda_{\Pi n} n \cdot I + \lambda_{\Pi\pi} \pi^{\mu\nu} \sigma_{\mu\nu}, \\ \mathcal{J}^\mu &= -\tau_n n_\nu \omega^{\nu\mu} - \delta_{nn} n^\mu \theta - \ell_{n\Pi} \nabla^\mu \Pi + \ell_{n\pi} \Delta^{\mu\nu} \nabla_\lambda \pi_\nu^\lambda + \tau_{n\Pi} \Pi F^\mu - \tau_{n\pi} \pi^{\mu\nu} F_\nu - \lambda_{nn} n_\nu \sigma^{\mu\nu} + \lambda_{n\Pi} \Pi I^\mu - \lambda_{n\pi} \pi^{\mu\nu} I_\nu, \\ \mathcal{J}^{\mu\nu} &= 2\tau_\pi \pi_\lambda^{(\mu} \omega^{\nu)\lambda} - \delta_{\pi\pi} \pi^{\mu\nu} \theta - \tau_{\pi\pi} \pi^{\lambda(\mu} \sigma_\lambda^{\nu)} + \lambda_{\pi\Pi} \Pi \sigma^{\mu\nu} - \tau_{\pi n} n^{(\mu} F^{\nu)} + \ell_{\pi n} \nabla^{(\mu} n^{\nu)} + \lambda_{\pi\pi} n^{(\mu} I^{\nu)}, \end{aligned} \quad (4)$$

where $\omega^{\mu\nu} = \frac{1}{2}(\nabla^\mu u^\nu - \nabla^\nu u^\mu)$ denotes the vorticity and we defined $F^\mu = \nabla^\mu P_0$ as the gradient of the thermodynamic pressure. The tensors \mathcal{K} , \mathcal{K}^μ , and $\mathcal{K}^{\mu\nu}$ contain all terms of second order in Knudsen number,

$$\begin{aligned} \mathcal{K} &= \tilde{\zeta}_1 \omega_{\mu\nu} \omega^{\mu\nu} + \tilde{\zeta}_2 \sigma_{\mu\nu} \sigma^{\mu\nu} + \tilde{\zeta}_3 \theta^2 + \tilde{\zeta}_4 I \cdot I + \tilde{\zeta}_5 F \cdot F + \tilde{\zeta}_6 I \cdot F + \tilde{\zeta}_7 \nabla \cdot I + \tilde{\zeta}_8 \nabla \cdot F, \\ \mathcal{K}^\mu &= \tilde{\kappa}_1 \sigma^{\mu\nu} I_\nu + \tilde{\kappa}_2 \sigma^{\mu\nu} F_\nu + \tilde{\kappa}_3 I^\mu \theta + \tilde{\kappa}_4 F^\mu \theta + \tilde{\kappa}_5 \omega^{\mu\nu} I_\nu + \tilde{\kappa}_6 \Delta_\lambda^\mu \nabla_\nu \sigma^{\lambda\nu} + \tilde{\kappa}_7 \nabla^\mu \theta, \\ \mathcal{K}^{\mu\nu} &= \tilde{\eta}_1 \omega_\lambda^{(\mu} \omega^{\nu)\lambda} + \tilde{\eta}_2 \theta \sigma^{\mu\nu} + \tilde{\eta}_3 \sigma^{\lambda(\mu} \sigma_\lambda^{\nu)} + \tilde{\eta}_4 \sigma_\lambda^{(\mu} \omega^{\nu)\lambda} + \tilde{\eta}_5 I^{(\mu} I^{\nu)} + \tilde{\eta}_6 F^{(\mu} F^{\nu)} \\ &\quad + \tilde{\eta}_7 I^{(\mu} F^{\nu)} + \tilde{\eta}_8 \nabla^{(\mu} I^{\nu)} + \tilde{\eta}_9 \nabla^{(\mu} F^{\nu)}. \end{aligned} \quad (5)$$

Note that, in contrast to Ref. [24], we now write the term proportional to $\tilde{\kappa}_6$ with a 3-gradient operator ∇_ν instead of a partial derivative ∂_ν . Finally, the tensors \mathcal{R} , \mathcal{R}^μ , and $\mathcal{R}^{\mu\nu}$ contain all terms of second order in inverse Reynolds number,

$$\mathcal{R} = \varphi_1 \Pi^2 + \varphi_2 n \cdot n + \varphi_3 \pi_{\mu\nu} \pi^{\mu\nu}, \quad (6)$$

$$\mathcal{R}^\mu = \varphi_4 n_\nu \pi^{\mu\nu} + \varphi_5 \Pi n^\mu, \quad (7)$$

$$\mathcal{R}^{\mu\nu} = \varphi_6 \Pi \pi^{\mu\nu} + \varphi_7 \pi^{\lambda(\mu} \pi_\lambda^{\nu)} + \varphi_8 n^{(\mu} n^{\nu)}. \quad (8)$$

These second-order terms follow from computing the collision integral beyond linear order in the dissipative quantities.

Observing the plethora of transport coefficients occurring in Eqs. (4) – (8), a natural question to ask is whether all of them are of the same order of magnitude or whether some coefficients are larger and thus more important than others. The goal of this paper is to answer this question. The coefficients in Eqs. (4) were explicitly computed in Ref. [24] for a massless Boltzmann gas with constant scattering cross section. We now supplement these results by computing the coefficients in Eqs. (5) – (8). We restrict ourselves to the 14-moment approximation. In this case, the coefficients in Eqs. (5) vanish identically (cf. Appendix I), and we only need to focus on the coefficients $\varphi_1, \dots, \varphi_8$ in Eqs. (6) – (8). The derivation and calculation of these coefficients is quite demanding and is presented in detail

in the remainder of this paper. For the rest of this introductory section, we simply quote the results and draw our conclusions.

For massless particles, the bulk viscous pressure vanishes identically, $\Pi = 0$, and we do not have to solve Eq. (1). Also, terms in Eqs. (4) – (8) proportional to Π can be neglected, such that we do not need to compute the corresponding coefficients. Moreover, as mentioned above, in the 14-moment approximation all coefficients in Eqs. (5) vanish. Dividing Eq. (2) by n_0 , and Eq. (3) by P_0 , these two equations can be written in the following form,

$$\begin{aligned} \tau_n \frac{\dot{n}^{(\mu)}}{n_0} + \frac{n^\mu}{n_0} &= \frac{\kappa}{n_0} \nabla^\mu \alpha_0 \\ &+ (\tau_n \omega^{\mu\nu} - \lambda_{nn} \sigma^{\mu\nu} - \delta_{nn} \theta g^{\mu\nu}) \frac{n_\nu}{n_0} \\ &+ \frac{\ell_{n\pi}}{\beta_0} \frac{\Delta^{\mu\nu} \nabla_\lambda \pi_\nu^\lambda}{P_0} - \left(\frac{\tau_{n\pi} P_0}{\beta_0} \frac{\nabla_\nu P_0}{P_0} + \frac{\lambda_{n\pi}}{\beta_0} \nabla_\nu \alpha_0 - \varphi_4 P_0 \frac{n_\nu}{n_0} \right) \frac{\pi^{\mu\nu}}{P_0}, \end{aligned} \quad (9)$$

$$\begin{aligned} \tau_\pi \frac{\dot{\pi}^{(\mu\nu)}}{P_0} + \frac{\pi^{\mu\nu}}{P_0} &= \frac{2\eta}{P_0} \sigma^{\mu\nu} \\ &+ \frac{\pi_\lambda^{(\mu}}{P_0} \left(2\tau_\pi \omega^{\nu)\lambda} - \tau_{\pi\pi} \sigma^{\nu)\lambda} - \delta_{\pi\pi} \theta g^{\nu)\lambda} + \varphi_7 P_0 \frac{\pi^{\nu)\lambda}}{P_0} \right) \\ &+ \ell_{\pi n} \beta_0 \frac{\nabla^{(\mu} n^{\nu)}}{n_0} + \frac{n^{(\mu}}{n_0} \left(\lambda_{\pi n} \beta_0 \nabla^{\nu)} \alpha_0 - \tau_{\pi n} n_0 \frac{\nabla^{\nu)} P_0}{P_0} + \varphi_8 \beta_0^2 P_0 \frac{n^{\nu)}}{n_0} \right). \end{aligned} \quad (10)$$

Here, we have made use of the equation of state of the massless Boltzmann gas, $P_0 = n_0/\beta_0$, with $\beta_0 = 1/T$. By dividing the dissipative quantities by n_0 or P_0 , respectively, we immediately identify terms which are proportional to inverse Reynolds number. Furthermore, the coefficients of terms involving gradients (or time derivatives) all have dimension of time (or mean-free path) and are thus proportional to the Knudsen number. In this form, it is easy to apply power-counting arguments to estimate the order of magnitude of the various terms. The Navier-Stokes terms appearing in the first lines are of first order in Knudsen number. The second lines contain terms proportional to the dissipative quantity that is evolved in the respective equation (n^μ in the first and $\pi^{\mu\nu}$ in the second equation), while the third lines contain cross terms proportional to the other dissipative quantity that is not evolved ($\pi^{\mu\nu}$ in the first and n^μ in the second equation). Terms in the second and third lines are of first order in the product of Knudsen and inverse Reynolds number as well as of second order in inverse Reynolds number.

At this point, one cannot draw any further conclusion without making assumptions about the relative magnitude of Knudsen and inverse Reynolds numbers. In the following, let us assume that all of them are of the *same* order of magnitude, $\text{Kn} \sim \text{R}_n^{-1} \sim \text{R}_\pi^{-1}$ [situations where this is no longer the case were studied, e.g., in Ref. [53]]. At least for asymptotically long times, when the values of the dissipative quantities approach their respective Navier-Stokes limits, this assumption is fulfilled.

Then we can simply assess the order of magnitude (and thus the importance) of the various terms by comparing the values of the coefficients accompanying them. These are listed in Tables I [for Eq. (9)] and II [for Eq. (10)]. Note that the coefficients of terms containing a power of Knudsen number are given in units of the relaxation times τ_n and τ_π , respectively.

$\frac{\kappa}{n_0} [\lambda_{\text{mfp}}]$	$\tau_n [\lambda_{\text{mfp}}]$	$\lambda_{nn} [\tau_n]$	$\delta_{nn} [\tau_n]$	$\frac{\ell_{n\pi}}{\beta_0} [\tau_n]$	$\frac{\tau_{n\pi} P_0}{\beta_0} [\tau_n]$	$\frac{\lambda_{n\pi}}{\beta_0} [\tau_n]$	$\varphi_4 P_0$
$\frac{3}{16}$	$\frac{9}{4}$	$\frac{3}{5}$	1	$\frac{1}{20}$	$\frac{1}{80}$	$\frac{1}{20}$	$\frac{1}{25}$

TABLE I. The coefficients in the particle diffusion equation for a massless Boltzmann gas with constant cross section σ_T in the 14-moment approximation [from Ref. [24]]. $\lambda_{\text{mfp}} = 1/(n_0 \sigma_T)$ is the mean-free path.

$\frac{2\eta}{P_0} [\lambda_{\text{mfp}}]$	$\tau_\pi [\lambda_{\text{mfp}}]$	$\tau_{\pi\pi} [\tau_\pi]$	$\delta_{\pi\pi} [\tau_\pi]$	$\varphi_7 P_0$	$\ell_{\pi n} \beta_0 [\tau_\pi]$	$\lambda_{\pi n} \beta_0 [\tau_\pi]$	$\tau_{\pi n} n_0 [\tau_\pi]$	$\varphi_8 \beta_0^2 P_0$
$\frac{8}{3}$	$\frac{5}{3}$	$\frac{10}{7}$	$\frac{4}{3}$	$\frac{9}{70}$	0	0	0	$\frac{8}{5}$

TABLE II. The coefficients in the shear-stress tensor equation for a massless Boltzmann gas with constant cross section σ_T in the 14-moment approximation [from Ref. [24]]. $\lambda_{\text{mfp}} = 1/(n_0 \sigma_T)$ is the mean-free path.

One immediately observes that the coefficients in the third line of Eq. (9) are (at least) an order of magnitude smaller than the ones in the second line. If Knudsen and inverse Reynolds numbers are of the same order of magnitude, it is then a reasonable assumption to drop these terms altogether, including the last term in Eq. (9) that arises from nonlinear terms in the collision integral.

In the 14-moment approximation, all terms in the third line of Eq. (10), except for the last one, vanish identically. We remark, however, that this is accidental; going beyond the 14-moment approximation [24], all terms are of the same order of magnitude. From the terms in the second line, the last one (arising from the nonlinear terms in the collision integral) is an order of magnitude smaller than the other second-order terms and hence can safely be neglected.

To summarize our results, we conclude that, in the 14-moment approximation, for a massless Boltzmann gas, and for situations where Knudsen and inverse Reynolds numbers are of the same order of magnitude, the equations of motion for the dissipative quantities read, to good approximation, as follows

$$\tau_n \dot{n}^{(\mu)} + n^\mu \simeq \kappa \nabla^\mu \alpha_0 + (\tau_n \omega^{\mu\nu} - \lambda_{nn} \sigma^{\mu\nu}) n_\nu - \delta_{nn} \theta n^\mu, \quad (11)$$

$$\tau_\pi \dot{\pi}^{(\mu\nu)} + \pi^{\mu\nu} \simeq 2\eta \sigma^{\mu\nu} + \pi_\lambda^{(\mu} \left(2\tau_\pi \omega^{\nu)\lambda} - \tau_{\pi\pi} \sigma^{\nu)\lambda} \right) - \delta_{\pi\pi} \theta \pi^{\mu\nu} + \varphi_8 n^{(\mu} n^{\nu)}. \quad (12)$$

In the application of relativistic dissipative fluid dynamics to describe the dynamics of nuclear collisions, this form of the equations should be applicable (far) above the QCD transition, i.e., (deep) in the quark-gluon plasma phase, where the velocity of sound is close to that of an ultrarelativistic gas, $c_s^2 = 1/3$, and the mass of the quasi-particles can be neglected. Nevertheless, the values of the transport coefficients may change from the values given in Tables I and II when accounting for proper quantum statistics, the correct number of degrees of freedom, as well as a more realistic (angular-dependent) scattering cross section. Closer to the QCD phase transition, however, the breaking of conformality and the effect from non-vanishing bulk viscous pressure Π can no longer be neglected.

The remainder of this paper is organized as follows. In Secs. II A and II B we briefly recapitulate the derivation of the equations of relativistic dissipative fluid dynamics from the Boltzmann equation using the method of moments [24]. In Sec. II C we give a general derivation of the second-order corrections to the collision integral. Explicit results will be derived in Sec. III by reducing the number of moments to 14. Details of our calculations are relegated to several Appendices.

We use natural units throughout this work, $\hbar = k_B = c = 1$. Covariant, $x_\mu = (t, -x, -y, -z)$, and contravariant, $x^\mu = (t, x, y, z)$, 4-vectors are related through the metric of flat space-time, $g^{\mu\nu} \equiv \text{diag}(1, -1, -1, -1) \equiv g_{\mu\nu}$, by $x_\mu = g_{\mu\nu} x^\nu$ and $x^\mu = g^{\mu\nu} x_\nu$. Furthermore, the symmetrization and anti-symmetrization operations are denoted by parentheses or brackets around indices, $A^{(\mu\nu)} = (A^{\mu\nu} + A^{\nu\mu})/2$ and $A^{[\mu\nu]} = (A^{\mu\nu} - A^{\nu\mu})/2$, respectively.

II. RELATIVISTIC FLUID DYNAMICS FROM KINETIC THEORY

A. General variables

In relativistic kinetic theory of dilute single-component gases, an ensemble of particles with mass m and 4-momenta $k^\mu = (k^0, \mathbf{k})$, with $k^0 = \sqrt{\mathbf{k}^2 + m^2}$, at a given space-time point $x^\mu = (t, \mathbf{x})$, is characterized by the invariant single-particle distribution function $f_{\mathbf{k}}(t, \mathbf{x}) \equiv f_{\mathbf{k}}$. In the absence of external forces or fields, the space-time evolution of $f_{\mathbf{k}}$ is described by the relativistic Boltzmann equation [54, 55],

$$k^\mu \partial_\mu f_{\mathbf{k}} = C[f], \quad (13)$$

where $C[f]$ is the collision term. In the case of binary elastic collisions it is given by

$$C[f] = \frac{1}{\nu} \int dK' dP dP' W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'} \left(f_{\mathbf{p}} f_{\mathbf{p}'} \tilde{f}_{\mathbf{k}} \tilde{f}_{\mathbf{k}'} - f_{\mathbf{k}} f_{\mathbf{k}'} \tilde{f}_{\mathbf{p}} \tilde{f}_{\mathbf{p}'} \right). \quad (14)$$

Here $\tilde{f}_{\mathbf{k}} = 1 - a f_{\mathbf{k}}$, with $a = 1$ (-1) for fermions (bosons) and $a = 0$ for classical particles, $dK = g d^3\mathbf{k} / [(2\pi)^3 k^0]$ is the Lorentz-invariant momentum-space volume, where g denotes the number of internal degrees of freedom, and $W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'}$ is the Lorentz-invariant transition rate. The factor $\nu = 2$ takes into account that the particles are indistinguishable, while the transition rate satisfies detailed balance, i.e., it is symmetric for time-reversed states $W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'} = W_{\mathbf{p}\mathbf{p}' \rightarrow \mathbf{k}\mathbf{k}'}$.

The conservation of particle number and of energy and momentum in individual collisions leads to the following continuity equations for the particle 4-current, N^μ , and energy-momentum tensor, $T^{\mu\nu}$ [54, 55],

$$\partial_\mu N^\mu \equiv \int dK C[f] = 0, \quad (15)$$

$$\partial_\mu T^{\mu\nu} \equiv \int dK k^\nu C[f] = 0. \quad (16)$$

The currents $N^\mu = N^\mu(t, \mathbf{x})$ and $T^{\mu\nu} = T^{\mu\nu}(t, \mathbf{x})$ are identified as the first and second moments of the single-particle distribution function, respectively. Without loss of generality, they can be tensor-decomposed in terms of the fluid 4-velocity as

$$N^\mu \equiv \langle k^\mu \rangle = \langle E_{\mathbf{k}} \rangle u^\mu + \langle k^{\langle \mu} \rangle, \quad (17)$$

$$T^{\mu\nu} \equiv \langle k^\mu k^\nu \rangle = \langle E_{\mathbf{k}}^2 \rangle u^\mu u^\nu + \frac{1}{3} \Delta^{\mu\nu} \langle \Delta^{\alpha\beta} k_\alpha k_\beta \rangle + \langle k^{\langle \mu} k^{\nu \rangle} \rangle, \quad (18)$$

where $\langle \dots \rangle = \int dK (\dots) f_{\mathbf{k}}$. Here, $E_{\mathbf{k}} = k^\mu u_\mu$ and $k^{\langle \mu} = \Delta^{\mu\nu} k_\nu$ correspond to the energy and the 3-momentum, respectively, of the particle in the local rest frame of the fluid, such that $k^\mu = E_{\mathbf{k}} u^\mu + k^{\langle \mu}$ [23, 24, 56–58]. Moreover, we denoted the orthogonal projection of a first-rank tensor as $A^{\langle \mu} \equiv \Delta^{\mu\nu} A_\nu$, while the symmetric, traceless, and orthogonal projection of second-rank tensors $A^{\mu\nu}$ is defined as $A^{\langle \mu\nu \rangle} \equiv \Delta^{\mu\nu} A^{\alpha\beta}$, with $\Delta^{\mu\nu\alpha\beta} = (\Delta^{\mu\alpha} \Delta^{\beta\nu} + \Delta^{\nu\alpha} \Delta^{\beta\mu})/2 - \Delta^{\mu\nu} \Delta^{\alpha\beta}/3$. In this work, the flow velocity u^μ is defined according to the Landau prescription [13] as the eigenvector of the energy-momentum tensor, i.e., $T^{\mu\nu} u_\nu = \langle E_{\mathbf{k}}^2 \rangle u^\mu$. As a consequence, the energy-momentum diffusion current vanishes, $W^\mu \equiv \langle E_{\mathbf{k}} k^{\langle \mu} \rangle = 0$.

Using Eqs. (17) – (18) we are able to identify the fundamental fluid-dynamical quantities as

$$n \equiv N^\mu u_\mu = \langle E_{\mathbf{k}} \rangle, \quad (19)$$

$$\varepsilon \equiv T^{\mu\nu} u_\mu u_\nu = \langle E_{\mathbf{k}}^2 \rangle, \quad (20)$$

$$P \equiv -\frac{1}{3} T^{\mu\nu} \Delta_{\mu\nu} = -\frac{1}{3} \langle \Delta^{\alpha\beta} k_\alpha k_\beta \rangle, \quad (21)$$

$$n^\mu \equiv N^\nu \Delta_\nu^\mu = \langle k^{\langle \mu} \rangle, \quad (22)$$

$$\pi^{\mu\nu} \equiv T^{\alpha\beta} \Delta_{\alpha\beta}^{\mu\nu} = \langle k^{\langle \mu} k^{\nu \rangle} \rangle, \quad (23)$$

where n is the particle number density, ε is the energy density, P is the local isotropic pressure, n^μ is the particle diffusion current, and $\pi^{\mu\nu}$ is the shear-stress tensor.

It is customary to separate the isotropic pressure into two components, $P = P_0 + \Pi$, with P_0 being the thermodynamic pressure and Π the bulk viscous pressure. The thermodynamic and bulk viscous pressures are defined with respect to the local equilibrium distribution function,

$$f_{0\mathbf{k}} = [\exp(\beta_0 E_{\mathbf{k}} - \alpha_0) + a]^{-1}, \quad (24)$$

and

$$P_0 = -\frac{1}{3} \langle \Delta^{\alpha\beta} k_\alpha k_\beta \rangle_0, \quad (25)$$

$$\Pi = -\frac{1}{3} \langle \Delta^{\alpha\beta} k_\alpha k_\beta \rangle_\delta, \quad (26)$$

where $\langle \dots \rangle_0 = \int dK (\dots) f_{0\mathbf{k}}$ and $\langle \dots \rangle_\delta = \int dK (\dots) \delta f_{\mathbf{k}}$, with $\delta f_{\mathbf{k}} \equiv f_{\mathbf{k}} - f_{0\mathbf{k}}$. The temperature and chemical potential, $\mu = T\alpha_0$, introduced in $f_{0\mathbf{k}}$ are defined by the so-called matching conditions which impose that the particle number density and energy density are given by their respective values in a fictitious local thermodynamic equilibrium state, i.e., $n_0 \equiv \langle E_{\mathbf{k}} \rangle_0 = n$ and $\varepsilon_0 \equiv \langle E_{\mathbf{k}}^2 \rangle_0 = \varepsilon$. In this state, there is an equation of state of the form $P_0(T, \mu)$, such that $n_0 = \partial P_0 / \partial \mu$, $s_0 = \partial P_0 / \partial T$, and the fundamental thermodynamical relation $\varepsilon_0 = T s_0 + \mu n_0 - P_0$ is fulfilled.

It is also convenient to introduce the irreducible moments of $\delta f_{\mathbf{k}}$,

$$\rho_r^{\mu_1 \dots \mu_\ell} = \langle E_{\mathbf{k}}^r k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} \rangle_\delta. \quad (27)$$

Such irreducible moments are constructed to be symmetric, traceless, and orthogonal to the 4-velocity, with the symmetrized, traceless, and orthogonal projections being defined as

$$k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} = \Delta_{\nu_1 \dots \nu_\ell}^{\mu_1 \dots \mu_\ell} k^{\nu_1} \dots k^{\nu_\ell}. \quad (28)$$

The details of construction and the properties of such tensors can be found in Appendix B. Note that the bulk viscous pressure, particle diffusion current, and shear-stress tensor are also irreducible moments of $\delta f_{\mathbf{k}}$,

$$\Pi = -\frac{m^2}{3} \rho_0, \quad n^\mu = \rho_0^\mu, \quad \pi^{\mu\nu} = \rho_0^{\mu\nu}. \quad (29)$$

Furthermore, the matching conditions and the definition of the local rest frame can also be expressed using irreducible moments. The matching conditions correspond to

$$\rho_1 \equiv \langle E_{\mathbf{k}} \rangle_{\delta} = 0, \quad (30)$$

$$\rho_2 \equiv \langle E_{\mathbf{k}}^2 \rangle_{\delta} = 0, \quad (31)$$

while the Landau definition of the fluid 4-velocity leads to

$$\rho_1^{\mu} \equiv \langle k^{(\mu)} E_{\mathbf{k}} \rangle_{\delta} = 0. \quad (32)$$

B. Moment expansion of $f_{\mathbf{k}}$ and the equations of motion for the irreducible moments

Following Ref. [24], $\delta f_{\mathbf{k}}$ is parametrized as

$$\delta f_{\mathbf{k}} = f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}} \phi_{\mathbf{k}}. \quad (33)$$

The function $\phi_{\mathbf{k}}$ is then expanded in terms of a series in the irreducible tensors given in Eq. (28),

$$\phi_{\mathbf{k}} = \sum_{\ell=0}^{\infty} \lambda_{\mathbf{k}}^{\langle \mu_1 \dots \mu_{\ell} \rangle} k_{\langle \mu_1} \dots k_{\mu_{\ell} \rangle}. \quad (34)$$

By expanding the tensor $\lambda_{\mathbf{k}}^{\langle \mu_1 \dots \mu_{\ell} \rangle}$ using a set of orthogonal polynomials, it is straightforward to prove that

$$\lambda_{\mathbf{k}}^{\langle \mu_1 \dots \mu_{\ell} \rangle} = \sum_{n=0}^{N_{\ell}} \mathcal{H}_{\mathbf{k}n}^{(\ell)} \rho_n^{\mu_1 \dots \mu_{\ell}}, \quad (35)$$

where N_{ℓ} denotes the order at which the expansion is truncated (for the coefficient of rank ℓ) and $\rho_n^{\mu_1 \dots \mu_{\ell}}$ is the irreducible moment defined in Eq. (27). The coefficients $\mathcal{H}_{\mathbf{k}n}^{(\ell)}$ are found to be [24]

$$\mathcal{H}_{\mathbf{k}n}^{(\ell)} = \frac{W^{(\ell)}}{\ell!} \sum_{m=n}^{N_{\ell}} a_{mn}^{(\ell)} P_{\mathbf{k}m}^{(\ell)}, \quad (36)$$

with $P_{\mathbf{k}m}^{(\ell)}$ being orthogonal polynomials in $E_{\mathbf{k}}$,

$$P_{\mathbf{k}m}^{(\ell)} = \sum_{r=0}^m a_{mr}^{(\ell)} E_{\mathbf{k}}^r. \quad (37)$$

The coefficients $a_{mr}^{(\ell)}$ are determined from the orthonormality condition

$$\int dK \omega^{(\ell)} P_{\mathbf{k}m}^{(\ell)} P_{\mathbf{k}n}^{(\ell)} = \delta_{mn}, \quad (38)$$

using Gram-Schmidt orthogonalization. The measure $\omega^{(\ell)}$ depends on the rank ℓ of the tensor being expanded and reads

$$\omega^{(\ell)} = \frac{W^{(\ell)}}{(2\ell+1)!!} (\Delta^{\alpha\beta} k_{\alpha} k_{\beta})^{\ell} f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}}, \quad (39)$$

where $W^{(\ell)}$ is a normalization constant defined as

$$W^{(\ell)} = (-1)^{\ell} (J_{2\ell, \ell})^{-1}. \quad (40)$$

For more details, see Refs. [24, 25].

Using the Boltzmann equation, one can derive the general equations of motion satisfied by $\rho_r^{\mu_1 \dots \mu_{\ell}}$. This is accomplished by explicitly taking the comoving derivative of the corresponding irreducible moment, i.e., $\dot{\rho}_r^{\langle \mu_1 \dots \mu_{\ell} \rangle} = \Delta_{\nu_1 \dots \nu_{\ell}}^{\mu_1 \dots \mu_{\ell}} D \int dK E_{\mathbf{k}}^r k^{\langle \nu_1} \dots k^{\nu_{\ell} \rangle} \delta f_{\mathbf{k}}$, and using the Boltzmann equation to express the comoving derivative of $\delta f_{\mathbf{k}}$ in

terms of the collision term, $f_{0\mathbf{k}}$ and its derivatives, and spatial derivatives of $\delta f_{\mathbf{k}}$. The details of this derivation as well as the general form of the resulting equations of motion is contained in Refs. [24, 25]. For the three lowest-rank moments, these equations of motion read

$$\dot{\rho}_r - C_{r-1} = \alpha_r^{(0)} \theta + (\text{nonlinear terms}), \quad (41)$$

$$\dot{\rho}_r^{(\mu)} - C_{r-1}^{(\mu)} = \alpha_r^{(1)} I^\mu + (\text{nonlinear terms}), \quad (42)$$

$$\dot{\rho}_r^{(\mu\nu)} - C_{r-1}^{(\mu\nu)} = 2\alpha_r^{(2)} \sigma^{\mu\nu} + (\text{nonlinear terms}). \quad (43)$$

Here we defined the following thermodynamic quantities,

$$\alpha_r^{(0)} = (1-r) I_{r1} - I_{r0} - \frac{n_0}{D_{20}} (h_0 G_{2r} - G_{3r}), \quad (44)$$

$$\alpha_r^{(1)} = J_{r+1,1} - h_0^{-1} J_{r+2,1}, \quad (45)$$

$$\alpha_r^{(2)} = I_{r+2,1} + (r-1) I_{r+2,2}, \quad (46)$$

where $h_0 = (\varepsilon_0 + P_0)/n_0$ denotes the enthalpy per particle and

$$G_{nm} = J_{n0} J_{m0} - J_{n-1,0} J_{m+1,0}, \quad (47)$$

$$D_{nq} = J_{n+1,q} J_{n-1,q} - (J_{nq})^2. \quad (48)$$

The variables $I_{n+r,q}(\alpha_0, \beta_0)$ and $J_{n+r,q}(\alpha_0, \beta_0)$ correspond to thermodynamic integrals defined as

$$I_{r+n,q} = \frac{(-1)^q}{(2q+1)!!} \left\langle E_{\mathbf{k}}^{n+r-2q} (\Delta^{\alpha\beta} k_\alpha k_\beta)^q \right\rangle_0, \quad (49)$$

$$J_{r+n,q} = \left. \frac{\partial I_{n+r,q}}{\partial \alpha_0} \right|_{\beta_0}. \quad (50)$$

More details can be found in Appendix A.

We also introduced the generalized irreducible collision integral $C_r^{\mu_1 \dots \mu_\ell}$ and its symmetric, traceless, and orthogonal projection,

$$C_r^{\langle \mu_1 \dots \mu_\ell \rangle} \equiv \Delta_{\nu_1 \dots \nu_\ell}^{\mu_1 \dots \mu_\ell} C_r^{\nu_1 \dots \nu_\ell} = \int dK E_{\mathbf{k}}^r k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} C[f]. \quad (51)$$

As was shown in Ref. [24], the moment equations (41) – (43) reduce to the fluid-dynamical equations for the dissipative variables when the fast-varying modes of the Boltzmann equation can be neglected and, simultaneously, the Knudsen number(s) and inverse Reynolds number(s) are small. In this case, the linear parts of the collision integrals introduced above determine the relaxation times for the dissipative variables, while their nonlinear parts give rise to the terms that are of second order in inverse Reynolds number(s), i.e., the tensors \mathcal{R} , \mathcal{R}^μ , and $\mathcal{R}^{\mu\nu}$ that appear in Eqs. (6) – (8). In Ref. [24] the existence of such nonlinear terms was pointed out but the explicit calculation of the corresponding transport coefficients was left for future work. In the next sections we shall complete this task.

C. Expansion of the collision integral

In this section we show how to express the collision integrals in terms of irreducible moments of $\delta f_{\mathbf{k}}$. Substituting the expression of the collision term for binary elastic collisions (14) into the expression for the irreducible collision integral (51), one obtains

$$C_{r-1}^{\langle \mu_1 \dots \mu_\ell \rangle} = \frac{1}{\nu} \int dK dK' dP dP' W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'} E_{\mathbf{k}}^{r-1} k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} \left(f_{\mathbf{p}} f_{\mathbf{p}'} \tilde{f}_{\mathbf{k}} \tilde{f}_{\mathbf{k}'} - f_{\mathbf{k}} f_{\mathbf{k}'} \tilde{f}_{\mathbf{p}} \tilde{f}_{\mathbf{p}'} \right). \quad (52)$$

Substituting the distribution function $f_{\mathbf{k}} = f_{0\mathbf{k}} + f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}} \phi_{\mathbf{k}}$ into the above formula and using

$$f_{\mathbf{p}} f_{\mathbf{p}'} = f_{0\mathbf{p}} f_{0\mathbf{p}'} \left(1 + \tilde{f}_{0\mathbf{p}'} \phi_{\mathbf{p}'} + \tilde{f}_{0\mathbf{p}} \phi_{\mathbf{p}} \right) + f_{0\mathbf{p}} f_{0\mathbf{p}'} \tilde{f}_{0\mathbf{p}} \tilde{f}_{0\mathbf{p}'} \phi_{\mathbf{p}} \phi_{\mathbf{p}'}, \quad (53)$$

$$\tilde{f}_{\mathbf{p}} \tilde{f}_{\mathbf{p}'} = \tilde{f}_{0\mathbf{p}} \tilde{f}_{0\mathbf{p}'} (1 - a f_{0\mathbf{p}'} \phi_{\mathbf{p}'} - a f_{0\mathbf{p}} \phi_{\mathbf{p}}) + a^2 f_{0\mathbf{p}} f_{0\mathbf{p}'} \tilde{f}_{0\mathbf{p}} \tilde{f}_{0\mathbf{p}'} \phi_{\mathbf{p}} \phi_{\mathbf{p}'}, \quad (54)$$

together with the equality $f_{0\mathbf{k}}f_{0\mathbf{k}'}\tilde{f}_{0\mathbf{p}}\tilde{f}_{0\mathbf{p}'} = f_{0\mathbf{p}}f_{0\mathbf{p}'}\tilde{f}_{0\mathbf{k}}\tilde{f}_{0\mathbf{k}'}$, the part that is linear in $\phi_{\mathbf{k}}$ reads

$$L_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle} = \frac{1}{\nu} \int_f E_{\mathbf{k}}^{r-1} k^{\langle\mu_1} \cdots k^{\mu_\ell\rangle} (\phi_{\mathbf{p}} + \phi_{\mathbf{p}'} - \phi_{\mathbf{k}} - \phi_{\mathbf{k}'}), \quad (55)$$

where we abbreviated $\int_f = \int dK dK' dP dP' W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'} f_{0\mathbf{k}} f_{0\mathbf{k}'} \tilde{f}_{0\mathbf{p}} \tilde{f}_{0\mathbf{p}'}$ and used the fact that the collision term vanishes for the local equilibrium distribution function $C_r^{\mu_1\cdots\mu_\ell} [f_0] = 0$. Inserting the expression for $\phi_{\mathbf{k}}$ from the moment expansion (34) into the previous equation, we obtain

$$\begin{aligned} L_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle} &= \frac{1}{\nu} \sum_{m=0}^{\infty} \sum_{n=0}^{N_m} \int_f E_{\mathbf{k}}^{r-1} k^{\langle\mu_1} \cdots k^{\mu_\ell\rangle} \rho_n^{\nu_1\cdots\nu_m} \\ &\times \left(\mathcal{H}_{\mathbf{p}n}^{(m)} p_{\langle\nu_1} \cdots p_{\nu_m\rangle} + \mathcal{H}_{\mathbf{p}'n}^{(m)} p'_{\langle\nu_1} \cdots p'_{\nu_m\rangle} - \mathcal{H}_{\mathbf{k}n}^{(m)} k_{\langle\nu_1} \cdots k_{\nu_m\rangle} - \mathcal{H}_{\mathbf{k}'n}^{(m)} k'_{\langle\nu_1} \cdots k'_{\nu_m\rangle} \right). \end{aligned} \quad (56)$$

It was shown in Ref. [24] that the linear part of the collision integral simplifies to

$$L_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle} = \sum_{m=0}^{\infty} \sum_{n=0}^{N_m} (\mathcal{A}_{rn})^{\mu_1\cdots\mu_\ell}_{\nu_1\cdots\nu_m} \rho_n^{\nu_1\cdots\nu_m} = \sum_{m=0}^{\infty} \sum_{n=0}^{N_m} \mathcal{A}_{rn}^{(\ell)} \rho_n^{\mu_1\cdots\mu_\ell}, \quad (57)$$

where

$$\begin{aligned} (\mathcal{A}_{rn})^{\mu_1\cdots\mu_\ell}_{\nu_1\cdots\nu_m} &= \frac{1}{\nu} \int_f E_{\mathbf{k}}^{r-1} k^{\langle\mu_1} \cdots k^{\mu_\ell\rangle} \\ &\times \left(\mathcal{H}_{\mathbf{p}n}^{(m)} p_{\langle\nu_1} \cdots p_{\nu_m\rangle} + \mathcal{H}_{\mathbf{p}'n}^{(m)} p'_{\langle\nu_1} \cdots p'_{\nu_m\rangle} - \mathcal{H}_{\mathbf{k}n}^{(m)} k_{\langle\nu_1} \cdots k_{\nu_m\rangle} - \mathcal{H}_{\mathbf{k}'n}^{(m)} k'_{\langle\nu_1} \cdots k'_{\nu_m\rangle} \right), \end{aligned} \quad (58)$$

while using the properties of the irreducible projection tensors one can show that

$$\mathcal{A}_{rn}^{(\ell)} = [\Delta_{\alpha_1\cdots\alpha_\ell}^{\mu_1\cdots\mu_\ell}]^{-1} \Delta_{\mu_1\cdots\mu_\ell}^{\nu_1\cdots\nu_\ell} (\mathcal{A}_{rn})^{\mu_1\cdots\mu_\ell}_{\nu_1\cdots\nu_\ell}. \quad (59)$$

The coefficient $\mathcal{A}_{rn}^{(\ell)}$ is the (rn) element of an $(N_\ell + 1) \times (N_\ell + 1)$ matrix, $\mathcal{A}^{(\ell)}$, and, in the linearized case, contains all the information of the underlying microscopic theory. We remark that, for $\ell = 0$, the second and third rows and columns ($r, n = 1, 2$) and, for $\ell = 1$, the second row and column ($r, n = 1$) are zero, because the moments ρ_1 , ρ_2 , and ρ_1^μ vanish due to the matching conditions and our choice of frame.

The computation of the nonlinear part of the collision integral is analogous. Inspecting the previous formulas we observe that the collision integral is a quartic function of $\phi_{\mathbf{k}}$. However, in this paper we shall restrict our calculations to the case of Boltzmann statistics ($a = 0$), in which case the dependence on $\phi_{\mathbf{k}}$ becomes quadratic. The collision integral can be written as

$$C_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle} = L_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle} + N_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle}, \quad (60)$$

where the quadratic contribution to the collision integral reads

$$\begin{aligned} N_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle} &\equiv \frac{1}{\nu} \int_f E_{\mathbf{k}}^{r-1} k^{\langle\mu_1} \cdots k^{\mu_\ell\rangle} (\phi_{\mathbf{p}} \phi_{\mathbf{p}'} - \phi_{\mathbf{k}} \phi_{\mathbf{k}'}) \\ &= \frac{1}{\nu} \sum_{m,m'=0}^{\infty} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m'}} \int_f E_{\mathbf{k}}^{r-1} k^{\langle\mu_1} \cdots k^{\mu_\ell\rangle} \rho_n^{\alpha_1\cdots\alpha_m} \rho_{n'}^{\beta_1\cdots\beta_{m'}} \\ &\times \left(\mathcal{H}_{\mathbf{p}n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m')} p_{\langle\alpha_1} \cdots p_{\alpha_m\rangle} p'_{\langle\beta_1} \cdots p'_{\beta_{m'}\rangle} - \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m')} k_{\langle\alpha_1} \cdots k_{\alpha_m\rangle} k'_{\langle\beta_1} \cdots k'_{\beta_{m'}\rangle} \right). \end{aligned} \quad (61)$$

This nonlinear contribution can be further simplified to

$$N_{r-1}^{\langle\mu_1\cdots\mu_\ell\rangle} = \sum_{m'=0}^{\infty} \sum_{m=0}^{m'} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m'}} (\mathcal{N}_{rnn'})^{\mu_1\cdots\mu_\ell}_{\alpha_1\cdots\alpha_m\beta_1\cdots\beta_{m'}} \rho_n^{\alpha_1\cdots\alpha_m} \rho_{n'}^{\beta_1\cdots\beta_{m'}}, \quad (62)$$

where we defined the following tensor of rank $\ell + m + m'$,

$$\begin{aligned}
(\mathcal{N}_{rnn'})^{\mu_1 \dots \mu_\ell}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} &= \frac{1}{\nu} \int_f E_{\mathbf{k}}^{r-1} k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} \\
&\times \left[\mathcal{H}_{\mathbf{p}n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m')} p_{\langle \alpha_1} \dots p_{\alpha_m \rangle} p'_{\langle \beta_1} \dots p'_{\beta_{m'} \rangle} + (1 - \delta_{mm'}) \mathcal{H}_{\mathbf{p}'n}^{(m)} \mathcal{H}_{\mathbf{p}n'}^{(m')} p'_{\langle \alpha_1} \dots p'_{\alpha_m \rangle} p_{\langle \beta_1} \dots p_{\beta_{m'} \rangle} \right. \\
&\left. - \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m')} k_{\langle \alpha_1} \dots k_{\alpha_m \rangle} k'_{\langle \beta_1} \dots k'_{\beta_{m'} \rangle} - (1 - \delta_{mm'}) \mathcal{H}_{\mathbf{k}'n}^{(m)} \mathcal{H}_{\mathbf{k}n'}^{(m')} k'_{\langle \alpha_1} \dots k'_{\alpha_m \rangle} k_{\langle \beta_1} \dots k_{\beta_{m'} \rangle} \right]. \quad (63)
\end{aligned}$$

In comparison with Eq. (61), we have split the double sum $\sum_{m=0}^{\infty} \sum_{m'=0}^{\infty}$ into a double sum $\sum_{m'=0}^{\infty} \sum_{m=0}^{m'}$ and a double sum $\sum_{m=0}^{\infty} \sum_{m'=0}^m$, and subtracted the superfluous term $m = m'$ in the last sum with the help of a Kronecker delta. Then we interchanged indices $m \leftrightarrow m'$, $n \leftrightarrow n'$ in the second sum.

The tensor $(\mathcal{N}_{rnn'})^{\mu_1 \dots \mu_\ell}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}}$ is symmetric under permutations of μ -type, α -type, and β -type indices, and depends solely on equilibrium distribution functions and the corresponding cross-section(s). The equilibrium distribution function contains only one 4-vector, i.e., the fluid 4-velocity u^μ . Therefore, $(\mathcal{N}_{rnn'})^{\mu_1 \dots \mu_\ell}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}}$ must be constructed from tensor structures made of u^μ and the metric tensor $g^{\mu\nu}$, or, equivalently, u^μ and $\Delta^{\mu\nu}$. Furthermore, $(\mathcal{N}_{rnn'})^{\mu_1 \dots \mu_\ell}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}}$ must be orthogonal to u^μ which implies that it can only be constructed from combinations of elementary projection operators, $\Delta^{\mu\nu}$. This already constrains the rank of the tensor, $\ell + m + m'$, to be an even number. Finally, it must satisfy the following property:

$$\Delta^{\mu'_1 \dots \mu'_\ell} \Delta^{\alpha_1 \dots \alpha_m}_{\alpha'_1 \dots \alpha'_m} \Delta^{\beta_1 \dots \beta_{m'}}_{\beta'_1 \dots \beta'_{m'}} (\mathcal{N}_{rnn'})^{\mu_1 \dots \mu_\ell}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} = (\mathcal{N}_{rnn'})^{\mu'_1 \dots \mu'_\ell}_{\alpha'_1 \dots \alpha'_m \beta'_1 \dots \beta'_{m'}}. \quad (64)$$

For our purposes it is sufficient to calculate terms that are of second order in inverse Reynolds number, i.e., the terms \mathcal{R} , \mathcal{R}^μ , and $\mathcal{R}^{\mu\nu}$. Therefore, we only need to consider the cases $\ell = 0$, $\ell = 1$, and $\ell = 2$. Since the actual deduction of the nonlinear collision integrals is complicated, this task is relegated to Appendix C and here we shall only give the final results.

The scalar nonlinear collision integral from Eq. (62) is given by

$$\begin{aligned}
N_{r-1} &\equiv \sum_{m'=0}^{\infty} \sum_{m=0}^{m'} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m'}} (\mathcal{N}_{rnn'})^{\mu_1 \dots \mu_\ell}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} \rho_n^{\alpha_1 \dots \alpha_m} \rho_{n'}^{\beta_1 \dots \beta_{m'}} \\
&= \sum_{n=0}^{N_0} \sum_{n'=0}^{N_0} \mathcal{C}_{rnn'}^{0(0,0)} \rho_n \rho_{n'} + \sum_{m=1}^{\infty} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_m} \mathcal{C}_{rnn'}^{0(m,m)} \rho_n^{\alpha_1 \dots \alpha_m} \rho_{n', \alpha_1 \dots \alpha_m}, \quad (65)
\end{aligned}$$

where $\mathcal{C}_{rnn'}^{0(m,m)}$ is the special case $\ell = 0$ of a more general coefficient

$$\begin{aligned}
\mathcal{C}_{rnn'}^{\ell(m,m+\ell)} &= \frac{1}{(2m+2\ell+1)\nu} \int_f E_{\mathbf{k}}^{r-1} k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} \\
&\times \left[\mathcal{H}_{\mathbf{p}n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m+\ell)} p^{\langle \nu_1} \dots p^{\nu_m \rangle} p'_{\langle \mu_1} \dots p'_{\mu_\ell} p'_{\nu_1} \dots p'_{\nu_m \rangle} \right. \\
&+ (1 - \delta_{m,m+\ell}) \mathcal{H}_{\mathbf{p}'n}^{(m)} \mathcal{H}_{\mathbf{p}n'}^{(m+\ell)} p'^{\langle \nu_1} \dots p'^{\nu_m \rangle} p_{\langle \mu_1} \dots p_{\mu_\ell} p_{\nu_1} \dots p_{\nu_m \rangle} \\
&- \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m+\ell)} k^{\langle \nu_1} \dots k^{\nu_m \rangle} k'_{\langle \mu_1} \dots k'_{\mu_\ell} k'_{\nu_1} \dots k'_{\nu_m \rangle} \\
&\left. - (1 - \delta_{m,m+\ell}) \mathcal{H}_{\mathbf{k}'n}^{(m)} \mathcal{H}_{\mathbf{k}n'}^{(m+\ell)} k'^{\langle \nu_1} \dots k'^{\nu_m \rangle} k_{\langle \mu_1} \dots k_{\mu_\ell} k_{\nu_1} \dots k_{\nu_m \rangle} \right]. \quad (66)
\end{aligned}$$

Similarly, the nonlinear collision term for $\ell = 1$ becomes,

$$\begin{aligned}
N_{r-1}^\mu &\equiv \sum_{m'=0}^{\infty} \sum_{m=0}^{m'} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m'}} (\mathcal{N}_{rnn'})^\mu_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} \rho_n^{\alpha_1 \dots \alpha_m} \rho_{n'}^{\beta_1 \dots \beta_{m'}} \\
&= \sum_{n=0}^{N_0} \sum_{n'=0}^{N_1} \mathcal{C}_{rnn'}^{1(0,1)} \rho_n \rho_{n'}^\mu + \sum_{m=1}^{\infty} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m+1}} \mathcal{C}_{rnn'}^{1(m,m+1)} \rho_n^{\alpha_1 \dots \alpha_m} \rho_{n', \alpha_1 \dots \alpha_m}^\mu. \quad (67)
\end{aligned}$$

where the coefficient $\mathcal{C}_{rnn'}^{1(m,m+1)}$ is the $\ell = 1$ case of the general coefficient $\mathcal{C}_{rnn'}^{\ell(m,m+\ell)}$ introduced in Eq. (66).

Finally, the rank-2 tensor terms are obtained taking $\ell = 2$,

$$\begin{aligned}
N_{r-1}^{\mu\nu} &\equiv \sum_{m'=0}^{\infty} \sum_{m=0}^{m'} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m'}} (\mathcal{N}_{rnn'})^{\mu\nu}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} \rho_n^{\alpha_1 \dots \alpha_m} \rho_{n'}^{\beta_1 \dots \beta_{m'}} \\
&= \sum_{m=0}^{\infty} \sum_{n=0}^{N_{m+2}} \sum_{n'=0}^{N_m} \mathcal{C}_{rnn'}^{2(m,m+2)} \rho_n^{\alpha_1 \dots \alpha_m} \rho_{n', \alpha_1 \dots \alpha_m}^{\mu\nu} \\
&\quad + \sum_{n=0}^{N_1} \sum_{n'=0}^{N_1} \mathcal{D}_{rnn'}^{2(11)} \rho_n^{\langle \mu} \rho_{n'}^{\nu \rangle} + \sum_{m=2}^{\infty} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_m} \mathcal{D}_{rnn'}^{2(mm)} \rho_n^{\alpha_2 \dots \alpha_m \langle \mu} \rho_{n', \alpha_2 \dots \alpha_m}^{\nu \rangle},
\end{aligned} \tag{68}$$

where $\mathcal{C}_{rnn'}^{2(m,m+2)}$ can be calculated from Eq. (66) and we introduced another coefficient,

$$\begin{aligned}
\mathcal{D}_{rnn'}^{2(mm)} &= \frac{1}{d^{(m)}_{\nu}} \int_f E_{\mathbf{k}}^{r-1} k^{\langle \mu} k^{\nu \rangle} \\
&\quad \times \left(\mathcal{H}_{\mathbf{p}'n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m)} p_{\langle \mu} p^{\beta_{q+1}} \dots p^{\beta_m \rangle} p'_{\langle \nu} p'^{\beta_{q+1}} \dots p'^{\beta_m \rangle} - \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m)} k_{\langle \mu} k^{\beta_{q+1}} \dots k^{\beta_m \rangle} k'_{\langle \nu} k^{\beta_{q+1}} \dots k'^{\beta_m \rangle} \right).
\end{aligned} \tag{69}$$

The normalization $d^{(m)}$ is complicated and is discussed in Appendix C together with other details of the derivation of the nonlinear collision term.

III. TRANSPORT COEFFICIENTS IN THE 14-MOMENT APPROXIMATION

In this section we calculate the previously introduced coefficients $\mathcal{A}_{rn}^{(\ell)}$, $\mathcal{C}_{rnn'}^{0(mm)}$, $\mathcal{C}_{rnn'}^{1(m,m+1)}$, $\mathcal{C}_{rnn'}^{2(m,m+2)}$, and $\mathcal{D}_{rnn'}^{2(m,m)}$ in the 14-moment approximation. As shown in Refs. [24, 25], this corresponds to the truncation $N_0 = 2$, $N_1 = 1$, $N_2 = 0$. This implies that the following irreducible moments appear: $\rho_0 = -3\Pi/m^2$, $\rho_1 = 0$, $\rho_2 = 0$, $\rho_0^\mu = n^\mu$, $\rho_1^\mu = 0$, and $\rho_0^{\mu\nu} = \pi^{\mu\nu}$. As one can see, they are uniquely related to the dissipative quantities.

Before proceeding and for the sake of later convenience, we re-express the coefficients $\mathcal{H}_{\mathbf{k}n}^{(\ell)}$ using Eqs. (36) and (37) as

$$\mathcal{H}_{\mathbf{k}n}^{(\ell)} \equiv \frac{W^{(\ell)}}{\ell!} \sum_{k=n}^{N_\ell} \sum_{r=0}^k a_{kr}^{(\ell)} a_{kn}^{(\ell)} E_{\mathbf{k}}^r = \sum_{r=n}^{N_\ell} A_{rn}^{(\ell)} E_{\mathbf{k}}^r + \sum_{r=0}^{n-1} A_{nr}^{(\ell)} E_{\mathbf{k}}^r, \tag{70}$$

where

$$A_{rn}^{(\ell)} = \frac{W^{(\ell)}}{\ell!} \sum_{k=r}^{N_\ell} a_{kr}^{(\ell)} a_{kn}^{(\ell)}. \tag{71}$$

Note that, for $n = 0$, the second sum in Eq. (70) identically vanishes, which greatly simplifies the calculation of the collision integral.

Furthermore, from the definition of the irreducible moments and using Eqs. (34) – (37) together with the orthogonality condition (B8) we obtain the following general result,

$$\begin{aligned}
\rho_r^{\mu_1 \dots \mu_\ell} &\equiv \frac{\ell!}{(2\ell+1)!!} \sum_{n=0}^{N_\ell} \rho_n^{\mu_1 \dots \mu_\ell} \int dK E_{\mathbf{k}}^r (\Delta^{\alpha\beta} k_\alpha k_\beta)^\ell \mathcal{H}_{\mathbf{k}n}^{(\ell)} f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}} \\
&= (-1)^\ell \ell! \sum_{n=0}^{N_\ell} \rho_n^{\mu_1 \dots \mu_\ell} \left(\sum_{r'=n}^{N_\ell} A_{r'n}^{(\ell)} J_{r+r'+2\ell, \ell} + \sum_{r'=0}^{n-1} A_{nr'}^{(\ell)} J_{r+r'+2\ell, \ell} \right).
\end{aligned} \tag{72}$$

where we used Eq. (70) in the last step. Therefore, truncating the above general result in the 14-moment approximation we obtain

$$\rho_r \equiv \gamma_r^\Pi \rho_0 = -\frac{3}{m^2} \left(A_{00}^{(0)} J_{r,0} + A_{10}^{(0)} J_{r+1,0} + A_{20}^{(0)} J_{r+2,0} \right) \Pi, \tag{73}$$

$$\rho_r^\mu \equiv \gamma_r^n \rho_0^\mu = - \left(A_{00}^{(1)} J_{r+2,1} + A_{10}^{(1)} J_{r+3,1} \right) n^\mu, \tag{74}$$

$$\rho_r^{\mu\nu} \equiv \gamma_r^\pi \rho_0^{\mu\nu} = \left(2A_{00}^{(2)} J_{r+4,2} \right) \pi^{\mu\nu}, \tag{75}$$

where for $r = 0$ we obviously have $\gamma_0^\Pi = \gamma_0^n = \gamma_0^\pi = 1$. The coefficients $A_{20}^{(0)}$, $A_{10}^{(1)}$, $A_{00}^{(2)}$, as well as $A_{00}^{(0)}$, $A_{10}^{(0)}$, $A_{20}^{(0)}$ are calculated from Eq. (71) and listed in Appendix D. These linear relations between the moments are the main result of the 14-moment approximation, which was also obtained in Ref. [25].

It is straightforward to show using Eqs. (58), (59), and (70) that the $\mathcal{A}_{rn}^{(\ell)}$ coefficients of the linear collision term can be expressed in terms of $A_{rn}^{(\ell)}$. For $\ell = 0$ where, in the 14-moment approximation, $N_0 = 2$, the coefficient is

$$\mathcal{A}_{r0}^{(0)} \equiv A_{20}^{(0)} \frac{1}{\nu} \int_f E_{\mathbf{k}}^{r-1} (E_{\mathbf{p}}^2 + E_{\mathbf{p}'}^2 - E_{\mathbf{k}}^2 - E_{\mathbf{k}'}^2) = A_{20}^{(0)} X_{(r-3)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta, \quad (76)$$

where the integrals proportional to $A_{00}^{(0)} \int_f (1 + 1 - 1 - 1) = 0$ and $A_{10}^{(0)} \int_f (E_{\mathbf{p}} + E_{\mathbf{p}'} - E_{\mathbf{k}} - E_{\mathbf{k}'}) = 0$ vanish due to particle number and energy conservation in binary collisions. Here, we introduced the following rank-4 tensor

$$X_{(r)}^{\mu\nu\alpha\beta} = \frac{1}{\nu} \int_f E_{\mathbf{k}}^r k^\mu k^\nu (p^\alpha p^\beta + p'^\alpha p'^\beta - k^\alpha k^\beta - k'^\alpha k'^\beta), \quad (77)$$

which is symmetric upon the interchange of indices (μ, ν) and (α, β) , i.e., $X_{(r)}^{\mu\nu\alpha\beta} = X_{(r)}^{(\mu\nu)(\alpha\beta)}$, and it is also traceless in the latter indices, $X_{(r)}^{\mu\nu\alpha\beta} g_{\alpha\beta} = 0$.

Similarly, for $\ell = 1$ we have

$$\mathcal{A}_{r0}^{(1)} \equiv A_{10}^{(1)} \frac{1}{3\nu} \int_f E_{\mathbf{k}}^{r-1} k^{(\mu)} (E_{\mathbf{p}} p_{(\mu)} + E_{\mathbf{p}'} p'_{(\mu)} - E_{\mathbf{k}} k_{(\mu)} - E_{\mathbf{k}'} k'_{(\mu)}) = A_{10}^{(1)} \frac{1}{3} X_{(r-2)}^{\mu\nu\alpha\beta} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)}, \quad (78)$$

where $A_{00}^{(1)} \int_f (p_{(\mu)} + p'_{(\mu)} - k_{(\mu)} - k'_{(\mu)}) = 0$ vanishes due to 3-momentum conservation.

Finally, for $\ell = 2$ we obtain

$$\mathcal{A}_{r0}^{(2)} \equiv A_{00}^{(2)} \frac{1}{5\nu} \int_f E_{\mathbf{k}}^{r-1} k^{(\mu} k^{\nu)} (p_{(\mu} p_{\nu)} + p'_{(\mu} p'_{\nu)} - k_{(\mu} k_{\nu)} - k'_{(\mu} k'_{\nu)}) = A_{00}^{(2)} \frac{1}{5} X_{(r-1)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu\alpha\beta}. \quad (79)$$

Recalling that $L_{r-1}^{(\mu_1 \dots \mu_\ell)} = \sum_{n=0}^{N_\ell} \mathcal{A}_{rn}^{(\ell)} \rho_n^{\mu_1 \dots \mu_\ell}$, these results lead to the linear collision terms in the 14-moment approximation,

$$L_{r-1} = \sum_{n=0}^2 \mathcal{A}_{rn}^{(0)} \rho_n \equiv \mathcal{A}_{r0}^{(0)} \rho_0 = -A_{20}^{(0)} X_{(r-3),1} \frac{3}{m^2} \Pi, \quad (80)$$

$$L_{r-1}^{(\mu)} = \sum_{n=0}^1 \mathcal{A}_{rn}^{(1)} \rho_n^\mu \equiv \mathcal{A}_{r0}^{(1)} \rho_0^\mu = A_{10}^{(1)} X_{(r-2),3} n^\mu, \quad (81)$$

$$L_{r-1}^{(\mu\nu)} = \mathcal{A}_{r0}^{(2)} \rho_0^{\mu\nu} = A_{00}^{(2)} X_{(r-1),4} \pi^{\mu\nu}. \quad (82)$$

Here, we denoted the different tensor projections as $X_{(r),1} = X_{(r)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta$, $X_{(r),3} = \frac{1}{3} X_{(r)}^{\mu\nu\alpha\beta} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)}$, and $X_{(r),4} = \frac{1}{5} X_{(r)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu\alpha\beta}$.

Now, with the help of these formulas and using Eqs. (73) – (75) the coefficients of bulk viscosity, particle diffusion coefficient, and shear viscosity, as well as the corresponding relaxation times can be calculated,

$$\zeta^r = \frac{\alpha_r^{(0)}}{A_{20}^{(0)} X_{(r-3),1}}, \quad \tau_\Pi^r = -\frac{\gamma_r^\Pi}{A_{20}^{(0)} X_{(r-3),1}}, \quad (83)$$

$$\kappa^r = -\frac{\alpha_r^{(1)}}{A_{10}^{(1)} X_{(r-2),3}}, \quad \tau_n^r = -\frac{\gamma_r^n}{A_{10}^{(1)} X_{(r-2),3}}, \quad (84)$$

$$\eta^r = -\frac{\alpha_r^{(2)}}{A_{00}^{(2)} X_{(r-1),4}}, \quad \tau_\pi^r = -\frac{\gamma_r^\pi}{A_{00}^{(2)} X_{(r-1),4}}, \quad (85)$$

where $\alpha_r^{(0)}$, $\alpha_r^{(1)}$, and $\alpha_r^{(2)}$ were defined in Eqs. (44) – (46) while γ_r^Π , γ_r^n , and γ_r^π are listed in Eqs. (73) – (75).

We now compute the nonlinear collision terms. With Eq. (70), the scalar contribution (66) is

$$\begin{aligned} \mathcal{C}_{rnn'}^{0(mm)} &= \frac{1}{(2m+1)\nu} \int_f E_{\mathbf{k}}^{r-1} \\ &\times \left[\left(\sum_{i=n}^{N_m} A_{in}^{(m)} E_{\mathbf{p}}^i + \sum_{i=0}^{n-1} A_{ni}^{(m)} E_{\mathbf{p}}^i \right) \left(\sum_{i'=n'}^{N_m} A_{i'n'}^{(m)} E_{\mathbf{p}'}^{i'} + \sum_{i'=0}^{n'-1} A_{n'i'}^{(m)} E_{\mathbf{p}'}^{i'} \right) p^{\langle \mu_1} \dots p^{\mu_m \rangle} p'_{\langle \mu_1} \dots p'_{\mu_m \rangle} \right. \\ &\left. - \left(\sum_{i=n}^{N_m} A_{in}^{(m)} E_{\mathbf{k}}^i + \sum_{i=0}^{n-1} A_{ni}^{(m)} E_{\mathbf{k}}^i \right) \left(\sum_{i'=n'}^{N_m} A_{i'n'}^{(m)} E_{\mathbf{k}'}^{i'} + \sum_{i'=0}^{n'-1} A_{n'i'}^{(m)} E_{\mathbf{k}'}^{i'} \right) k^{\langle \mu_1} \dots k^{\mu_m \rangle} k'_{\langle \mu_1} \dots k'_{\mu_m \rangle} \right]. \end{aligned} \quad (86)$$

As noted before, in the 14-moment approximation terms proportional to $\sum_{i=0}^{n-1} A_{ni}^{(m)}$ vanish, hence Eq. (65) leads to

$$N_{r-1} = \frac{9}{m^4} \mathcal{C}_{r00}^{0(00)} \Pi^2 + \mathcal{C}_{r00}^{0(11)} n^\mu n_\mu + \mathcal{C}_{r00}^{0(22)} \pi^{\mu\nu} \pi_{\mu\nu}, \quad (87)$$

where we used Eqs. (73) – (75) for the $\rho_n^{\alpha_1 \dots \alpha_m}$'s. The coefficients in the above equation are

$$\mathcal{C}_{r00}^{0(00)} = \left(A_{10}^{(0)} \right)^2 Y_{2(r-3),1} + \left(A_{20}^{(0)} \right)^2 Y_{5(r-3),1} + \left(A_{00}^{(0)} A_{20}^{(0)} \right) X_{(r-3),1} + \left(A_{10}^{(0)} A_{20}^{(0)} \right) Y_{3(r-3),1}, \quad (88)$$

$$\mathcal{C}_{r00}^{0(11)} = -\frac{1}{3} \left(A_{00}^{(1)} \right)^2 Y_{2(r-3),1} + \left(A_{00}^{(1)} A_{10}^{(1)} \right) Y_{3(r-3),5} + \left(A_{10}^{(1)} \right)^2 Y_{5(r-3),5}, \quad (89)$$

$$\mathcal{C}_{r00}^{0(22)} = 2 \left(A_{00}^{(2)} \right)^2 Y_{5(r-3),9}, \quad (90)$$

where the detailed derivation is given in Appendix E. Comparing the above result to Eq. (6) and taking into account the corresponding relaxation time from Eq. (1), we obtain

$$\varphi_1 = \frac{9}{m^4} \frac{\tau_{\Pi}^r}{\gamma_{\Pi}^r} \mathcal{C}_{r00}^{0(00)}, \quad (91)$$

$$\varphi_2 = \frac{\tau_{\Pi}^r}{\gamma_{\Pi}^r} \mathcal{C}_{r00}^{0(11)}, \quad (92)$$

$$\varphi_3 = \frac{\tau_{\Pi}^r}{\gamma_{\Pi}^r} \mathcal{C}_{r00}^{0(22)}. \quad (93)$$

Similarly, the vector term (67) in the 14-moment approximation leads to the following formula

$$N_{r-1}^\mu = \mathcal{C}_{r00}^{1(12)} n^\nu \pi_\nu^\mu - \frac{3}{m^2} \mathcal{C}_{r00}^{1(01)} \Pi n^\mu, \quad (94)$$

where

$$\mathcal{C}_{r00}^{1(12)} = 2 \left[\left(A_{00}^{(1)} A_{00}^{(2)} \right) Y_{3(r-2),8} + \left(A_{10}^{(1)} A_{00}^{(2)} \right) Y_{4(r-2),8} \right], \quad (95)$$

$$\begin{aligned} \mathcal{C}_{r00}^{1(01)} &= \left(A_{00}^{(0)} A_{10}^{(1)} \right) X_{(r-2),3} + \left(A_{10}^{(0)} A_{00}^{(1)} \right) Y_{1(r-2),3} + \left(A_{10}^{(0)} A_{10}^{(1)} \right) Y_{3(r-2),3} \\ &- \left(A_{20}^{(0)} A_{00}^{(1)} \right) (3Y_{3(r-2),6} + 2Y_{3(r-2),8}) + \left(A_{20}^{(0)} A_{10}^{(1)} \right) Y_{4(r-2),3}. \end{aligned} \quad (96)$$

Now, recalling Eq. (7) we obtain

$$\varphi_4 = \frac{\tau_n^r}{\gamma_n^r} \mathcal{C}_{r00}^{1(12)}, \quad (97)$$

$$\varphi_5 = -\frac{3}{m^2} \frac{\tau_n^r}{\gamma_n^r} \mathcal{C}_{r00}^{1(01)}. \quad (98)$$

Finally, the tensor term (68) is

$$N_{r-1}^{\mu\nu} = -\frac{3}{m^2} \mathcal{C}_{r00}^{2(0,2)} \Pi \pi^{\mu\nu} + \mathcal{D}_{r00}^{2(2,2)} \pi^{\lambda\langle\mu} \pi_{\lambda}^{\nu\rangle} + \mathcal{D}_{r00}^{2(1,1)} n^{\langle\mu} n^{\nu\rangle}, \quad (99)$$

where

$$\mathcal{C}_{r00}^{2(02)} = A_{00}^{(2)} \left[A_{00}^{(0)} X_{(r-1),4} + A_{10}^{(0)} Y_{3(r-1),4} + A_{20}^{(0)} Y_{4(r-1),4} \right], \quad (100)$$

$$\mathcal{D}_{r00}^{2(22)} = 8 \left(A_{00}^{(2)} \right)^2 Y_{5(r-1),11}, \quad (101)$$

$$\mathcal{D}_{r00}^{2(11)} = \left(A_{00}^{(1)} \right)^2 Y_{2(r-1),4} + 2 \left(A_{00}^{(1)} A_{10}^{(1)} \right) Y_{3(r-1),7} + 2 \left(A_{10}^{(1)} \right)^2 Y_{5(r-1),7}, \quad (102)$$

and a comparison with Eq. (8) yields

$$\varphi_6 = -\frac{3}{m^2} \frac{\tau_\pi^r}{\gamma_\pi^r} \mathcal{C}_{r00}^{2(0,2)}, \quad (103)$$

$$\varphi_7 = \frac{\tau_\pi^r}{\gamma_\pi^r} \mathcal{D}_{r00}^{2(2,2)}, \quad (104)$$

$$\varphi_8 = \frac{\tau_\pi^r}{\gamma_\pi^r} \mathcal{D}_{r00}^{2(1,1)}. \quad (105)$$

In order to calculate these coefficients we have introduced five new tensors, which are similar to $X_{(r)}^{\mu\nu\alpha\beta}$ and are given as follows,

$$Y_{1(r)}^{\mu\nu\alpha\beta} = \frac{1}{\nu} \int_f E_{\mathbf{k}}^r k^\mu k^\nu (p^\alpha p'^\beta + p'^\alpha p^\beta - k^\alpha k'^\beta - k'^\alpha k^\beta), \quad (106)$$

$$Y_{2(r)}^{\mu\nu\alpha\beta} = \frac{1}{\nu} \int_f E_{\mathbf{k}}^r k^\mu k^\nu [p^\alpha p'^\beta - k^\alpha k'^\beta], \quad (107)$$

$$Y_{3(r)}^{\mu\nu\alpha\beta\kappa} = \frac{1}{\nu} \int_f E_{\mathbf{k}}^r k^\mu k^\nu (p^\alpha p'^\beta p'^\kappa + p'^\alpha p^\beta p^\kappa - k^\alpha k'^\beta k'^\kappa - k'^\alpha k^\beta k^\kappa), \quad (108)$$

$$Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} = \frac{1}{\nu} \int_f E_{\mathbf{k}}^r k^\mu k^\nu (p^\alpha p^\beta p'^\kappa p'^\lambda + p'^\alpha p'^\beta p^\kappa p^\lambda - k^\alpha k^\beta k'^\kappa k'^\lambda - k'^\alpha k'^\beta k^\kappa k^\lambda), \quad (109)$$

$$Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} = \frac{1}{\nu} \int_f E_{\mathbf{k}}^r k^\mu k^\nu (p^\alpha p^\beta p'^\kappa p'^\lambda - k^\alpha k^\beta k'^\kappa k'^\lambda). \quad (110)$$

Note that the $Y_{i(r),j}$ terms in the previous equations are different contractions of these five tensors. Our notation is such that the i -index specifies the tensor while the j -index labels a particular contraction. More details are given in Appendix F.

We have shown earlier that the coefficients in the equations of motion depend explicitly on the choice of the moment, i.e., the index r . Therefore, once the 14-moment approximation is enforced any moment of the Boltzmann equation leads to a closed set of equations, but when calculating the coefficients of the nonlinear collision integrals one has to account for the exact form of the relaxation equations which follow from Eqs. (41) – (43) using Eqs. (73) – (75). As an example we quote the equations for the particle diffusion current and shear-stress tensor for arbitrary r ,

$$\tau_n^r \dot{n}^{\langle\mu\rangle} + n^\mu + \frac{3}{m^2} \frac{\tau_n^r \mathcal{C}_{r00}^{1(01)}}{\gamma_n^r} \Pi n^\mu - \frac{\tau_n^r \mathcal{C}_{r00}^{1(12)}}{\gamma_n^r} n^\nu \pi_\nu^\mu = \kappa^r \nabla^\mu \alpha_0 + \dots \quad (111)$$

$$\tau_\pi^r \pi^{\langle\mu\nu\rangle} + \pi^{\mu\nu} + \frac{3}{m^2} \frac{\tau_\pi^r \mathcal{C}_{r00}^{2(0,2)}}{\gamma_\pi^r} \Pi \pi^{\mu\nu} - \frac{\tau_\pi^r \mathcal{D}_{r00}^{2(2,2)}}{\gamma_\pi^r} \pi^{\lambda\langle\mu} \pi_{\lambda}^{\nu\rangle} - \frac{\tau_\pi^r \mathcal{D}_{r00}^{2(1,1)}}{\gamma_\pi^r} n^{\langle\mu} n^{\nu\rangle} = 2\eta^r \sigma^{\mu\nu} + \dots \quad (112)$$

Note that in order to recover Eqs. (1) – (3) one must take $r = 0$, i.e., $\zeta = \zeta^0$, $\kappa = \kappa^0$, $\eta = \eta^0$, $\tau_\Pi = \tau_\Pi^0$, $\tau_n = \tau_n^0$ and $\tau_\pi = \tau_\pi^0$.

Using the above relations it was already shown in Ref. [25] how to derive the equations of motion and calculate the transport coefficients for different choices of the moments corresponding to the traditional method by Israel and Stewart [22] and to the one proposed by Denicol, Koide, and Rischke (DKR) [23]. Here, we shall follow this recipe and calculate the coefficients of the nonlinear collision integral in both cases.

The equations of motion derived by Israel and Stewart [22] can be obtained by the choice $\rho_3 = -\frac{3}{m^2} \gamma_3^\Pi \Pi$, $\rho_2^\mu = \gamma_2^n n^\mu$, $\rho_1 = \gamma_1^\pi \pi^{\mu\nu}$ in Eqs. (73) – (75) and substituting these values into the equations of motion (41) – (43). Therefore, in the IS theory all coefficients need to be calculated with $r = 3$ for the scalar moments, $r = 2$ for vector moments, and $r = 1$ for second-rank tensor moments. In contrast, the choice of DKR is to use $r = 0$ for all equations and coefficients.

We explicitly compute some of these coefficients in the ultrarelativistic limit, $m\beta_0 \rightarrow 0$, for a classical gas ($a = 0$) with fixed cross section. Since in this limit $\Pi = 0$, we do not need to compute the coefficients φ_1, φ_2 , and φ_3 appearing in the term \mathcal{R} , Eq. (6), which enters the equation of motion (1) for the bulk viscous pressure. Furthermore, φ_5, φ_6 are coefficients in terms which are proportional to Π , and thus also need not be computed. The remaining, non-vanishing coefficients corresponding to the DKR choice $r = 0$ are simply denoted as φ_4, φ_7 , and φ_8 , while the ones corresponding to the IS choice are denoted by φ_4^{IS} for $r = 2$, while φ_7^{IS} and φ_8^{IS} for $r = 1$. They read

$$\varphi_4 = \frac{1}{25}P_0^{-1}, \quad \varphi_4^{IS} = \frac{1}{4}P_0^{-1}, \quad (113)$$

$$\varphi_7 = \frac{9}{70}P_0^{-1}, \quad \varphi_7^{IS} = \frac{1}{5}P_0^{-1}, \quad (114)$$

$$\varphi_8 = \frac{8}{5\beta_0^2}P_0^{-1}, \quad \varphi_8^{IS} = -\frac{4}{5\beta_0^2}P_0^{-1}. \quad (115)$$

We observe that only φ_8 (multiplying $n^{\langle\mu}n^{\nu\rangle}$) differs in sign between the DKR and IS choices, with the absolute magnitude of the latter being half as large. The coefficients φ_7 (multiplying $\pi^{\lambda\langle\mu}\pi^{\nu\rangle}$) are approximately of the same magnitude for both choices, while φ_4 (multiplying $n_\nu\pi^{\mu\nu}$) is more than six times smaller in DKR than in IS theory. The implications of these results have already been discussed in the Introduction and Conclusions for the DKR choice.

Finally, for further reference, we also quote the coefficients of particle diffusion and shear viscosity,

$$\kappa = \frac{3}{16}n_0\lambda_{\text{mfp}}, \quad \kappa^{IS} = \frac{1}{8}n_0\lambda_{\text{mfp}}, \quad (116)$$

$$\eta = \frac{4}{3}P_0\lambda_{\text{mfp}}, \quad \eta^{IS} = \frac{6}{5}P_0\lambda_{\text{mfp}}, \quad (117)$$

where $\lambda_{\text{mfp}} = 1/(n_0\sigma_T)$ is the mean-free path and σ_T is the total cross section. Note that all remaining transport coefficients from Eq. (4) were already computed in Ref. [25] for both the DKR and IS choices and it was shown that the differences are of the order of 10 – 30%. Therefore, the only coefficients that change considerably from one formalism to the other are φ_4 and φ_8 .

In closing we remark that the numerical solutions of both the IS and DKR theories were compared to the numerical solutions of the Boltzmann equation in various cases [23, 53, 59–62]. These investigations showed the advantages of the DKR choice for the corresponding coefficients, which leads to a far better agreement with numerical solutions of the Boltzmann equation than the IS theory.

ACKNOWLEDGMENTS

This work was supported by the Helmholtz International Center for FAIR within the framework of the LOEWE program launched by the State of Hesse. G.S. Denicol is supported by a Banting Fellowship of the Natural Sciences and Engineering Research Council of Canada. The work of H. Niemi was supported by Academy of Finland, Project No. 133005. E. Molnár was partially supported by the European Union and the European Social Fund through project Supercomputer, the national virtual lab (grant no.: TAMOP-4.2.2.C-11/1/KONV-2012-0010), as well as by TAMOP 4.2.4. A/2-11-1-2012-0001 National Excellence Program (A2-MZPDÖ-13-0042).

The authors thank R. Paatelainen for his help with FeynCalc.

Appendix A: Thermodynamic integrals

Following Refs. [22, 28] we introduced the following equilibrium moments of tensor rank n ,

$$I_n^{\mu_1 \dots \mu_n} \equiv \langle k^{\mu_1} \dots k^{\mu_n} \rangle_0 = \sum_{q=0}^{[n/2]} (-1)^q b_{nq} I_{nq} \Delta^{(\mu_1 \mu_2} \dots \Delta^{\mu_{2q-1} \mu_{2q}} u^{\mu_{2q+1}} \dots u^{\mu_n)}, \quad (A1)$$

where n, q are natural numbers and $[n/2]$ is the largest integer not exceeding $n/2$, cf. Eq. (A8) in Ref. [22]. The parentheses (...) around indices denote symmetrization. For an arbitrary tensor of rank n , this operation is defined by $A^{(\mu_1 \dots \mu_n)} = \frac{1}{n!} \sum_{\varphi_\mu} A^{\mu_1 \mu_2 \dots \mu_n}$, where φ_μ denotes all possible permutations of the μ -indices.

The b_{nq} coefficient is equal to the number of permutations in the set \wp_μ , which lead to identical tensor products of the u^μ and $\Delta^{\mu\nu}$ projectors [22],

$$b_{nq} \equiv \frac{n!}{2^q q! (n-2q)!} = \frac{n! (2q-1)!!}{(2q)! (n-2q)!}, \quad (\text{A2})$$

see Eq. (A2) of Ref. [22].

The thermodynamic integrals I_{nq} and J_{nq} were defined in Eqs. (49), (50):

$$I_{nq} = \frac{(-1)^q}{(2q+1)!!} \int dK (E_{\mathbf{k}})^{n-2q} (\Delta^{\alpha\beta} k_\alpha k_\beta)^q f_{0\mathbf{k}}, \quad (\text{A3})$$

$$J_{nq} = \frac{(-1)^q}{(2q+1)!!} \int dK (E_{\mathbf{k}})^{n-2q} (\Delta^{\alpha\beta} k_\alpha k_\beta)^q f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}}. \quad (\text{A4})$$

Replacing $(\Delta^{\alpha\beta} k_\alpha k_\beta)^q = (m^2 - E_{\mathbf{k}}^2)^q$ we get the following recursion relations for $0 \leq q \leq n/2$,

$$I_{n+2,q} = m^2 I_{n,q} + (2q+3) I_{n+2,q+1}, \quad (\text{A5})$$

$$J_{n+2,q} = m^2 J_{n,q} + (2q+3) J_{n+2,q+1}, \quad (\text{A6})$$

while an integration by parts of Eq. (A3) leads to the following relation,

$$\beta_0 J_{nq} = I_{n-1,q-1} + (n-2q) I_{n-1,q}. \quad (\text{A7})$$

Furthermore,

$$\begin{aligned} dI_{nq}(\alpha_0, \beta_0) &\equiv \frac{\partial I_{nq}}{\partial \alpha_0} d\alpha_0 + \frac{\partial I_{nq}}{\partial \beta_0} d\beta_0, \\ &= J_{nq} d\alpha_0 - J_{n+1,q} d\beta_0, \end{aligned} \quad (\text{A8})$$

with a similar relation for $dJ_{nq}(\alpha_0, \beta_0)$.

Appendix B: Irreducible tensors

We define the following projection operator [54]

$$\Delta^{\mu_1 \dots \mu_n \nu_1 \dots \nu_n} = \sum_{q=0}^{[n/2]} c_{nq} \Phi_{(nq)}^{\mu_1 \dots \mu_n \nu_1 \dots \nu_n}, \quad (\text{B1})$$

where the coefficients are given by

$$c_{nq} = (-1)^q \frac{(n!)^2}{(2n)!} \frac{(2n-2q)!}{q! (n-q)! (n-2q)!}, \quad (\text{B2})$$

and

$$\Phi_{(nq)}^{\mu_1 \dots \mu_n \nu_1 \dots \nu_n} = (n-2q)! \left(\frac{2^q q!}{n!} \right)^2 \sum_{\wp_\mu \wp_\nu} \Delta^{\mu_1 \mu_2} \dots \Delta^{\mu_{2q-1} \mu_{2q}} \Delta^{\nu_1 \nu_2} \dots \Delta^{\nu_{2q-1} \nu_{2q}} \Delta^{\mu_{2q+1} \nu_{2q+1}} \dots \Delta^{\mu_n \nu_n}. \quad (\text{B3})$$

The summation is taken over all *distinct* permutations of μ - and ν -type indices (without mutual exchange of these types of indices). The prefactor is the inverse of the number of distinct permutations. This can be seen as follows: the *total* number of permutations of μ - and ν -type indices is $(n!)^2$. In order to obtain the number of *distinct* permutations, we have to divide this by the following three numbers: $(2^q)^2$ permutations lead to terms which only differ by a trivial permutation of indices on the same projector, e.g. $\Delta^{\mu_1 \mu_2} = \Delta^{\mu_2 \mu_1}$; $(q!)^2$ terms just correspond to a pairwise exchange of indices between projectors, e.g. $\Delta^{\mu_1 \mu_2} \Delta^{\mu_3 \mu_4} = \Delta^{\mu_3 \mu_4} \Delta^{\mu_1 \mu_2}$; for any given distribution of μ -type indices in the product $\Delta^{\mu_{2q+1} \nu_{2q+1}} \dots \Delta^{\mu_n \nu_n}$, there are $(n-2q)!$ possible ways to distribute the ν -type indices, which lead to the same product of projectors.

The projection operator has the following properties [for details, see Ref. [54]]:

(i) It is separately symmetric upon interchange of μ - or ν -type indices,

$$\Delta^{\mu_1 \dots \mu_n \nu_1 \dots \nu_n} = \Delta^{(\mu_1 \dots \mu_n)(\nu_1 \dots \nu_n)}. \quad (\text{B4})$$

(ii) It is traceless upon contraction of μ - or ν -type indices,

$$\Delta^{\mu_1 \dots \mu_n \nu_1 \dots \nu_n} g_{\mu_i \mu_j} = \Delta^{\mu_1 \dots \mu_n \nu_1 \dots \nu_n} g_{\nu_i \nu_j} = 0, \quad (\text{B5})$$

for any pair of indices μ_i, μ_j or ν_i, ν_j , where $1 \leq i, j \leq n$.

(iii) The complete contraction is

$$\Delta^{\mu_1 \dots \mu_\ell}_{\mu_1 \dots \mu_\ell} = 2\ell + 1. \quad (\text{B6})$$

The irreducible tensors $k^{\langle \mu_1} \dots k^{\mu_\ell \rangle}$ defined in Eq. (28) are

$$k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} = \Delta^{\mu_1 \dots \mu_\ell}_{\nu_1 \dots \nu_\ell} k^{\nu_1} \dots k^{\nu_\ell}, \quad (\text{B7})$$

where $\Delta^{\mu_1 \dots \mu_\ell}_{\nu_1 \dots \nu_\ell} \equiv \Delta^{\mu_1 \dots \mu_\ell \alpha_1 \dots \alpha_\ell} g_{\alpha_1 \nu_1} \dots g_{\alpha_\ell \nu_\ell}$. Furthermore, the tensors $k^{\langle \mu_1} \dots k^{\mu_m \rangle}$ satisfy the following orthogonality condition,

$$\int dK F_{\mathbf{k}} k^{\langle \mu_1} \dots k^{\mu_m \rangle} k^{\langle \nu_1} \dots k^{\nu_n \rangle} = \frac{m! \delta_{mn}}{(2m+1)!!} \Delta^{\mu_1 \dots \mu_m \nu_1 \dots \nu_m} \int dK F_{\mathbf{k}} (\Delta^{\alpha\beta} k_\alpha k_\beta)^m, \quad (\text{B8})$$

where $F_{\mathbf{k}}$ is an arbitrary scalar function of $E_{\mathbf{k}}$.

Let us explicitly write down the projection operators (B1) which are needed for our calculations. The first one follows from Eq. (B1) for $n = 1$, which defines the elementary projection operator, $\Delta^{\mu_1 \nu_1}$, and hence for any 4-vector we have

$$A^{\langle \mu_1 \rangle} = \Delta^{\mu_1 \nu_1} A_{\nu_1}. \quad (\text{B9})$$

The next one is given for $n = 2$, which defines the symmetric, traceless, and orthogonal projection in case of arbitrary second-rank tensors,

$$\Delta^{\mu_1 \mu_2 \nu_1 \nu_2} = \Delta^{\mu_1 (\nu_1} \Delta^{\mu_2) \nu_2} - \frac{1}{3} \Delta^{\mu_1 \mu_2} \Delta^{\nu_1 \nu_2}, \quad (\text{B10})$$

hence for any second-rank tensor formed from the dyadic product of two 4-vectors, A^{μ_1} and A^{μ_2} , we obtain

$$A^{\langle \mu_1} A^{\mu_2 \rangle} = A^{\langle \mu_1 \rangle} A^{\langle \mu_2 \rangle} - \frac{1}{3} \Delta^{\mu_1 \mu_2} (\Delta^{\alpha\beta} A_\alpha A_\beta). \quad (\text{B11})$$

The case $n = 3$ leads to

$$\begin{aligned} \Delta^{\mu_1 \mu_2 \mu_3 \nu_1 \nu_2 \nu_3} &= \frac{1}{3} \left(\Delta^{\mu_1 \nu_1} \Delta^{\mu_2 (\nu_2} \Delta^{\mu_3) \nu_3} + \Delta^{\mu_1 \nu_2} \Delta^{\mu_2 (\nu_1} \Delta^{\mu_3) \nu_3} + \Delta^{\mu_1 \nu_3} \Delta^{\mu_2 (\nu_1} \Delta^{\mu_3) \nu_2} \right) \\ &\quad - \frac{3}{5} \Delta^{\mu_1 \mu_2} \Delta^{\mu_3 (\nu_1} \Delta^{\nu_2) \nu_3}, \end{aligned} \quad (\text{B12})$$

and so for any rank-3 tensor formed from the dyadic product of the 4-vectors A^{μ_1} , A^{μ_2} , and A^{μ_3} we obtain

$$A^{\langle \mu_1} A^{\mu_2} A^{\mu_3 \rangle} = A^{\langle \mu_1 \rangle} A^{\langle \mu_2 \rangle} A^{\langle \mu_3 \rangle} - \frac{1}{5} \left(\Delta^{\mu_1 \mu_2} A^{\langle \mu_3 \rangle} + \Delta^{\mu_1 \mu_3} A^{\langle \mu_2 \rangle} + \Delta^{\mu_2 \mu_3} A^{\langle \mu_1 \rangle} \right) (\Delta^{\alpha\beta} A_\alpha A_\beta). \quad (\text{B13})$$

Finally, for $n = 4$ Eq. (B1) leads to

$$\begin{aligned} \Delta^{\mu_1 \mu_2 \mu_3 \mu_4 \nu_1 \nu_2 \nu_3 \nu_4} &= \frac{1}{4!} \sum_{\wp_\mu \wp_\nu} \Delta^{\mu_1 \nu_1} \Delta^{\mu_2 \nu_2} \Delta^{\mu_3 \nu_3} \Delta^{\mu_4 \nu_4} - \frac{3}{14} \Delta^{\mu_1 \mu_2} \Delta^{\mu_3 (\nu_1} \Delta^{\nu_2) \nu_3} \Delta^{\mu_4 \nu_4} \\ &\quad - \frac{3}{14} \Delta^{\mu_1 \mu_2} \Delta^{\mu_4 (\nu_1} \Delta^{\nu_2) \nu_3} \Delta^{\mu_3 \nu_4} - \frac{3}{14} \Delta^{\mu_1 \mu_3} \Delta^{\mu_4 (\nu_1} \Delta^{\nu_2) \nu_3} \Delta^{\mu_2 \nu_4} \\ &\quad - \frac{3}{14} \Delta^{\mu_2 \mu_3} \Delta^{\mu_4 (\nu_1} \Delta^{\nu_2) \nu_3} \Delta^{\mu_1 \nu_4} + \frac{3}{35} \Delta^{\mu_1 \mu_2} \Delta^{\mu_3 \mu_4} \Delta^{\nu_1 \nu_2} \Delta^{\nu_3 \nu_4}, \end{aligned} \quad (\text{B14})$$

and

$$\begin{aligned}
A^{\langle\mu_1\rangle} A^{\mu_2} A^{\mu_3} A^{\mu_4} &= A^{\langle\mu_1\rangle} A^{\langle\mu_2\rangle} A^{\langle\mu_3\rangle} A^{\langle\mu_4\rangle} - \frac{3}{14} \Delta^{(\mu_1\mu_2} A^{\langle\mu_3\rangle}) A^{\langle\mu_4\rangle} (\Delta^{\alpha\beta} A_\alpha A_\beta) \\
&\quad - \frac{3}{14} \Delta^{(\mu_1\mu_2} A^{\langle\mu_4\rangle}) A^{\langle\mu_3\rangle} (\Delta^{\alpha\beta} A_\alpha A_\beta) - \frac{3}{14} \Delta^{(\mu_1\mu_4} A^{\langle\mu_3\rangle}) A^{\langle\mu_2\rangle} (\Delta^{\alpha\beta} A_\alpha A_\beta) \\
&\quad - \frac{3}{14} \Delta^{(\mu_4\mu_2} A^{\langle\mu_3\rangle}) A^{\langle\mu_1\rangle} (\Delta^{\alpha\beta} A_\alpha A_\beta) + \frac{3}{35} \Delta^{(\mu_1\mu_2} \Delta^{\mu_3\mu_4}) (\Delta^{\alpha\beta} A_\alpha A_\beta)^2.
\end{aligned} \tag{B15}$$

Note that we also use the notation with mixed indices such as,

$$A^{\langle\mu_1\rangle} A^{\mu_2} \Delta_{\mu_2\nu_1} = A^{\langle\mu_1\rangle} A_{\nu_1}, \tag{B16}$$

$$A^{\langle\mu_1\rangle} A^{\mu_2} A^{\mu_3} A^{\mu_4} \Delta_{\mu_3\mu_4\nu_1\nu_2} = A^{\langle\mu_1\rangle} A^{\mu_2} A_{\nu_1} A_{\nu_2}. \tag{B17}$$

Appendix C: Reduction of collision tensors

In this Appendix, we show how to derive the general structure of the collision integrals introduced in the main text. As already discussed, the tensor structure of $(\mathcal{N}_{rnn'})_{\alpha_1\cdots\alpha_m\beta_1\cdots\beta_{m'}}^{\mu_1\cdots\mu_\ell}$ can only be constructed from tensors formed using projection operators $\Delta^{\mu\nu}$. We start by collecting all possible combinations of projection operators that can appear in $(\mathcal{N}_{rnn'})_{\alpha_1\cdots\alpha_m\beta_1\cdots\beta_{m'}}^{\mu_1\cdots\mu_\ell}$:

- (i) Terms where all μ -type indices pair up on projectors, all α -type indices pair up on projectors, and all β -type indices pair up on projectors, e.g.

$$\Delta^{\mu_1\mu_2} \cdots \Delta^{\mu_{\ell-1}\mu_\ell} \Delta_{\alpha_1\alpha_2} \cdots \Delta_{\alpha_{m-1}\alpha_m} \Delta_{\beta_1\beta_2} \cdots \Delta_{\beta_{m'-1}\beta_{m'}}. \tag{C1}$$

All possible permutations of the μ, α, β -type indices among themselves are allowed. In this case, ℓ, m , and m' must all be even.

- (ii) Terms where at least one μ -type index pairs with an α -type index on a projector, or one μ -type index pairs with a β -type index on a projector, or one α -type index pairs with a β -type index on a projector, e.g.

$$\Delta_{\alpha_1}^{\mu_1} \Delta^{\mu_2\mu_3} \cdots \Delta^{\mu_{\ell-1}\mu_\ell} \Delta_{\alpha_2\alpha_3} \cdots \Delta_{\alpha_{m-1}\alpha_m} \Delta_{\beta_1\beta_2} \cdots \Delta_{\beta_{m'-1}\beta_{m'}} , \tag{C2}$$

$$\Delta_{\beta_1}^{\mu_1} \Delta^{\mu_2\mu_3} \cdots \Delta^{\mu_{\ell-1}\mu_\ell} \Delta_{\alpha_1\alpha_2} \cdots \Delta_{\alpha_{m-1}\alpha_m} \Delta_{\beta_2\beta_3} \cdots \Delta_{\beta_{m'-1}\beta_{m'}} , \tag{C3}$$

$$\Delta_{\alpha_1\beta_1} \Delta^{\mu_1\mu_2} \cdots \Delta^{\mu_{\ell-1}\mu_\ell} \Delta_{\alpha_2\alpha_3} \cdots \Delta_{\alpha_{m-1}\alpha_m} \Delta_{\beta_2\beta_3} \cdots \Delta_{\beta_{m'-1}\beta_{m'}} . \tag{C4}$$

Again, all possible permutations of the μ -type, α -type, and β -type indices are allowed.

- (iii) Terms where *each* μ -type, α -type, and β -type index pairs up with an index of another type. To guarantee that the μ -type indices have sufficiently many partners among the other two types of indices, one must have $\ell \leq m + m'$. Similarly, in order for the α -type indices to pair up in this way, we have to require $m \leq \ell + m'$. Finally, for the β -type indices we need the condition $m' \leq \ell + m$. In this case, only projectors of the type $\Delta_{\alpha_j}^{\mu_i}$, $\Delta_{\beta_j}^{\mu_i}$ or $\Delta_{\alpha_i\beta_j}$ exist, with no left-over projectors containing indices of the same type. Such terms have the form

$$\Delta_{\alpha_p}^{\mu_i} \Delta_{\beta_q}^{\mu_j} \Delta_{\alpha_r\beta_s} \cdots . \tag{C5}$$

Again, all permutations of the μ, α, β -type indices among themselves are allowed.

It is important to emphasize that terms of the type (i) and (ii) by themselves do not satisfy the property (64), since they are not traceless. This can also be seen from the fact that any term which contains at least one projector of the type $\Delta^{\mu_i\mu_j}$, $\Delta_{\alpha_p\alpha_q}$, or $\Delta_{\beta_r\beta_s}$ vanishes when contracted with $\Delta_{\mu_1'\cdots\mu_\ell'}^{\mu_1\cdots\mu_\ell} \Delta_{\alpha_1'\cdots\alpha_m'}^{\alpha_1\cdots\alpha_m} \Delta_{\beta_1'\cdots\beta_{m'}}^{\beta_1\cdots\beta_{m'}}$. Thus, $(\mathcal{N}_{rnn'})_{\alpha_1\cdots\alpha_m\beta_1\cdots\beta_{m'}}^{\mu_1\cdots\mu_\ell}$ cannot be solely constructed from terms of type (i) and (ii) and there must be at least one term of type (iii).

Therefore, terms of type (iii) are of special importance in this derivation and it is convenient to further discuss some of their properties. The inequalities that constrain terms of type (iii), i.e., $\ell \leq m + m'$, $m \leq \ell + m'$, $m' \leq \ell + m$, can be solved and lead to

$$\ell = q + r, \quad m = p + r, \quad m' = p + q, \tag{C6}$$

with $p, q, r = 0, 1, 2, \dots$. Since the ℓ index is always fixed in the summations appearing in Eq. (62), one can re-express the above equations as

$$m = p - q + \ell, \quad m' = p + q, \quad q \leq \ell. \quad (\text{C7})$$

For our purposes it is sufficient to calculate terms of second order in inverse Reynolds number in the terms \mathcal{R} , \mathcal{R}^μ , and $\mathcal{R}^{\mu\nu}$. Therefore, we only need to consider the cases $\ell = 0$, $\ell = 1$, and $\ell = 2$:

1. $\ell = 0$

If $\ell = 0$, the equalities (C7) imply that $m' = m = 0, 1, \dots$ and, consequently, one must have

$$(\mathcal{N}_{rnn'})_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} = \delta_{mm'} \mathcal{C}_{(0)} \Delta_{(\alpha_1 \beta_1} \dots \Delta_{\alpha_m \beta_m)} + [\text{terms of type (i) and (ii)}]. \quad (\text{C8})$$

Contracting Eq. (C8) with $\Delta_{\alpha'_1 \dots \alpha'_m}^{\alpha_1 \dots \alpha_m} \Delta_{\beta'_1 \dots \beta'_{m'}}^{\beta_1 \dots \beta_m}$ and using Eq. (64), we prove that

$$(\mathcal{N}_{rnn'})_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} = \delta_{mm'} \mathcal{C}_{(0)} \Delta_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_m}, \quad (\text{C9})$$

where $\mathcal{C}_{(0)}$ is the trace of $(\mathcal{N}_{rnn'})_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}}$,

$$\begin{aligned} \mathcal{C}_{(0)} &\equiv \left[\Delta_{\beta_1 \dots \beta_m}^{\alpha_1 \dots \alpha_m} \Delta_{\alpha_1 \dots \alpha_m}^{\beta_1 \dots \beta_m} \right]^{-1} \Delta_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_m} (\mathcal{N}_{rnn'})_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_m} \\ &= \frac{1}{(2m+1)\nu} \int_f E_{\mathbf{k}}^{r-1} \left(\mathcal{H}_{\mathbf{p}n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m)} p^{\langle \mu_1} \dots p^{\mu_m \rangle} p'_{\langle \mu_1} \dots p'_{\mu_m \rangle} - \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m)} k^{\langle \mu_1} \dots k^{\mu_m \rangle} k'_{\langle \mu_1} \dots k'_{\mu_m \rangle} \right). \end{aligned} \quad (\text{C10})$$

The coefficient $\mathcal{C}_{(0)} = \mathcal{C}_{rnn'}^{0(mm)}$, the $\ell = 0$ case of Eq. (66). Thus, we obtain Eq. (65) for the scalar nonlinear collision integral.

2. $\ell = 1$

For $\ell = 1$, Eqs. (C7) imply that $m' = m + 1$, and, consequently,

$$(\mathcal{N}_{rnn'})_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}}^\mu = \delta_{m+1, m'} \mathcal{C}_{(1)} \Delta_{(\beta_1}^\mu \Delta_{\beta_2 \alpha_1} \dots \Delta_{\beta_{m+1} \alpha_m)} + [\text{terms of type (i) and (ii)}]. \quad (\text{C11})$$

All permutations of the α -indices and β -indices among themselves are allowed, while permutations of the α -indices with the β -indices are forbidden. Contracting Eq. (C11) with $\Delta_{\mu'}^{\mu} \Delta_{\alpha'_1 \dots \alpha'_m}^{\alpha_1 \dots \alpha_m} \Delta_{\beta'_1 \dots \beta'_{m+1}}^{\beta_1 \dots \beta_{m+1}}$ and using Eq. (64), we prove that

$$(\mathcal{N}_{rnn'})_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+1}}^\mu = \mathcal{C}_{(1)} \Delta_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+1}}^\mu. \quad (\text{C12})$$

The coefficient $\mathcal{C}_{(1)}$ is obtained from the trace of $(\mathcal{N}_{rnn'})_{\beta_1 \dots \beta_{m+1}}^{\mu \alpha_1 \dots \alpha_m}$, i.e.,

$$\begin{aligned} \mathcal{C}_{(1)} &\equiv \left[\Delta_{\beta_1 \dots \beta_{m+1}}^{\mu \alpha_1 \dots \alpha_m} \Delta_{\mu \alpha_1 \dots \alpha_m}^{\beta_1 \dots \beta_{m+1}} \right]^{-1} \Delta_{\mu}^{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+1}} (\mathcal{N}_{rnn'})_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+1}}^\mu \\ &= \frac{1}{[2(m+1)+1]\nu} \int_f E_{\mathbf{k}}^{r-1} k_\mu \\ &\quad \times \left(\mathcal{H}_{\mathbf{p}n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m+1)} p^{\langle \mu} p^{\alpha_1} \dots p^{\alpha_m \rangle} p'_{\langle \alpha_1} \dots p'_{\alpha_m \rangle} p'^{\mu} p^{\alpha_1} \dots p^{\alpha_m \rangle} \right. \\ &\quad \left. - \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m+1)} k^{\langle \mu} k^{\alpha_1} \dots k^{\alpha_m \rangle} k'_{\langle \alpha_1} \dots k'_{\alpha_m \rangle} k'^{\mu} k^{\alpha_1} \dots k^{\alpha_m \rangle} \right). \end{aligned} \quad (\text{C13})$$

Note that $\mathcal{C}_{(1)} = \mathcal{C}_{rnn'}^{1(m, m+1)}$, defined in Eq. (66) in the main text. Thus, we obtain N_{r-1}^μ as given in Eq. (67).

3. $\ell = 2$

For terms with $\ell = 2$, two solutions are possible: $m' = m = 0, 1, \dots$, and $m' = m + 2 = 2, 3, \dots$. Therefore, two different type (iii) tensors can be constructed, leading to

$$\begin{aligned} (\mathcal{N}_{rnn'})^{\mu\nu}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} &= \delta_{m+2, m'} \mathcal{C}_{(2)} \Delta_{(\beta_1}^{\mu} \Delta_{\beta_2}^{\nu} \Delta_{\beta_3 \alpha_1} \dots \Delta_{\beta_{m+2} \alpha_m)} \\ &+ \delta_{mm'} \mathcal{D}_{(2)} \Delta_{(\beta_1}^{(\mu} \Delta_{\alpha_1}^{\nu)} \Delta_{\beta_2 \alpha_2} \dots \Delta_{\beta_m \alpha_m)} \\ &+ [\text{terms of type (i) and (ii)}]. \end{aligned} \quad (\text{C14})$$

All permutations of the α -indices and β -indices among themselves are allowed, while permutations of the α -indices with the β -indices are forbidden. Contracting Eq. (C14) with $\Delta_{\mu\nu}^{\mu' \nu'} \Delta_{\alpha_1' \dots \alpha_m'}^{\alpha_1 \dots \alpha_m} \Delta_{\beta_1' \dots \beta_{m'}}^{\beta_1 \dots \beta_{m'}}$ and using Eq. (64), we prove that

$$\begin{aligned} (\mathcal{N}_{rnn'})^{\mu\nu}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}} &= \delta_{m+2, m'} \mathcal{C}_{(2)} \Delta_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+2}}^{\mu\nu} \\ &+ \delta_{mm'} \mathcal{D}_{(2)} \Delta_{\lambda_1 \sigma}^{\mu\nu} \Delta_{\lambda_2 \dots \lambda_m \alpha_1 \dots \alpha_m}^{\sigma} \Delta_{\beta_1 \dots \beta_m}^{\lambda_1 \dots \lambda_m}, \end{aligned} \quad (\text{C15})$$

with the coefficients $\mathcal{C}_{(2)}$ and $\mathcal{D}_{(2)}$ being obtained from the corresponding trace of $(\mathcal{N}_{rnn'})^{\mu\nu}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m'}}$ when $m' = m + 2$ and $m' = m$, respectively. That is, the coefficient $\mathcal{C}_{(2)}$ is given by

$$\begin{aligned} \mathcal{C}_{(2)} &\equiv \left[\Delta_{\beta_1 \dots \beta_{m+2}}^{\mu\nu \alpha_1 \dots \alpha_m} \Delta_{\mu\nu \alpha_1 \dots \alpha_m}^{\beta_1 \dots \beta_{m+2}} \right]^{-1} \Delta_{\mu\nu}^{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+2}} (\mathcal{N}_{rnn'})^{\mu\nu}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+2}} \\ &= \frac{1}{[2(m + \ell) + 1] \nu} \Delta^{\mu\nu \alpha_1 \dots \alpha_m \beta_1 \dots \beta_{m+2}} \int_f E_{\mathbf{k}}^{r-1} k_{\langle \mu} k_{\nu \rangle} \\ &\times \left(\mathcal{H}_{\mathbf{p}n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m+2)} p_{\langle \alpha_1} \dots p_{\alpha_m \rangle} p'^{\langle \mu} p'^{\nu} p'^{\alpha_1} \dots p'^{\alpha_m \rangle} + \mathcal{H}_{\mathbf{p}'n}^{(m)} \mathcal{H}_{\mathbf{p}n'}^{(m+2)} p'_{\langle \alpha_1} \dots p'_{\alpha_m \rangle} p^{\langle \mu} p^{\nu} p^{\alpha_1} \dots p^{\alpha_m \rangle} \right. \\ &\quad \left. - \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m+2)} k_{\langle \alpha_1} \dots k_{\alpha_m \rangle} k'^{\langle \mu} k'^{\nu} k'^{\alpha_1} \dots k'^{\alpha_m \rangle} - \mathcal{H}_{\mathbf{k}'n}^{(m)} \mathcal{H}_{\mathbf{k}n'}^{(m+2)} k'_{\langle \alpha_1} \dots k'_{\alpha_m \rangle} k^{\langle \mu} k^{\nu} k^{\alpha_1} \dots k^{\alpha_m \rangle} \right), \end{aligned} \quad (\text{C16})$$

while $\mathcal{D}_{(2)}$ is

$$\begin{aligned} \mathcal{D}_{(2)} &\equiv \left[d^{(m)} \right]^{-1} \Delta_{\mu\nu}^{\lambda_1 \sigma} \Delta_{\sigma}^{\lambda_2 \dots \lambda_m \alpha_1 \dots \alpha_m} \Delta_{\lambda_1 \dots \lambda_m}^{\beta_1 \dots \beta_m} (\mathcal{N}_{rnn'})^{\mu\nu}_{\alpha_1 \dots \alpha_m \beta_1 \dots \beta_m} \\ &= \left[d^{(m)} \right]^{-1} \frac{1}{\nu} \int_f E_{\mathbf{k}}^{r-1} k^{\langle \lambda_1} k_{\sigma \rangle} \\ &\times \left(\mathcal{H}_{\mathbf{p}n}^{(m)} \mathcal{H}_{\mathbf{p}'n'}^{(m)} p^{\langle \sigma} p^{\lambda_2} \dots p^{\lambda_m \rangle} p'_{\langle \lambda_1} \dots p'_{\lambda_m \rangle} - \mathcal{H}_{\mathbf{k}n}^{(m)} \mathcal{H}_{\mathbf{k}'n'}^{(m)} k^{\langle \sigma} k^{\lambda_2} \dots k^{\lambda_m \rangle} k'_{\langle \lambda_1} \dots k'_{\lambda_m \rangle} \right), \end{aligned} \quad (\text{C17})$$

where we defined $d^{(m)} = \Delta_{\lambda_1 \sigma}^{\rho_1 \psi} \Delta_{\psi}^{\rho_2 \dots \rho_m \alpha_1 \dots \alpha_m} \Delta_{\sigma}^{\lambda_2 \dots \lambda_m \alpha_1 \dots \alpha_m} \Delta_{\rho_1 \dots \rho_m}^{\lambda_1 \dots \lambda_m}$. The coefficients $\mathcal{C}_{(2)}$ and $\mathcal{D}_{(2)}$ can be identified with the coefficients $\mathcal{C}_{rnn'}^{2(m, m+2)}$ and $\mathcal{D}_{rnn'}^{2(m, m)}$, respectively, defined in the main text in Eqs. (66) and (69), respectively. Thus,

$$\begin{aligned} N_{r-1}^{\mu\nu} &= \sum_{m=0}^{\infty} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m'}} \delta_{m+2, m'} \mathcal{C}_{rnn'}^{2(m, m+2)} \rho_n^{\alpha_1 \dots \alpha_m} \rho_{n'}^{\mu\nu \alpha_1 \dots \alpha_m} \\ &+ \sum_{m=1}^{\infty} \sum_{n=0}^{N_m} \sum_{n'=0}^{N_{m'}} \delta_{mm'} \mathcal{D}_{rnn'}^{2(m, m)} \rho_{n, \lambda_2 \dots \lambda_m}^{\langle \mu} \rho_{n'}^{\nu \rangle \lambda_2 \dots \lambda_m}, \end{aligned} \quad (\text{C18})$$

which was already presented in the main text in Eq. (68). Calculating the trace $d^{(m)}$ for an arbitrary m can be very complicated. In this paper, we shall only do it for the cases $m = 1, 2$, which are actually needed. It follows that, for $m = 1$,

$$d^{(1)} \equiv \Delta_{\lambda_1 \sigma}^{\rho_1 \psi} \Delta_{\psi}^{\alpha_1} \Delta_{\alpha_1}^{\sigma} \Delta_{\rho_1}^{\lambda_1} = \Delta_{\lambda_1 \sigma}^{\lambda_1 \sigma} = 5, \quad (\text{C19})$$

while for $m = 2$ one obtains

$$d^{(2)} \equiv \Delta_{\lambda_1 \sigma}^{\rho_1 \psi} \Delta_{\psi}^{\rho_2 \alpha_1 \alpha_2} \Delta_{\sigma}^{\lambda_2 \alpha_1 \alpha_2} \Delta_{\rho_1 \rho_2}^{\lambda_1 \lambda_2} = \frac{35}{12}. \quad (\text{C20})$$

Appendix D: Expansion coefficients

In this appendix, we construct the polynomials $P_{\mathbf{k}n}^{(\ell)}$, see Eq. (37). For any $\ell \geq 0$, we set

$$P_{\mathbf{k}0}^{(\ell)} \equiv a_{00}^{(\ell)} = 1, \quad (\text{D1})$$

and obtain

$$P_{\mathbf{k}1}^{(0)} = a_{11}^{(0)} E_{\mathbf{k}} + a_{10}^{(0)}, \quad (\text{D2})$$

$$P_{\mathbf{k}1}^{(1)} = a_{11}^{(1)} E_{\mathbf{k}} + a_{10}^{(1)}, \quad (\text{D3})$$

$$P_{\mathbf{k}2}^{(0)} = a_{22}^{(0)} E_{\mathbf{k}}^2 + a_{21}^{(0)} E_{\mathbf{k}} + a_{20}^{(0)}. \quad (\text{D4})$$

From the orthonormality condition $\int dK \omega^{(\ell)} P_{\mathbf{k}i}^{(\ell)} P_{\mathbf{k}j}^{(\ell)} = \delta_{ij}$, it follows that the measure, $\omega^{(\ell)}$, and the normalization constant, $W^{(\ell)}$, are given in Eqs. (39,40). Therefore, using the above equations together with the orthonormality conditions we obtain

$$\frac{a_{10}^{(0)}}{a_{11}^{(0)}} = -\frac{J_{10}}{J_{00}}, \quad \left(a_{11}^{(0)}\right)^2 = \frac{J_{00}^2}{D_{10}}, \quad (\text{D5})$$

$$\frac{a_{21}^{(0)}}{a_{22}^{(0)}} = \frac{G_{12}}{D_{10}}, \quad \frac{a_{20}^{(0)}}{a_{22}^{(0)}} = \frac{D_{20}}{D_{10}}, \quad (\text{D6})$$

$$\left(a_{22}^{(0)}\right)^2 = \frac{J_{00}D_{10}}{J_{20}D_{20} + J_{30}G_{12} + J_{40}D_{10}}, \quad (\text{D7})$$

$$\frac{a_{10}^{(1)}}{a_{11}^{(1)}} = -\frac{J_{31}}{J_{21}}, \quad \left(a_{11}^{(1)}\right)^2 = \frac{J_{21}^2}{D_{31}}, \quad (\text{D8})$$

where the G_{nm} and D_{nq} functions were defined in Eqs. (47), (48).

The coefficients $\mathcal{H}_{\mathbf{k}n}^{(\ell)}$ are defined in Eq. (36). In the 14-moment approximation we only need $\rho_0 = -3\Pi/m^2$, $\rho_0^\mu = n^\mu$ and $\rho_0^{\mu\nu} = \pi^{\mu\nu}$ with $N_0 = 2$, $N_1 = 1$, and $N_2 = 0$. Furthermore $\rho_1 = 0$ and $\rho_2 = 0$ due to the matching conditions (30), while $\rho_1^\mu = 0$ by the choice (32) of the local rest frame. Hence,

$$\mathcal{H}_{\mathbf{k}0}^{(0)} \equiv W^{(0)} \left(a_{00}^{(0)} P_{\mathbf{k}0}^{(0)} + a_{10}^{(0)} P_{\mathbf{k}1}^{(0)} + a_{20}^{(0)} P_{\mathbf{k}2}^{(0)} \right) = A_{00}^{(0)} + A_{10}^{(0)} E_{\mathbf{k}} + A_{20}^{(0)} E_{\mathbf{k}}^2, \quad (\text{D9})$$

$$\mathcal{H}_{\mathbf{k}0}^{(1)} \equiv W^{(1)} \left(a_{00}^{(1)} P_{\mathbf{k}0}^{(1)} + a_{10}^{(1)} P_{\mathbf{k}1}^{(1)} \right) = A_{00}^{(1)} + A_{10}^{(1)} E_{\mathbf{k}}, \quad (\text{D10})$$

$$\mathcal{H}_{\mathbf{k}0}^{(2)} \equiv \frac{W^{(2)}}{2} a_{00}^{(2)} P_{\mathbf{k}0}^{(2)} = A_{00}^{(2)}, \quad (\text{D11})$$

where the $A_{rn}^{(\ell)}$ were introduced in Eq. (71) and, in the 14-moment approximation, are given by

$$A_{00}^{(0)} \equiv W^{(0)} \left[1 + \left(a_{10}^{(0)}\right)^2 + \left(a_{20}^{(0)}\right)^2 \right] = \frac{D_{30}}{J_{20}D_{20} + J_{30}G_{12} + J_{40}D_{10}}, \quad (\text{D12})$$

$$A_{10}^{(0)} \equiv W^{(0)} \left(a_{10}^{(0)} a_{11}^{(0)} + a_{20}^{(0)} a_{21}^{(0)} \right) = \frac{G_{23}}{J_{20}D_{20} + J_{30}G_{12} + J_{40}D_{10}}, \quad (\text{D13})$$

$$A_{20}^{(0)} \equiv W^{(0)} \left(a_{20}^{(0)} a_{22}^{(0)} \right) = \frac{D_{20}}{J_{20}D_{20} + J_{30}G_{12} + J_{40}D_{10}}, \quad (\text{D14})$$

and

$$A_{00}^{(1)} \equiv W^{(1)} \left[1 + \left(a_{10}^{(1)}\right)^2 \right] = -\frac{J_{41}}{D_{31}}, \quad (\text{D15})$$

$$A_{10}^{(1)} \equiv W^{(1)} \left(a_{10}^{(1)} a_{11}^{(1)} \right) = \frac{J_{31}}{D_{31}}, \quad (\text{D16})$$

$$A_{00}^{(2)} \equiv \frac{W^{(2)}}{2} = \frac{1}{2J_{42}}. \quad (\text{D17})$$

Note that these coefficients closely resemble the ones given in Eqs. (108) – (113) of Ref. [25].

Appendix E: The coefficients of the collision terms

In this appendix, we calculate the coefficients of the nonlinear collision integral in the 14-moment approximation. The nonlinear scalar term N_r from Eq. (65) is expanded with the help of the following coefficients,

$$\begin{aligned} \mathcal{C}_{r00}^{0(00)} &\equiv \frac{1}{\nu} \int_f E_{\mathbf{k}}^{r-1} \left(\mathcal{H}_{\mathbf{p}0}^{(0)} \mathcal{H}_{\mathbf{p}'0}^{(0)} - \mathcal{H}_{\mathbf{k}0}^{(0)} \mathcal{H}_{\mathbf{k}'0}^{(0)} \right) \\ &= \frac{1}{\nu} \left(A_{10}^{(0)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} (E_{\mathbf{p}} E_{\mathbf{p}'} - E_{\mathbf{k}} E_{\mathbf{k}'}) + \frac{1}{\nu} \left(A_{20}^{(0)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} (E_{\mathbf{p}}^2 E_{\mathbf{p}'}^2 - E_{\mathbf{k}}^2 E_{\mathbf{k}'}^2) \\ &\quad + \frac{1}{\nu} \left(A_{00}^{(0)} A_{20}^{(0)} \right) \int_f E_{\mathbf{k}}^{r-1} (E_{\mathbf{p}}^2 + E_{\mathbf{p}'}^2 - E_{\mathbf{k}}^2 - E_{\mathbf{k}'}^2) + \frac{1}{\nu} \left(A_{10}^{(0)} A_{20}^{(0)} \right) \int_f E_{\mathbf{k}}^{r-1} (E_{\mathbf{p}} E_{\mathbf{p}'}^2 + E_{\mathbf{p}'} E_{\mathbf{p}}^2 - E_{\mathbf{k}} E_{\mathbf{k}'}^2 - E_{\mathbf{k}'} E_{\mathbf{k}}^2), \end{aligned} \quad (\text{E1})$$

where terms proportional to $\left(A_{00}^{(0)} \right)^2$ and $A_{00}^{(0)} A_{10}^{(0)}$ vanish on account of energy conservation in binary collisions. The above result can be re-expressed using the $X_{(r)}$ and $Y_{i(r)}$ tensors given in Eq. (77) and Eqs. (106) – (110). Thus, after some calculation we obtain

$$\begin{aligned} \mathcal{C}_{r00}^{0(00)} &= \left(A_{10}^{(0)} \right)^2 Y_{2(r-3)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta + \left(A_{20}^{(0)} \right)^2 Y_{5(r-3)}^{\mu\nu\alpha\beta\kappa\lambda} u_\mu u_\nu u_\alpha u_\beta u_\kappa u_\lambda \\ &\quad + \left(A_{00}^{(0)} A_{20}^{(0)} \right) X_{(r-3)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta + \left(A_{10}^{(0)} A_{20}^{(0)} \right) Y_{3(r-3)}^{\mu\nu\alpha\beta\kappa} u_\mu u_\nu u_\alpha u_\beta u_\kappa. \end{aligned} \quad (\text{E2})$$

After this step we still need to evaluate the terms $Y_{2(r-3)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta$, $Y_{5(r-3)}^{\mu\nu\alpha\beta\kappa\lambda} u_\mu u_\nu u_\alpha u_\beta u_\kappa u_\lambda$ etc. This is relegated to Appendix F, for example $Y_{2(r-3)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta = Y_{2(r),1}$ as shown in Eq. (F9). In a similar fashion we repeat the calculation for all components and later, in Appendix H, we calculate them in the massless limit.

The next coefficient is

$$\begin{aligned} \mathcal{C}_{r00}^{0(11)} &\equiv \frac{1}{3\nu} \int_f E_{\mathbf{k}}^{r-1} \left(\mathcal{H}_{\mathbf{p}0}^{(1)} \mathcal{H}_{\mathbf{p}'0}^{(1)} p_{\langle\mu} p'_{\rangle} - \mathcal{H}_{\mathbf{k}0}^{(1)} \mathcal{H}_{\mathbf{k}'0}^{(1)} k_{\langle\mu} k'_{\rangle} \right) \\ &= \frac{1}{3\nu} \left(A_{00}^{(1)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} \left(p_{\langle\mu} p'_{\rangle} - k_{\langle\mu} k'_{\rangle} \right) \\ &\quad + \frac{1}{3\nu} \left(A_{00}^{(1)} A_{10}^{(1)} \right) \int_f E_{\mathbf{k}}^{r-1} \left(E_{\mathbf{p}} p_{\langle\mu} p'_{\rangle} + E_{\mathbf{p}'} p_{\langle\mu} p'_{\rangle} - E_{\mathbf{k}} k_{\langle\mu} k'_{\rangle} - E_{\mathbf{k}'} k_{\langle\mu} k'_{\rangle} \right) \\ &\quad + \frac{1}{3\nu} \left(A_{10}^{(1)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} \left(E_{\mathbf{p}} E_{\mathbf{p}'} p_{\langle\mu} p'_{\rangle} - E_{\mathbf{k}} E_{\mathbf{k}'} k_{\langle\mu} k'_{\rangle} \right), \end{aligned} \quad (\text{E3})$$

so that

$$\mathcal{C}_{r00}^{0(11)} = \frac{1}{3} \left[\left(A_{00}^{(1)} \right)^2 Y_{2(r-3)}^{\mu\nu\alpha\beta} u_\mu u_\nu \Delta_{\alpha\beta} + \left(A_{00}^{(1)} A_{10}^{(1)} \right) Y_{3(r-3)}^{\mu\nu\alpha\beta\kappa} u_\mu u_\nu u_{(\beta} \Delta_{\kappa)\alpha} + \left(A_{10}^{(1)} \right)^2 Y_{5(r-3)}^{\mu\nu\alpha\beta\kappa\lambda} u_\mu u_\nu u_{(\alpha} \Delta_{\beta)(\kappa} u_{\lambda)} \right]. \quad (\text{E4})$$

The last scalar coefficient is,

$$\begin{aligned} \mathcal{C}_{r00}^{0(22)} &\equiv \frac{1}{5\nu} \int_f E_{\mathbf{k}}^{r-1} \left(\mathcal{H}_{\mathbf{p}0}^{(2)} \mathcal{H}_{\mathbf{p}'0}^{(2)} p_{\langle\mu} p^{\nu\rangle} p'_{\langle\mu} p'_{\nu\rangle} - \mathcal{H}_{\mathbf{k}0}^{(2)} \mathcal{H}_{\mathbf{k}'0}^{(2)} k_{\langle\mu} k^{\nu\rangle} k'_{\langle\mu} k'_{\nu\rangle} \right) \\ &= \frac{1}{5\nu} \left(A_{00}^{(2)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} \left(p_{\langle\mu} p^{\nu\rangle} p'_{\langle\mu} p'_{\nu\rangle} - k_{\langle\mu} k^{\nu\rangle} k'_{\langle\mu} k'_{\nu\rangle} \right), \end{aligned} \quad (\text{E5})$$

therefore,

$$\mathcal{C}_{r00}^{0(22)} = \frac{1}{5} \left(A_{00}^{(2)} \right)^2 Y_{5(r-3)}^{\mu\nu\alpha\beta\kappa\lambda} u_\mu u_\nu \Delta_{\alpha\beta\kappa\lambda}. \quad (\text{E6})$$

The vector coefficients are given by

$$\begin{aligned}
\mathcal{C}_{r00}^{1(01)} &\equiv \frac{1}{3\nu} \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(\mathcal{H}_{\mathbf{p}0}^{(0)} \mathcal{H}_{\mathbf{p}'0}^{(1)} p'^{\langle\mu} + \mathcal{H}_{\mathbf{p}'0}^{(0)} \mathcal{H}_{\mathbf{p}0}^{(1)} p^{\langle\mu} - \mathcal{H}_{\mathbf{k}0}^{(0)} \mathcal{H}_{\mathbf{k}'0}^{(1)} k'^{\langle\mu} - \mathcal{H}_{\mathbf{k}'0}^{(0)} \mathcal{H}_{\mathbf{k}0}^{(1)} k^{\langle\mu} \right) \\
&= \frac{1}{3\nu} \left(A_{00}^{(0)} A_{10}^{(1)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(E_{\mathbf{p}} p^{\langle\mu} + E_{\mathbf{p}'} p'^{\langle\mu} - E_{\mathbf{k}} k^{\langle\mu} - E_{\mathbf{k}'} k'^{\langle\mu} \right) \\
&+ \frac{1}{3\nu} \left(A_{10}^{(0)} A_{00}^{(1)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(E_{\mathbf{p}} p'^{\langle\mu} + E_{\mathbf{p}'} p^{\langle\mu} - E_{\mathbf{k}} k'^{\langle\mu} - E_{\mathbf{k}'} k^{\langle\mu} \right) \\
&+ \frac{1}{3\nu} \left(A_{10}^{(0)} A_{10}^{(1)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(E_{\mathbf{p}} E_{\mathbf{p}'} p'^{\langle\mu} + E_{\mathbf{p}'} E_{\mathbf{p}} p^{\langle\mu} - E_{\mathbf{k}} E_{\mathbf{k}'} k'^{\langle\mu} - E_{\mathbf{k}'} E_{\mathbf{k}} k^{\langle\mu} \right) \\
&+ \frac{1}{3\nu} \left(A_{20}^{(0)} A_{00}^{(1)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(E_{\mathbf{p}}^2 p^{\langle\mu} + E_{\mathbf{p}'}^2 p'^{\langle\mu} - E_{\mathbf{k}}^2 k^{\langle\mu} - E_{\mathbf{k}'}^2 k'^{\langle\mu} \right) \\
&+ \frac{1}{3\nu} \left(A_{20}^{(0)} A_{10}^{(1)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(E_{\mathbf{p}}^2 E_{\mathbf{p}'} p'^{\langle\mu} + E_{\mathbf{p}'}^2 E_{\mathbf{p}} p^{\langle\mu} - E_{\mathbf{k}}^2 E_{\mathbf{k}'} k'^{\langle\mu} - E_{\mathbf{k}'}^2 E_{\mathbf{k}} k^{\langle\mu} \right), \tag{E7}
\end{aligned}$$

hence

$$\begin{aligned}
\mathcal{C}_{r00}^{1(01)} &= \frac{1}{3} \left[\left(A_{00}^{(0)} A_{10}^{(1)} \right) X_{(r-2)}^{\mu\nu\alpha\beta} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)} + \left(A_{10}^{(0)} A_{00}^{(1)} \right) Y_{1(r-2)}^{\mu\nu\alpha\beta} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)} \right. \\
&+ \left(A_{10}^{(0)} A_{10}^{(1)} \right) Y_{3(r-2)}^{\mu\nu\alpha\beta\kappa} u_{(\mu} \Delta_{\nu)(\beta} u_{\kappa)} u_{\alpha} + \left(A_{20}^{(0)} A_{00}^{(1)} \right) Y_{3(r-2)}^{\mu\nu\alpha\beta\kappa} \Delta_{\alpha(\mu} u_{\nu)} u_{\beta} u_{\kappa} \\
&\left. + \left(A_{20}^{(0)} A_{10}^{(1)} \right) Y_{4(r-2)}^{\mu\nu\alpha\beta\kappa\lambda} u_{(\mu} \Delta_{\nu)(\kappa} u_{\lambda)} u_{\alpha} u_{\beta} \right]. \tag{E8}
\end{aligned}$$

The second term is

$$\begin{aligned}
\mathcal{C}_{r00}^{1(12)} &\equiv \frac{1}{5\nu} \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(\mathcal{H}_{\mathbf{p}0}^{(1)} \mathcal{H}_{\mathbf{p}'0}^{(2)} p'^{\langle\mu} p'^{\nu\rangle} + \mathcal{H}_{\mathbf{p}'0}^{(1)} \mathcal{H}_{\mathbf{p}0}^{(2)} p'^{\langle\mu} p^{\nu\rangle} - \mathcal{H}_{\mathbf{k}0}^{(1)} \mathcal{H}_{\mathbf{k}'0}^{(2)} k'^{\langle\mu} k'^{\nu\rangle} - \mathcal{H}_{\mathbf{k}'0}^{(1)} \mathcal{H}_{\mathbf{k}0}^{(2)} k'^{\langle\mu} k^{\nu\rangle} \right) \\
&= \frac{1}{5\nu} \left(A_{00}^{(1)} A_{00}^{(2)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(p_{\langle\nu} p'^{\langle\mu} p'^{\nu\rangle} + p'_{\langle\nu} p^{\langle\mu} p^{\nu\rangle} - k_{\langle\nu} k'^{\langle\mu} k'^{\nu\rangle} - k'_{\langle\nu} k^{\langle\mu} k^{\nu\rangle} \right) \\
&+ \frac{1}{5\nu} \left(A_{10}^{(1)} A_{00}^{(2)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} \left(E_{\mathbf{p}} p_{\langle\nu} p'^{\langle\mu} p'^{\nu\rangle} + E_{\mathbf{p}'} p'_{\langle\nu} p^{\langle\mu} p^{\nu\rangle} - E_{\mathbf{k}} k_{\langle\nu} k'^{\langle\mu} k'^{\nu\rangle} - E_{\mathbf{k}'} k'_{\langle\nu} k^{\langle\mu} k^{\nu\rangle} \right), \tag{E9}
\end{aligned}$$

hence

$$\mathcal{C}_{r00}^{1(12)} = \frac{1}{5} A_{00}^{(2)} \left[A_{00}^{(1)} Y_{3(r-2)}^{\mu\nu\alpha\beta\kappa} u_{(\mu} \Delta_{\nu)\alpha\beta\kappa} + \frac{1}{2} A_{10}^{(1)} Y_{4(r-2)}^{\mu\nu\alpha\beta\kappa\lambda} (u_{(\mu} \Delta_{\nu)\alpha\kappa\lambda} u_{\beta} + u_{(\mu} \Delta_{\nu)\beta\kappa\lambda} u_{\alpha}) \right]. \tag{E10}$$

The first tensor coefficient is given by

$$\begin{aligned}
\mathcal{C}_{r00}^{2(02)} &\equiv \frac{1}{5\nu} \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu} \left(\mathcal{H}_{\mathbf{p}0}^{(0)} \mathcal{H}_{\mathbf{p}'0}^{(2)} p'^{\langle\mu} p'^{\nu\rangle} + \mathcal{H}_{\mathbf{p}'0}^{(0)} \mathcal{H}_{\mathbf{p}0}^{(2)} p'^{\langle\mu} p^{\nu\rangle} - \mathcal{H}_{\mathbf{k}0}^{(0)} \mathcal{H}_{\mathbf{k}'0}^{(2)} k'^{\langle\mu} k'^{\nu\rangle} - \mathcal{H}_{\mathbf{k}'0}^{(0)} \mathcal{H}_{\mathbf{k}0}^{(2)} k'^{\langle\mu} k^{\nu\rangle} \right) \\
&= \frac{1}{5\nu} \left(A_{00}^{(0)} A_{00}^{(2)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu} \left(p'^{\langle\mu} p'^{\nu\rangle} + p'^{\langle\mu} p^{\nu\rangle} - k'^{\langle\mu} k'^{\nu\rangle} - k'^{\langle\mu} k^{\nu\rangle} \right) \\
&+ \frac{1}{5\nu} \left(A_{10}^{(0)} A_{00}^{(2)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu} \left(E_{\mathbf{p}} p'^{\langle\mu} p'^{\nu\rangle} + E_{\mathbf{p}'} p'^{\langle\mu} p^{\nu\rangle} - E_{\mathbf{k}} k'^{\langle\mu} k'^{\nu\rangle} - E_{\mathbf{k}'} k'^{\langle\mu} k^{\nu\rangle} \right) \\
&+ \frac{1}{5\nu} \left(A_{20}^{(0)} A_{00}^{(2)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu} \left(E_{\mathbf{p}}^2 p'^{\langle\mu} p'^{\nu\rangle} + E_{\mathbf{p}'}^2 p'^{\langle\mu} p^{\nu\rangle} - E_{\mathbf{k}}^2 k'^{\langle\mu} k'^{\nu\rangle} - E_{\mathbf{k}'}^2 k'^{\langle\mu} k^{\nu\rangle} \right), \tag{E11}
\end{aligned}$$

and so,

$$\mathcal{C}_{r00}^{2(0,2)} = \frac{1}{5} A_{00}^{(2)} \left[A_{00}^{(0)} X_{(r-1)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu\alpha\beta} + A_{10}^{(0)} Y_{3(r-1)}^{\mu\nu\alpha\beta\kappa} u_{\alpha} \Delta_{\mu\nu\beta\kappa} + A_{20}^{(0)} Y_{4(r-1)}^{\mu\nu\alpha\beta\kappa\lambda} u_{\alpha} u_{\beta} \Delta_{\mu\nu\kappa\lambda} \right]. \tag{E12}$$

The $\mathcal{D}_{r00}^{2(11)}$ term is

$$\begin{aligned}
\mathcal{D}_{r00}^{2(11)} &\equiv \frac{1}{5\nu} \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu\rangle} \left(\mathcal{H}_{\mathbf{p}0}^{(1)} \mathcal{H}_{\mathbf{p}'0}^{(1)} p^{\langle\mu\rangle} p'^{\langle\nu\rangle} - \mathcal{H}_{\mathbf{k}0}^{(1)} \mathcal{H}_{\mathbf{k}'0}^{(1)} k^{\langle\mu\rangle} k'^{\langle\nu\rangle} \right) \\
&= \frac{1}{5\nu} \left(A_{00}^{(1)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu\rangle} \left(p^{\langle\mu\rangle} p'^{\langle\nu\rangle} - k^{\langle\mu\rangle} k'^{\langle\nu\rangle} \right) \\
&+ \frac{1}{5\nu} \left(A_{00}^{(1)} A_{10}^{(1)} \right) \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu\rangle} \left(E_{\mathbf{p}} p^{\langle\mu\rangle} p'^{\langle\nu\rangle} + E_{\mathbf{p}'} p'^{\langle\mu\rangle} p'^{\langle\nu\rangle} - E_{\mathbf{k}} k^{\langle\mu\rangle} k'^{\langle\nu\rangle} - E_{\mathbf{k}'} k'^{\langle\mu\rangle} k'^{\langle\nu\rangle} \right) \\
&+ \frac{1}{5\nu} \left(A_{10}^{(1)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu\rangle} \left(E_{\mathbf{p}} E_{\mathbf{p}'} p^{\langle\mu\rangle} p'^{\langle\nu\rangle} - E_{\mathbf{k}} E_{\mathbf{k}'} k^{\langle\mu\rangle} k'^{\langle\nu\rangle} \right), \tag{E13}
\end{aligned}$$

hence,

$$\begin{aligned}
\mathcal{D}_{r00}^{2(11)} &= \frac{1}{5} \left[\left(A_{00}^{(1)} \right)^2 Y_{2(r-1)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu\alpha\beta} + \left(A_{00}^{(1)} A_{10}^{(1)} \right) Y_{3(r-1)}^{\mu\nu\alpha\beta\kappa} \Delta_{\mu\nu\alpha(\beta} u_{\kappa)} \right. \\
&\quad \left. + \frac{1}{2} \left(A_{10}^{(1)} \right)^2 Y_{5(r-1)}^{\mu\nu\alpha\beta\kappa\lambda} (\Delta_{\mu\nu\alpha(\kappa} u_{\lambda)} u_{\beta)} + \Delta_{\mu\nu\beta(\kappa} u_{\lambda)} u_{\alpha)} \right], \tag{E14}
\end{aligned}$$

The last coefficient we need is

$$\begin{aligned}
\mathcal{D}_{r00}^{2(22)} &\equiv \frac{12}{35\nu} \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu\rangle} \left(\mathcal{H}_{\mathbf{p}0}^{(2)} \mathcal{H}_{\mathbf{p}'0}^{(2)} p^{\langle\mu} p_{\lambda\rangle} p'^{\langle\nu} p'^{\lambda\rangle} - \mathcal{H}_{\mathbf{k}0}^{(2)} \mathcal{H}_{\mathbf{k}'0}^{(2)} k^{\langle\mu} k_{\lambda\rangle} k'^{\langle\nu} k'^{\lambda\rangle} \right) \\
&= \frac{12}{35\nu} \left(A_{00}^{(2)} \right)^2 \int_f E_{\mathbf{k}}^{r-1} k_{\langle\mu} k_{\nu\rangle} \left(p^{\langle\mu} p_{\lambda\rangle} p'^{\langle\nu} p'^{\lambda\rangle} - k^{\langle\mu} k_{\lambda\rangle} k'^{\langle\nu} k'^{\lambda\rangle} \right), \tag{E15}
\end{aligned}$$

and so,

$$\mathcal{D}_{r00}^{2(22)} = \frac{12}{35} \left(A_{00}^{(2)} \right)^2 Y_{5(r-1)}^{\mu\nu\alpha\beta\kappa\lambda} \left[\frac{1}{2} \left(\Delta_{\mu\nu\mu_1\kappa} \Delta_{\alpha\beta\lambda}^{\mu_1} + \Delta_{\mu\nu\mu_1\lambda} \Delta_{\alpha\beta\kappa}^{\mu_1} \right) - \frac{1}{3} \Delta_{\mu\nu\alpha\beta} \Delta_{\kappa\lambda} \right]. \tag{E16}$$

Appendix F: Tensor decompositions

In this appendix, we discuss the decompositions of all collision tensors required in the 14-moment approximation. The collision tensor $X_{(r)}^{\mu\nu\alpha\beta} = X_{(r)}^{(\mu\nu)(\alpha\beta)}$ from Eq. (77) is symmetric upon the interchange of indices (μ, ν) and (α, β) , and it is also traceless in the latter indices, $X_{(r)}^{\mu\nu\alpha\beta} g_{\alpha\beta} = 0$, which follows from the mass-shell condition, $k^\mu k_\mu \equiv p^\mu p_\mu = m^2$. Using these properties one can show that $X_{(r)}^{\mu\nu\alpha\beta}$ is a spatially isotropic tensor which can be constructed using the 4-velocity u^μ , the projector $\Delta^{\mu\nu}$, and different scalar coefficients x_{ij} . The most general decomposition of $X_{(r)}^{\mu\nu\alpha\beta}$ which is symmetric upon the interchange of indices (μ, ν) and (α, β) is

$$X^{(\mu\nu)(\alpha\beta)} \equiv x_{10} u^\mu u^\nu u^\alpha u^\beta + x_{11} u^\mu u^\nu \Delta^{\alpha\beta} + x_{21} u^\alpha u^\beta \Delta^{\mu\nu} + 4x_{31} u^{(\mu} \Delta^{\nu)(\alpha} u^{\beta)} + x_{12} \Delta^{\mu\nu} \Delta^{\alpha\beta} + 2x_{22} \Delta^{\mu(\alpha} \Delta^{\beta)\nu}. \tag{F1}$$

The indices of the scalar coefficients x_{ij} are chosen such that the second index (j) denotes the number of projection tensors belonging to the respective coefficient while the first index (i) counts the number of such coefficients. For example x_{10} is the coefficient without any projection tensor, while x_{12} is the first coefficient which contains two elementary projection tensors.

From the tracelessness relation, $X_{(r)}^{\mu\nu\alpha\beta} g_{\alpha\beta} = 0$, follows that we only have four independent coefficients, since

$$x_{11} = -\frac{x_{10}}{3}, \quad x_{22} = -\frac{1}{2} (x_{21} + 3x_{12}). \tag{F2}$$

Thus we obtain

$$X_{(r)}^{\mu\nu\alpha\beta} \equiv (x_{10} u^\mu u^\nu + x_{21} \Delta^{\mu\nu}) \left(u^\alpha u^\beta - \frac{1}{3} \Delta^{\alpha\beta} \right) + 4x_{31} u^{(\mu} \Delta^{\nu)(\alpha} u^{\beta)} - (x_{21} + 3x_{12}) \Delta^{\mu\nu\alpha\beta}. \tag{F3}$$

Introducing the notation $X_{(r),1} = x_{10}$, $X_{(r),2} = x_{21}$, $X_{(r),3} = x_{31}$, and $X_{(r),4} = 2x_{22}$, one can easily show that these coefficients are the result of the following contractions,

$$X_{(r),1} \equiv X_{(r)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta = -X_{(r)}^{\mu\nu\alpha\beta} u_\mu u_\nu \Delta_{\alpha\beta}, \quad (\text{F4})$$

$$X_{(r),2} \equiv \frac{1}{3} X_{(r)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu} u_\alpha u_\beta = -\frac{1}{3} X_{(r)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu} \Delta_{\alpha\beta}, \quad (\text{F5})$$

$$X_{(r),3} = \frac{1}{3} X_{(r)}^{\mu\nu\alpha\beta} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)}, \quad (\text{F6})$$

$$X_{(r),4} = \frac{1}{5} X_{(r)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu\alpha\beta}. \quad (\text{F7})$$

Similarly as shown here, the other rank-4 collision tensors $Y_{1(r)}^{\mu\nu\alpha\beta}$ and $Y_{2(r)}^{\mu\nu\alpha\beta}$ from Eqs. (106), (107) can be decomposed taking into account the symmetry properties $Y_{1(r)}^{\mu\nu\alpha\beta} = Y_{1(r)}^{(\mu\nu)(\alpha\beta)}$, $Y_{2(r)}^{\mu\nu\alpha\beta} = Y_{2(r)}^{(\mu\nu)\alpha\beta}$ and the tracelessness relations, $Y_{1(r)}^{\mu\nu\alpha\beta} g_{\alpha\beta} = 0$, $Y_{2(r)}^{\mu\nu\alpha\beta} g_{\alpha\beta} = 0$. Here we only list the coefficients which are needed for later calculations,

$$Y_{1(r),3} = \frac{1}{3} Y_{1(r)}^{\mu\nu\alpha\beta} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)}, \quad (\text{F8})$$

$$Y_{2(r),1} \equiv Y_{2(r)}^{\mu\nu\alpha\beta} u_\mu u_\nu u_\alpha u_\beta = -Y_{2(r)}^{\mu\nu\alpha\beta} u_\mu u_\nu \Delta_{\alpha\beta}, \quad (\text{F9})$$

$$Y_{2(r),4} = \frac{1}{5} Y_{2(r)}^{\mu\nu\alpha\beta} \Delta_{\mu\nu\alpha\beta}. \quad (\text{F10})$$

In the following we will decompose the rank-5 collision tensor $Y_{3(r)}^{\mu\nu\alpha\beta\kappa} = Y_{3(r)}^{(\mu\nu)(\alpha\beta)\kappa}$ from Eq. (108),

$$\begin{aligned} Y_3^{(\mu\nu)(\alpha\beta)\kappa} &= y_{10} u^\mu u^\nu u^\beta u^\kappa u^\alpha + y_{11} u^\mu u^\nu \Delta^{\beta\kappa} u^\alpha + y_{21} u^\beta u^\kappa \Delta^{\mu\nu} u^\alpha + 4y_{31} u^{(\mu} \Delta^{\nu)(\beta} u^{\kappa)} u^\alpha \\ &+ 2y_{41} u^\mu u^\nu u^{(\beta} \Delta^{\kappa)\alpha} + 2y_{51} u^\beta u^\kappa u^{(\mu} \Delta^{\nu)\alpha} + y_{12} \Delta^{\mu\nu} \Delta^{\beta\kappa} u^\alpha + 2y_{22} \Delta^{(\mu\beta} \Delta^{\kappa)\nu} u^\alpha \\ &+ 2y_{32} \Delta^{\mu\nu} u^{(\beta} \Delta^{\kappa)\alpha} + 2y_{42} \Delta^{\beta\kappa} u^{(\mu} \Delta^{\nu)\alpha} + 4y_{52} \Delta^{\alpha(\mu} \Delta^{\nu)(\beta} u^{\kappa)} + 4y_{62} \Delta^{\alpha(\beta} \Delta^{\kappa)(\mu} u^{\nu)}. \end{aligned} \quad (\text{F11})$$

Using the tracelessness relation $Y_3^{\mu\nu\alpha\beta\kappa} g_{\beta\kappa} = 0$,

$$y_{11} = -\frac{y_{10}}{3}, \quad y_{22} = -\frac{1}{2} (y_{21} + 3y_{12}), \quad y_{51} = -3y_{42} - 2y_{62}, \quad (\text{F12})$$

thus the number of unknown coefficients is nine. We can express them similarly as in the previous cases, however, here we only list the ones which will be used later

$$Y_{3(r),1} \equiv y_{10} = Y_{3(r)}^{\mu\nu\alpha\beta\kappa} u_\mu u_\nu u_\alpha u_\beta u_\kappa, \quad (\text{F13})$$

$$Y_{3(r),3} \equiv y_{31} = \frac{1}{3} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} u_{(\mu} \Delta_{\nu)(\beta} u_{\kappa)} u_\alpha, \quad (\text{F14})$$

$$Y_{3(r),4} \equiv 2y_{22} = \frac{1}{5} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} \Delta_{\mu\nu\beta\kappa} u_\alpha, \quad (\text{F15})$$

$$Y_{3(r),5} \equiv y_{41} = \frac{1}{3} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} u_\mu u_\nu u_{(\beta} \Delta_{\kappa)\alpha}, \quad (\text{F16})$$

$$Y_{3(r),6} \equiv y_{42} = -\frac{1}{15} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} [\Delta_{\alpha(\beta} \Delta_{\kappa)(\mu} u_{\nu)} + 2u_\beta u_\kappa u_{(\mu} \Delta_{\nu)\alpha}], \quad (\text{F17})$$

$$Y_{3(r),7} \equiv y_{52} = \frac{1}{30} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} [3\Delta_{\alpha(\mu} \Delta_{\nu)(\beta} u_{\kappa)} - \Delta_{\mu\nu} u_{(\beta} \Delta_{\kappa)\alpha}] = \frac{1}{10} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} \Delta_{\mu\nu\alpha(\beta} u_{\kappa)}, \quad (\text{F18})$$

$$Y_{3(r),8} \equiv y_{62} = \frac{1}{30} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} [3\Delta_{\alpha(\beta} \Delta_{\kappa)(\mu} u_{\nu)} + u_\beta u_\kappa u_{(\mu} \Delta_{\nu)\alpha}]. \quad (\text{F19})$$

Note that in Eq. (96) we made use of

$$y_{51} \equiv \frac{1}{3} Y_{3(r)}^{\mu\nu\alpha\beta\kappa} u_{(\mu} \Delta_{\nu)\alpha} u_\beta u_\kappa = -3Y_{3(r),6} - 2Y_{3(r),8}. \quad (\text{F20})$$

The last two tensors we need are $Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} \equiv Y_{4(r)}^{(\mu\nu)(\alpha\beta)(\kappa\lambda)} = Y_{4(r)}^{(\mu\nu)(\kappa\lambda)(\alpha\beta)}$ and $Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} = Y_{5(r)}^{(\mu\nu)(\alpha\beta)(\kappa\lambda)}$, where the trace relations are for example $Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\kappa\lambda} = m^2 X_{(r)}^{\mu\nu\alpha\beta}$, $Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\beta\lambda} = m^2 Y_{1(r)}^{\mu\nu\alpha\kappa}$, and $Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\beta\lambda} = m^2 Y_{2(r)}^{\mu\nu\alpha\kappa}$, as

well as $Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\kappa\lambda} g_{\alpha\beta} \equiv Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\kappa\lambda} g_{\alpha\beta} = 0$. Although both rank-6 tensor are very similar, there are important differences due to symmetry, and so we write down both decompositions,

$$\begin{aligned}
Y_{4(r)}^{(\mu\nu)(\alpha\beta)(\kappa\lambda)} &= w_{10} u^\mu u^\nu u^\alpha u^\beta u^\kappa u^\lambda + w_{11} u^\mu u^\nu [\Delta^{\alpha\beta} u^\kappa u^\lambda + \Delta^{\kappa\lambda} u^\alpha u^\beta] + w_{21} u^\alpha u^\beta \Delta^{\mu\nu} u^\kappa u^\lambda \\
&+ 4w_{31} [u^{(\mu} \Delta^{\nu)(\alpha} u^{\beta)} u^\kappa u^\lambda + u^{(\mu} \Delta^{\nu)(\kappa} u^{\lambda)} u^\alpha u^\beta] + 4w_{41} u^\mu u^\nu u^{(\alpha} \Delta^{\beta)(\kappa} u^{\lambda)} \\
&+ w_{12} \Delta^{\mu\nu} [\Delta^{\alpha\beta} u^\kappa u^\lambda + \Delta^{\kappa\lambda} u^\alpha u^\beta] + 2w_{22} [\Delta^{\mu(\alpha} \Delta^{\beta)\nu} u^\kappa u^\lambda + \Delta^{\mu(\kappa} \Delta^{\lambda)\nu} u^\alpha u^\beta] \\
&+ 4w_{32} \Delta^{\mu\nu} u^{(\alpha} \Delta^{\beta)(\kappa} u^{\lambda)} + 4w_{42} [\Delta^{\alpha\beta} u^{(\mu} \Delta^{\nu)(\kappa} u^{\lambda)} + \Delta^{\kappa\lambda} u^{(\mu} \Delta^{\nu)(\alpha} u^{\beta)}] \\
&+ 8w_{52} u^{(\kappa} \Delta^{\lambda)(\mu} \Delta^{\nu)(\alpha} u^{\beta)} + 8w_{62} [u^{(\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)(\kappa} u^{\lambda)} + u^{(\mu} \Delta^{\nu)(\kappa} \Delta^{\lambda)(\alpha} u^{\beta)}] \\
&+ 2w_{72} u^\mu u^\nu \Delta^{\kappa(\alpha} \Delta^{\beta)\lambda} + w_{82} u^\mu u^\nu \Delta^{\alpha\beta} \Delta^{\kappa\lambda} \\
&+ w_{13} \Delta^{\mu\nu} \Delta^{\alpha\beta} \Delta^{\kappa\lambda} + 8w_{23} \Delta^{(\kappa} \Delta^{\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)\lambda)} \\
&+ 2w_{33} [\Delta^{\alpha\beta} \Delta^{\mu(\kappa} \Delta^{\lambda)\nu} + \Delta^{\kappa\lambda} \Delta^{\mu(\alpha} \Delta^{\beta)\nu}] + 2w_{43} \Delta^{\mu\nu} \Delta^{\alpha(\kappa} \Delta^{\lambda)\beta}, \tag{F21}
\end{aligned}$$

where $\Delta^{(\kappa} \Delta^{\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)\lambda)} = \frac{1}{2} (\Delta^{\kappa(\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)\lambda} + \Delta^{\lambda(\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)\kappa})$ and

$$\begin{aligned}
Y_{5(r)}^{(\mu\nu)(\alpha\beta)(\kappa\lambda)} &= z_{10} u^\mu u^\nu u^\alpha u^\beta u^\kappa u^\lambda + z_{11} u^\mu u^\nu \Delta^{\alpha\beta} u^\kappa u^\lambda + z'_{11} u^\mu u^\nu \Delta^{\kappa\lambda} u^\alpha u^\beta + z_{21} u^\alpha u^\beta \Delta^{\mu\nu} u^\kappa u^\lambda \\
&+ 4z_{31} u^{(\mu} \Delta^{\nu)(\alpha} u^{\beta)} u^\kappa u^\lambda + 4z'_{31} u^{(\mu} \Delta^{\nu)(\kappa} u^{\lambda)} u^\alpha u^\beta + 4z_{41} u^\mu u^\nu u^{(\alpha} \Delta^{\beta)(\kappa} u^{\lambda)} \\
&+ z_{12} \Delta^{\mu\nu} \Delta^{\alpha\beta} u^\kappa u^\lambda + z'_{12} \Delta^{\mu\nu} \Delta^{\kappa\lambda} u^\alpha u^\beta + 2z_{22} \Delta^{\mu(\alpha} \Delta^{\beta)\nu} u^\kappa u^\lambda + 2z'_{22} \Delta^{\mu(\kappa} \Delta^{\lambda)\nu} u^\alpha u^\beta \\
&+ 4z_{32} \Delta^{\mu\nu} u^{(\alpha} \Delta^{\beta)(\kappa} u^{\lambda)} + 4z_{42} \Delta^{\alpha\beta} u^{(\mu} \Delta^{\nu)(\kappa} u^{\lambda)} + 4z'_{42} \Delta^{\kappa\lambda} u^{(\mu} \Delta^{\nu)(\alpha} u^{\beta)} \\
&+ 8z_{52} u^{(\kappa} \Delta^{\lambda)(\mu} \Delta^{\nu)(\alpha} u^{\beta)} + 8z_{62} u^{(\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)(\kappa} u^{\lambda)} + 8z'_{62} u^{(\mu} \Delta^{\nu)(\kappa} \Delta^{\lambda)(\alpha} u^{\beta)} \\
&+ 2z_{72} u^\mu u^\nu \Delta^{\kappa(\alpha} \Delta^{\beta)\lambda} + z_{82} u^\mu u^\nu \Delta^{\alpha\beta} \Delta^{\kappa\lambda} \\
&+ z_{13} \Delta^{\mu\nu} \Delta^{\alpha\beta} \Delta^{\kappa\lambda} + 8z_{23} \Delta^{(\kappa} \Delta^{\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)\lambda)} + 2z_{33} \Delta^{\alpha\beta} \Delta^{\mu(\kappa} \Delta^{\lambda)\nu} \\
&+ 2z'_{33} \Delta^{\kappa\lambda} \Delta^{\mu(\alpha} \Delta^{\beta)\nu} + 2z_{43} \Delta^{\mu\nu} \Delta^{\alpha(\kappa} \Delta^{\lambda)\beta}. \tag{F22}
\end{aligned}$$

There are only a few coefficients which we will need to recall later, these are

$$Y_{4(r),3} \equiv w_{31} = \frac{1}{3} Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)} u_{\kappa} u_{\lambda} = \frac{1}{3} Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} u_{(\mu} \Delta_{\nu)(\kappa} u_{\lambda)} u_{\alpha} u_{\beta}, \tag{F23}$$

$$Y_{4(r),4} \equiv 2w_{22} = \frac{1}{5} Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} \Delta_{\mu\nu\alpha\beta} u_{\kappa} u_{\lambda}, \tag{F24}$$

$$\begin{aligned}
Y_{4(r),8} \equiv w_{62} &= \frac{1}{30} Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} [3u_{(\mu} \Delta_{\nu)(\kappa} \Delta_{\lambda)(\alpha} u_{\beta)} - \Delta_{\kappa\lambda} u_{(\mu} \Delta_{\nu)(\alpha} u_{\beta)}], \\
&= \frac{1}{30} Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} [3u_{(\mu} \Delta_{\nu)(\alpha} \Delta_{\beta)(\kappa} u_{\lambda)} - \Delta_{\alpha\beta} u_{(\mu} \Delta_{\nu)(\kappa} u_{\lambda)}], \tag{F25}
\end{aligned}$$

and

$$Y_{5(r),1} \equiv z_{10} = Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} u_{\mu} u_{\nu} u_{\alpha} u_{\beta} u_{\kappa} u_{\lambda}, \tag{F26}$$

$$Y_{5(r),5} \equiv z_{41} = \frac{1}{3} Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} u_{\mu} u_{\nu} u_{(\alpha} \Delta_{\beta)(\kappa} u_{\lambda)}, \tag{F27}$$

$$Y_{5(r),7} \equiv z_{52} = \frac{1}{30} Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} [3u_{(\kappa} \Delta_{\lambda)(\mu} \Delta_{\nu)(\alpha} u_{\beta)} - \Delta_{\mu\nu} u_{(\alpha} \Delta_{\beta)(\kappa} u_{\lambda)}], \tag{F28}$$

$$Y_{5(r),9} \equiv z_{72} = \frac{1}{30} Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} [3\Delta_{\kappa(\alpha} \Delta_{\beta)\lambda} u_{\mu} u_{\nu} - \Delta_{\alpha\beta} \Delta_{\kappa\lambda} u_{\mu} u_{\nu}] = \frac{1}{10} Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} u_{\mu} u_{\nu} \Delta_{\alpha\beta\kappa\lambda}, \tag{F29}$$

$$\begin{aligned}
Y_{5(r),11} \equiv z_{23} &= \frac{1}{210} Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} [2\Delta_{\mu\nu} \Delta_{\alpha\beta} \Delta_{\kappa\lambda} + 9\Delta_{(\kappa} \Delta^{\mu} \Delta^{\nu)(\alpha} \Delta^{\beta)\lambda)}] \\
&- \frac{1}{70} Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} [\Delta_{\alpha\beta} \Delta_{\mu(\kappa} \Delta_{\lambda)\nu} + \Delta_{\mu\nu} \Delta_{\alpha(\kappa} \Delta_{\lambda)\beta} + \Delta_{\kappa\lambda} \Delta_{\mu(\alpha} \Delta_{\beta)\nu}]. \tag{F30}
\end{aligned}$$

It is easy to realize that in case the particles have the same mass, $Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\kappa\lambda} = m^2 X_{(r)}^{\mu\nu\alpha\beta}$, $Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\beta\lambda} = m^2 Y_{1(r)}^{\mu\nu\alpha\kappa}$, also $Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} g_{\beta\lambda} = Y_{2(r)}^{\mu\nu\alpha\kappa}$, and so on. This means that we only have rank-5 and -6 tensors to evaluate, $Y_{3(r)}^{\mu\nu\alpha\beta\kappa}$,

$Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda}$. However, since it is much simpler to decompose and project rank-4 tensors and then use them to check the results through the trace relations for the rank-6 tensors, we will keep using all of them.

Appendix G: Collision integrals in the Boltzmann limit

In this appendix we calculate the previously defined collision tensors Eqs. (77) – (110) in the massless Boltzmann limit. These tensors are based on the collision integral $C[f]$ defined in Eq. (14), where the Lorentz-invariant transition rate, $W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'}$, may only depend on the following collision invariant

$$s \equiv (k^\mu + k'^\mu)^2 = (p^\mu + p'^\mu)^2. \quad (\text{G1})$$

In the center of mass (CM) frame, where the sum of 3-momenta vanishes $\mathbf{k} + \mathbf{k}' = \mathbf{p} + \mathbf{p}' = 0$ and so $k^0 = k'^0 = p^0 = p'^0$, the scattering angle θ_s is given with the help of another collision invariant, $t \equiv (k^\mu - p^\mu)^2 = (k'^\mu - p'^\mu)^2$, such that

$$\cos \theta_s \equiv 1 + \frac{2t}{s - 4m^2} = \frac{(k^\mu - k'^\mu)(p_\mu - p'_\mu)}{(k^\mu - k'^\mu)^2}. \quad (\text{G2})$$

The above defined collision invariants together with $u \equiv (k^\mu - p'^\mu)^2 = (k'^\mu - p^\mu)^2$ are the so-called Mandelstam variables, satisfying $s + t + u = 4m^2$, where m is the mass of the particles.

Therefore, the total momentum involved in a binary collision $P_T^\mu \equiv k^\mu + k'^\mu = p^\mu + p'^\mu$ is related to the total energy $P_T^\mu = (\sqrt{s}, 0, 0, 0)$, where $s \equiv 2(m^2 + k^\mu k'_\mu) = (2k^0)^2$, in the CM frame. The Lorentz-invariant transition rate is defined as

$$g^2 W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'} = (2\pi)^6 s \sigma(s, \theta_s) \delta^4(k^\mu + k'^\mu - p^\mu - p'^\mu). \quad (\text{G3})$$

The quantity $\sigma(s, \theta_{CM})$ is the differential cross section and $\delta^4(k^\mu + k'^\mu - p^\mu - p'^\mu)$ enforces energy and momentum conservation in binary collisions. Furthermore, we also define the total cross section $\sigma_T(s)$ as the integral of the differential cross section over the solid angle, $d\Omega = \int_0^{2\pi} d\varphi \int_0^\pi \sin \theta_s d\theta_s$,

$$\sigma_T(s) = \frac{1}{\nu} \int_0^\pi d\Omega \sigma(s, \theta_s). \quad (\text{G4})$$

For later purposes, let us define the rank- n tensor

$$\begin{aligned} \Theta^{\mu_1 \dots \mu_n} &\equiv \frac{1}{\nu} \int \frac{d^3 \mathbf{p}}{p^0} \frac{d^3 \mathbf{p}'}{p'^0} s \sigma(s, \theta_s) p^{\mu_1} \dots p^{\mu_n} \delta^4(k^\mu + k'^\mu - p^\mu - p'^\mu), \\ &= \sum_{q=0}^{[n/2]} (-1)^q b_{nq} \mathcal{B}_{nq} \Delta_{P_T}^{(\mu_1 \mu_2} \dots \Delta_{P_T}^{\mu_{2q-1} \mu_{2q}} P_T^{\mu_{2q+1}} \dots P_T^{\mu_n}), \end{aligned} \quad (\text{G5})$$

where we assumed that the differential cross section is isotropic, i.e., $\sigma(s, \theta_s) = \sigma(s)$. The b_{nq} coefficients are given in Eq. (A2) and

$$\begin{aligned} \mathcal{B}_{nq} &\equiv \frac{(-1)^q}{(2q+1)!!} \frac{1}{\nu} \int \frac{d^3 \mathbf{p}}{p^0} \frac{d^3 \mathbf{p}'}{p'^0} s^{-(n-2q)/2} \sigma(s, \theta_s) (P_T^\mu p_\mu)^{n-2q} \left(\Delta_{P_T}^{\alpha\beta} p_\alpha p_\beta \right)^q, \\ &= \frac{1}{(2q+1)!!} \frac{\sigma_T(s)}{2^{n+1}} s^{1/2} (s - 4m^2)^{(2q+1)/2}. \end{aligned} \quad (\text{G6})$$

Here we defined the projection orthogonal to the total momentum, $\Delta_P^{\alpha\beta} = g^{\alpha\beta} - P_T^\mu P_T^\nu / s$, hence the particle 4-momentum can be decomposed as $p^\mu = P_T^\mu (P_T^\alpha p_\alpha) / s + p_\alpha \Delta_{P_T}^{\mu\alpha}$ and thus $\Delta_{P_T}^{\alpha\beta} p_\alpha p_\beta = m^2 - s/4$. Similarly as in the previous integral we define a slightly different rank- n tensor $\Gamma_{(n,m)}^{\mu_1 \dots \mu_n - m \nu_{n-m+1} \dots \nu_n}$

$$\begin{aligned} \Gamma_{(n,m)}^{\mu_1 \dots \mu_n - m \nu_{n-m+1} \dots \nu_n} &\equiv \frac{1}{\nu} \int \frac{d^3 \mathbf{p}}{p^0} \frac{d^3 \mathbf{p}'}{p'^0} s \sigma(s, \theta_{CM}) p^{\mu_1} \dots p^{\mu_n - m} p'^{\nu_{n-m+1}} \dots p'^{\nu_n} \delta^4(k^\mu + k'^\mu - p^\mu - p'^\mu), \\ &= (-1)^{n-m} \Theta^{\mu_1 \dots \mu_n - m \nu_{n-m+1} \dots \nu_n} + (-1)^{n-m-1} P_T^{\nu_n} \Theta^{\mu_1 \dots \mu_n - m \nu_{n-m+1} \dots \nu_{n-1}} \\ &\quad + \dots + P_T^{\nu_{n-m+1}} \dots P_T^{\nu_n} \Theta^{\mu_1 \dots \mu_{n-m}}, \end{aligned} \quad (\text{G7})$$

where we replaced $p'^\nu = P_T^\nu - p^\nu$ to obtain the above result.

Appendix H: Coefficients in the massless limit

Using the results from Appendix G we can express the collision integrals from Eqs. (77) – (110) as a function of particle momenta and the metric tensor. As an example, here we discuss the $X_{(r)}^{\mu\nu\alpha\beta}$ tensor, which we separate into gain $\mathcal{G}_{(r)}^{\mu\nu\alpha\beta}$ and loss $\mathcal{L}_{(r)}^{\mu\nu\alpha\beta}$ parts as $X_{(r)}^{\mu\nu\alpha\beta} = \mathcal{G}_{(r)}^{\mu\nu\alpha\beta} - \mathcal{L}_{(r)}^{\mu\nu\alpha\beta}$, where

$$\begin{aligned}\mathcal{G}_{(r)}^{\mu\nu\alpha\beta} &\equiv \frac{1}{\nu} \int dK dK' dP dP' f_{0\mathbf{k}} f_{0\mathbf{k}'} W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'} E_{\mathbf{k}}^r k^\mu k^\nu (p^\alpha p^\beta + p'^\alpha p'^\beta), \\ &= 2 \int dK dK' f_{0\mathbf{k}} f_{0\mathbf{k}'} E_{\mathbf{k}}^r k^\mu k^\nu \Theta^{\alpha\beta},\end{aligned}\quad (\text{H1})$$

and

$$\begin{aligned}\mathcal{L}_{(r)}^{\mu\nu\alpha\beta} &\equiv \frac{1}{\nu} \int dK dK' dP dP' f_{0\mathbf{k}} f_{0\mathbf{k}'} W_{\mathbf{k}\mathbf{k}' \rightarrow \mathbf{p}\mathbf{p}'} E_{\mathbf{k}}^r k^\mu k^\nu (k^\alpha k^\beta + k'^\alpha k'^\beta), \\ &= \int dK dK' f_{0\mathbf{k}} f_{0\mathbf{k}'} E_{\mathbf{k}}^r k^\mu k^\nu (k^\alpha k^\beta + k'^\alpha k'^\beta) \Theta.\end{aligned}\quad (\text{H2})$$

Here $\Theta \equiv \frac{\sigma_T}{2} \sqrt{s(s-4m^2)} = \sigma_T \sqrt{(k^\mu k'_\mu)^2 - m^4}$ is the invariant flux and $\Theta^{\mu\nu} = \frac{\Theta}{4} [P_T^\mu P_T^\nu - \frac{1}{3}(s-4m^2) \Delta_{PT}^{\mu\nu}]$, cf. Eq. (G5). The invariant flux can be expressed using the relative velocity between particles, $\Theta = v_{rel} k^\mu k'_\mu$ where $v_{rel} = (1 - \mathbf{v} \cdot \mathbf{v}')^{-1} \sqrt{(\mathbf{v} - \mathbf{v}')^2 - (\mathbf{v} \times \mathbf{v}')^2}$ and $\mathbf{v} = \mathbf{k}/k^0$, $\mathbf{v}' = \mathbf{k}'/k'^0$. Hence the total cross section is $\sigma_T = v_{rel} \sigma_T(s)$. However, in the massless limit, $m \rightarrow 0$, the relative velocity is $v_{rel} = 1$.

From now on we will work in the massless limit $m \rightarrow 0$. For example the solution for $\mathcal{G}_{(r)}^{\mu\nu\alpha\beta}$ can be written formally as $\mathcal{G}_{(r)}^{\mu\nu\alpha\beta} = \int (A_X - B_X/s) P_T^\mu P_T^\nu + B_X g^{\mu\nu}$. After finding the coefficients A_X and B_X we replace $P_T^\mu = k^\mu + k'^\mu$. Thus using the above definitions and relations we obtain

$$\mathcal{G}_{(r)}^{\mu\nu\alpha\beta} = \frac{2\sigma_T}{3} I_{(r+5)}^{\mu\nu\alpha\beta\kappa} I_{(1)\kappa} + \frac{4\sigma_T}{3} I_{(r+4)}^{\mu\nu\kappa(\alpha} I_{(2)\kappa}^{\beta)} + \frac{2\sigma_T}{3} I_{(r+3)}^{\mu\nu\kappa} I_{(3)\kappa}^{\alpha\beta} - \frac{\sigma_T}{3} g^{\alpha\beta} I_{(r+4)}^{\mu\nu\kappa\lambda} I_{(2)\kappa\lambda}, \quad (\text{H3})$$

$$\mathcal{L}_{(r)}^{\mu\nu\alpha\beta} = \sigma_T I_{(r+5)}^{\mu\nu\alpha\beta\kappa} I_{(1)\kappa} + \sigma_T I_{(r+3)}^{\mu\nu\kappa} I_{(3)\kappa}^{\alpha\beta}, \quad (\text{H4})$$

and hence,

$$X_{(r)}^{\mu\nu\alpha\beta} = -\frac{\sigma_T}{3} I_{(r+5)}^{\mu\nu\alpha\beta\kappa} I_{(1)\kappa} + \frac{4\sigma_T}{3} I_{(r+4)}^{\mu\nu\kappa(\alpha} I_{(2)\kappa}^{\beta)} - \frac{\sigma_T}{3} I_{(r+3)}^{\mu\nu\kappa} I_{(3)\kappa}^{\alpha\beta} - \frac{\sigma_T}{3} g^{\alpha\beta} I_{(r+4)}^{\mu\nu\kappa\lambda} I_{(2)\kappa\lambda}, \quad (\text{H5})$$

where the $I_{(r+n)}^{\mu_1 \dots \mu_n}$ tensors were defined in Eq. (A1).

Similarly, we obtain for the next rank-4 collision tensor

$$Y_{1(r)}^{\mu\nu\alpha\beta} = -X_{(r)}^{\mu\nu\alpha\beta}. \quad (\text{H6})$$

The last rank-4 collision tensor we need is

$$Y_{2(r)}^{\mu\nu\alpha\beta} = \frac{\sigma_T}{6} I_{(r+5)}^{\mu\nu\alpha\beta\kappa} I_{(1)\kappa} + \frac{\sigma_T}{3} I_{(r+4)}^{\mu\nu\kappa(\alpha} I_{(2)\kappa}^{\beta)} + \frac{\sigma_T}{6} I_{(r+3)}^{\mu\nu\kappa} I_{(3)\kappa}^{\alpha\beta} - \sigma_T I_{(r+4)}^{\mu\nu\kappa\alpha} I_{(2)\kappa}^\beta + \frac{\sigma_T}{6} g^{\alpha\beta} I_{(r+4)}^{\mu\nu\kappa\lambda} I_{(2)\kappa\lambda}. \quad (\text{H7})$$

The only rank-5 tensor we have is

$$\begin{aligned}Y_{3(r)}^{(\mu\nu)\alpha(\beta\kappa)} &= \frac{\sigma_T}{6} \left[I_{(r+6)}^{\mu\nu\alpha\beta\kappa\lambda} I_{(1)\lambda} + 2 I_{(r+5)}^{\mu\nu\lambda\alpha(\beta} I_{(2)\lambda}^{\kappa)} + I_{(r+5)}^{\mu\nu\lambda\kappa\beta} I_{(2)\lambda}^\alpha \right] + \frac{\sigma_T}{6} \left[2 I_{(r+4)}^{\mu\nu\lambda(\beta} I_{(3)\lambda}^{\kappa)\alpha} + I_{(r+4)}^{\mu\nu\lambda\alpha} I_{(3)\lambda}^{\beta\kappa} + I_{(r+3)}^{\mu\nu\lambda} I_{(4)\lambda}^{\alpha\beta\kappa} \right] \\ &+ \frac{\sigma_T}{6} 2 g^{\alpha(\beta} \left[I_{(r+5)}^{\mu\nu\lambda\sigma} I_{(2)\lambda\sigma} + I_{(3)\lambda\sigma} I_{(r+4)}^{\mu\nu\lambda\sigma} \right] - \frac{\sigma_T}{6} g^{\beta\kappa} \left[I_{(r+5)}^{\alpha\mu\nu\lambda\sigma} I_{(2)\lambda\sigma} + I_{(3)\lambda\sigma} I_{(r+4)}^{\alpha\mu\nu\lambda\sigma} \right] \\ &- \sigma_T \left[I_{(r+4)}^{\mu\nu\alpha\lambda} I_{(3)\lambda}^{\beta\kappa} + I_{(r+5)}^{\mu\nu\beta\kappa\lambda} I_{(2)\lambda}^\alpha \right],\end{aligned}\quad (\text{H8})$$

where the solution $Y_{3(r)}^{(\mu\nu)\alpha(\beta\kappa)} \sim \int (A_3 - 2B_3/s - C_3/s) P_T^\alpha P_T^\beta P_T^\kappa + 2B_3 g^{\alpha(\beta} P_T^{\kappa)} + C_3 g^{\beta\kappa} P_T^\alpha$.

The rank-6 tensors are

$$\begin{aligned}
Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} = & \frac{\sigma_T}{15} \left[I_{(r+7)}^{\mu\nu\alpha\beta\kappa\lambda\sigma} I_{(1)\sigma} + 4I_{(r+6)}^{\mu\nu\sigma(\alpha\beta\kappa} I_{(2)\sigma}^{\lambda)} + 6I_{(r+5)}^{\mu\nu\sigma(\alpha\beta} I_{(3)\sigma}^{\kappa\lambda)} + 4I_{(r+4)}^{\mu\nu\sigma(\alpha} I_{(4)\sigma}^{\beta\kappa\lambda)} + I_{(r+3)}^{\mu\nu\sigma} I_{(5)\sigma}^{\alpha\beta\kappa\lambda} \right] \\
& + \frac{\sigma_T}{15} 2g^{\beta(\kappa} \left[I_{(r+6)}^{\lambda)\alpha\mu\nu\sigma\rho} I_{(2)\sigma\rho} + I_{(3)\sigma\rho}^{\lambda)} I_{(r+5)}^{\alpha\mu\nu\sigma\rho} + I_{(r+5)}^{\lambda)\mu\nu\sigma\rho} I_{(3)\sigma\rho}^{\alpha} + I_{(4)\sigma\rho}^{\lambda)\alpha} I_{(r+4)}^{\mu\nu\sigma\rho} \right] \\
& + \frac{\sigma_T}{15} 2g^{\alpha(\kappa} \left[I_{(r+6)}^{\lambda)\beta\mu\nu\sigma\rho} I_{(2)\sigma\rho} + I_{(3)\sigma\rho}^{\lambda)} I_{(r+5)}^{\beta\mu\nu\sigma\rho} + I_{(r+5)}^{\lambda)\mu\nu\sigma\rho} I_{(3)\sigma\rho}^{\beta} + I_{(4)\sigma\rho}^{\lambda)\beta} I_{(r+4)}^{\mu\nu\sigma\rho} \right] \\
& + \frac{\sigma_T}{30} \left[2g^{\alpha(\kappa} g^{\lambda)\beta} + g^{\alpha\beta} g^{\kappa\lambda} \right] I_{(r+5)}^{\mu\nu\sigma\rho\tau} I_{(3)\sigma\rho\tau} \\
& - \frac{\sigma_T}{10} g^{\alpha\beta} \left[I_{(r+6)}^{\kappa\lambda\mu\nu\sigma\rho} I_{(2)\sigma\rho} + I_{(3)\sigma\rho}^{\lambda)} I_{(r+5)}^{\kappa\mu\nu\sigma\rho} + I_{(3)\sigma\rho}^{\kappa} I_{(r+5)}^{\lambda\mu\nu\sigma\rho} + I_{(4)\sigma\rho}^{\kappa\lambda} I_{(r+4)}^{\mu\nu\sigma\rho} \right] \\
& - \frac{\sigma_T}{10} g^{\kappa\lambda} \left[I_{(r+6)}^{\alpha\beta\mu\nu\sigma\rho} I_{(2)\sigma\rho} + I_{(3)\sigma\rho}^{\beta)} I_{(r+5)}^{\alpha\mu\nu\sigma\rho} + I_{(3)\sigma\rho}^{\alpha} I_{(r+5)}^{\beta\mu\nu\sigma\rho} + I_{(4)\sigma\rho}^{\alpha\beta} I_{(r+4)}^{\mu\nu\sigma\rho} \right] \\
& - \sigma_T I_{(r+5)}^{\mu\nu\alpha\beta\sigma} I_{(3)\sigma}^{\kappa\lambda} - \sigma_T I_{(r+5)}^{\mu\nu\kappa\lambda\sigma} I_{(3)\sigma}^{\alpha\beta},
\end{aligned} \tag{H9}$$

and

$$Y_{5(r)}^{\mu\nu\alpha\beta\kappa\lambda} = \frac{1}{2} \left[Y_{4(r)}^{\mu\nu\alpha\beta\kappa\lambda} - \sigma_T I_{(r+5)}^{\mu\nu\alpha\beta\sigma} I_{(3)\sigma}^{\kappa\lambda} + \sigma_T I_{(r+5)}^{\mu\nu\kappa\lambda\sigma} I_{(3)\sigma}^{\alpha\beta} \right]. \tag{H10}$$

Here the solution has the form $Y^{\mu\nu\alpha\beta\kappa\lambda} \sim \int A_Y P_T^\alpha P_T^\beta P_T^\kappa P_T^\lambda + 2B_Y \left(P_T^\alpha g^{\beta(\kappa} P^\lambda) + P_T^\beta g^{\alpha(\kappa} P^\lambda) \right) + 2C_Y g^{\alpha(\kappa} g^{\lambda)\beta} + D_Y g^{\alpha\beta} P_T^\lambda P_T^\kappa + E_Y g^{\kappa\lambda} P_T^\alpha P_T^\beta + F_Y g^{\alpha\beta} g^{\kappa\lambda}$.

In order to calculate the above collision tensors, we need the following $I_{(r+n)}^{\mu_1 \dots \mu_n}$ tensors from Eq. (A1),

$$I_{(r+1)}^\mu = I_{r+1,0} u^\mu, \tag{H11}$$

$$I_{(r+2)}^{\mu\nu} = I_{r+2,0} u^\mu u^\nu - I_{r+2,1} \Delta^{\mu\nu}, \tag{H12}$$

$$I_{(r+3)}^{\mu\nu\alpha} = I_{r+3,0} u^\mu u^\nu u^\alpha - 3I_{r+3,1} u^{(\mu} \Delta^{\nu\alpha)}, \tag{H13}$$

$$I_{(r+4)}^{\mu\nu\alpha\beta} = I_{r+4,0} u^\mu u^\nu u^\alpha u^\beta - 6I_{r+4,1} u^{(\mu} u^\nu \Delta^{\alpha\beta)} + 3I_{r+4,2} \Delta^{(\mu\nu} \Delta^{\alpha\beta)}, \tag{H14}$$

$$I_{(r+5)}^{\mu\nu\alpha\beta\kappa} = I_{r+5,0} u^\mu u^\nu u^\alpha u^\beta u^\kappa - 10I_{r+5,1} u^{(\mu} u^\nu u^\alpha \Delta^{\beta\kappa)} + 15I_{r+5,2} u^{(\mu} \Delta^{\nu\alpha} \Delta^{\beta\kappa)}, \tag{H15}$$

$$\begin{aligned}
I_{(r+6)}^{\mu\nu\alpha\beta\kappa\lambda} = & I_{r+6,0} u^\mu u^\nu u^\alpha u^\beta u^\kappa u^\lambda - 15I_{r+6,1} u^{(\mu} u^\nu u^\alpha u^\beta \Delta^{\kappa\lambda)} \\
& + 45I_{r+6,2} u^{(\mu} u^\nu \Delta^{\alpha\beta} \Delta^{\kappa\lambda)} - 15I_{r+6,3} \Delta^{(\mu\nu} \Delta^{\alpha\beta} \Delta^{\kappa\lambda)},
\end{aligned} \tag{H16}$$

$$\begin{aligned}
I_{(r+7)}^{\mu\nu\alpha\beta\kappa\lambda\tau} = & I_{r+7,0} u^\mu u^\nu u^\alpha u^\beta u^\kappa u^\lambda u^\tau - 21I_{r+7,1} u^{(\mu} u^\nu u^\alpha u^\beta u^\kappa \Delta^{\lambda\tau)} \\
& + 105I_{r+7,2} u^{(\mu} u^\nu u^\alpha \Delta^{\beta\kappa} \Delta^{\lambda\tau)} - 105I_{r+7,3} \Delta^{(\mu\nu} \Delta^{\alpha\beta} \Delta^{\kappa\lambda} u^\tau).
\end{aligned} \tag{H17}$$

In the massless Boltzmann limit, we use the following formula for the thermodynamic integrals,

$$I_{n+r,q}(\alpha_0, \beta_0) \underset{m \rightarrow 0}{=} \frac{P_0 (r+n+1)!}{2\beta_0^{r+n-2} (2q+1)!!}, \tag{H18}$$

where $P_0 = ge^{\alpha_0} \beta_0^{-4}/\pi^2$, and we extensively use the recursion relation (A5) in the massless limit, i.e., $I_{r+n,q} = (2q+3) I_{r+n,q+1}$.

The coefficients for the linear part of the collision integral are

$$X_{(r),1} \equiv x_{10} = -\frac{\sigma_T}{3} [I_{r+5,0} I_{1,0} + I_{r+3,0} I_{3,0} - 24I_{r+4,1} I_{2,1}] = -\frac{\sigma_T P_0^2 (r+4)!}{6\beta_0^{r+2}} (r^2 + 3r + 2), \tag{H19}$$

$$X_{(r),3} \equiv x_{31} = \frac{\sigma_T}{3} [I_{r+5,1} I_{1,0} - I_{r+3,1} I_{3,0} - 4I_{r+4,1} I_{2,1}] = \frac{\sigma_T P_0^2 (r+4)!}{18\beta_0^{r+2}} (r^2 + 7r + 6), \tag{H20}$$

$$X_{(r),4} \equiv 2x_{22} = -\frac{2\sigma_T}{3} [I_{r+5,2} I_{1,0} + 4I_{r+4,2} I_{2,1}] = -\frac{\sigma_T P_0^2 (r+5)!}{45\beta_0^{r+2}} (r+10). \tag{H21}$$

The other coefficients resulting from the rank-4 tensor are

$$Y_{1(r),3} = -X_{(r),3}, \quad (\text{H22})$$

$$Y_{2(r),1} = -\frac{1}{2}X_{(r),1}, \quad (\text{H23})$$

$$Y_{2(r),4} = -\frac{1}{2}X_{(r),4}. \quad (\text{H24})$$

The coefficients of the rank-5 tensor are

$$\begin{aligned} Y_{3(r),1} &\equiv y_{10} = \frac{\sigma_T}{6} [I_{r+6,0}I_{1,0} - 15I_{r+5,1}I_{2,1} - 15I_{r+4,1}I_{3,1} + I_{r+3,0}I_{4,0}] \\ &= \frac{\sigma_T P_0^2 (r+4)!}{12\beta_0^{r+3}} (r+1)(r+2)(r+10), \end{aligned} \quad (\text{H25})$$

$$\begin{aligned} Y_{3(r),3} &\equiv y_{31} = -\frac{\sigma_T}{6} [I_{r+6,1}I_{1,0} - 9I_{r+5,1}I_{2,1} + 5I_{r+4,1}I_{3,1} - I_{r+3,1}I_{4,1}] \\ &= -\frac{\sigma_T P_0^2 (r+4)!}{36\beta_0^{r+3}} (r+1)(r^2 + 8r + 20), \end{aligned} \quad (\text{H26})$$

$$\begin{aligned} Y_{3(r),4} &\equiv 2y_{22} = \frac{\sigma_T}{3} [I_{r+6,2}I_{1,0} - 17I_{r+5,2}I_{2,1} - 2I_{r+4,2}I_{3,1}] \\ &= \frac{\sigma_T P_0^2 (r+5)!}{90\beta_0^{r+3}} (r^2 - 4r - 68), \end{aligned} \quad (\text{H27})$$

$$Y_{3(r),5} \equiv y_{41} = -\frac{1}{3}Y_{3(r),1}, \quad (\text{H28})$$

$$\begin{aligned} Y_{3(r),6} &\equiv y_{42} = \frac{\sigma_T}{30} [I_{r+6,1}I_{1,0} + 23I_{r+5,1}I_{2,1} - 35I_{r+4,1}I_{3,1} - I_{r+3,1}I_{4,1}] \\ &= \frac{\sigma_T P_0^2 (r+4)!}{180\beta_0^{r+3}} (r+1)(r^2 + 40r + 180), \end{aligned} \quad (\text{H29})$$

$$\begin{aligned} Y_{3(r),7} &\equiv y_{52} = \frac{\sigma_T}{30} [I_{r+6,1}I_{1,0} + 7I_{r+5,1}I_{2,1} + 4I_{r+4,1}I_{3,1}] \\ &= \frac{\sigma_T P_0^2 (r+5)!}{180\beta_0^{r+3}} (r+10)^2. \end{aligned} \quad (\text{H30})$$

$$\begin{aligned} Y_{3(r),8} &\equiv y_{62} = \frac{\sigma_T}{30} [I_{r+6,1}I_{1,0} - 17I_{r+5,1}I_{2,1} + 15I_{r+4,1}I_{3,1} - I_{r+3,1}I_{4,1}] \\ &= \frac{\sigma_T P_0^2 (r+4)!}{180\beta_0^{r+3}} (r+1)(r^2 - 20). \end{aligned} \quad (\text{H31})$$

The coefficients from the rank-6 tensors are

$$\begin{aligned} Y_{4(r),3} &\equiv w_{31} = -\frac{\sigma_T}{15} [I_{r+7,1}I_{1,0} + 10I_{r+6,1}I_{2,1} - 23I_{r+5,1}I_{3,1} - I_{r+4,1}I_{4,1} - I_{r+3,1}I_{5,1}] \\ &= -\frac{\sigma_T P_0^2 (r+4)!}{90\beta_0^{r+4}} (r+1)(r^3 + 35r^2 + 304r + 800), \end{aligned} \quad (\text{H32})$$

$$\begin{aligned} Y_{4(r),4} &\equiv 2w_{22} = \frac{\sigma_T}{75} [10I_{r+7,2}I_{1,0} - 20I_{r+6,2}I_{2,1} - 400I_{r+5,2}I_{3,1} - 26I_{r+4,2}I_{4,1}] \\ &= \frac{\sigma_T P_0^2 (r+5)!}{225\beta_0^{r+4}} (r+19)(r^2 - 40), \end{aligned} \quad (\text{H33})$$

$$\begin{aligned} Y_{4(r),8} &\equiv w_{62} = \frac{\sigma_T}{150} [10I_{r+7,2}I_{1,0} - 40I_{r+6,1}I_{2,1} + 14I_{r+5,1}I_{3,1} + 28I_{r+4,1}I_{4,1} - 2I_{r+3,1}I_{5,1}] \\ &= \frac{\sigma_T P_0^2 (r+4)!}{450\beta_0^{r+4}} (r+1)(r^3 + 5r^2 - 86r - 400), \end{aligned} \quad (\text{H34})$$

and

$$\begin{aligned} Y_{5(r),1} &\equiv z_{10} = \frac{\sigma_T}{30} [I_{r+7,0}I_{1,0} + 48I_{r+6,1}I_{2,1} - 165I_{r+5,1}I_{3,1} + 48I_{r+4,1}I_{4,1} + 9I_{r+3,1}I_{5,1}] \\ &= \frac{\sigma_T P_0^2 (r+4)!}{60\beta_0^{r+4}} (r+1)(r+2)(r^2 + 39r + 200), \end{aligned} \quad (\text{H35})$$

$$\begin{aligned} Y_{5(r),5} &\equiv z_{41} = -\frac{\sigma_T}{30} [I_{r+7,1}I_{1,0} - 4I_{r+6,1}I_{2,1} - 5I_{r+5,1}I_{3,1} - 4I_{r+4,1}I_{4,1} + 3I_{r+3,1}I_{5,1}] \\ &= -\frac{\sigma_T P_0^2 (r+4)!}{180\beta_0^{r+4}} (r+1)(r+2)(r^2 + 19r + 100), \end{aligned} \quad (\text{H36})$$

$$\begin{aligned} Y_{5(r),7} &\equiv z_{52} = \frac{\sigma_T}{150} [5I_{r+7,2}I_{1,0} + 40I_{r+6,2}I_{2,1} + 125I_{r+5,2}I_{3,1} - 8I_{r+4,2}I_{4,1}] \\ &= \frac{\sigma_T P_0^2 (r+5)!}{900\beta_0^{r+4}} (r^3 + 29r^2 + 350r + 1240), \end{aligned} \quad (\text{H37})$$

$$\begin{aligned} Y_{5(r),9} &\equiv z_{72} = \frac{\sigma_T}{150} [5I_{r+7,2}I_{1,0} - 44I_{r+6,1}I_{2,1} + 95I_{r+5,1}I_{3,1} - 44I_{r+4,1}I_{4,1} + 3I_{r+3,1}I_{5,1}] \\ &= \frac{\sigma_T P_0^2 (r+4)!}{900\beta_0^{r+4}} (r^4 - 18r^3 - 161r^2 - 342r - 200), \end{aligned} \quad (\text{H38})$$

$$\begin{aligned} Y_{5(r),11} &\equiv z_{23} = -\frac{\sigma_T}{210} [I_{r+7,2}I_{1,0} - 32I_{r+6,2}I_{2,1} + 35I_{r+5,2}I_{3,1} - 28I_{r+4,2}I_{4,2}] \\ &= -\frac{\sigma_T P_0^2 (r+5)!}{6300\beta_0^{r+4}} (r^3 - 11r^2 - 130r - 280). \end{aligned} \quad (\text{H39})$$

We remark that to crosscheck some of our calculations found in this appendix we made use of the symbolic computer algebra software Cadabra by K. Peeters [63, 64], as well as of the Mathematica package FeynCalc developed and maintained by Rolf Mertig and Frederik Orellana [65].

Appendix I: Coefficients of the \mathcal{K} tensors

In this appendix, we list the coefficients of the tensors \mathcal{K} , \mathcal{K}^μ , and $\mathcal{K}^{\mu\nu}$ in Eqs. (5), in the notation of Ref. [24]. They can be derived following the derivation of the equations of motion for the dissipative quantities presented in that reference. For the sake of brevity, they were not explicitly given in Ref. [24], so we decided to list them here for further reference. Note, however, that in the 14-moment approximation, where $N_0 = 2$, $N_1 = 1$, and $N_2 = 0$, they vanish identically.

$$\tilde{\zeta}_1 = \sum_{r=3}^{N_0} \tau_{0r}^{(0)} \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right), \quad (\text{I1})$$

$$\tilde{\zeta}_2 = -\frac{2m^2}{3} \sum_{r=1}^{N_0-2} \tau_{0,r+2}^{(0)} (r+1) \left(\eta_r - \Omega_{r0}^{(2)} \eta \right) - \tilde{\zeta}_1, \quad (\text{I2})$$

$$\tilde{\zeta}_3 = \sum_{r=3}^{N_0} \tau_{0r}^{(0)} \bar{F}_r(\alpha_0, \beta_0) + \frac{1}{3} \sum_{r=3}^{N_0} \tau_{0r}^{(0)} (r+1) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) - \frac{m^2}{3} \sum_{r=3}^{N_0-2} \tau_{0,r+2}^{(0)} (r+1) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right), \quad (\text{I3})$$

$$\tilde{\zeta}_4 = \frac{m^2}{3} \sum_{r=2}^{N_0-1} \tau_{0,r+1}^{(0)} \left(\frac{\partial}{\partial \alpha_0} + \frac{n_0}{\varepsilon_0 + P_0} \frac{\partial}{\partial \beta_0} \right) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right), \quad (\text{I4})$$

$$\tilde{\zeta}_5 = -\frac{2(\varepsilon_0 + P_0) + \beta_0 J_{30}}{(\varepsilon_0 + P_0)^3} \tilde{\zeta}_1, \quad (\text{I5})$$

$$\tilde{\zeta}_6 = -\frac{(\varepsilon_0 + P_0) J_{20} - n_0 J_{30}}{(\varepsilon_0 + P_0)^3} \tilde{\zeta}_1 - \frac{m^2}{3} \frac{1}{\varepsilon_0 + P_0} \sum_{r=2}^{N_0-1} \tau_{0,r+1}^{(0)} \left[(r+1) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) + \frac{\partial \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right)}{\partial \ln \beta_0} \right], \quad (\text{I6})$$

$$\tilde{\zeta}_7 = \frac{m^2}{3} \sum_{r=2}^{N_0-1} \tau_{0,r+1}^{(0)} \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right), \quad (\text{I7})$$

$$\tilde{\zeta}_8 = \frac{\tilde{\zeta}_1}{\varepsilon_0 + P_0}. \quad (\text{I8})$$

where $\zeta = \zeta^0$, $\kappa = \kappa^0$, $\eta = \eta^0$ in consistency with Eqs. (1) – (3).

In Eq. (I3), we defined the function $\bar{F}_r(\alpha_0, \beta_0)$ through the relation

$$D \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) = \bar{F}_r(\alpha_0, \beta_0) \theta. \quad (\text{I9})$$

Furthermore,

$$\begin{aligned} \tilde{\kappa}_1 = & -2 \sum_{r=1}^{N_1-1} \tau_{0,r+1}^{(1)} \left(\frac{\partial}{\partial \alpha_0} + \frac{n_0}{\varepsilon_0 + P_0} \frac{\partial}{\partial \beta_0} \right) \left(\eta_r - \Omega_{r0}^{(2)} \eta \right) \\ & + \frac{2m^2}{5} \sum_{r=2}^{N_1-2} \tau_{0,r+2}^{(1)} (r+1) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) - \frac{2}{5} \sum_{r=2}^{N_1} \tau_{0r}^{(1)} (r-1) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right), \end{aligned} \quad (\text{I10})$$

$$\tilde{\kappa}_2 = \frac{2}{\varepsilon_0 + P_0} \sum_{r=1}^{N_1-1} \tau_{0,r+1}^{(1)} \left[(r+1) \left(\eta_r - \Omega_{r0}^{(2)} \eta \right) + \frac{\partial \left(\eta_r - \Omega_{r0}^{(2)} \eta \right)}{\partial \ln \beta_0} \right], \quad (\text{I11})$$

$$\begin{aligned} \tilde{\kappa}_3 = & -\sum_{r=2}^{N_1} \tau_{0r}^{(1)} \left[\bar{G}_r(\alpha_0, \beta_0) + \left(\frac{r+2}{3} + \frac{\partial \mathcal{H}(\alpha_0, \beta_0)}{\partial \alpha_0} + \frac{n_0}{\varepsilon_0 + P_0} \frac{\partial \mathcal{H}(\alpha_0, \beta_0)}{\partial \beta_0} \right) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) \right] \\ & + \frac{m^2}{3} \sum_{r=2}^{N_1-2} \tau_{0,r+2}^{(1)} (r+1) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) - \sum_{r=3}^{N_1-1} \tau_{0,r+1}^{(1)} \left(\frac{\partial}{\partial \alpha_0} + \frac{n_0}{\varepsilon_0 + P_0} \frac{\partial}{\partial \beta_0} \right) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) \\ & + \frac{1}{m^2} \sum_{r=3}^{N_1+1} \tau_{0,r-1}^{(1)} \left(\frac{\partial}{\partial \alpha_0} + \frac{n_0}{\varepsilon_0 + P_0} \frac{\partial}{\partial \beta_0} \right) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right), \end{aligned} \quad (\text{I12})$$

$$\begin{aligned} \tilde{\kappa}_4 = & \frac{1}{\varepsilon_0 + P_0} \left\{ \left[\mathcal{H}(\alpha_0, \beta_0) + \frac{\partial \mathcal{H}(\alpha_0, \beta_0)}{\partial \ln \beta_0} \right] \sum_{r=2}^{N_1} \tau_{0r}^{(1)} \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) + \sum_{r=3}^{N_1-1} \tau_{0,r+1}^{(1)} \left(r+1 + \frac{\partial}{\partial \ln \beta_0} \right) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) \right. \\ & \left. - \frac{1}{m^2} \sum_{r=3}^{N_1+1} \tau_{0,r-1}^{(1)} \left(r+2 + \frac{\partial}{\partial \ln \beta_0} \right) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) \right\}, \end{aligned} \quad (\text{I13})$$

$$\tilde{\kappa}_5 = 2 \sum_{r=2}^{N_1} \tau_{0r}^{(1)} \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right), \quad (\text{I14})$$

$$\tilde{\kappa}_6 = -2 \sum_{r=1}^{N_1-1} \tau_{0,r+1}^{(1)} \left(\eta_r - \Omega_{r0}^{(2)} \eta \right), \quad (\text{I15})$$

$$\tilde{\kappa}_7 = -\mathcal{H}(\alpha_0, \beta_0) \frac{\tilde{\kappa}_5}{2} - \sum_{r=3}^{N_1-1} \tau_{0,r+1}^{(1)} \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) + \frac{1}{m^2} \sum_{r=3}^{N_1+1} \tau_{0,r-1}^{(1)} \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right). \quad (\text{I16})$$

Here, we defined the function $\bar{G}_r(\alpha_0, \beta_0)$ through the relation

$$D\left(\kappa_r - \Omega_{r0}^{(1)}\kappa\right) = \bar{G}_r(\alpha_0, \beta_0)\theta. \quad (\text{I17})$$

Moreover,

$$\mathcal{H}(\alpha_0, \beta_0) = \frac{(\varepsilon_0 + P_0)J_{20} - n_0J_{30}}{D_{20}}. \quad (\text{I18})$$

Finally,

$$\tilde{\eta}_1 = 2 \sum_{r=1}^{N_2} \tau_{0r}^{(2)} \left(\eta_r - \Omega_{r0}^{(2)} \eta \right), \quad (\text{I19})$$

$$\begin{aligned} \tilde{\eta}_2 = & 2 \left\{ - \sum_{r=1}^{N_2} \tau_{0r}^{(2)} \left[\bar{H}_r(\alpha_0, \beta_0) + \frac{1}{3}(r+2) \left(\eta_r - \Omega_{r0}^{(2)} \eta \right) \right] \right. \\ & + \frac{m^2}{3} \sum_{r=1}^{N_2-2} \tau_{0,r+2}^{(2)}(r+1) \left(\eta_r - \Omega_{r0}^{(2)} \eta \right) + \frac{m^2}{5} \sum_{r=3}^{N_2-2} \tau_{0,r+2}^{(2)}(r+1) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) \\ & \left. - \frac{1}{5} \sum_{r=3}^{N_2} \tau_{0r}^{(2)}(2r+3) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) + \frac{1}{5m^2} \sum_{r=3}^{N_2+2} \tau_{0,r-2}^{(2)}(r+2) \left(\zeta_r - \Omega_{r0}^{(0)} \zeta \right) \right\}, \end{aligned} \quad (\text{I20})$$

$$\tilde{\eta}_3 = \frac{2}{7} \left[- \sum_{r=1}^{N_2} \tau_{0r}^{(2)}(4r+3) \left(\eta_r - \Omega_{r0}^{(2)} \eta \right) + 4m^2 \sum_{r=1}^{N_2-2} \tau_{0,r+2}^{(2)}(r+1) \left(\eta_r - \Omega_{r0}^{(2)} \eta \right) \right], \quad (\text{I21})$$

$$\tilde{\eta}_4 = 2\tilde{\eta}_1, \quad (\text{I22})$$

$$\tilde{\eta}_5 = \frac{2}{5} \left[\sum_{r=2}^{N_2+1} \tau_{0,r-1}^{(2)} \left(\frac{\partial}{\partial \alpha_0} + \frac{n_0}{\varepsilon_0 + P_0} \frac{\partial}{\partial \beta_0} \right) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) - m^2 \sum_{r=2}^{N_2-1} \tau_{0,r+1}^{(2)} \left(\frac{\partial}{\partial \alpha_0} + \frac{n_0}{\varepsilon_0 + P_0} \frac{\partial}{\partial \beta_0} \right) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) \right], \quad (\text{I23})$$

$$\tilde{\eta}_6 = \frac{2(\varepsilon_0 + P_0) + \beta_0 J_{30}}{(\varepsilon_0 + P_0)^3} \tilde{\eta}_1, \quad (\text{I24})$$

$$\begin{aligned} \tilde{\eta}_7 = & \frac{2}{5(\varepsilon_0 + P_0)} \left\{ \frac{5(\varepsilon_0 + P_0)J_{20} - n_0J_{30}}{(\varepsilon_0 + P_0)^2} \tilde{\eta}_1 - \sum_{r=2}^{N_2+1} \tau_{0,r-1}^{(2)} \left(r + 4 + \frac{\partial}{\partial \ln \beta_0} \right) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) \right. \\ & \left. + m^2 \sum_{r=2}^{N_2-1} \tau_{0,r+1}^{(2)} \left(r + 1 + \frac{\partial}{\partial \ln \beta_0} \right) \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) \right\}, \end{aligned} \quad (\text{I25})$$

$$\tilde{\eta}_8 = \frac{2}{5} \left[\sum_{r=2}^{N_2+1} \tau_{0,r-1}^{(2)} \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) - m^2 \sum_{r=2}^{N_2-1} \tau_{0,r+1}^{(2)} \left(\kappa_r - \Omega_{r0}^{(1)} \kappa \right) \right], \quad (\text{I26})$$

$$\tilde{\eta}_9 = -\frac{\tilde{\eta}_1}{\varepsilon_0 + P_0}. \quad (\text{I27})$$

Here, we defined the function $\bar{H}_r(\alpha_0, \beta_0)$ through the relation

$$D\left(\eta_r - \Omega_{r0}^{(2)}\eta\right) = \bar{H}_r(\alpha_0, \beta_0)\theta. \quad (\text{I28})$$

-
- [1] U. W. Heinz and R. Snellings, Annu. Rev. Nucl. Part. Sci. **63**, 123 (2013) [arXiv:1301.2826 [nucl-th]].
[2] C. Gale, S. Jeon and B. Schenke, Int. J. Mod. Phys. A **28**, 1340011 (2013) [arXiv:1301.5893 [nucl-th]].
[3] L. Baiotti, B. Giacomazzo and L. Rezzolla, Phys. Rev. D **78**, 084033 (2008) [arXiv:0804.0594 [gr-qc]].
[4] N. Andersson and G. L. Comer, Living Rev. Relativity **10**, 1 (2007); <http://www.livingreviews.org/lrr-2007-1>.
[5] R. Maartens, [astro-ph/9609119].

- [6] R. Maartens, *Class. Quant. Grav.* **12**, 1455 (1995).
- [7] W. Zimdahl, D. Pavon and R. Maartens, *Phys. Rev. D* **55**, 4681 (1997) [astro-ph/9611147].
- [8] X. Chen and E. A. Spiegel, *Class. Quant. Grav.* **28**, 175003 (2011) [arXiv:1107.0319 [gr-qc]].
- [9] P. Danielewicz and M. Gyulassy, *Phys. Rev. D* **31**, 53 (1985).
- [10] P. Kovtun, D. T. Son, A. O. Starinets, *Phys. Rev. Lett.* **94**, 111601 (2005) [hep-th/0405231].
- [11] L. P. Csernai, J. I. Kapusta, L. D. McLerran, *Phys. Rev. Lett.* **97**, 152303 (2006) [nucl-th/0604032].
- [12] C. Eckart, *Phys. Rev.* **58**, 919 (1940).
- [13] L. D. Landau and E. M. Lifshitz, *Fluid Dynamics*, Second Edition, Butterworth-Heinemann (1987).
- [14] W. A. Hiscock and L. Lindblom, *Ann. Phys. (N.Y.)* **151**, 466 (1983); *Phys. Rev. D* **31**, 725 (1985); *Phys. Rev. D* **35**, 3723 (1987); *Phys. Lett. A* **131**, 509 (1988); *Phys. Lett. A* **131**, 509 (1988).
- [15] S. Pu, T. Koide and D. H. Rischke, *Phys. Rev. D* **81**, 114039 (2010) [arXiv:0907.3906 [hep-ph]].
- [16] H. Grad, *Comm. Pure Appl. Math.* **2**, 331 (1949).
- [17] I. Müller, *Z. Physik* **198**, 329 (1967).
- [18] I-Shih Liu, I. Müller and T. Ruggeri, *Annals Phys.* **169**, 191 (1986).
- [19] I. Müller, *Living Rev. Relativity* **2**, 1 (1999); <http://www.livingreviews.org/lrr-1999-1>.
- [20] J. M. Stewart, *Lecture Notes in Physics* (Springer-Verlag, Berlin, 1971), Vol. 10.
- [21] J. M. Stewart, *Proc. Roy. Soc. A* **357**, 59 (1977); W. Israel and J. M. Stewart, *ibid. A* **365**, 43 (1979).
- [22] W. Israel and J. M. Stewart, *Annals Phys.* **118**, 341 (1979).
- [23] G. S. Denicol, T. Koide and D. H. Rischke, *Phys. Rev. Lett.* **105**, 162501 (2010) [arXiv:1004.5013 [nucl-th]].
- [24] G. S. Denicol, H. Niemi, E. Molnar and D. H. Rischke, *Phys. Rev. D* **85**, 114047 (2012) [arXiv:1202.4551 [nucl-th]].
- [25] G. S. Denicol, E. Molnar, H. Niemi and D. H. Rischke, *Eur. Phys. J. A* **48**, 170 (2012) [arXiv:1206.1554 [nucl-th]].
- [26] B. Betz, D. Henkel and D. H. Rischke, *Prog. Part. Nucl. Phys.* **62**, 556 (2009) [arXiv:0812.1440 [nucl-th]]; B. Betz, D. Henkel and D. H. Rischke, *J. Phys. G* **36**, 064029 (2009).
- [27] B. Betz, G. S. Denicol, T. Koide, E. Molnar, H. Niemi and D. H. Rischke, *EPJ Web Conf.* **13**, 07005 (2011) [arXiv:1012.5772 [nucl-th]].
- [28] A. Muronga, *Phys. Rev. C* **76**, 014909 (2007) [arXiv:nucl-th/0611090].
- [29] T. Tsumura, T. Kunihiro and K. Ohnishi, *Phys. Lett. B* **646**, 134 (2007).
- [30] K. Tsumura and T. Kunihiro, *Phys. Lett. B* **690**, 255 (2010) [arXiv:0906.0079 [hep-ph]].
- [31] K. Tsumura and T. Kunihiro, *Prog. Theor. Phys. Suppl.* **195**, 19 (2012) [arXiv:1205.5843 [nucl-th]].
- [32] M. A. York and G. D. Moore, *Phys. Rev. D* **79**, 054011 (2009) [arXiv:0811.0729 [hep-ph]].
- [33] A. Monnai and T. Hirano, *Phys. Rev. C* **80**, 054906 (2009) [arXiv:0903.4436 [nucl-th]].
- [34] A. Monnai, T. Hirano, *Nucl. Phys. A* **847**, 283 (2010) [arXiv:1003.3087 [nucl-th]].
- [35] W. Florkowski and R. Ryblewski, *Phys. Rev. C* **83**, 034907 (2011) [arXiv:1007.0130 [nucl-th]].
- [36] M. Martinez and M. Strickland, *Nucl. Phys. A* **848**, 183 (2010) [arXiv:1007.0889 [nucl-th]].
- [37] A. Jaiswal, R. S. Bhalerao and S. Pal, *Phys. Lett. B* **720**, 347 (2013) [arXiv:1204.3779 [nucl-th]].
- [38] A. Jaiswal, *Phys. Rev. C* **87**, 051901 (2013) [arXiv:1302.6311 [nucl-th]].
- [39] A. Jaiswal, *Phys. Rev. C* **88**, 021903 (2013) [arXiv:1305.3480 [nucl-th]].
- [40] D. Teaney and L. Yan, *Phys. Rev. C* **89**, 014901 (2014) [arXiv:1304.3753 [nucl-th]].
- [41] D. Jou, J. Casas-Vázquez and G. Lebon, *Rep. Prog. Phys.* **51**, 1105 (1988).
- [42] D. Jou, J. Casas-Vázquez and G. Lebon, *Rep. Prog. Phys.* **62**, 1035 (1999).
- [43] J. Gariel and G. Le Denmat, *Phys. Rev. D* **50**, 2560 (1994).
- [44] T. Koide, G. S. Denicol, P. Mota and T. Kodama, *Phys. Rev. C* **75**, 034909 (2007) [hep-ph/0609117].
- [45] A. El, Z. Xu and C. Greiner, *Phys. Rev. C* **81**, 041901 (2010) [arXiv:0907.4500 [hep-ph]].
- [46] P. Van and T. S. Biro, *Eur. Phys. J. ST* **155**, 201 (2008) [arXiv:0704.2039 [nucl-th]].
- [47] T. S. Biro, E. Molnar and P. Van, *Phys. Rev. C* **78**, 014909 (2008) [arXiv:0805.1061 [nucl-th]].
- [48] P. Van and T. S. Biro, *Phys. Lett. B* **709**, 106 (2012) [arXiv:1109.0985 [nucl-th]].
- [49] A. Muronga, *J. Phys. G* **37**, 094008 (2010).
- [50] T. Osada, *Phys. Rev. C* **85**, 014906 (2012) [arXiv:1111.1276 [nucl-th]].
- [51] A. Jaiswal, R. S. Bhalerao and S. Pal, *Phys. Rev. C* **87**, 021901 (2013) [arXiv:1302.0666 [nucl-th]].
- [52] G. S. Denicol, J. Noronha, H. Niemi and D. H. Rischke, *Phys. Rev. D* **83**, 074019 (2011) [arXiv:1102.4780 [hep-th]].
- [53] G. S. Denicol, H. Niemi, I. Bouras, E. Molnar, Z. Xu, D. H. Rischke and C. Greiner, arXiv:1207.6811 [nucl-th].
- [54] S. R. de Groot, W. A. van Leeuwen and Ch. G. van Weert, *Relativistic Kinetic Theory - Principles and applications*, North Holland (1980).
- [55] C. Cercignani and G. M. Kremer, *The Relativistic Boltzmann Equation: Theory and Applications*, Birkhäuser (2002).
- [56] J. L. Anderson, *J. Math. Phys.* **15** No.7, 1116 (1974).
- [57] H. Struchtrup *Annals Phys.* **257**, 111 (1997).
- [58] H. Struchtrup *Physica A* **253**, 555 (1998).
- [59] P. Huovinen and D. Molnar, *Phys. Rev. C* **79**, 014906 (2009) [arXiv:0808.0953 [nucl-th]].
- [60] I. Bouras, E. Molnar, H. Niemi, Z. Xu, A. El, O. Fochler, C. Greiner and D. H. Rischke, *Nucl. Phys. A* **830**, 741C (2009) [arXiv:0907.4519 [hep-ph]].
- [61] I. Bouras, E. Molnar, H. Niemi, Z. Xu, A. El, O. Fochler, C. Greiner and D. H. Rischke, *Phys. Rev. C* **82**, 024910 (2010) [arXiv:1006.0387 [hep-ph]].
- [62] W. Florkowski, R. Ryblewski and M. Strickland, *Phys. Rev. C* **88**, 024903 (2013) [arXiv:1305.7234 [nucl-th]].
- [63] K. Peeters, *Comput. Phys. Commun.* **176**, 550 (2007) [cs/0608005 [cs.SC]].

- [64] K. Peeters, [hep-th/0701238 [HEP-TH]].
- [65] R. Mertig, M. Bohm and A. Denner, Comput. Phys. Commun. **64**, 345 (1991).