Large-N Structure of Tetraquark Mesons

Richard F. Lebed*

Department of Physics, Arizona State University, Tempe, AZ 85287-1504

(Dated: August 2013)

Abstract

Following Weinberg's argument that narrow tetraquark mesons are not precluded in large N_c QCD, we explore the flow of N_c factors needed for the consistency of this picture, and show that they must arise in a novel way, not simply through the usual combinatoric counting.

PACS numbers: 11.15.Pg,14.40.Rt

arXiv:1308.2657v2 [hep-ph] 24 Sep 2013

*Electronic address: Richard.Lebed@asu.edu

Recently, Weinberg [1] re-evaluated the well-known argument of Coleman [2], that color-singlet operators of the form $q\bar{q}q\bar{q}$ create nothing but meson pairs, to propose that such operators can consistently create narrow tetraquarks [widths of $O(1/N_c)$, the same as ordinary mesons] in the limit of a large number N_c of QCD colors. Subsequently, Knecht and Peris [3] showed that some tetraquarks of this type would mix with mesons at $O(N_c^0)$ and hence be very difficult to disentangle in experimental data, but they also showed that tetraquarks of certain flavor structures do not suffer this fate and may be discernible in data. Our purpose in this short note is to show that narrow tetraquark states, if they indeed exist, exhibit an N_c behavior at the quark level that is novel compared to that of conventional meson or baryon states.

In either approach, one begins with the color-singlet quartic quark operator

$$\mathcal{Q}(x) = \sum_{ij} C_{ij} \mathcal{B}_i(x) \mathcal{B}_j(x) \,, \tag{1}$$

where $\mathcal{B}_i(x)$ are color-singlet quark bilinears,

$$\mathcal{B}_i(x) = \sum_{a=1}^{N_c} \overline{q^a}(x) \Gamma_i \, q^a(x) \,, \tag{2}$$

which (if needed) may be redefined to have zero vacuum expectation values, the index a sums over all N_c colors, Γ^i represents the N_c-independent spin-flavor connections defining $\mathcal{B}_i$, and $C_{ij} = C_{ji}$ are a collection of numerical coefficients (taken N_c-independent in Ref. [1]) that describe the correlation of the bilinears $\mathcal{B}_i$, $\mathcal{B}_j$ at the spacetime point x. It is a feature of SU(N_c) group theory (operationally manifested by Fierz reordering) that any color-singlet quartic $\mathcal{Q}$ can be decomposed into such a product. The interesting quantity to study is the $\mathcal{Q}$ two-point function, which has disconnected and connected pieces:

$$\langle \mathcal{Q}(x) \mathcal{Q}^\dagger(y) \rangle \;=\; \sum_{ijkl} C_{ij} C_{kl} \left[\langle \mathcal{B}_i(x) \mathcal{B}_k^\dagger(y) \rangle \langle \mathcal{B}_j(x) \mathcal{B}_l^\dagger(y) \rangle + \langle \mathcal{B}_i(x) \mathcal{B}_j(x) \mathcal{B}_k^\dagger(y) \mathcal{B}_l^\dagger(y) \rangle_{\rm conn} \right] . \tag{3}$$

Suppose that Eq. (1) is generalized to a nonlocal form by allowing the spacetime point x appearing in the two bilinears to separate into distinct values $x_1 \neq x_2$, and that such bilinears are used in the definition of Eq. (3) with $y \to y_1 \neq y_2$ as well. The coefficients in this case generalize to $C_{ij}(x_1 - x_2) C_{kl}(y_1 - y_2)$. Then, by conventional large N_c counting, the first term is $O(N_c^2)$ and is saturated by sums over the propagators of two noninteracting mesons, each of which (by LSZ reduction) has a decay constant $f_{\mathcal{B}} = O(N_c^{1/2})$ appearing for each

asymptotic state, thus neatly accounting for the leading-order N_c behavior. The second term is $O(N_c^1)$ and represents, at least in part, two-meson scattering diagrams; once the four $f_{\mathcal{B}}$ factors are amputated, one finds the well-known result that meson-meson scattering amplitudes are only $O(1/N_c)$.

Now allow the points $x_{1,2}$ and $y_{1,2}$ to again coalesce to x and y, respectively, as in Eq. (1). Then the N_c counting remains the same as before, but poles due to localized one-tetraquark states may now be considered. However, true propagating tetraquark states can only appear in the connected term, which is subleading in $1/N_c$. From this fact, Ref. [2] concludes that the two-point function of Eq. (3) produces only two-meson states, but Ref. [1] concludes that nothing precludes the production of tetraquark states in its subleading, connected piece. Moreover, since this $O(N_c^1)$ term contains the full amplitude for the free $O(N_c^0)$ propagation of a tetraquark, LSZ reduction shows that the tetraquark decay constant $f_{\mathcal{Q}} = O(N_c^{1/2})$, and by using this result on the $O(N_c^1)$ connected three-point function $\langle \mathcal{Q}(x)\mathcal{B}_i(y)\mathcal{B}_j(z)\rangle_{\rm conn}$, one finds the trilinear coupling to be $O(N_c^{-1/2})$, and hence the tetraquark decay width to be $O(1/N_c)$. These are the central results of Ref. [1].

The point of interest in this note is the interpretation of the tetraquark decay constant $f_{\mathcal{Q}}$ in terms of the underlying quark degrees of freedom. In the case of mesons, the fact that $f_{\mathcal{B}} = O(N_c^{1/2})$ is equivalent to the fact that the LSZ reduction identifies the operator $\mathcal{B}/\sqrt{N_c}$ as the one that creates or destroys a properly normalized meson state. Viewed in terms of the quark color degrees of freedom in Eq. (2), this prefactor is simply the one that produces a normalized wave function over the color degrees of freedom c_a, $\sum_{a=1}^{N_c} \bar{c}_a c_a/\sqrt{N_c}$. In the case of the tetraquark, however, Eq. (3) indicates that the color wave function of the operator $\mathcal{Q}$ has a norm-square of N_c^2, and that of the operator $\mathcal{Q}/\sqrt{N_c}$, which according to Ref. [1] creates or destroys normalized tetraquark states, actually has norm-square of N_c^1 rather than unity. How can these facts be made to agree?

In fact, while the operator $\mathcal{Q}/\sqrt{N_c}$ creates and destroys normalized tetraquarks, it need not produce tetraquarks alone. Indeed, our argument that the connected nonlocal four-bilinear correlation function is dominated by two-meson scattering suggests that a dominant component of the $\mathcal{Q}/\sqrt{N_c}$ two-point function is represented by coincident (propagating from y to x) pairs of mesons, including meson molecules. The authors of Refs. [1, 3] assert that narrow tetraquarks *can* exist; our purpose is to find a condition that causes them to arise naturally. In order to obtain the additional $1/\sqrt{N_c}$ suppression for creating properly

normalized tetraquarks, we note that pointlike tetraquark states appear only in the local limits $x_2 \to x_1$, $y_2 \to y_1$, or at least when $x_1 - x_2$ and $y_1 - y_2$ are sufficiently small that the tetraquark wave function has appreciable amplitude at all four of these points; but then, the use of LSZ reduction (which requires well-separated sources) to identify four distinct meson sources is no longer justified. One may, for example, model the required behavior by allowing the coefficients C_{ij} of Eq. (1) in the nonlocal case to have a contribution of the Gaussian form

$$\delta C_{ij} \sim \exp[-N_c^m \Lambda_{\rm QCD}^2 (x_1 - x_2)^2] \,, \tag{4}$$

where $m > 0$. For any finite separation $(x_1 - x_2)^2 \gtrsim 1/\Lambda_{\rm QCD}^2$, one finds $\delta C_{ij} \to 0$ as $N_c \to \infty$, so that tetraquark poles are features of the connected four-bilinear correlator only in the local limit and do not modify the naive $O(N_c^0)$ counting of C_{ij}. However, integrating δC_{ij} over an arbitrarily small range of separation between x_1 and x_2—that is to say, averaging over the $O(1/\Lambda_{\rm QCD})$ spatial extent of the tetraquark wave function—gives a contribution of $O(N_c^{-m/2})$ for each integral taken over C_{ij}. For the special case of 3 spatial integrals and $m = \frac{1}{3}$, one obtains precisely the desired $1/\sqrt{N_c}$ behavior to obtain a narrow tetraquark with $f_{\mathcal{Q}} = O(\sqrt{N_c})$. Any particular choice for carrying out this averaging is suitable, as long as it integrates over the whole functional peak.

The proposed profile δC_{ij} presented here is something both new and familiar in large N_c counting. It arises through the noncommutativity of the $(x_1 - x_2)^2 \to 0$ and $N_c \to \infty$ limits. δC_{ij} is not obtained from any first-principles counting of factors of the QCD coupling constant $g_s \sim 1/\sqrt{N_c}$ and combinatoric N_c color factors [4], which are sufficient to explain the basic N_c counting for ordinary mesons and glueballs, hybrid mesons [5], baryons [6], and baryon resonances [7]. Indeed, δC_{ij} is nonperturbative in N_c counting, and since $1/g_s^2 \sim N_c$, the profile is reminiscent of an instanton structure. Nevertheless, we do not assert the presence of any specific instanton-like physics at work here, but rather only the need for a contribution to C_{ij} with distinct local and nonlocal limits with respect to N_c; any functional form that produces the $1/\sqrt{N_c}$ for $(x_1 - x_2)^2 \to 0$ is suitable for this purpose. Finally, the appearance of a $1/\sqrt{N_c}$-suppressed four-quark state arising in the limit in which two $\mathcal{B}$ operators converge in the $O(N_c^1)$ three-point function $\langle \mathcal{B}_i(x) \mathcal{B}_j(y) \mathcal{B}_k(z) \rangle_{\rm conn}$ provides a mechanism through which meson-tetraquark mixing can occur at $O(N_c^0)$ [3].

In summary, we propose a resolution to the problem of N_c counting for the proposed narrow tetraquark states of Ref. [1], and are led to a novel application of inducing the

required N_c factors via a term nonperturbative in N_c.

Acknowledgments

The author is indebted to S. Weinberg and T. Cohen for helpful discussions. This work was supported by the National Science Foundation under Grant No. PHY-1068286.

---

[1] S. Weinberg, Phys. Rev. Lett. 110, **261601** (2013) [arXiv:1303.0342 [hep-ph]].

[2] S.R. Coleman, *Aspects of Symmetry* (Cambridge Univ. Press, Cambridge, England, 1985), pp. 377-378.

[3] M. Knecht and S. Peris, Phys. Rev. D **88**, 036016 (2013) [arXiv:1307.1273 [hep-ph]].

[4] G. 't Hooft, Nucl. Phys. B **72**, 461 (1974).

[5] T.D. Cohen, Phys. Lett. B **427**, 348 (1998) [hep-ph/9801316].

[6] E. Witten, Nucl. Phys. B **160**, 57 (1979).

[7] T.D. Cohen and R. F. Lebed, Phys. Rev. Lett. **91**, 012001 (2003) [hep-ph/0301167].