

PRECISION MEASURES OF THE PRIMORDIAL ABUNDANCE OF DEUTERIUM*

RYAN J. COOKE^{1,7}, MAX PETTINI^{2,3}, REGINA A. JORGENSEN⁴, MICHAEL T. MURPHY⁵, CHARLES C. STEIDEL⁶

(Dated: February 26, 2022)

Draft version February 26, 2022

ABSTRACT

We report the discovery of deuterium absorption in the very metal-poor ([Fe/H] = −2.88) damped Lyman- α system at $z_{\text{abs}} = 3.06726$ toward the QSO SDSS J1358+6522. On the basis of 13 resolved D_I absorption lines and the damping wings of the H_I Lyman α transition, we have obtained a new, precise measure of the primordial abundance of deuterium. Furthermore, to bolster the present statistics of precision D/H measures, we have reanalyzed all of the known deuterium absorption-line systems that satisfy a set of strict criteria. We have adopted a blind analysis strategy (to remove human bias), and developed a software package that is specifically designed for precision D/H abundance measurements. For this reanalyzed sample of systems, we obtain a weighted mean of $(\text{D}/\text{H})_{\text{p}} = (2.53 \pm 0.04) \times 10^{-5}$, corresponding to a universal baryon density $100 \Omega_{\text{b},0} h^2 = 2.202 \pm 0.046$ for the standard model of big bang nucleosynthesis (BBN). By combining our measure of $(\text{D}/\text{H})_{\text{p}}$ with observations of the cosmic microwave background (CMB), we derive the effective number of light fermion species, $N_{\text{eff}} = 3.28 \pm 0.28$. We therefore rule out the existence of an additional (sterile) neutrino (i.e. $N_{\text{eff}} = 4.046$) at 99.3 per cent confidence (2.7σ), provided that the values of N_{eff} and of the baryon-to-photon ratio (η_{10}) did not change between BBN and recombination. We also place a strong bound on the neutrino degeneracy parameter, independent of the ^4He primordial mass fraction, Y_{p} : $\xi_{\text{D}} = +0.05 \pm 0.13$ based only on the CMB+(D/H)_p observations. Combining this value of ξ_{D} with the current best literature measure of Y_{p} , we find a 2σ upper bound on the neutrino degeneracy parameter, $|\xi| \leq +0.062$.

Subject headings: quasars: absorption lines – quasars: individual: J1358+6522 – cosmology: observations.

1. INTRODUCTION

We are currently in an exciting era of high-precision cosmology, with most of the “standard” cosmological model parameters now known to within a few percent. In particular, the recent analysis of the cosmic microwave background (CMB) temperature fluctuations recorded by the *Planck* satellite has found that the present day density of baryons, $\Omega_{\text{b},0}$, contributes just $(2.205 \pm 0.028)/h^2$ per cent of the critical density (Planck Collaboration 2013), where h is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$. *Planck*’s determination of $\Omega_{\text{b},0} h^2$, which is now limited by cosmic variance, is the most precise measure of this fundamental physical quantity for the foreseeable future (when derived from the CMB).

For a long time (e.g. Wagoner, Fowler, & Hoyle 1967), it has been appreciated that a complementary measurement of

$\Omega_{\text{b},0} h^2$ can be deduced from the relative abundances of the light elements that were created during big bang nucleosynthesis (BBN). Aside from protons, the only stable nuclei that were produced in astrophysically accessible abundances are ^2H (a.k.a. deuterium, D), ^3He , ^4He , and trace amounts of ^7Li (see Steigman 2007 for a comprehensive review). In particular, considerable effort has been devoted to measuring the abundance of deuterium (D/H), the mass fraction of ^4He (Y_{p}), and the abundance of ^7Li . Of these, the primordial abundance of deuterium is generally accepted as the best ‘baryometer’, owing to its sensitivity and monotonic relationship to $\Omega_{\text{b},0} h^2$.

The mass fraction of ^4He , on the other hand, is relatively insensitive to $\Omega_{\text{b},0} h^2$, but depends quite strongly on the expansion rate of the early Universe (typically parameterized in non-standard models of BBN as an effective number of neutrino species, N_{eff}). At present, this measurement is primarily limited by systematic uncertainties in converting the He and H emission lines of metal-poor H II regions into an estimate of Y_{p} (see e.g. Izotov, Stasinska, & Guseva 2013; Aver et al. 2013). Unlike Y_{p} , the $^7\text{Li}/\text{H}$ ratio depends modestly on both $\Omega_{\text{b},0} h^2$ and N_{eff} . However, the observationally inferred value for the “primordial” level, derived from metal-poor Galactic halo stars (Spite & Spite 1982; Meléndez et al. 2010), is discrepant by a factor of ~ 3 compared with the standard model predictions, becoming even more discrepant at the lowest metallicities (Sbordone et al. 2010).

As pointed out recently by Nollett & Holder (2011) (see also Cyburt 2004), *precise* measures of the primordial D/H ratio can provide interesting bounds on N_{eff} , when combined with a measure of $\Omega_{\text{b},0} h^2$ from the CMB. The promise of this method was recently demonstrated by Pettini & Cooke (2012), who obtained the most precise measure of the D/H ratio to date, using a metal-poor ([Fe/H] = −2.33) damped Lyman α system (DLA), seen in absorption against a bright

* Based on observations collected at the European Organisation for Astronomical Research in the Southern Hemisphere, Chile [VLT program IDs: 68.B-0115(A), 70.A-0425(C), 078.A-0185(A), 085.A-0109(A)], and at the W.M. Keck Observatory which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation. Keck telescope time was partially granted by NAOJ, through the Telescope System Instrumentation Program (TSIP). TSIP is funded by NSF.

¹ Department of Astronomy and Astrophysics, UCO/Lick Observatory, University of California, Santa Cruz, CA 95064, USA

² Institute of Astronomy, Madingley Road, Cambridge, UK, CB3 0HA

³ Kavli Institute for Cosmology, Madingley Road, Cambridge, UK, CB3 0HA

⁴ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

⁵ Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Hawthorn, Victoria 3122, Australia

⁶ California Institute of Technology, MS 249-17, Pasadena, CA 91125, USA

⁷ Morrison Fellow; email: rcooke@ucolick.org

background quasar.

The prospect of measuring the D/H ratio in gas clouds seen in absorption along the line-of-sight to a high-redshift quasar was first pointed out by Adams (1976). This vision was only realized much later, with the advent of 8-10 m class telescopes equipped with high resolution echelle spectrographs (Burles & Tytler 1998a,b). Since these first discoveries, a handful of additional cases have been identified (see Pettini & Cooke 2012 for the ‘*top-ten*’ list), including one system that appears to be chemically pristine (Fumagalli, O’Meara, & Prochaska 2011). One lingering concern with this prime set of measurements is that the dispersion in the reported D/H measures is significantly larger than the quoted errors (first noted by Steigman 2001). Indeed, a simple χ^2 test reveals that the errors for all D/H measurements would need to be scaled upwards by $\sim 33\%$, if the observed dispersion were due to chance alone. An alternative possibility, if the quoted random errors are in fact realistic, is that the analyses may suffer from small systematic biases that are now important to recognize and correct for.

This concern has prompted us to identify the rare handful of systems which afford *precise* measurements of the primordial abundance of deuterium together with realistic error estimates. This effort is part of our ongoing survey to study the chemistry of the most metal-poor DLAs, described in more detail in Pettini et al. (2008a) and Cooke et al. (2011).

In this paper, we present the *Precision Sample* of primordial deuterium abundance measurements, consisting of the current best estimates of D/H in QSO absorption line systems. We also report a DLA where a new, precise measurement of the primordial deuterium abundance could be obtained. All of these systems have been reanalyzed in a self-consistent manner, taking into account the dominant sources of systematic uncertainty. In the following section, we provide the details of the observations and data reduction procedures for the newly observed DLA, and discuss the selection criteria we have adopted to define the Precision Sample. In Section 3, we describe the analysis of the new system with deuterium absorption, and estimate its D/H ratio. In Section 4 we derive the value of $(D/H)_p$ from the Precision Sample, discuss its cosmological implications, and consider current limitations in using D/H abundance measurements for cosmology. Section 5 deals with the implications of our results for physics beyond the standard model, considering in particular bounds on the effective number of neutrino families and on the lepton asymmetry. Finally, in Section 6, we summarize the main findings from this work.

2. OBSERVATIONS AND DATA REDUCTION

2.1. The DLA towards J1358+6522

Most newly discovered DLAs are now identified using the low-resolution spectra provided by the Sloan Digital Sky Survey (SDSS). Recent searches for DLAs have yielded several thousand systems (Prochaska & Wolfe 2009; Noterdaeme et al. 2012). At the modest resolution ($R \sim 2000$) and signal-to-noise ratio ($S/N \sim 20$) of the SDSS spectra, the metal absorption lines of the most metal-poor DLAs are unresolved and undetected. Follow-up, high spectral resolution observations are thus required to pin-down their chemical abundances (see Cooke et al. 2011).

Penprase et al. (2010) were the first to recognize the very metal-poor DLA at redshift $z_{\text{abs}} \approx 3.0674$ towards the $z_{\text{em}} \approx 3.173$ quasar SDSS J1358+6522. On the basis of their in-

termediate resolution observations (60 km s^{-1} full-width at half maximum; FWHM) with the Keck Echelle Spectrograph and Imager, these authors concluded that this DLA has a metallicity of $[\text{Fe}/\text{H}] \leq -3.02$, and is thus among the most metal-poor known (Cooke et al. 2011). We re-observed this QSO with the Keck High Resolution Echelle Spectrometer (HIRES; Vogt et al. 1994) using the red cross-disperser to perform an accurate chemical abundance analysis (Cooke, Pettini, & Murphy 2012). The HIRES data pinned down the metallicity of the DLA ($[\text{Fe}/\text{H}] = -2.84 \pm 0.03$) and provided the first indication for the presence of a handful of resolved D I absorption lines.

Taking advantage of the high efficiency of the HIRES UV cross-disperser at blue wavelengths, we then performed dedicated HIRES observations with the goal of obtaining a precise measure of the deuterium abundance. Our observations included a total of 24 300 s on 2012, May 11, (divided equally into nine exposures) and an additional 29 750 s on 2013, May 3 and 5. An identical instrument setup was employed for both runs; we used the C1 decker ($7'' \times 0.861''$, well-matched to the seeing conditions of $0.6'' - 0.8''$), which delivers a nominal spectral resolution of $R \sim 48\,000$ ($\equiv 6.3 \text{ km s}^{-1}$ FWHM) for a uniformly illuminated slit. Our setup covers the wavelength range $3650 - 6305 \text{ \AA}$, with small gaps near 4550 \AA and 5550 \AA due to the gaps between the HIRES detector chips. The exposures were binned on-chip 2×2 .

The data were reduced following the standard procedures of bias subtraction, flat-fielding, order definition and extraction. The data were mapped onto a vacuum, heliocentric wavelength scale from observations of a ThAr lamp which bracketed the science exposures. To test the quality of the data reduction, we reduced the data using two pipelines: MAKEE which is maintained by T. Barlow⁹, and HIREDEX which is maintained by J. X. Prochaska¹⁰. We found that the skyline subtraction performed by HIREDEX was superior to that of MAKEE, and substantially improved the S/N of the spectra at blue wavelengths. This was very important for the removal of telluric features near the redshifted D I and H I transitions of the DLA. All data analyzed and presented herein for J1358+6522 were reduced with HIREDEX. The individual orders were combined with UVES_POPLER, which is maintained by M. T. Murphy¹¹. As a final step, we flux-calibrated the echelle data by comparison to the SDSS data. The combined S/N ratio of the final spectrum per 2.5 km s^{-1} pixel is ~ 30 near 4000 \AA , ~ 45 at 5000 \AA and ~ 20 near 6000 \AA . The reduced, flux-calibrated spectrum is presented in Figure 1, where the red tick marks indicate the wavelengths of the redshifted H I Lyman series transitions.

2.2. Literature D/H QSO Absorption Line Systems

2.2.1. Selection Criteria

In this section, we outline the strict set of rules that we have used to define the Precision Sample. Our goal is to identify the small handful of systems currently known where the most accurate and precise measures of D/H can potentially be obtained. The set of restrictions we applied are as follows:

⁹ MAKEE is available at:

<http://www.astro.caltech.edu/~tb/makee/>

¹⁰ HIREDEX can be obtained from:

<http://www.ucolick.org/~xavier/HIREDEX/index.html>

¹¹ UVES_POPLER can be downloaded from:

http://astronomy.swin.edu.au/~mmurphy/UVES_popler/




FIG. 1.— Flux-calibrated spectrum of J1358+6522 (black histogram). The zero-level and error spectrum are shown as the green dashed and solid blue lines respectively. The H I Lyman series lines of the DLA are indicated by red tick marks. Since the SDSS data do not extend below 3800 Å, we extrapolated the response curve for fluxing the data a further 100 Å to the blue.

- We require that the H I Ly α absorption line must exhibit Lorentzian damping wings. This criterion is satisfied for absorption line systems with $\log N(\text{H I})/\text{cm}^{-2} \geq 19$. Such absorption lines lie on the damping regime of the curve-of-growth, where $N(\text{H I})$ can be derived independently of the cloud model; the damping wings uniquely determine the total H I column density.
- Due to the power of the H I Ly α transition in establishing the total H I column density, we require that the wings of this transition are not strongly blended with nearby, unrelated, strong H I absorption systems. If there exist additional strong H I absorption systems nearby, we impose that the edges of their Ly α absorption troughs should be separated by at least 500 km s^{-1} , and that all such absorption systems should be modelled simultaneously.
- At least two, resolved and apparently unblended, optically thin transitions of D I must be available. This ensures that the total D I column density can be measured accurately, and independently of the cloud model.
- The data must have been obtained with a high resolution echelle spectrograph, with $R \geq 30\,000$ (i.e. $v_{\text{FWHM}} \leq 10 \text{ km s}^{-1}$), to resolve the broadening of the D I lines, and recorded at $S/N \gtrsim 10$ per pixel ($\sim 3 \text{ km s}^{-1}$) at both Ly α and the weakest D I absorption line used in the analysis.
- Several unblended metal lines with a range of oscillator strengths for a given species (ideally O I and Si II) must be present if there is only one optically thin D I transition in order to determine the velocity structure of the absorbing cloud, and ensure that the presence of any partly ionized gas (which may contribute to the H I column density) is accurately modelled. However, if there exist at least two optically thin D I transitions, metal lines are not strictly required.

Of course, we are also limited by data that we, and others, have access to. Data that are not publicly available on archives could not be used in this study. In total, there are four systems in the literature that meet the above criteria. We now give a brief description of these systems.

2.2.2. *HS0105+1619*, $z_{\text{abs}} = 2.53651$

Keck+HIRES observations of the QSO HS 0105+1619 were obtained as follows: 1800 s in 1999 (Program ID: U05H, PI: A. Wolfe), 85 000 s in 1999–2000 by O’Meara et al. (2001), and 21 500 s in 2005 (Program ID: G10H, PI: D. Tytler). The data obtained by O’Meara et al. (2001) are not available from the Keck Observatory Archive. Thus, we have reanalysed only the last dataset which, in any case, was obtained with a much improved HIRES detector. As reported by O’Meara et al. (2001), the spectrum shows nine clean H I transitions in the Lyman series together with four optically thin D I absorption lines; the absorption is confined to a single velocity component at $z_{\text{abs}} = 2.536509$. The first ions exhibit slightly asymmetric profiles, arising from ionized gas blueshifted by 7.1 km s^{-1} relative to the main component (see Figure 8). The H I column density of this ionized component contributes just 0.01 dex to the total column density $\log N(\text{H I})/\text{cm}^{-2} = 19.42 \pm 0.01$. The data and best-fitting model are shown in Figure 8.

2.2.3. *Q0913+072*, $z_{\text{abs}} = 2.61829$

The quasar Q0913+072 intersects a DLA with $\log N(\text{H I})/\text{cm}^{-2} = 20.34 \pm 0.04$ at $z_{\text{abs}} = 2.61829$. Q0913+072 was observed with the Ultraviolet and Visual Echelle Spectrograph (UVES) on the European Southern Observatory’s (ESO) Very Large Telescope (VLT) facility in 2002 (Program ID: 68.B-0115(A), PI: P. Molaro), and then again in 2007 to obtain UV coverage down to the Lyman limit (Program ID: 078.A-0185(A), PI: M. Pettini), resulting in a total exposure time of 77 500 s. The details of this system’s chemical composition were reported by Pettini et al. (2008a) (see also Erni et al. 2006), and the analysis of the D I lines was presented by Pettini et al. (2008b); with $[\text{O}/\text{H}] = -2.40$ this is currently the most metal-poor DLA known with resolved D I lines. There is a small column density of ionized gas at velocities near that of the DLA; we have carefully modelled its contribution to the overall absorption in the metal lines and the H I Lyman series (see Figure 9).

Since Q0913+072 does not have an SDSS spectrum from which we could perform a flux calibration, we obtained a set of VLT+FOR2 (Focal Reducer/low dispersion Spectrograph 2) exposures from the VLT archive to flux calibrate the UVES spectrum. (Program ID: 70.A-0425(C), PI: L. Wisotzki). The FOR2 data were reduced following standard procedures, using the software routines from the ESO FOR2 data reduction pipeline¹². The UVES data and the corresponding model fits are shown in Figure 9.

2.2.4. *J1419+0829*, $z_{\text{abs}} = 3.04973$

The chemical properties of this DLA were discussed by Cooke et al. (2011), while Pettini & Cooke (2012) reported the analysis of the deuterium lines. This system currently holds the record for the most precise determination of the D/H ratio. The analysis technique developed in Pettini & Cooke (2012) is the same as that employed here (see Section 3.2), except for two aspects: (1) in the present work all QSO spectra were analysed “blind,” i.e., without knowledge of the value of D/H until the line fitting procedure was concluded; and (2) we now allow for an arbitrary continuum fitting with Legendre polynomials, rather than the combination of power-law continuum and Gaussian emission lines used by Pettini & Cooke (2012). To maintain consistency with the other DLAs investigated here, we have reanalyzed this system with the new prescription for the continuum definition, and the analysis was performed blind. We found that the final value of D/H from this new analysis deviated by just 0.005 dex from that reported by Pettini & Cooke (2012), with an identical estimate for the error. This small shift is due to our new approach for dealing with the systematic uncertainty in the continuum level.

2.2.5. *J1558–0031*, $z_{\text{abs}} = 2.70242$

The data for J1558–0031 consist of a 11 300 s Keck+HIRES spectrum obtained in 2006 (Program ID: U152Hb, PI: J. X. Prochaska), first reported by O’Meara et al. (2006). With $\log N(\text{H I})/\text{cm}^{-2} = 20.67 \pm 0.05$, this $z_{\text{abs}} = 2.70242$ DLA is the highest H I column density system currently known with at least one optically thin D I absorption feature. There exists another metal-poor DLA along this sightline, at $z_{\text{abs}} = 2.629756$, that also exhibits damped Ly α absorption ($\log N(\text{H I})/\text{cm}^{-2} = 19.726 \pm 0.007$).

¹² We used version 3.9.6, obtained from:
<http://www.eso.org/sci/software/pipelines/>

This second system contains a single absorption component, as evidenced by several Si II absorption lines (see bottom-right panel of Figure 10). Although D I absorption is only seen in the higher redshift DLA, we include the H I and metal lines of both DLAs in the fitting procedure, since the damping wings of the two DLAs overlap slightly. Portions of the HIRES spectrum, together with the best-fitting model, are reproduced in Figure 10. Unlike the O’Meara et al. (2006) analysis, which uses data acquired with both Magellan+MIKE and Keck+HIRES, our analysis exclusively uses the Keck+HIRES data described above.

3. PROFILE ANALYSIS

The analysis we have adopted to measure the D/H ratio is an improvement over previous studies, with two main differences: (1) the profile fitting and χ^2 minimization process was performed with new, purpose-built software routines; and (2) we have tried to identify the dominant systematics, and have attempted to account for these affects. Before we discuss the details of our line profile analysis, we briefly describe the important aspects of the new software we have written to measure D/H in metal-poor DLAs. We will focus our discussion on the newly discovered system with a precise value of the D/H abundance (J1358+6522), but we stress that all five DLAs were analyzed in an identical manner.

3.1. Line Profile Fitting Software

To account for the dominant systematics that could potentially affect the measurement of D/H, we have developed a new suite of software routines written in the PYTHON software environment. Our Absorption Line Software (ALIS; described in more detail by R. Cooke 2014, in preparation) uses a modified version of the MPFIT package (Markwardt 2009). MPFIT employs a Levenberg-Marquardt least-squares minimization algorithm to derive the model parameters that best fit the data (i.e. the parameters that minimize the difference between the data and the model, weighted by the error in the data). Unlike the software packages that have previously been used to measure D/H, ALIS has the advantage of being able to fit an arbitrary emission profile for the quasar whilst simultaneously fitting to the absorption from the DLA; any uncertainty in the continuum is therefore automatically folded into the final uncertainty in the D/H ratio. For all transitions, we use the atomic data compiled by Morton (2003).

3.2. Analysis Technique for Measuring D/H

We first highlight one of the main strengths of our analysis which, to our knowledge, has not been implemented in previous D/H studies. In order to remove human bias from the results, we have adopted a blind analysis strategy, whereby the only information revealed during the profile fitting process is the best-fitting χ^2 value, and plots of the corresponding model fits to the data; all of the parameter fitting results are entirely hidden from view. Only after the analysis converged on the best-fitting model were the full results of the model fit uncovered, and these are the final values quoted here. No further changes were made to the model fit nor data reduction after the results were revealed.

To allow for an efficient computation, a small wavelength interval (typically $\pm 300 \text{ km s}^{-1}$) around each absorption line of interest was extracted. For each of these intervals, we manually identified the regions to be used in the fitting process by

selecting pixels that we attribute to either the quasar continuum or the DLA’s absorption lines. In some cases, it was also necessary to model unrelated absorption lines in the fitting procedure if there was significant line-blending.

We have also attempted to account for the main systematics and limitations that might affect the reported D/H value. The dominant systematics likely include the instrumental broadening function, the cloud model, and the placement of the quasar continuum and zero-level. We marginalize over the first of these systematics by assuming that the instrumental profile is a Gaussian¹³, where the Gaussian FWHM was allowed to be a free model parameter during the χ^2 minimization process.

Any uncertainty or bias in the choice of the cloud model is overcome by our selection criteria listed in Section 2.2.1. Specifically, DLAs have two advantages over other absorption line systems for the determination of the D/H ratio, both stemming from their high H I column densities: (1) The H I Ly α line exhibits well-defined Lorentzian damping wings¹⁴ from which the total H I column density can be deduced precisely and independently of the cloud model; and (2) many D I transitions in the Lyman series are normally detectable. With their widely varying oscillator strengths these multiple D I lines can pin down the D I column density very precisely; the availability of unsaturated, high-order lines ensures that the value of D/H thus deduced does not depend on the detailed kinematic structure of the gas (i.e. the cloud model). Neither of these two advantages applies to Lyman limit systems whose H I column density can be three orders of magnitude lower than that of DLAs.

Finally, if the background level is not accurately determined, a small offset can systematically affect the derived D/H value. We have overcome this concern by allowing the zero level to be fit during the minimization process (assuming the offset to be the same at all wavelengths).

Arguably, the above systematic effects are sub-dominant compared to the choice of the continuum level. One of the improvements over previous analyses is that we have not specified *a priori* the level of the quasar continuum. Instead, we have marginalized over the continuum level during the fitting process by fitting Legendre polynomials to the flux-calibrated quasar continuum. The order of the polynomial was chosen to be sufficiently large to allow flexibility in arbitrarily defining the continuum level, whilst ensuring that the continuum was not being over-fit to the noise (examples are shown in the top panels of Figs. 2, 8, 9, and 10).

We emphasize that by including the above systematics into the fitting process, the uncertainties in the continuum, zero level, and instrumental broadening are incorporated into the final uncertainty in the derived D/H value. To summarize, we have marginalized over the continuum, zero level and instrumental FWHM, whilst simultaneously fitting for all of the DLA’s unbled metal lines, the DLA’s Ly α transition, and the entire H I and D I Lyman series. Our analysis therefore includes all of the available information that can help pin down the D/H value, and it accounts for the systematics that are currently thought to contribute the most to the overall error budget. We finally point out that every H I component used to fit

¹³ We confirmed that emission lines from the ThAr wavelength calibration spectrum were well-fit by a Gaussian profile.

¹⁴ Lee (2013) recently suggested that the wings of the damped Ly α absorption line exhibit an asymmetric deviation from a Lorentzian. Such a deviation from a Lorentzian profile would be marginalized over during the arbitrary continuum fitting process described below, and is thus expected to have a negligible impact on our results.




FIG. 2.— The top panel displays a portion of the flux-calibrated HIRES spectrum near the damped Ly α line at $z_{\text{abs}} = 3.06726$ toward J1358+6522 (black histogram) together with the error spectrum (continuous blue line). The dashed green line marks the best-fitting zero level of the data and the dashed blue line shows the best-fitting continuum level. The solid red line shows the overall best-fitting model to the DLA. The bottom panel shows a zoom-in of the data and model; the weak absorption feature that we have modelled on the blue wing of Ly α is Si III $\lambda 1206.5$ at the redshift of the DLA (~ 4907 Å in the observed frame).

the DLA was forced to have the same D I/H I ratio. We have therefore fit directly to the D I/H I value, rather than deriving separately the column densities of D I and H I. All absorption lines were modeled as Voigt profiles where the line broadening has both thermal and turbulent contributions. All of the DLAs in our survey were first modeled with a single absorption component. If the metal lines exhibited an asymmetric profile, an additional component was included to improve the quality of the fit.

To confirm that the model parameters had converged to their best-fitting values, we performed a convergence test; once the difference in the χ^2 between successive iterations was < 0.01 , the parameter values were stored and the χ^2 minimization recommenced with a tolerance of 10^{-3} . Once a successive iteration reduced the χ^2 by $< 10^{-3}$, the minimization was terminated and the parameter values from the two convergence criteria were compared. If all parameter values differed by $< 0.2\sigma$ (i.e. 20 per cent of the parameter error) then the model fit has converged.

As a final step, we repeated the χ^2 minimization process 20 times, perturbing the starting parameters of each run by the covariance matrix. This exercise ensures that our choice of starting parameters does not influence the final result. We

found that the choice of starting parameters has a negligible contribution to the error on D I/H I (typically 0.002 dex), but can introduce a small bias (again, typically 0.002 dex). We have accounted for this small bias in all of the results quoted herein.

3.3. Component Structure

Most of the narrow, low-ionization metal lines of the DLA toward J1358+6522 consist of a single component at $z_{\text{abs}} = 3.067259$. A second weaker component, blueshifted by 17.4 km s^{-1} ($z_{\text{abs}} = 3.06702$) contributes to Si III $\lambda 1206.5$ and to the strongest C II and Si II lines. Evidently, this weaker absorption arises in nearby ionized gas.

In fitting the absorption lines, we tied the redshift, turbulent Doppler parameter and kinetic temperature of the gas to be the same for the metal, D I and H I absorption lines. We allowed all of the cloud model parameters to vary, whilst simultaneously fitting for the continuum near every absorption line. Relevant parameters of the best-fitting cloud model so determined are collected in Table 1. Figures 2, 3 and 4 compare the data and model fits for, respectively, the damped Ly α line, the full Lyman series, and selected metal lines. [Since the metal lines analyzed here are the same as those shown in




FIG. 3.— A montage of the full Lyman series absorption in the DLA at $z_{\text{abs}} = 3.067259$ toward J1358+6522. The black histogram shows the data, fully adjusted to the best-fitting continuum and zero levels, while the red continuous line is the model fit. The minimum χ^2/dof for this fit is 6282.3/6401. Tick marks above the spectrum indicate the location of the velocity components (red ticks for H I, green ticks for D I).

TABLE 1
BEST-FITTING MODEL PARAMETERS FOR THE DLA AT $z_{\text{abs}} = 3.067259$ TOWARD THE QSO J1358+6522

Comp.	z_{abs}	T (K)	b_{turb} (km s $^{-1}$)	$\log N(\text{H I})$ (cm $^{-2}$)	$\log (\text{D I}/\text{H I})$	$\log N(\text{C II})$ (cm $^{-2}$)	$\log N(\text{N I})$ (cm $^{-2}$)
1a	3.0672594	5 700	2.4	20.21	−4.587 ^a	14.32	12.86
	± 0.0000005	$\pm 1\,000$	± 0.2	± 0.06	± 0.012	± 0.06	± 0.06
1b	3.0672591	4 000	8.8	20.16	−4.587 ^a	... ^b	... ^b
	± 0.0000018	± 500	± 0.5	± 0.07	± 0.012		
2	3.06702	13 100	8.5	17.4	−4.587 ^a	12.30	... ^b
	± 0.00001	$\pm 1\,500$	± 1.4	± 0.03	± 0.012	± 0.15	

Comp.	$\log N(\text{O I})$ (cm $^{-2}$)	$\log N(\text{Al II})$ (cm $^{-2}$)	$\log N(\text{Si II})$ (cm $^{-2}$)	$\log N(\text{Si III})$ (cm $^{-2}$)	$\log N(\text{S II})$ (cm $^{-2}$)	$\log N(\text{Fe II})$ (cm $^{-2}$)
1a	14.85	11.93	13.44	11.87	13.00	13.09
	± 0.02	± 0.03	± 0.02	± 0.04	± 0.08	± 0.02
1b	... ^b	... ^b	... ^b	... ^b	... ^b	... ^b
2	... ^b	... ^b	11.53 ± 0.06	11.80 ± 0.06	... ^b	... ^b

^aForced to be the same for all components.

^bAbsorption is undetected for this ion in this component.




FIG. 4.— A selection of metal absorption lines for the DLA at $z_{\text{abs}} = 3.067259$ toward the QSO J1358+6522 (black histogram, cf. Cooke, Pettini, & Murphy 2012). The best-fitting model is shown as the solid red line, while fitted blends are shown as the thick orange lines. In the top panels, the red tick marks above the spectrum indicate the location of the two absorption components, while the bottom panels only display the tick mark for the main absorption component (i.e. component 1a+1b). The long-dashed blue and dashed green lines show the continuum and zero level respectively. The absorption feature near +25 km s $^{-1}$ in the Si II $\lambda 1260$ panel is Fe II $\lambda 1260.5$.

Figure 1 of Cooke, Pettini, & Murphy (2012), albeit now with a higher S/N ratio, we only present a small selection of them in Figure 4 to avoid repetition]. The best-fitting chi-squared value for this fit is also provided for completeness¹⁵.

Returning to Table 1, it can be seen that we found it necessary to separate the main absorption into two separate components, labeled 1a and 1b in the Table. A statistically acceptable fit¹⁶ to the metal, D I and H I lines could not be achieved with a single absorbing cloud in which the turbulent broadening is the same for all species and the thermal broadening is proportional to the square root of the ion mass (i.e. $b_{\text{th}}^2 = 2KT/m$, where K is the Boltzmann constant). The main absorption component of this DLA appears to consist of two ‘clouds’ with very similar redshifts, temperatures and H I column densities, but with significantly different turbulence parameters (see Table 1). The turbulent broadening for component 1a is bounded by the metal-lines, whereas the thermal broadening is bounded by the relatively narrow H I line profiles. This combination of turbulent and thermal broadening is unable to reproduce the observed widths of the strongest D I lines, which require an additional component with a larger contribution of turbulent broadening. Surprisingly, metal absorption is only seen in the low-turbulence cloud (component 1a); component 1b has less than 1/100 of the metallicity of component 1a. We propose three possible interpretations for this unusual finding: (1) there exists an essentially pristine cloud of gas near the very metal-poor DLA; (2) the metals see a different potential to D I and H I (i.e. the metals have not been entirely mixed into the H I); (3) the assumption of a constant kinetic temperature for all elements through the entire sightline is incorrect.

While any of these explanations is interesting in its own right, we stress here that *the choice of the cloud model does not affect the derived value of D/H* in this system, for the reasons discussed in Section 3.2. Provided that there exists a series of optically thin D I lines, and H I Ly α exhibits damping wings (as indeed is the case for the DLA towards J1358+6522—see Figures 2 and 3), the derived total D I and H I column densities will be independent of the cloud model. We therefore only require that the model provides a good fit to the data.

The final, best-fitting value of the deuterium abundance for the DLA towards J1358+6522 is (expressed as a logarithmic and linear quantity):

$$\log (\text{D I}/\text{H I}) = -4.588 \pm 0.012 \quad (1)$$

$$10^5 \text{D I}/\text{H I} = 2.58 \pm 0.07 \quad (2)$$

where the quoted error term includes the random (observational) error in addition to the systematic uncertainties that were marginalized over during the fitting process.

4. THE PRIMORDIAL ABUNDANCE OF DEUTERIUM FROM THE PRECISION SAMPLE

¹⁵ We caution that the quoted chi-squared value is likely underestimated in our analysis because: (1) there is some degree of correlation between neighbouring pixels that is not accounted for in the error spectrum, and (2) the continuum regions selected tend to have lower fluctuations about the mean than average.

¹⁶ The addition of an extra absorption component (i.e. three components as opposed to two, and four additional free parameters) reduces the minimum chi-squared value by $\Delta\chi_{\text{min}}^2 \simeq 660$, which is *highly* significant (see e.g. Lampton, Margon, & Bowyer 1976).

Including the new D/H case reported herein, there are just five metal-poor absorption line systems in which a *precise* measure of the primordial abundance of deuterium can be obtained, based on the criteria outlined in Section 2.2. Relevant details of these systems are collected in Table 2. The four DLAs from the literature were all reanalyzed in an identical manner to that described above (Section 3) for J1358+6522. In particular, we adopted the same blind analysis strategy and marginalized over the important systematic uncertainties. We refer to this sample of five high quality measurements as the *Precision Sample*.

In Table 2, we provide a measure of the *total* H I column density, along with the associated error. Many of our systems contain more than one component in H I and the column density estimates for these multiple components are correlated with one another. To calculate the error on the total H I column density, we have drawn 10 000 realizations of the component column densities from the covariance matrix. We then calculated the total column density for each realization; in Table 2, we provide the mean and 1σ error derived from this Monte Carlo analysis.

We consider the five measures of D I/H I in these DLAs as five independent determinations of the primordial abundance of deuterium, $(\text{D}/\text{H})_{\text{p}}$, for the following reasons: (1) we are not aware of any physical mechanism that would alter the ionization balance of D compared to H. Thus, to our knowledge, $\text{D I}/\text{H I} \equiv \text{D}/\text{H}$; (2) the degree of astration of D (i.e. its destruction when gas is turned into stars) is expected to be negligible at the low metallicities ($[\text{O}/\text{H}] < -1.5$) of the DLAs considered here (e.g., see Figure 2 of Romano et al. 2006); thus $(\text{D}/\text{H})_{\text{DLA}} = (\text{D}/\text{H})_{\text{p}}$. (3) the lack of dust in metal-poor DLAs makes it extremely unlikely that selective depletion of D onto grains occurs in the cases considered here [it has been proposed that such a mechanism may be responsible for the local variations in $(\text{D}/\text{H})_{\text{ISM}}$ —see Linsky et al. 2006]; (4) the five DLAs sample entirely independent sites in the distant Universe.

As can be seen from Table 2 and Figure 5, the five measures of D/H in the Precision Sample are in very good mutual agreement and the dispersion of the measurements is consistent with the errors estimated with our improved analysis. A χ^2 test indeed confirms that the five measurements are consistent within 2σ of being drawn from a single value of D/H. We can therefore combine the five independent determinations of $(\text{D}/\text{H})_{\text{DLA}}$ to deduce the weighted mean value of the primordial abundance of deuterium:

$$\log (\text{D}/\text{H})_{\text{p}} = -4.597 \pm 0.006 \quad (3)$$

$$10^5 (\text{D}/\text{H})_{\text{p}} = 2.53 \pm 0.04 \quad (4)$$

This value of $(\text{D}/\text{H})_{\text{p}}$ is not markedly different from other recent estimates (Pettini et al. 2008b; Fumagalli, O’Meara, & Prochaska 2011; Pettini & Cooke 2012), but its precision is significantly better than achieved in earlier papers that considered a more heterogeneous set of $(\text{D}/\text{H})_{\text{DLA}}$ determinations. For completeness, we have recalculated the weighted mean for all the known D/H measurements listed in Table 2 of Pettini & Cooke (2012), after updating the D/H values of the systems we have reanalyzed here. The resulting weighted mean value of the primordial deuterium abundance is $(\text{D}/\text{H})_{\text{p}} = -4.596 \pm 0.006$. This compares very well with the value derived from the Precision Sample (eqs. 3 and 4). Perhaps this is not surprising, since




FIG. 5.— Values of D/H for the Precision Sample of DLA measurements analyzed in this paper. The orange point represents the new case reported here (J1358+6522). The left and right panels show respectively the D/H measures as a function of the DLA oxygen abundance and H I column density. The dark and light green bands are the 1σ and 2σ determinations of $\Omega_{b,0} h^2$ from the analysis of the CMB temperature fluctuations recorded by the *Planck* satellite (Planck Collaboration 2013) assuming the standard model of physics. The conversion from D/H to $\Omega_{b,0} h^2$ is given by eqs. 5 and 6.

TABLE 2
THE PRECISION SAMPLE OF D/H MEASUREMENTS IN QSO ABSORPTION LINE SYSTEMS

QSO	z_{em}	z_{abs}	$[O/H]^a$	Literature		This work		Ref. ^b
				$\log N(\text{H I})$ (cm^{-2})	$\log(D/H)$	$\log N(\text{H I})$ (cm^{-2})	$\log(D/H)$	
HS 0105+1619	2.652	2.53651	-1.77	19.42 ± 0.01	-4.60 ± 0.04	19.426 ± 0.006	-4.589 ± 0.026	1, 2
Q0913+072	2.785	2.61829	-2.40	20.34 ± 0.04	-4.56 ± 0.04	20.312 ± 0.008	-4.597 ± 0.018	1, 3, 4
SDSS J1358+6522	3.173	3.06726	-2.33	20.495 ± 0.008	-4.588 ± 0.012	1
SDSS J1419+0829	3.030	3.04973	-1.92	20.391 ± 0.008	-4.596 ± 0.009	20.392 ± 0.003	-4.601 ± 0.009	1, 5, 6
SDSS J1558-0031	2.823	2.70242	-1.55	20.67 ± 0.05	-4.48 ± 0.06	20.75 ± 0.03	-4.619 ± 0.026	1, 7

^aWe adopt the solar value $\log(O/H)_{\odot} + 12 = 8.69$ (Asplund et al. 2009).

^bReferences – (1) This work, (2) O’Meara et al. (2001), (3) Pettini et al. (2008a), (4) Pettini et al. (2008b), (5) Pettini & Cooke (2012), (6) Cooke et al. (2011), (7) O’Meara et al. (2006).

the literature systems that did not meet our selection criteria (see Section 2.2.1) have larger uncertainties, and thus their contribution to the weighted mean value of D/H is relatively low.

4.1. The Cosmic Density of Baryons

Using the most up-to-date calculations of the network of nuclear reactions involved in BBN, the primordial abundance of deuterium is related to the cosmic density of baryons (in units of the critical density), $\Omega_{b,0}$, via the following relations (Steigman 2012; G. Steigman 2013, private communication):

$$(D/H)_p = 2.55 \times 10^{-5} (6/\eta_D)^{1.6} \times (1 \pm 0.03) \quad (5)$$

$$\eta_D = \eta_{10} - 6(S - 1) + 5\xi/4 \quad (6)$$

where $\eta_{10} = 273.9 \Omega_{b,0} h^2$, $S = [1 + 7(N_{\text{eff}} - 3.046)/43]^{1/2}$ is the expansion factor and ξ is the neutrino degeneracy parameter (related to the lepton asymmetry by Equation 14 from Steigman 2012). The rightmost term in eq. 5 represents the current 3% uncertainty in the conversion of $(D/H)_p$ to η_D due to the uncertainties in the relevant nuclear reactions rates (see Section 4.2). For the standard model, $N_{\text{eff}} \approx 3.046$ and $\xi = 0$. In this case, the Precision Sample of D/H measurements implies a cosmic density of baryons:

$$100 \Omega_{b,0} h^2(\text{BBN}) = 2.202 \pm 0.020 \text{ (random)} \pm 0.041 \text{ (systematic)} \quad (7)$$

where we have decoupled the error terms from our measurement (i.e. the random error term) and the systematic uncertainty in converting the D abundance into the baryon density parameter.

As can be seen from Figure 5, this value of $\Omega_{b,0} h^2$ is in excellent agreement with that derived from the analysis of the CMB temperature fluctuations measured by the *Planck* satellite (Planck Collaboration 2013):

$$100 \Omega_{b,0} h^2(\text{CMB}) = 2.205 \pm 0.028. \quad (8)$$

4.2. The Current Limitation

In the era of high-precision cosmology, we feel that it is important to highlight the main limitations affecting the use of $(D/H)_p$ in the estimation of cosmological parameters. As can be seen from eq. 7, the main source of error is in the conversion of $(D/H)_p$ to the baryon density parameter (η_D , and hence $\Omega_{b,0} h^2$). In large part, this systematic uncertainty is due to the relative paucity of experimental measures for several nuclear cross-sections that are important in the network of BBN reactions, particularly deuteron–deuteron reactions and the $d(p, \gamma)^3\text{He}$ reaction rate at the relevant energies (Fiorentini et al. 1998; Nollett & Burles 2000; Cyburt 2004; Serpico et al. 2004). Since these studies, estimates for the deuteron–deuteron reaction cross-sections (Leonard et al. 2006) have improved and their contribution to the error budget has been reduced. The main lingering concern involves the reaction rate $d(p, \gamma)^3\text{He}$, for which only a single reliable dataset

of the S -factor is currently available in the relevant energy range (Ma et al. 1997).¹⁷ The concern for $d(p, \gamma)^3\text{He}$ is made worse by the fact that the theoretical and experimental values of the S -factor do not agree. This paucity of data, in addition to the difficulties of obtaining an accurate and precise measure of $d(p, \gamma)^3\text{He}$ at BBN energies, led Nollett & Holder (2011) to adopt a theoretical curve for the S -factor between 50–500 keV. This has resulted in the improved conversion given in eq. 5.

It is worth pointing out here that a further reduction by a modest factor of two in the uncertainty of the conversion from $(D/H)_p$ to η_D would make the precision of $\Omega_{b,0} h^2$ (BBN) from the DLA Precision Sample comparable to that of $\Omega_{b,0} h^2$ (CMB) achieved by the *Planck* mission (cf. eqs. 7 and 8).

5. BEYOND THE STANDARD MODEL

In this section we combine the results presented here with those of the CMB analysis by the *Planck* Collaboration to place bounds on some parameters denoting new physics beyond the standard model of cosmology and particle physics. In particular, we use the Markov-Chain Monte Carlo chains from the combined analysis of the *Planck* temperature fluctuations (Planck Collaboration 2013), the low-multipole *WMAP* polarization data ($l < 23$; Bennett et al. 2012), and high-resolution CMB data (see Planck Collaboration 2013 for an appropriate list of references). Throughout, we refer to this combined dataset as *Planck*+WP+highL, for consistency with the work by the Planck Collaboration (2013). In what follows, we have assumed that N_{eff} and the baryon-to-photon ratio remained unchanged from the epoch of BBN to the epoch of recombination.

5.1. Combined CMB+BBN Measure of N_{eff} and $\Omega_{b,0} h^2$

It has long been known that the mass fraction of ^4He , Y_p , is potentially a very sensitive probe of additional light degrees of freedom (Steigman, Schramm, & Gunn 1977). Unfortunately, systematic uncertainties in the determination of Y_p have limited its use as an effective probe of N_{eff} (see Figure 8 of Steigman 2012). However, Nollett & Holder (2011) (see also Cyburt 2004) have recently highlighted the potential of using precise measurements of the primordial deuterium abundance in conjunction with observations of the CMB to place a strong, independent bound on N_{eff} .

In the left panel of Figure 6, we show the current 1σ and 2σ confidence contours for N_{eff} and $\Omega_{b,0} h^2$ derived from the *Planck*+WP+highL CMB analysis¹⁸ (green), and from the BBN-derived $(D/H)_p$ reported here (blue). The combined confidence bounds are overlaid as red contours. In what follows, it is instructive to recall that the CMB-only bounds are (Planck Collaboration 2013):

$$100 \Omega_{b,0} h^2 = 2.23 \pm 0.04 \quad (9)$$

$$N_{\text{eff}} = 3.36 \pm 0.34. \quad (10)$$

(Note that solving simultaneously for $\Omega_{b,0} h^2$ and N_{eff} leads to a slightly different best-fitting value of $\Omega_{b,0} h^2$ than that obtained for the standard model; cf. eqs. 8 and 9). For compari-

son, from the joint BBN+CMB analysis we deduce:

$$100 \Omega_{b,0} h^2 = 2.23 \pm 0.04 \quad (11)$$

$$N_{\text{eff}} = 3.28 \pm 0.28. \quad (12)$$

Thus, combining $(D/H)_p$ with the CMB does not significantly change the uncertainty in $\Omega_{b,0} h^2$, but does reduce the error on N_{eff} by ~ 20 per cent. The uncertainty on N_{eff} could be reduced further by an improvement in the cross-section of the $d(p, \gamma)^3\text{He}$ (see right panel of Figure 6, and Section 4.2). Based on the current bound on N_{eff} from CMB+ $(D/H)_p$, we can nevertheless rule out the existence of an additional (sterile) neutrino (i.e. $N_{\text{eff}} = 4.046$) at 99.3% confidence (i.e. $\sim 2.7\sigma$), provided that N_{eff} and η_{10} remained unchanged between BBN and recombination. However, as noted recently by Steigman (2013), if the CMB photons are heated after the neutrinos have decoupled [for example, by a weakly interacting massive particle that annihilates to photons], N_{eff} will be less than 3.046 for three standard model neutrinos; a sterile neutrino can in principle exist even when $N_{\text{eff}} < 4.046$.

Looking to the future, Y_p has contours that are almost orthogonal to those of the CMB and $(D/H)_p$ (see e.g. Steigman 2007). Thus, measures of Y_p that are not limited by systematic uncertainties could potentially provide a very strong bound, when combined with $(D/H)_p$, on the number of equivalent neutrinos during the epoch of BBN, independently of CMB observations.

Following improvements in the He I emissivity calculations (Porter et al. 2012, 2013), there have been two recent reassessments of Y_p , by Izotov, Stasinska, & Guseva (2013) and by Aver et al. (2013) respectively. Izotov, Stasinska, & Guseva (2013) find $Y_p = 0.253 \pm 0.003$; this value includes a small correction of -0.001 to reflect the fact that their *Cloudy* modelling overpredicts Y_p by this amount at the lowest metallicities — see their Figure 7(a). Aver et al. (2013) find $Y_p = 0.254 \pm 0.004$ from the average of all the low metallicity H II regions in their sample. Thus, the values deduced by these two teams are in good mutual agreement; in the analysis that follows we adopt $Y_p = 0.253 \pm 0.003$ from Izotov, Stasinska, & Guseva (2013). Using the following conversion relation for Y_p (Steigman 2012; G. Steigman 2013, private communication):

$$Y_p = 0.2469 \pm 0.0006 + 0.0016 (\eta_{\text{He}} - 6) \quad (13)$$

$$\eta_{\text{He}} = \eta_{10} + 100(S - 1) - 575\xi/4 \quad (14)$$

combined with the Izotov, Stasinska, & Guseva (2013) measure of Y_p , we derive the following BBN-only bound on the baryon density and the effective number of neutrino species:

$$100 \Omega_{b,0} h^2 = 2.28 \pm 0.05 \quad (15)$$

$$N_{\text{eff}} = 3.50 \pm 0.20. \quad (16)$$

The corresponding contours are shown in Figure 7. Thus, it appears that even with the most recent reappraisals of the primordial abundance of ^4He by Izotov, Stasinska, & Guseva (2013) and Aver et al. (2013), there is better agreement (within the standard model) between $(D/H)_p$ and the CMB, than between $(D/H)_p$ and Y_p .

5.2. Deuterium and the Lepton Asymmetry

In the past, the primordial deuterium abundance has been commonly used as a tool for measuring the present-day universal density of baryons (see e.g. Steigman 2007), and more

¹⁷ To avoid possible confusion caused by the unfortunate nomenclature, we point out that the S -factor in question is directly related to the reaction cross-section as a function of energy, $\sigma(E)$. It has nothing to do with the expansion factor S in eq. 6.

¹⁸ We used the base cosmology set with N_{eff} added as a free parameter (see Section 6.4.4 of Planck Collaboration 2013).




FIG. 6.— The 1σ and 2σ confidence contours (dark and light shades respectively) for N_{eff} and $\Omega_{\text{b},0} h^2$ derived from the primordial deuterium abundance (blue), the CMB (green), and the combined confidence contours (red). The left panel illustrates the current situation, while the right panel shows the effect of reducing the uncertainty in the conversion from $(\text{D}/\text{H})_{\text{p}}$ to $\Omega_{\text{b},0} h^2$ by a factor of two (see discussion in Section 4.2). Dashed and dotted lines indicate the hidden contour lines for BBN and CMB bounds respectively.




FIG. 7.— The 1σ and 2σ confidence contours (dark and light shades respectively) for N_{eff} and $\Omega_{\text{b},0} h^2$ derived from the primordial deuterium abundance (blue), the primordial He mass fraction (green), and the combined confidence contours (red). Dashed and dotted lines indicate the hidden contour lines for $(\text{D}/\text{H})_{\text{p}}$ and Y_{p} bounds respectively.

recently as a probe of the effective number of neutrino families (Cyburt 2004; Nollett & Holder 2011; Pettini & Cooke 2012, see also Section 5.1). Here, we demonstrate that *precise* measures of the primordial deuterium abundance (in combination with the CMB) can also be used to estimate the neutrino degeneracy parameter, ξ , which is related to the lepton asymmetry by Equation 14 from Steigman (2012).

Steigman (2012) recently suggested that combined estimates for $(\text{D}/\text{H})_{\text{p}}$, Y_{p} , and a measure of N_{eff} from the CMB, can provide interesting limits on the neutrino degeneracy parameter ($\xi \leq 0.079$, 2σ ; see also, Serpico & Raffelt 2005; Popa & Vasile 2008; and Simha & Steigman 2008). By combining $(\text{D}/\text{H})_{\text{p}}$ and Y_{p} , this approach effectively removes the dependence on $\Omega_{\text{b},0} h^2$. Using the conversion relations for $(\text{D}/\text{H})_{\text{p}}$ and Y_{p} (eqs. 5–6 and 13–14) and the current best determination of Y_{p} (0.253 ± 0.003 ; Izotov, Stasinska, & Guseva 2013), in addition to the *Planck*+WP+highL¹⁹ constraint on N_{eff} and the precise determination of $(\text{D}/\text{H})_{\text{p}}$ reported here, we derive a 2σ upper limit on the neutrino degeneracy parameter, $|\xi| \leq 0.064$, based on the approach by Steigman (2012).

We propose that an equally powerful technique for estimat-

ing ξ does *not* involve removing the dependence on $\Omega_{\text{b},0} h^2$ by combining $(\text{D}/\text{H})_{\text{p}}$ and Y_{p} , as in Steigman (2012). Instead, one can obtain a measure of both $\Omega_{\text{b},0} h^2$ and N_{eff} from the CMB, and use *either* $(\text{D}/\text{H})_{\text{p}}$ or Y_{p} to obtain two separate measures of ξ . This has the clear advantage of decoupling $(\text{D}/\text{H})_{\text{p}}$ and Y_{p} ; any systematic biases in either of these two values could potentially bias the measure of ξ . Separating $(\text{D}/\text{H})_{\text{p}}$ and Y_{p} also allows one to check that the two estimates agree with one another.

Our calculation involved a Monte Carlo technique, whereby we generated random values from the Gaussian-distributed primordial D/H abundance measurements, whilst simultaneously drawing random values from the (correlated) distribution between $\Omega_{\text{b},0} h^2$ and N_{eff} from the *Planck*+WP+highL CMB data (Planck Collaboration 2013)²⁰. Using Equation 19 from Steigman (2012, equivalent to eq. 6 here), we find $\xi_{\text{D}} = +0.05 \pm 0.13$ for $(\text{D}/\text{H})_{\text{p}}$, leading to a 2σ upper limit of $|\xi_{\text{D}}| \leq 0.31$.

With the technique outlined above, we have also computed the neutrino degeneracy parameter from the current observational bound on Y_{p} . For this calculation, we have used the MCMC chains from the *Planck*+WP+highL CMB base cosmology with N_{eff} and Y_{p} added as free parameters. In this case, the CMB distribution was weighted by the observational bound on Y_{p} ($Y_{\text{p}} = 0.253 \pm 0.003$; Izotov, Stasinska, & Guseva 2013). Using Equations 19–20 from Steigman (2012, equivalent to eqs. 6 and 14 here), we find $\xi_{\text{D}} = +0.04 \pm 0.15$ for $(\text{D}/\text{H})_{\text{p}}$ and $\xi_{\text{He}} = -0.010 \pm 0.027$ for Y_{p} . These values translate into corresponding 2σ upper limits $|\xi_{\text{D}}| \leq 0.34$ and $|\xi_{\text{He}}| \leq 0.064$. Combining these two constraints then gives $\xi = -0.008 \pm 0.027$, or $|\xi| \leq 0.062$ (2σ).

Alternatively, if we assume that the effective number of neutrino species is consistent with three standard model neutrinos (i.e. $N_{\text{eff}} \approx 3.046$), we obtain the following BBN-only bound on the neutrino degeneracy parameter by combining $(\text{D}/\text{H})_{\text{p}}$ and Y_{p} , $\xi = -0.026 \pm 0.015$, or $|\xi| \leq 0.056$ (2σ). We therefore conclude that all current estimates of the neutrino degeneracy parameter, and hence the lepton asymmetry, are consistent with the standard model value, $\xi = 0$.

¹⁹ We used the base cosmology set with N_{eff} and Y_{p} added as free parameters (see Section 6.4.5 of Planck Collaboration 2013).

²⁰ Rather than drawing values of $\Omega_{\text{b},0} h^2$ and N_{eff} from the appropriate distribution, we instead used the Markov-Chain Monte Carlo chains provided by the *Planck* science team, which are available at: <http://www.sciops.esa.int/wikiSI/planckpla/index.php?title=CosmologicalParameters&instance=PlanckPublic.PLA>

From the above calculations, it is clear that Y_p is the more sensitive probe of any lepton asymmetry, as is already appreciated. However, $(D/H)_p$ offers an additional bound on ξ that is complementary to that obtained from Y_p . We note further that, if the uncertainty in the conversion of $(D/H)_p$ to η_D could be reduced by a factor of two, the bound on ξ_D would be reduced by 15%, corresponding to a 1σ uncertainty on ξ of ~ 0.11 from the CMB and deuterium measurements alone.

6. SUMMARY AND CONCLUSIONS

We have reported a new precise measurement of the primordial abundance of deuterium in a very metal-poor damped Lyman α system at $z_{\text{abs}} = 3.06726$ towards the QSO SDSS J1358+6522. Furthermore, we have reanalyzed self-consistently all literature systems that meet a set of strict criteria (four cases). These criteria were imposed to identify the systems where both accurate and precise measures of D/H could potentially be obtained. The new system, plus the four from the literature, constitute the *Precision Sample* of DLAs that are best suited for a precise determination of $(D/H)_p$. Our reanalysis was performed blind (to remove human bias), and took advantage of a new software package that we have specifically developed for the measurement of D/H in QSO absorption line systems. From the analysis of this sample, we draw the following conclusions.

(i) The very metal-poor DLA ($[Fe/H] = -2.88 \pm 0.03$) towards SDSS J1358+6522 provides a strong bound on the primordial abundance of deuterium, $(D/H)_p = (2.58 \pm 0.07) \times 10^{-5}$. A weighted mean of the five systems in the Precision Sample gives the most stringent limit on $(D/H)_p$ to date: $(D/H)_p = (2.53 \pm 0.04) \times 10^{-5}$. The corresponding baryon density is $100 \Omega_{b,0} h^2 = 2.202 \pm 0.046$, assuming the standard model of particle physics with three families of neutrinos and no lepton asymmetry. This value is in excellent agreement with $100 \Omega_{b,0} h^2$ (CMB) = 2.205 ± 0.028 deduced from the analysis of *Planck* observations of the CMB.

(ii) The main limitation in using $(D/H)_p$ for cosmological parameter estimation is the conversion of $(D/H)_p$ into the ratio of baryons-to-photons. In particular, modern measurements of the cross-section for the reaction $d(p, \gamma)^3\text{He}$ at energies between 50–500 keV, where there currently exists a paucity of reliable data, are highly desirable. We estimate that a factor of two improvement in the conversion from $(D/H)_p$ to $\Omega_{b,0} h^2$ would provide a measurement of the universal baryon density from D/H as precise as that derived from the published *Planck* data.

(iii) By combining our D/H determination with *Planck* ob-

servations of the CMB, we can place a tight bound on both $\Omega_{b,0} h^2$ and N_{eff} . The best-fitting parameters so derived are $100 \Omega_{b,0} h^2 = 2.23 \pm 0.04$ and $N_{\text{eff}} = 3.28 \pm 0.28$. We therefore rule out the existence of an additional (sterile) neutrino at 99.3% confidence, provided that N_{eff} and η_{10} remained unchanged from BBN to recombination. The uncertainty on N_{eff} could be reduced further by a more accurate set of reaction cross-sections for $d(p, \gamma)^3\text{He}$ and, to a lesser degree, $d + d$.

(iv) For the first time, we have combined the *Planck* CMB results with our measure of $(D/H)_p$ to place a bound on the neutrino degeneracy parameter, $\xi_D = +0.05 \pm 0.13$. When including the current best-estimate of the ^4He mass fraction, Y_p , derived from metal-poor H II regions, the combined bound on the neutrino degeneracy parameter is $\xi = -0.008 \pm 0.027$, or $|\xi| < 0.062$ at 2σ .

We conclude by re-emphasizing that the most metal-poor DLAs are potentially the best environments where both accurate and precise measures of the primordial abundance of deuterium can be obtained. A combined effort to measure anew several important nuclear reaction cross-sections, in addition to dedicated searches for the rare metal-poor DLAs that exhibit resolved deuterium absorption, are the next steps necessary to pin down further the value of $(D/H)_p$. This approach offers a promising and exciting avenue to test for departures from the standard model of cosmology and particle physics.

ACKNOWLEDGMENTS

We are grateful to the staff astronomers at the ESO VLT and Keck Observatory for their assistance with the observations. We are indebted to Gary Steigman for kindly communicating ahead of publication the latest fitting formulae used in this work, and for providing valuable comments on the manuscript. We also thank an anonymous referee who provided valuable comments that improved the presentation of this work. Discussions, advice and help with various aspects of the work described in this paper were provided by Antony Lewis, Jordi Miralda-Escudé, Paolo Molaro, Ken Nollett, Pasquier Noterdaeme, John O'Meara, Jason X. Prochaska, Signe Riemer-Sørensen, Donatella Romano, and John Webb. We thank the Hawaiian people for the opportunity to observe from Mauna Kea; without their hospitality, this work would not have been possible. R. J. C. is partially supported by NSF grant AST-1109447. R. A. J. is supported by an NSF Astronomy and Astrophysics Postdoctoral Fellowship under award AST-1102683. M. T. M. thanks the Australian Research Council for a QEII Research Fellowship (DP0877998) and a Discovery Project grant (DP130100568).

APPENDIX

MODEL FITS TO LITERATURE SYSTEMS

In this Appendix we show portions of the spectra of the metal-poor DLAs from the literature that were reanalyzed in the present work. Further details are given in Section 2.2.

REFERENCES

- Adams T. F., 1976, *A&A*, 50, 461
- Asplund M., Grevesse N., Sauval A. J., Scott P., 2009, *ARA&A*, 47, 481
- Aver E., Olive K. A., Porter R. L., Skillman E. D., 2013, arXiv e-prints, arXiv:1309.0047
- Bennett, C. L., et al., 2012, arXiv e-prints, arXiv:1212.5225
- Burles S., Tytler D., 1998a, *ApJ*, 499, 699
- Burles S., Tytler D., 1998, *ApJ*, 507, 732
- Cooke R., Pettini M., Steidel C. C., Rudie G. C., Nissen P. E., 2011, *MNRAS*, 417, 1534
- Cooke R., Pettini M., Murphy M. T., 2012, *MNRAS*, 425, 347
- Cybur R. H., 2004, *PhRvD*, 70, 023505
- Erni, P., Richter, P., Ledoux, C., & Petitjean, P. 2006, *A&A*, 451, 19
- Fiorentini G., Lisi E., Sarkar S., Villante F. L., 1998, *PhRvD*, 58, 063506
- Fumagalli M., O'Meara J. M., Prochaska J. X., 2011, *Sci*, 334, 1245
- Izotov Y. I., Stasinska G., Guseva N. G., 2013, arXiv, arXiv:1308.2100




FIG. 8.— The data (black histogram) and best-fitting model (continuous red line) for the DLA at $z_{\text{abs}} = 2.536509$ toward the QSO HS0105+1619 (cf. O’Meara et al. 2001). The long-dashed blue line in the top panel marks the best-fitting continuum level near the Ly α absorption line. In the remaining panels, the data have been normalized to the continuum. The minimum χ^2/dof for this fit is 2088.1/3038. The red tick marks above the spectrum indicate the absorption components in this DLA. The green tick marks in the Lyman series panels show the corresponding location of D I absorption. The absorption feature near $\lambda \approx 4267 \text{ \AA}$ in the top panel is Si III $\lambda 1206.5$ at the redshift of the DLA. The feature near $+25 \text{ km s}^{-1}$ in the Si II $\lambda 1260$ panel is Fe II $\lambda 1260.5$.




FIG. 9.— Same as Fig. 8 for the DLA at $z_{\text{abs}} = 2.61829$ toward the QSO Q0913+072 (cf. Pettini et al. 2008b). The minimum χ^2/dof for this fit is 5539.8/4473.




FIG. 10.— Same as Fig. 8 for the DLA at $z_{\text{abs}} = 2.70242$ toward the QSO J1558–0031 (cf. O’Meara et al. 2006). The DLA near 4500 Å is the system where D/H can be measured. The lower redshift system (at $z_{\text{abs}} = 2.62976$) is included in the simultaneous fit to the higher redshift system (see the bottom-right panel for an example Si II line profile in the low-redshift system). The minimum χ^2/dof for this fit is 4754.4/4700. In the top panel, several absorption features from metal lines at $z_{\text{abs}} = 2.70242$ (Si II $\lambda\lambda 1190.4, 1193.3$ near 4407 Å and 4418 Å respectively, N I $\lambda 1200$ and including some additional blends near 4440 Å , and Si II $\lambda 1206.5$ near 4467 Å) are indicated in red. The ‘extra’ absorption in the trough of H I Ly10 is due to H I Ly ϵ from the system at redshift $z_{\text{abs}} = 2.62976$. The absorption feature at +100 km s^{-1} in this same panel corresponds to O I $\lambda 919.6$ at $z_{\text{abs}} = 2.70242$.

- Lampton M., Margon B., Bowyer S., 1976, *ApJ*, 208, 177
- Lee H.-W., (2013) *ApJ*, 772, 123
- Leonard D. S., Karwowski H. J., Brune C. R., Fisher B. M., Ludwig E. J., (2006) *Phys. Rev. C*, 73, 045801
- Linsky, J. L., Draine, B. T., Moos, H. W., et al. 2006, *ApJ*, 647, 1106
- Ma L., et al., 1997, *Phys. Rev. C*, 55, 588
- Markwardt C. B., 2009, *ASPC*, 411, 251
- Meléndez, J., Casagrande, L., Ramírez, I., Asplund, M., & Schuster, W. J. 2010, *A&A*, 515, L3
- Morton, D. C. 2003, *ApJS*, 149, 205
- Nollett K. M., Burles S., 2000, *PhRvD*, 61, 123505
- Nollett K. M., Holder G. P., 2011, arXiv e-prints, arXiv:1112.2683
- Noterdaeme P., et al., 2012, *A&A*, 547, L1
- O’Meara J. M., Tytler D., Kirkman D., Suzuki N., Prochaska J. X., Lubin D., Wolfe A. M., 2001, *ApJ*, 552, 718
- O’Meara J. M., Burles S., Prochaska J. X., Prochter G. E., Bernstein R. A., Burgess K. M., 2006, *ApJ*, 649, L61
- Penprase B. E., Prochaska J. X., Sargent W. L. W., Toro-Martinez I., Beeler D. J., 2010, *ApJ*, 721, 1
- Pettini M. & Cooke R., 2012, *MNRAS*, 425, 2477
- Pettini M., Zych B. J., Steidel C. C., Chaffee F. H., 2008a, *MNRAS*, 385, 2011
- Pettini M., Zych B. J., Murphy M. T., Lewis A., Steidel C. C., 2008b, *MNRAS*, 391, 1499
- Planck Collaboration, 2013, arXiv e-prints, arXiv:1303.5076
- Porter R. L., Ferland G. J., Storey P. J., Detisch M. J., 2012, *MNRAS*, 425, L28
- Porter R. L., Ferland G. J., Storey P. J., Detisch M. J., 2013, *MNRAS*, 433, L89
- Popa L. A., Vasile A., 2008, *JCAP*, 6, 28
- Prochaska J. X., Wolfe A. M., 2009, *ApJ*, 696, 1543
- Romano, D., Tosi, M., Chiappini, C., Matteucci, F., 2006, *MNRAS*, 369, 295
- Sbordone L., et al., 2010, *A&A*, 522, A26
- Serpico P. D., Esposito S., Iocco F., Mangano G., Miele G., Pisanti O., 2004, *JCAP*, 12, 10
- Serpico P. D., Raffelt G. G., 2005, *PhRvD*, 71, 127301
- Simha V., Steigman G., 2008, *JCAP*, 8, 11
- Spite F., Spite M., 1982, *A&A*, 115, 357
- Steigman G., 2001, in *Proc. of the STScI Symp. Ser. 15, The Dark Universe: Matter, energy and gravity*, ed. M. Livio (Baltimore, MD: Cambridge University Press), 46
- Steigman G., 2007, *ARNPS*, 57, 463
- Steigman G., 2012, arXiv e-prints, arXiv:1208.0032
- Steigman G., 2013, *PhRvD*, 87, 103517
- Steigman G., Schramm D. N., Gunn J. E., 1977, *Phys. Lett. B*, 66, 202
- Vogt S. S., et al., 1994, *SPIE*, 2198, 362
- Wagoner R. V., Fowler W. A., Hoyle F., 1967, *ApJ*, 148, 3