

# Heavy Scalar, Vector and Axial-Vector Mesons in Hot and Dense Nuclear Medium

Arvind Kumar\*

*Department of Physics, Dr. B R Ambedkar National Institute  
of Technology Jalandhar, Jalandhar – 144011, Punjab, India*

## Abstract

In this work we shall investigate the mass modifications of scalar mesons ( $D_0, B_0$ ), vector mesons ( $D^*, B^*$ ) and axial-vector mesons ( $D_1, B_1$ ) at finite density and temperature of the nuclear medium. The above mesons are modified in the nuclear medium through the modification of quark and gluon condensates. We shall find the medium modification of quark and gluon condensates within chiral SU(3) model through the medium modification of scalar-isoscalar fields  $\sigma$  and  $\zeta$  at finite density and temperature. These medium modified quark and gluon condensates will further be used through QCD sum rules for the evaluation of in-medium properties of above mentioned scalar, vector and axial vector mesons. We shall also discuss the effects of density and temperature of the nuclear medium on the scattering lengths of above scalar, vector and axial-vector mesons. The study of the medium modifications of above mesons may be helpful for understanding their production rates in heavy-ion collision experiments. The results of present investigations of medium modifications of scalar, vector and axial-vector mesons at finite density and temperature can be verified in the Compressed Baryonic Matter (CBM) experiment of FAIR facility at GSI, Germany.

**Keywords:** Dense hadronic matter, heavy-ion collisions, effective chiral model, QCD sum rules, heavy mesons.

PACS numbers : -14.40.Lb , -14.40.Nd, 13.75.Lb

---

\*Electronic address: iitd.arvind@gmail.com, kumara@nitj.ac.in

## I. INTRODUCTION

The motive behind the heavy-ion collision experiments at different experimental facilities is to explore the different phases of QCD phase diagram. These experiments help us to understand the nuclear matter properties for different values of temperatures and densities. The hadronic matter produced in heavy-ion collisions may undergo different phase transitions e.g. liquid-gas phase transition, the kaon condensation, the restoration of chiral symmetry and may be the formation of quark gluon plasma [1–4]. The Compressed Baryonic Matter (CBM) experiment of the FAIR project at GSI, Germany may explore the phase of hadronic matter at high baryon densities and moderate temperatures. These kind of phases may exist in the compact astrophysical objects e.g. neutron stars. The property of restoration of chiral symmetry is closely related to the medium modifications of hadrons [4]. The medium modifications of Kaons, D mesons and light vector mesons had been studied using different theoretical approaches e.g. chiral model [5–12], QCD sum rules [13–18] and coupled channel approach [19–21]. Due to interactions the properties of hadrons in the medium are found to be different as compared to their free space properties.

The medium modifications of heavy scalar, vector and axial-vector mesons at finite density and temperature of the medium had been studied very rarely [22–25]. In the present investigation we shall study the mass modifications of heavy scalar mesons ( $D_0, B_0$ ), vector mesons ( $D^*, B^*$ ) and axial vector mesons ( $D_1, B_1$ ) at finite densities and temperatures. The study of in-medium properties of scalar, vector and axial-vector mesons will be helpful to understand their experimental production rates. The medium modification of charmed mesons may modify the experimental production of ground state charmonium  $J/\psi$  and the excited charmonium states ( $\psi'$  and  $\chi_c$ ). The charmonium,  $J/\psi$  may be produced due to the decay of the higher charmonium states. However the vacuum threshold value of heavy meson pairs lies above the vacuum mass of the excited charmonium states. Now if these heavy mesons get modified (undergo mass drop in the medium) then the excited charmonium states may decay to the open charmed meson pairs instead of decaying to the ground state charmonium. Thus to understand the production of charmonium states in heavy-ion collisions it is very necessary to study the medium modification of the heavy scalar, vector and axial-vector mesons. The medium modifications of heavy vector mesons may also help us in understanding the dilepton spectra produced in heavy-ion collision experiments [26–28].

The dileptons are considered as interesting probe to study the evolution of matter produced in heavy ion collision experiments as they do not undergo strong interactions in the medium. In ref. [29] the production of open charm and charmonium in hot hadronic medium had been investigated using the statistical hadronization model at SPS/FAIR energies. In this work it was observed that the medium modifications of charmed hadrons do not lead to appreciable changes in cross-section for D mesons production. This is because of large charm quark mass and different times scales for charm quark and charm hadron production. However, the charmonia yield is effected appreciably due to in-medium modifications.

The properties of scalar charm resonances  $D_{s0}(2317)$  and  $D_0(2400)$  and hidden charm resonance,  $X(3700)$  had been studied in ref. [24] using coupled channel approach. In these studies the  $D_{s0}(2317)$  and  $X(3700)$  were found to undergo a width of about 100 and 200 MeV respectively at nuclear matter density. However, for the  $D_0(2400)$  mesons there was already large width of resonance in the free space and the medium effect were found to be weak as compared to  $D_{s0}(2317)$  and  $X(3700)$ . In ref.[17] the mass splitting of  $D$ - $\bar{D}$  and  $B$ - $\bar{B}$  mesons had been studied using the QCD sum rules in the cold nuclear matter and the calculated values of mass splitting at nuclear saturation density were 60 and 130 MeV respectively. The Borel transformed QCD sum rules had also been used to study the properties of pseudoscalar  $D$  mesons [18] and vector mesons,  $\rho$ ,  $\omega$  and  $\phi$  [16]. The properties of the scalar mesons ( $D_0, B_0$ ) in the cold nuclear matter using QCD sum rules have been investigated in ref. [22]. The vectors mesons ( $D^*, B^*$ ) and axial vector mesons ( $D_1, B_1$ ) had also been studied using QCD sum rules in cold nuclear matter in ref. [23]. Note that in ref. [22] and [23] the properties of the meson were investigated at zero temperature and at normal nuclear matter density. However in the present investigation we shall find the in-medium masses of the scalar ( $D_0, B_0$ ) and vector ( $D^*, B^*$ ) and axial vector ( $D_1, B_1$ ) mesons at finite temperatures as well as at the densities greater than the nuclear saturation density.

In the present work to investigate the properties of scalar, vector and axial-vector mesons we shall use the QCD sum rules and chiral SU(3) model [5]. Within QCD sum rules, the in-medium properties of mesons are related to the in-medium properties of quark and gluon condensates. We shall investigate the in-medium properties of quark and gluon condensates using the chiral SU(3) model. Using chiral SU(3) model we shall find the values of quark and gluon condensates at finite values of temperatures and baryonic densities. These values of condensates will further be used to find the medium modification of mesons using QCD sum

rules. The chiral SU(3) model along with QCD sum rules had been used in the literature to investigate the in-medium modification of the charmonium states  $J/\psi$  and  $\eta_c$  [30].

The present article is organized as follows: In section II we shall give a brief review of chiral SU(3) model. Then in section III we shall discuss that how we will evaluate the in-medium modifications of the scalar, vector and axial-vector mesons within QCD sum rules and using the properties of quark and gluon condensates as evaluated in the chiral SU(3) model. In section IV we shall discuss the results of the present investigation and finally in section V we shall give a brief summary of present work.

## II. CHIRAL SU(3) MODEL

In this section we shall briefly review the chiral SU(3) model used in the present investigation for the in-medium properties of heavy mesons. The chiral SU(3) model is based on the broken scale invariance and non-linear realization of chiral symmetry [31–35]. The model involve the Lagrangian densities describing e.g. kinetic energy terms, baryon-meson interactions, self interactions of scalar mesons, vector mesons, symmetry breaking terms and also the scale invariance breaking logarithmic potential terms.

For the investigation of hadron properties at finite temperature and densities we use the mean field approximation. Under this approximation all the meson fields are treated as classical fields and only the scalar and the vector fields contribute to the baryon-meson interactions. From the interaction Lagrangian densities, using the mean-field approximation, we derive the equations of motions for the scalar fields  $\sigma$  and  $\zeta$  and the dilaton field,  $\chi$  in isospin symmetric nuclear medium. We solve these coupled equations to obtain the density and temperature dependence of scalar fields  $\sigma$  and  $\zeta$  and the dilaton field,  $\chi$  in isospin symmetric nuclear medium [11]. The concept of broken scale invariance leading to the trace anomaly in (massless) QCD,  $\theta_\mu^\mu = \frac{\beta_{QCD}}{2g} G_{\mu\nu}^a G^{\mu\nu a}$ , where  $G_{\mu\nu}^a$  is the gluon field strength tensor of QCD, is simulated in the effective Lagrangian at tree level [36] through the introduction of the scale breaking terms [11]. Within chiral SU(3) model the scale breaking terms are written in terms of the dilaton field  $\chi$  and also the scalar fields  $\sigma$  and  $\zeta$ . From this we obtain the energy momentum tensor and this is compared with the energy momentum tensor of QCD which is written in terms of gluon condensates. In this way we extract the value of gluon condensates in terms of the scalar fields  $\sigma$  and  $\zeta$  and the dilaton

field,  $\chi$  and is given by the following equation [11],

$$\left\langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{a\mu\nu} \right\rangle = \frac{8}{9} \left[ (1-d)\chi^4 + \left( \frac{\chi}{\chi_0} \right)^2 \left( m_\pi^2 f_\pi \sigma + (\sqrt{2} m_K^2 f_K - \frac{1}{\sqrt{2}} m_\pi^2 f_\pi) \zeta \right) \right], \quad (1)$$

where the value of parameter  $d$  is 0.064 [9],  $m_\pi$  and  $m_K$  denote the masses of pions and kaons and have values 139 and 498 MeV respectively.  $f_\pi$  and  $f_K$  are the decay constants having values 93.3 and 122 MeV respectively. The symbols  $\sigma$ ,  $\zeta$  and  $\chi$  denote the non-strange scalar-isoscalar field, strange scalar-isoscalar field and the dilaton field respectively.  $\chi_0$  denotes the value of the dilaton field in vacuum. The vacuum values of  $\sigma$ ,  $\zeta$  and  $\chi$  are -93.3, -106.6 and 409.8 MeV respectively. Note that in above equation the gluon condensate is written considering finite quark masses. If we have massless QCD, then only first term written in terms of dilaton field  $\chi$  contributes to the gluon condensates. Using above equation we obtain the values of scalar gluon condensates at different values of densities and temperatures of the nuclear medium.

### III. QCD SUM RULES FOR SCALAR ( $D_0, B_0$ ), VECTOR ( $D^*, B^*$ ) AND AXIAL VECTOR ( $D_1, B_1$ ) MESONS

In this section we shall discuss the QCD sum rules [22, 23] which will be used later along with the chiral SU(3) model for the evaluation of in-medium properties of scalar, vector and axial vector mesons. To find the mass modification of above discussed heavy mesons we shall use the two-point correlation function  $\Pi_{\mu\nu}(q)$ ,

$$\Pi_{\mu\nu}(q) = i \int d^4x e^{iq \cdot x} \langle T \{ J_\mu(x) J_\nu^\dagger(0) \} \rangle_{\rho_B, T}. \quad (2)$$

In above equation  $J_\mu(x)$  denotes the isospin averaged current,  $x = x^\mu = (x^0, \mathbf{x})$  is the four coordinate,  $q = q^\mu = (q^0, \mathbf{q})$  is four momentum and  $T$  denotes the time ordered operation on the product of quantities in the brackets. From above definition it is clear that the two point correlation function is actually a Fourier transform of the expectation value of the time ordered product of two currents. The two-point correlation function for the scalar mesons is defined as,

$$\Pi(q) = i \int d^4x e^{iq \cdot x} \langle T \{ J(x) J^\dagger(0) \} \rangle_{\rho_B, T}. \quad (3)$$

For the scalar, vector and axial vector mesons isospin average currents are given by the expressions

$$J(x) = J^\dagger(x) = \frac{\bar{c}(x)q(x) + \bar{q}(x)c(x)}{2}, \quad (4)$$

$$J_\mu(x) = J_\mu^\dagger(x) = \frac{\bar{c}(x)\gamma_\mu q(x) + \bar{q}(x)\gamma_\mu c(x)}{2}, \quad (5)$$

and

$$J_{5\mu}(x) = J_{5\mu}^\dagger(x) = \frac{\bar{c}(x)\gamma_\mu\gamma_5 q(x) + \bar{q}(x)\gamma_\mu\gamma_5 c(x)}{2}, \quad (6)$$

respectively. Note that in above equations  $q$  denotes the light  $u$  or  $d$  quark whereas  $c$  denotes the heavy charm quark. Note that in the present work, instead of considering the mass splitting between particles and antiparticles, we emphasize on the mass shift of iso-doublet  $D$  and  $B$  mesons as a whole and therefore we consider the average in the definitions of scalar, vector and iso-vector currents which is referred as centroid [18]. To find the mass splitting of particles and antiparticles in the nuclear medium one has to consider the even and odd part of QCD sum rules [17]. For example, in ref. [17] the mass splitting between pseudoscalar  $D$  and  $\bar{D}$  mesons was investigated using the even and odd QCD sum rules whereas in [18, 22, 23] the mass-shift of  $D$  mesons was investigated under centroid approximation.

The two point correlation function can be decomposed into the vacuum part, a static one-nucleon part and pion bath contribution i.e. we can write

$$\Pi_{\mu\nu}(q) = \Pi_{\mu\nu}^0(q) + \frac{\rho_B}{2M_N} T_{\mu\nu}^N(q) + \Pi_{\mu\nu}^{P.B.}(q), \quad (7)$$

where

$$T_{\mu\nu}^N(\omega, \mathbf{q}) = i \int d^4x e^{iq \cdot x} \langle N(p) | T \{ J_\mu(x) J_\nu^\dagger(0) \} | N(p) \rangle. \quad (8)$$

In above equation  $|N(p)\rangle$  denotes the isospin and spin averaged static nucleon state with the four-momentum  $p = (M_N, 0)$ . The state is normalized as  $\langle N(\mathbf{p}) | N(\mathbf{p}') \rangle = (2\pi)^3 2p_0 \delta^3(\mathbf{p} - \mathbf{p}')$ . The third term,  $\Pi_{\mu\nu}^{P.B.}(q)$  in equation (7) gives the contribution from pion bath at finite

temperature. Note that in the present work instead of considering the contribution of pion bath the effects of finite temperature of the nuclear matter on the properties of  $D$  and  $B$  mesons will be evaluated through the temperature dependence of scalar fields  $\sigma$ ,  $\zeta$  and  $\chi$ . The temperature dependence of scalar fields  $\sigma$ ,  $\zeta$  and  $\chi$  modify the nucleon properties in the medium and these modified nucleons further modify the in-medium properties of  $D$  and  $B$  mesons at finite temperature and density. In literature the properties of kaons and antikaons,  $D$  mesons and charmonium had been studied at finite temperature of the nuclear matter using the above mentioned scalar fields  $\sigma$ ,  $\zeta$  and  $\chi$  [7, 11, 12, 30].

As discussed in Ref. [23], in the limit of the 3-vector  $\mathbf{q} \rightarrow \mathbf{0}$ , the correlation functions  $T_N(\omega, \mathbf{q})$  can be related to the  $D^*N$  and  $D_1N$  scattering T-matrices. Thus we write [23]

$$\begin{aligned}\mathcal{T}_{D^*N}(M_{D^*}, 0) &= 8\pi(M_N + M_{D^*})a_{D^*} \\ \mathcal{T}_{D_1N}(M_{D_1}, 0) &= 8\pi(M_N + M_{D_1})a_{D_1}\end{aligned}\quad (9)$$

In above equation  $a_{D^*}$  and  $a_{D_1}$  are the scattering lengths of  $D^*N$  and  $D_1N$  respectively. Similarly, we can also write the scattering  $T$  matrix corresponding to  $D_0N$  ( $D_0$  is a scalar meson) in terms of the scattering lengths [22],

$$\mathcal{T}_{D_0N}(M_{D_0}, 0) = 8\pi(M_N + M_{D_0})a_{D_0}. \quad (10)$$

Near the pole positions of the scalar, vector and axial vector mesons the phenomenological spectral densities can be parameterized with three unknown parameters  $a, b$  and  $c$  i.e. we write [18, 22, 23]

$$\begin{aligned}\rho(\omega, 0) &= -\frac{f_{D_0/D^*/D_1}^2 M_{D_0/D^*/D_1}^2}{\pi} \text{Im} \left[ \frac{\mathcal{T}_{D_0/D^*/D_1N}(\omega, \mathbf{0})}{(\omega^2 - M_{D_0/D^*/D_1}^2 + i\varepsilon)^2} \right] \\ &+ \dots = a \frac{d}{d\omega^2} \delta'(\omega^2 - M_{D_0/D^*/D_1}^2) + b \delta(\omega^2 - M_{D_0/D^*/D_1}^2) + c \theta(\omega^2 - s_0). \quad (11)\end{aligned}$$

The term denoted by ... represent the continuum contributions. The first term denotes the double-pole term and corresponds to the on-shell effects of the T-matrices,

$$a = -8\pi(M_N + M_{D_0/D^*/D_1})a_{D_0/D^*/D_1}f_{D_0/D^*/D_1}^2 M_{D_0/D^*/D_1}^2. \quad (12)$$

Now we shall write the relation between the scattering length of mesons and their in-medium mass-shift. For this first we note that the shift of squared mass of mesons can be

written in terms of the parameter  $a$  appearing in equation (11) through relation [16],

$$\begin{aligned}\Delta m_{D_0/D^*/D_1}^2 &= \frac{\rho_N}{2M_N} \frac{a}{f_{D_0/D^*/D_1}^2 M_{D_0/D^*/D_1}^2} \\ &= \frac{\rho_N}{2M_N} 8\pi(M_N + M_{D_0/D^*/D_1}) a_{D_0/D^*/D_1},\end{aligned}\quad (13)$$

where in the last term we used equation (12). The mass shift is now defined by the relation

$$\delta M_{D_0/D^*/D_1} = \sqrt{m_{D_0/D^*/D_1}^2 + \Delta m_{D_0/D^*/D_1}^2} - m_{D_0/D^*/D_1}. \quad (14)$$

The second term in equation (11) denotes the single-pole term, and corresponds to the off-shell (i.e.  $\omega^2 \neq M_{D_0/D^*/D_1}^2$ ) effects of the  $T$ -matrices. The third term denotes the continuum term or the remaining effects, where,  $s_0$ , is the continuum threshold. The continuum threshold parameter  $s_0$  define the scale below which the continuum contribution vanishes [37].

It can be observed from equations (13) and (14) that if we want to find the value of mass shift of mesons then we first need to find the value of unknown parameter  $a$ . For this we proceed as follows: we note that in the low energy limit,  $\omega \rightarrow 0$ , the  $T_N(\omega, \mathbf{0})$  is equivalent to the Born term  $T_{D_0/D^*/D_1 N}^{\text{Born}}(\omega, \mathbf{0})$ . We take into account the Born term at the phenomenological side,

$$\begin{aligned}T_N(\omega^2) &= T_{D_0/D^*/D_1 N}^{\text{Born}}(\omega^2) + \frac{a}{(M_{D_0/D^*/D_1}^2 - \omega^2)^2} \\ &+ \frac{b}{M_{D_0/D^*/D_1}^2 - \omega^2} + \frac{c}{s_0 - \omega^2},\end{aligned}\quad (15)$$

with the constraint

$$\frac{a}{M_{D_0/D^*/D_1}^4} + \frac{b}{M_{D_0/D^*/D_1}^2} + \frac{c}{s_0} = 0. \quad (16)$$

Note that in Eq. (15) the phenomenological side of scattering amplitude for  $q_\mu \neq 0$  is not exactly equal to Born term but there are contributions from other terms. However, for  $\omega \rightarrow 0$ ,  $T_N$  on left should be equal to  $T^{\text{Born}}$  on right side of Eq. (15) and this requirement results in constraint given in Eq. (16). As we shall discuss below the constraint (16) help in eliminating the parameter  $c$  and scattering amplitude will be function of parameters  $a$  and  $b$  only. The Born terms to be used in equation (15) for scalar, vector and axial-vector mesons are given by following relations [22, 23]

$$T_{D_0 N}^{\text{Born}}(\omega, \mathbf{0}) = \frac{2f_{D_0}^2 M_{D_0}^2 M_N (M_H - M_N) g_{D_0 N H}^2}{[\omega^2 - (M_H - M_N)^2] [\omega^2 - M_{D_0}^2]^2},$$



$$\begin{aligned}
T_{D^*N}^{\text{Born}}(\omega, \mathbf{0}) &= \frac{2f_{D^*}^2 M_{D^*}^2 M_N (M_H + M_N) g_{D^*NH}^2}{[\omega^2 - (M_H + M_N)^2] [\omega^2 - M_{D^*}^2]^2}, \\
T_{D_1N}^{\text{Born}}(\omega, \mathbf{0}) &= \frac{2f_{D_1}^2 M_{D_1}^2 M_N (M_H - M_N) g_{D_1NH}^2}{[\omega^2 - (M_H - M_N)^2] [\omega^2 - M_{D_1}^2]^2}.
\end{aligned} \tag{17}$$

In the above equations  $g_{D_0NH}$ ,  $g_{D^*NH}$  and  $g_{D_1NH}$  are the coupling constants.  $M_H$  is the mass of the hadron e.g. corresponding to charm mesons we have  $\Lambda_c$  and  $\Sigma_c$  whereas corresponding to bottom mesons we have the hadrons  $\Lambda_b$  and  $\Sigma_b$ . Corresponding to charm mesons we take the average value of the masses of  $M_{\Lambda_c}$  and  $M_{\Sigma_c}$  and is equal to 2.4 GeV. For the case of mesons having bottom quark,  $b$ , we consider the average value of masses of  $\Lambda_b$  and  $\Sigma_b$  and it is equal to 5.7 GeV.

Now we write the equation for the Borel transformation of the scattering matrix on the phenomenological side and equate that to the Borel transformation of the scattering matrix for the operator expansion side. For the scalar meson,  $D_0$ , the Borel transformation equation is written as [22],

$$\begin{aligned}
& a \left\{ \frac{1}{M^2} e^{-\frac{M_{D_0}^2}{M^2}} - \frac{s_0}{M_{D_0}^4} e^{-\frac{s_0}{M^2}} \right\} + b \left\{ e^{-\frac{M_{D_0}^2}{M^2}} - \frac{s_0}{M_{D_0}^2} e^{-\frac{s_0}{M^2}} \right\} \\
& + A \left[ \frac{1}{(M_H - M_N)^2 - M_{D_0}^2} - \frac{1}{M^2} \right] e^{-\frac{M_{D_0}^2}{M^2}} - \frac{A}{(M_H - M_N)^2 - M_{D_0}^2} e^{-\frac{(M_H - M_N)^2}{M^2}} \\
& = \left\{ \frac{m_c \langle \bar{q}q \rangle_N}{2} - \langle q^\dagger i D_0 q \rangle_N + \frac{m_c^2 \langle q^\dagger i D_0 q \rangle_N}{M^2} \right\} e^{-\frac{m_c^2}{M^2}} \\
& - \left\{ \frac{2m_c \langle \bar{q} i D_0 i D_0 q \rangle_N}{M^2} - \frac{m_c^3 \langle \bar{q} i D_0 i D_0 q \rangle_N}{M^4} \right\} e^{-\frac{m_c^2}{M^2}} \\
& + \frac{1}{16} \langle \frac{\alpha_s GG}{\pi} \rangle_N \int_0^1 dx \left( 1 + \frac{\tilde{m}_c^2}{M^2} \right) e^{-\frac{\tilde{m}_c^2}{M^2}} - \frac{1}{48M^4} \langle \frac{\alpha_s GG}{\pi} \rangle_N \int_0^1 dx \frac{1-x}{x} \tilde{m}_c^4 e^{-\frac{\tilde{m}_c^2}{M^2}}, \tag{18}
\end{aligned}$$

where,  $A = \frac{2f_{D_0}^2 M_{D_0}^2 M_N (M_H - M_N) g_{D_0NH}^2}{(M_H - M_N)^2 - M_{D_0}^2}$ .

Note that in equation (18) we have two unknown parameters  $a$  and  $b$ . We differentiate equation (18) w.r.t.  $\frac{1}{M^2}$  so that we could have two equations and two unknowns. By solving those two coupled equations we will be able to get the values of parameters  $a$  and  $b$ . Same procedure will be applied to obtain the values of parameters  $a$  and  $b$  corresponding to vector and axial-vector mesons. For vector meson,  $D^*$ , the Borel transformation equation is given by [23],

$$a \left\{ \frac{1}{M^2} e^{-\frac{M_{D^*}^2}{M^2}} - \frac{s_0}{M_{D^*}^4} e^{-\frac{s_0}{M^2}} \right\} + b \left\{ e^{-\frac{M_{D^*}^2}{M^2}} - \frac{s_0}{M_{D^*}^2} e^{-\frac{s_0}{M^2}} \right\}$$

$$\begin{aligned}
& + B \left[ \frac{1}{(M_H + M_N)^2 - M_{D^*}^2} - \frac{1}{M^2} \right] e^{-\frac{M_{D^*}^2}{M^2}} - \frac{B}{(M_H + M_N)^2 - M_{D^*}^2} e^{-\frac{(M_H + M_N)^2}{M^2}} \\
& = \left\{ -\frac{m_c \langle \bar{q}q \rangle_N}{2} - \frac{2 \langle q^\dagger i D_0 q \rangle_N}{3} + \frac{m_c^2 \langle q^\dagger i D_0 q \rangle_N}{M^2} \right\} e^{-\frac{m_c^2}{M^2}} + \frac{m_c \langle \bar{q} g_s \sigma G q \rangle_N}{3M^2} e^{-\frac{m_c^2}{M^2}} \\
& + \left\{ \frac{8m_c \langle \bar{q} i D_0 i D_0 q \rangle_N}{3M^2} - \frac{m_c^3 \langle \bar{q} i D_0 i D_0 q \rangle_N}{M^4} \right\} e^{-\frac{m_c^2}{M^2}} \\
& - \frac{1}{24} \langle \frac{\alpha_s GG}{\pi} \rangle_N \int_0^1 dx \left( 1 + \frac{\widetilde{m}_c^2}{2M^2} \right) e^{-\frac{\widetilde{m}_c^2}{M^2}} \\
& + \frac{1}{48M^2} \langle \frac{\alpha_s GG}{\pi} \rangle_N \int_0^1 dx \frac{1-x}{x} \left( \widetilde{m}_c^2 - \frac{\widetilde{m}_c^4}{M^2} \right) e^{-\frac{\widetilde{m}_c^2}{M^2}}. \tag{19}
\end{aligned}$$

where,  $B = \frac{2f_{D^*}^2 M_{D^*}^2 M_N (M_H + M_N) g_{D^* N H}^2}{(M_H + M_N)^2 - M_{D^*}^2}$ . For the axial-vector mesons,  $D_1$ , the Borel transformation equation is given by [23],

$$\begin{aligned}
& a \left\{ \frac{1}{M^2} e^{-\frac{M_{D_1}^2}{M^2}} - \frac{s_0}{M_{D_1}^4} e^{-\frac{s_0}{M^2}} \right\} + b \left\{ e^{-\frac{M_{D_1}^2}{M^2}} - \frac{s_0}{M_{D_1}^2} e^{-\frac{s_0}{M^2}} \right\} \\
& + C \left[ \frac{1}{(M_H - M_N)^2 - M_{D_1}^2} - \frac{1}{M^2} \right] e^{-\frac{M_{D_1}^2}{M^2}} - \frac{C}{(M_H - M_N)^2 - M_{D_1}^2} e^{-\frac{(M_H - M_N)^2}{M^2}} \\
& = \left\{ \frac{m_c \langle \bar{q}q \rangle_N}{2} - \frac{2 \langle q^\dagger i D_0 q \rangle_N}{3} + \frac{m_c^2 \langle q^\dagger i D_0 q \rangle_N}{M^2} \right\} e^{-\frac{m_c^2}{M^2}} - \frac{m_c \langle \bar{q} g_s \sigma G q \rangle_N}{3M^2} e^{-\frac{m_c^2}{M^2}} \\
& - \left\{ \frac{8m_c \langle \bar{q} i D_0 i D_0 q \rangle_N}{3M^2} - \frac{m_c^3 \langle \bar{q} i D_0 i D_0 q \rangle_N}{M^4} \right\} e^{-\frac{m_c^2}{M^2}} \\
& - \frac{1}{24} \langle \frac{\alpha_s GG}{\pi} \rangle_N \int_0^1 dx \left( 1 + \frac{\widetilde{m}_c^2}{2M^2} \right) e^{-\frac{\widetilde{m}_c^2}{M^2}} \\
& + \frac{1}{48M^2} \langle \frac{\alpha_s GG}{\pi} \rangle_N \int_0^1 dx \frac{1-x}{x} \left( \widetilde{m}_c^2 - \frac{\widetilde{m}_c^4}{M^2} \right) e^{-\frac{\widetilde{m}_c^2}{M^2}}, \tag{20}
\end{aligned}$$

where,  $C = \frac{2f_{D_1}^2 M_{D_1}^2 M_N (M_H - M_N) g_{D_1 N H}^2}{(M_H - M_N)^2 - M_{D_1}^2}$ . In the above equations  $\widetilde{m}_c^2 = \frac{m_c^2}{x}$ .

As discussed earlier, in determining the properties of hadrons from QCD sum rules, we shall use the values of quark and gluon condensates as calculated using chiral SU(3) model. Any operator  $\mathcal{O}$  on OPE side can be written as [16, 37, 38],

$$\begin{aligned}
\mathcal{O}_{\rho_B} &= \mathcal{O}_{vacuum} + 4 \int \frac{d^3 p}{(2\pi)^3 2E_p} n_F \langle N(p) | \mathcal{O} | N(p) \rangle + 3 \int \frac{d^3 k}{(2\pi)^3 2E_k} n_B \langle \pi(k) | \mathcal{O} | \pi(k) \rangle \\
&= \mathcal{O}_{vacuum} + \frac{\rho_B}{2M_N} \mathcal{O}_N + \mathcal{O}_{P.B.} \tag{21}
\end{aligned}$$

In above equation,  $\mathcal{O}_{\rho_B}$ , gives us the expectation value of the operator at finite baryonic density. The term  $\mathcal{O}_{vacuum}$  stands for the vacuum expectation value of the operator,  $\mathcal{O}_N$  gives us the nucleon expectation value of the operator and  $\mathcal{O}_{P.B.}$  denotes the contribution

from the pion bath at finite temperature.  $n_B = [e^{E_k/T} - 1]^{-1}$  and  $n_F = [e^{(E_p - \mu_N)/T} - 1]^{-1}$  are the thermal Boson and Fermion distribution functions. Within the chiral SU(3) model the quark and gluon condensates can be expressed in terms of scalar fields  $\sigma$ ,  $\zeta$  and  $\chi$ . As discussed earlier, the finite temperature effects in the present investigation will be evaluated through the scalar fields and therefore contribution of third term will not be considered. However, for completeness we shall compare the temperature dependence of scalar quark and scalar gluon condensates at zero baryon density as evaluated in the present work with the situation when the temperature dependence is evaluated using only pion bath contribution [37]. Thus within chiral SU(3) model, we can find the values of  $\mathcal{O}_{\rho_B}$  at finite density of the nuclear medium and hence can find  $\mathcal{O}_N$  using

$$\mathcal{O}_N = [\mathcal{O}_{\rho_B} - \mathcal{O}_{vacuum}] \frac{2M_N}{\rho_B}. \quad (22)$$

The quark condensate,  $\bar{q}q$ , can be extracted from the explicit symmetry breaking term of the Lagrangian density and is given by,

$$\sum_i m_i \bar{q}_i q_i = -\mathcal{L}_{SB} = \left( \frac{\chi}{\chi_0} \right)^2 \left[ m_\pi^2 f_\pi \sigma + (\sqrt{2} m_k^2 f_k - \frac{1}{\sqrt{2}} m_\pi^2 f_\pi) \zeta \right]. \quad (23)$$

In our present investigation of hadron properties, we are interested in light quark condensates,  $\bar{u}u$  and  $\bar{d}d$ , which are proportional to the non-strange scalar field  $\sigma$  within chiral SU(3) model. Considering equal mass of light quarks,  $u$  and  $d$  i.e.  $m_u = m_d = m_q = 0.006$  GeV, we can write,

$$\langle \bar{q}q \rangle_{\rho_B} = \frac{1}{2m_q} \left( \frac{\chi}{\chi_0} \right)^2 [m_\pi^2 f_\pi \sigma]. \quad (24)$$

The condensate  $\langle \bar{q}g_s \sigma Gq \rangle_{\rho_B}$  is given by the equation [39],

$$\langle \bar{q}g_s \sigma Gq \rangle_{\rho_B} = \lambda^2 \langle \bar{q}q \rangle_{\rho_B} + 3.0 GeV^2 \rho_B. \quad (25)$$

Also we write [39],

$$\langle \bar{q}iD_0iD_0q \rangle_{\rho_B} + \frac{1}{8} \langle \bar{q}g_s \sigma Gq \rangle_{\rho_B} = 0.3 GeV^2 \rho_B. \quad (26)$$

As discussed above the quark condensate,  $\langle \bar{q}q \rangle_{\rho_B}$ , can be calculated within the chiral SU(3) model. This value of  $\langle \bar{q}q \rangle_{\rho_B}$  can be used through equations (25) and (26) to calculate the value of condensates  $\langle \bar{q}g_s \sigma Gq \rangle_{\rho_B}$  and  $\langle \bar{q}iD_0iD_0q \rangle_{\rho_B}$  within chiral SU(3) model. The value of condensate  $\langle q^\dagger iD_0q \rangle$  is equal to  $0.18 GeV^2 \rho_B$  [39].

At finite temperature and zero baryon density we can write the expectation values of quark condensates and scalar gluon condensates as [37],

$$\langle \bar{q}q \rangle_T = \langle \bar{q}q \rangle_{vacuum} \left[ 1 - \frac{T^2}{8f_\pi^2} B_1 \left( \frac{m_\pi}{T} \right) \right], \quad (27)$$

and

$$\left\langle \frac{\alpha_s}{\pi} G^a_{\mu\nu} G^{a\mu\nu} \right\rangle_T = \left\langle \frac{\alpha_s}{\pi} G^a_{\mu\nu} G^{a\mu\nu} \right\rangle_{vacuum} - \frac{1}{9} m_\pi^2 T^2 B_1 \left( \frac{m_\pi}{T} \right). \quad (28)$$

respectively. In equations (27) and (28)  $B_1(x) = \frac{6}{\pi^2} \int_x^\infty dy \frac{\sqrt{y^2 - x^2}}{e^y - 1}$ . From equations (27) and (28) we observe that the contribution from pion bath to the expectation values of operator arises only at finite temperature. Note that in equations (18), (19) and (20) we need the nucleon expectation values of various condensates which can be evaluated in general using equation(22). In equations (1), (24), (25) and (26) the values of condensates are given at finite value of baryonic density. To find the corresponding nucleon expectation values of various condensates we use the values of condensates at finite baryonic density from equations (1), (24), (25) and (26) in equation(22). The equation (22) is then further used in equations (18), (19) and (20) for calculation of medium modifications of  $D$  mesons.

It may be noted that the QCD sum rules for the evaluation of in-medium properties of scalar mesons,  $B_0$ , vector mesons,  $B^*$  and axial vector mesons  $B_1$ , can be written by replacing masses of charmed mesons  $D_0$ ,  $D^*$  and  $D_1$ , by corresponding masses of bottom mesons  $B_0$ ,  $B^*$  and  $B_1$  in equations (18), (19) and (20) respectively. Also the bare charm quark mass,  $m_c$  will be replaced by the mass of bottom quark,  $m_b$ .

#### IV. RESULTS AND DISCUSSIONS

In this section we shall present the results of our investigation of in-medium properties of scalar ( $D_0, B_0$ ), vector ( $D^*, B^*$ ) and axial vector mesons ( $D_1, B_1$ ). The nuclear matter saturation density used in the present investigation is  $0.15 \text{ fm}^{-3}$ . The values of various coupling constants  $g_{D_0 N \Lambda_c} \approx g_{D_0 N \Sigma_c} \approx g_{B_0 N \Lambda_b} \approx g_{B_0 N \Sigma_b}$  are approximated to 6.74 [22]. The coupling constants  $g_{D^* N \Lambda_c} \approx g_{D^* N \Sigma_c} \approx g_{D_1 N \Lambda_c} \approx g_{D_1 N \Sigma_c} \approx g_{B^* N \Lambda_b} \approx g_{B^* N \Sigma_b} \approx g_{B_1 N \Lambda_b} \approx g_{B_1 N \Sigma_b}$  are approximated to 3.86 [23]. The masses of mesons  $M_{D_0}$ ,  $M_{B_0}$ ,  $M_{D^*}$ ,  $M_{B^*}$ ,  $M_{D_1}$  and  $M_{B_1}$  to be used in present investigation are 2.355, 5.74, 2.01, 5.325, 2.42 and 5.75 GeV respectively. The values of decay constants  $f_{D_0}$ ,  $f_{B_0}$ ,  $f_{D^*}$ ,  $f_{B^*}$ ,  $f_{D_1}$  are  $f_{B_1}$  are 0.334, 0.28, 0.270, 0.195, 0.305 and 0.255 GeV respectively. The values of threshold parameters,  $s^0$ ,

corresponding to  $D_0$ ,  $B_0$ ,  $D^*$ ,  $B^*$ ,  $D_1$  and  $B_1$  mesons are 8, 39, 6.5, 35, 8.5 and 39  $\text{GeV}^2$  respectively [22, 23]. As discussed earlier the mass-shift of scalar, vector and axial-vector mesons is calculated through the parameter  $a$  which is related to scattering length through equations (12) to (14). This parameter  $a$ , for example, for  $D^0$  is calculated by solving the coupled equations as discussed after Eq. (18) and is subjected to the medium modifications through the medium dependence of condensates. The medium dependence of condensates is further evaluated through the scalar fields  $\sigma$ ,  $\zeta$  and  $\chi$ . The various coupling constants listed above, the decay constants of  $D$  and  $B$  mesons, the threshold parameter  $s_0$  are not subjected to the medium modifications. In the present work we shall show the variation of mass shift as a function of squared Borel mass parameter,  $M$ . The Borel window is chosen such that there is almost no change in the mass of  $D$  and  $B$  mesons w.r.t variation in Borel mass parameter. For the charmed scalar,  $D_0$ , vector,  $D^*$ , and axial-vector,  $D_1$  mesons Borel windows are found to be  $(6.1 - 7.4)$ ,  $(4.5 - 5.4)$  and  $(6.5 - 7.6)$   $\text{GeV}^2$  respectively. The Borel windows for bottom scalar,  $B_0$ , vector,  $B^*$ , and axial-vector,  $B_1$  mesons are  $(33 - 39)$ ,  $(22 - 24)$  and  $(34 - 37)$   $\text{GeV}^2$  respectively.

In the present work we are studying the in-medium masses of scalar, vector and axial vector mesons using QCD sum rules and chiral SU(3) model. Since we are evaluating the quark and gluon condensates within the chiral SU(3) model through the modification of scalar fields  $\sigma$  and  $\zeta$  and the scalar dilaton field  $\chi$ , so first we shall discuss in short the effect of temperature and density of the medium on the values of scalar fields  $\sigma$  and  $\zeta$  and the scalar dilaton field  $\chi$ . We observe that as a function of density of the nuclear medium the magnitude of scalar fields decreases. For example at nuclear saturation density,  $\rho_B = \rho_0$ , the drop in magnitude of the scalar fields  $\sigma$  and  $\zeta$  and the dilaton field  $\chi$  is observed to be 33.49, 9.54 and 3.39 MeV respectively. At baryon densities,  $4\rho_0$ , these values changes to 63.41, 14.51 and 13.06 MeV respectively.

At zero baryon density, we observe that the magnitude of the scalar fields  $\sigma$  and  $\zeta$  and the dilaton field  $\chi$  decreases with increase in the temperature. However, the change in the values of scalar fields with temperature of the medium is observed to be very small. The reason for the non zero values of scalar fields at finite temperature and zero baryon density of the medium is the formation of baryon-antibaryon pairs [32, 40]. At finite baryon densities, the magnitude of the scalar fields increases with increase in the temperature of symmetric nuclear medium. This also leads to increase in the masses of the nucleons with

the temperature of the nuclear medium for finite baryon densities [11]. At nuclear saturation density,  $\rho_0$ , the magnitude of the scalar fields,  $\sigma$  and  $\zeta$  and the dilaton field  $\chi$  increases by 5.47, 1.28 and 0.96 MeV respectively as we move from  $T = 0$  to  $T = 150$  MeV respectively.

In figures (1) and (2) we show the variation of the light scalar quark condensate  $\bar{q}q$ , given by equation (24) and the scalar gluon condensate,  $G_0 = \langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{a\mu\nu} \rangle$ , given by equation (1), respectively as a function of density of the symmetric nuclear medium. We show the results for temperatures,  $T = 0, 50, 100$  and  $150$  MeV respectively. From equation (24) we observe that the value of the scalar quark condensate  $\bar{q}q$  is directly proportional to the scalar-isoscalar field,  $\sigma$ . Therefore, the behavior of the  $\bar{q}q$  as a function of temperature and density of the nuclear medium will be same as that of  $\sigma$  field. For given value of temperature of the nuclear medium the magnitude of the light quark condensate decreases with increase in the density. For example, at  $T = 0$ , the values of light quark condensate are observed to be  $-0.8829 \times 10^{-2}$  and  $-0.4200 \times 10^{-2}$   $\text{GeV}^3$  at baryon densities,  $\rho_B = \rho_0$  and  $4\rho_0$  respectively. At temperature,  $T = 150$  MeV, these values of quark condensates changes to  $-0.9681 \times 10^{-2}$  and  $-0.4993 \times 10^{-2}$   $\text{GeV}^3$  respectively. At zero baryon density the magnitude of the  $\bar{q}q$  decreases with increase in the temperature of the nuclear medium. At  $\rho_B = 0$ , the values of  $\bar{q}q$  are observed to be  $-1.4014 \times 10^{-2}$ ,  $-1.4014 \times 10^{-2}$ ,  $-1.4006 \times 10^{-2}$  and  $-1.3628 \times 10^{-2}$   $\text{GeV}^3$  at temperatures  $T = 0, 50, 100$  and  $150$  MeV respectively.

From figure (2), we observe that the values of the scalar gluon condensates decreases with increase in the density of the nuclear medium. At baryon density,  $\rho_B = \rho_0$ , the values of  $G_0$  are observed to be  $1.90646 \times 10^{-2}$   $\text{GeV}^4$ ,  $1.9119 \times 10^{-2}$   $\text{GeV}^4$ ,  $1.91755 \times 10^{-2}$   $\text{GeV}^4$ , and  $1.92 \times 10^{-2}$   $\text{GeV}^4$  for temperatures,  $T = 0, 50, 100$  and  $150$  MeV respectively. For the same values of the temperature, in the absence of finite quark masses, the values of  $G_0$  are observed to be  $2.269 \times 10^{-2}$   $\text{GeV}^4$ ,  $2.2771 \times 10^{-2}$   $\text{GeV}^4$ ,  $2.2857 \times 10^{-2}$   $\text{GeV}^4$  and  $2.29 \times 10^{-2}$   $\text{GeV}^4$  for  $\rho_B = \rho_0$ . For baryon density  $\rho_B = 4\rho_0$ , the values of  $G_0$  are given as  $1.7367 \times 10^{-2}$   $\text{GeV}^4$  ( $2.06 \times 10^{-2}$   $\text{GeV}^4$ ),  $1.74612 \times 10^{-2}$   $\text{GeV}^4$  ( $2.0713 \times 10^{-2}$   $\text{GeV}^4$ ),  $1.7656 \times 10^{-2}$   $\text{GeV}^4$  ( $2.094 \times 10^{-2}$   $\text{GeV}^4$ ) and  $1.78 \times 10^{-2}$   $\text{GeV}^4$  ( $2.112 \times 10^{-2}$   $\text{GeV}^4$ ) for values of temperature,  $T = 0, 50, 100$  and  $150$  MeV respectively, for the cases of the finite (zero) quark masses in the trace anomaly.

It may be noted that in the above discussion the finite temperature effects of the nuclear medium on the values of quark and gluon condensates are evaluated through the temperature dependence of scalar fields  $\sigma$ ,  $\zeta$  and the dilaton field  $\chi$ . In literature the scalar quark and




FIG. 1: (Color online) The light quark condensate  $\bar{q}q$  plotted as a function of density of the nuclear medium, in units of nuclear saturation density, for different values of temperatures ( $T = 0, 50, 100$  and  $150 \text{ MeV}$ ).

gluon condensates at finite temperature are evaluated due to contribution from pion bath using equations (27) and (28) respectively [37]. Using  $m_q \langle \bar{q}q \rangle_0 = - (0.11 \text{ GeV})^4$  and  $\langle \frac{\alpha_s}{\pi} G^2 \rangle = 0.005 \text{ GeV}^4$  [37] we calculate the quark and gluon condensates at finite temperature and zero baryon density using equations (27) and (28) respectively. The values of scalar quark condensates are observed to be  $-1.824 \times 10^{-2}$ ,  $-1.751 \times 10^{-2}$  and  $-1.566 \times 10^{-2} \text{ GeV}^3$




FIG. 2: (Color online) The scalar gluon condensate  $G_0$  plotted as a function of density of the nuclear medium, in units of nuclear saturation density, for different values of temperatures ( $T = 0, 50, 100$  and  $150$  MeV).

at temperatures  $T = 50, 100$  and  $150$  MeV respectively. These values of scalar quark condensates can be compared to the values  $-1.4014 \times 10^{-2}$ ,  $-1.4006 \times 10^{-2}$  and  $-1.3628 \times 10^{-2}$   $\text{GeV}^3$  at  $T = 50, 100$  and  $150$  MeV calculated in our present approach where finite temperature effects were considered through scalar fields  $\sigma$ ,  $\zeta$  and  $\chi$ . Similarly, the values of scalar gluon condensates calculated using equation (28) are found to be  $0.5 \times 10^{-2}$ ,  $0.4994 \times$



$10^{-2}$  and  $0.497 \times 10^{-2} \text{ GeV}^4$  at  $T = 50, 100$  and  $150 \text{ MeV}$  respectively. However, for zero baryon density, using the chiral  $SU(3)$  model the values of scalar gluon condensates for finite (zero) quark mass term, are observed to be  $1.94264 \times 10^{-2}$  ( $2.34550 \times 10^{-2}$ ),  $1.94269 \times 10^{-2}$  ( $2.3454 \times 10^{-2}$ ) and  $1.94456 \times 10^{-2}$  ( $2.3437 \times 10^{-2}$ )  $\text{GeV}^4$  at temperatures  $T = 50, 100$  and  $150 \text{ MeV}$  respectively. From above discussion we observe that as a function of temperature there are very small variations in the values of scalar quark and gluon condensates in the nuclear medium. This observation agree e.g. with ref. [41] where gluon and chiral condensates were studied at finite temperature with an effective Lagrangian of pseudoscalar mesons coupled to a scalar glueball. The gluon condensates were found to be very stable up to temperatures of  $200 \text{ MeV}$ , where the chiral sector of the theory reaches its limit of validity [41]. Actually scalar quark and gluon condensates are found to vary effectively with temperature above critical temperature but in hadronic medium, for zero baryon density, these are not much sensitive to temperature effects [42, 43].

Note that in the above discussion of scalar gluon and quark condensates, calculated using equations (1) and (24) respectively, we considered the vacuum values of decay constants  $f_\pi$  and  $f_K$  as well as masses  $m_\pi$  and  $m_K$  of pions and kaons respectively. Now we shall discuss the effect of medium modified values of  $f_\pi$ ,  $f_K$ ,  $m_\pi$  and  $m_K$  on the scalar condensates. In the chiral effective model the pion and kaon decay constants are related to the scalar fields  $\sigma$  and  $\zeta$  through relations [5, 11]

$$f_\pi = -\sigma \quad (29)$$

and

$$f_K = -\frac{1}{2}(\sigma + \sqrt{2}\zeta) \quad (30)$$

respectively. From the above relations it is clear that the medium dependence of scalar fields  $\sigma$  and  $\zeta$  can be used to study the density and temperature dependence of decay constants of pions and kaons. For example, using equation (29), at zero temperature the values of pion decay constant are observed to be  $93.3, 59.75$  and  $42.99 \text{ MeV}$  at baryonic densities  $\rho_B = 0, \rho_0$  and  $2\rho_0$  respectively. The values of kaons decay constants, using equation (30), are observed to be  $122, 98.6$  and  $87.93 \text{ MeV}$  at  $\rho_B = 0, \rho_0$  and  $2\rho_0$  respectively. In reference [44] the masses of pions were calculated in linear density approximation using the chiral perturbation theory. At baryon density  $\rho_B = \rho_0$  and zero temperature the mass of pion changes from the vacuum value  $139 \text{ MeV}$  to  $145.8 \text{ MeV}$ . The properties of kaons and antikaons had been

investigated in the literature using the chiral model [8] and coupled channel approach [45–48]. In the present work, to study the effect of medium modified masses of kaons on the values of quark and gluon condensates we shall use the chiral SU(3) model [8]. In the nuclear medium the kaons feel repulsive interactions and their in-medium mass increases as a function of density whereas the antikaons feel attractive interactions and their in-medium masses drop as we move to higher baryon density. For the present purpose considering the average mass of kaons and antikaons, the in-medium mass at temperature  $T = 0$  is observed to be 494, 488.88 and 466.08 MeV at  $\rho_B = 0$ ,  $\rho_0$  and  $2\rho_0$  respectively. Taking into account the above discussed in-medium properties of  $f_\pi$ ,  $f_K$ ,  $m_\pi$  and  $m_K$ , at baryonic density  $\rho_B = \rho_0$  and zero temperature the values of scalar quark and gluon condensates are observed to be  $-0.6222 \times 10^{-2} \text{ GeV}^3$  and  $1.9869 \times 10^{-2} \text{ GeV}^4$  respectively. These values of scalar quark and gluon condensates can be compared to  $-0.8829 \times 10^{-2} \text{ GeV}^3$  and  $1.9065 \times 10^{-2} \text{ GeV}^4$  respectively which were calculated without the medium modification of  $f_\pi$ ,  $f_K$ ,  $m_\pi$  and  $m_K$ . We conclude that the medium modification of  $f_\pi$ ,  $f_K$ ,  $m_\pi$  and  $m_K$  causes more decrease in the values of scalar quark and gluon condensates at finite baryonic density.

Now we shall calculate the in-medium masses of scalar, vector and axial vector mesons using the values of condensates from chiral SU(3) model. In figure (3), the subfigures (a), (c) and (e) show the variation of the mass shift of scalar mesons  $D_0$  as a function of square of the Borel mass parameter,  $M$ , at nuclear matter densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. The subfigures (b), (d) and (f) show the variation of the scalar mesons  $B_0$  as a function of square of the Borel mass parameter,  $M$ , at nuclear matter densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. In each subplot we have shown the results at temperatures  $T = 0, 50, 100$  and  $150$  MeV. We observe that for scalar mesons,  $D_0$ , at temperature,  $T = 0$ , the values of mass shift are found to be 76, 114 and 148 MeV at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. For temperature  $T = 50$  MeV the values of mass shift are observed to be 71, 109 and 144 MeV at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. At temperature  $T = 100$  MeV the above values of mass shift are observed to be 66, 103 and 139 MeV, whereas at  $T = 150$  MeV the values of mass shift changes to 58, 94 and 131 MeV at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. From the above discussion we conclude that for a given value of temperature the mass shift of scalar mesons,  $D_0$ , increases as a function of density of the nuclear medium. On the other hand as a function of temperature of the nuclear medium, for a constant value of density, the mass shift of scalar mesons,  $D_0$ , decreases.

As we can see from figure (3) the values of mass shift of scalar  $B_0$  mesons also increases as a function of density of the nuclear medium. At temperature  $T = 0$ , the values of mass shift are observed to 224, 334 and 420 MeV at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. At temperature  $T = 50$  MeV the above values of mass shift are found to be 211, 321 and 413 MeV at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. For temperature  $T = 100$  MeV the values of mass shift are found to be 196, 302 and 399 MeV whereas at  $T = 150$  MeV these values of mass shift changes to 171, 274 and 372 MeV at  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$  respectively. For a given value of nuclear matter density the values of mass shift for the  $B_0$  mesons are found to decrease with increase in the temperature of the medium.

We also calculate the values of scattering lengths of scalar mesons using equation (12) for different values of density and temperature of the medium. At temperature  $T = 0$  and baryon densities,  $\rho_B = \rho_0$  and  $4\rho_0$  the values of scattering lengths for scalar mesons  $D_0$  ( $B_0$ ) are observed to be 1.42 (5.05) and 0.70 (2.41) fm respectively. For temperature  $T = 100$  MeV and baryon densities,  $\rho_B = \rho_0$  and  $4\rho_0$  the values of scattering lengths for  $D_0$  ( $B_0$ ) are observed to be 1.23 (4.40) and 0.66 (2.28) fm respectively. We note that the value of scattering length decreases as a function of density and temperature of the nuclear medium. As discussed above the scattering lengths are evaluated using equation (12). In this equation we have the parameter  $a$  which is directly proportional to the scattering length of mesons. As discussed earlier the value of parameter  $a$  is evaluated by solving simultaneously equation (18) and the equation obtained by differentiate equation (18) w.r.t.  $\frac{1}{M^2}$ . The magnitude of parameter  $a$  is found to decrease as we move from low to higher value of baryonic density or from zero to finite value of temperature of the medium. This behavior of parameter  $a$  as a function of density and temperature of the medium results in the similar changes in the values of scattering lengths of scalar mesons.

Figure (4) shows the variation of the mass shift of vector mesons  $D^*$  and  $B^*$  as a function of square of the Borel mass parameter. Here also we have shown the results at nuclear matter densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ . We observe that at nuclear matter density  $\rho_B = \rho_0$  the values of mass shift for vector mesons,  $D^*$  are observed to be -76, -71, -65 and -56 MeV at temperatures  $T = 0, 50, 100$  and  $150$  MeV respectively. For baryon density  $\rho_B = 2\rho_0$  the values of mass shift are found to be -111, -106, -98, and -87 MeV whereas for  $\rho_B = 4\rho_0$  the values of the mass shift changes to -128, -125, -120 and -108 MeV at temperatures  $T = 0, 50, 100$  and  $150$  MeV respectively. Note that for a given value of density the magnitude




FIG. 3: (Color online) Figure shows the variation of the mass shift of Scalar mesons  $D_0$  (subplots (a), (c), (e)) and  $B_0$  (subplots (b), (d), (f)) as a function of the squared Borel mass Parameter,  $M^2$ . We show the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ . For each value of baryon density the results are shown at temperatures,  $T = 0, 50, 100$  and  $150$  MeV. Also the medium modifications of decay constants and masses of pions and kaons are considered while evaluating the above shown mass-shift of scalar mesons.

of the mass shift of vector mesons decreases as a function of temperature of the nuclear medium. On the other hand as a function of density of the medium the magnitude of the mass shift of the vector mesons,  $D^*$  increases. For nuclear matter saturation density the values of mass shift for the  $B^*$  mesons are found to be  $-366, -344, -311$  and  $-275$  MeV at temperatures  $T = 0, 50, 100$  and  $150$  MeV respectively. At baryon density  $2\rho_0$  the above

values of mass shift changes to  $-557$ ,  $-534$ ,  $-498$  and  $-447$  MeV at temperatures  $T = 0, 50, 100$  and  $150$  MeV respectively whereas for baryon density  $4\rho_0$  the above values of mass shift found to be  $-701$ ,  $-689$ ,  $-662$  and  $-609$  MeV at temperatures  $T = 0, 50, 100$  and  $150$  MeV respectively. At temperature  $T = 0$  and baryon densities,  $\rho_B = \rho_0$  and  $4\rho_0$  the values of scattering lengths for vector mesons  $D^*$  ( $B^*$ ) are observed to be  $-1.31$  ( $-7.72$ ) and  $-0.54$  ( $-3.58$ ) fm respectively. For temperature  $T = 100$  MeV and baryon densities,  $\rho_B = \rho_0$  and  $4\rho_0$  the values of scattering lengths for  $D^*$  ( $B^*$ ) are observed to be  $-1.12$  ( $-6.72$ ) and  $-0.51$  ( $-3.54$ ) fm respectively. We observe that the magnitude of the scattering lengths of vector mesons decreases in moving from low to higher value of density or temperature of the medium.

In figure (5), for given values of temperatures and densities we have shown the variation of mass shift of axial-vector mesons  $D_1$  and  $B_1$  as a function of square of the Borel mass parameter. We observe that at baryon density  $\rho_B = \rho_0$ , the values of mass shift for axial vector meson  $D_1$  are observed to be  $73, 69, 63$  and  $55$  MeV at temperatures,  $T = 0, 50, 100$  and  $150$  MeV. At baryon density  $2\rho_0$  ( $4\rho_0$ ) the values of mass shift are found to be  $108$  ( $131$ ),  $104$ ( $128$ ),  $97$ ( $123$ ) and  $87$ ( $113$ ) at temperatures,  $T = 0, 50, 100$  and  $150$  MeV. For the axial vector meson  $B_1$  at baryon density  $\rho_0$  ( $2\rho_0$ ) the values of mass shift are found to be  $267$  ( $396$ ),  $251$ ( $381$ ),  $233$ ( $357$ ) and  $203$ ( $324$ ) MeV at temperatures,  $T = 0, 50, 100$  and  $150$  MeV respectively. At baryon density  $4\rho_0$  the values of mass shift are found to be  $492, 485, 467$  and  $434$  MeV at temperatures,  $T = 0, 50, 100$  and  $150$  MeV respectively. The values of scattering lengths for axial-vector mesons  $D_1$  ( $B_1$ ) at temperature  $T = 0$  and baryon densities,  $\rho_B = \rho_0$  and  $4\rho_0$  are observed to be  $1.38$  ( $6.02$ ) and  $0.62$  ( $2.83$ ) fm respectively. For temperature  $T = 100$  MeV and baryon densities,  $\rho_B = \rho_0$  and  $4\rho_0$  the values of scattering lengths for  $D^*$  ( $B^*$ ) changes to  $1.19$  ( $5.25$ ) and  $0.58$  ( $2.69$ ) fm respectively. From the above discussions we observe a positive value of mass shift for scalar ( $D_0, B_0$ ) and axial vector mesons ( $D_1, B_1$ ) in the nuclear medium. However the values of mass shift for vector mesons ( $D^*, B^*$ ) are found to be negative. It means the masses of above scalar and axial-vector mesons in the nuclear medium may be large compared to the value in free space and this may lead to a decrease in the yield of these mesons in heavy-ion collisions.

Now we shall discuss the effect of different individual condensates on the in-medium modification of scalar ( $D_0, B_0$ ), vector ( $D^*, B^*$ ) and axial vector ( $D_1, B_1$ ) mesons. In figures (6), (7) and (8) we compare the contributions of individual condensates to the mass shift of scalar mesons,  $D_0$ , vector mesons,  $D^*$  and axial vector mesons,  $D_1$  respectively. The




FIG. 4: (Color online) Figure shows the variation of the mass shift of vector mesons  $D^*$  (subplots (a), (c), (e)) and  $B^*$  (subplots (b), (d), (f)) as a function of the squared Borel mass Parameter,  $M^2$ . We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ . For each value of the baryon density the results are shown at temperatures,  $T = 0, 50, 100$  and  $150$  MeV.

subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ . Note that in figures (6) and in the subsequent figures of this paper the word “*Total*” is for the contribution of all condensates to the properties of mesons. Similarly, “*Quark<sub>1</sub>*”, “*Quark<sub>2</sub>*”, “*Quark<sub>4</sub>*” and “*Gluon<sub>1</sub>*” are denoting the contribution of  $\langle \bar{q}q \rangle$ ,  $\langle q^\dagger i D_0 q \rangle$ ,  $\langle \bar{q} i D_0 i D_0 q \rangle$  and  $\langle \frac{\alpha_s G G}{\pi} \rangle$  respectively to the in-medium properties of mesons.




FIG. 5: (Color online) Figure shows the variation of the mass shift of axial vector mesons  $D_1$  (subplots (a), (c), (e)) and  $B_1$  (subplots (b), (d), (f)) as a function of the squared Borel mass Parameter,  $M^2$ . We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ . For each value of the baryon density the results are shown at temperatures,  $T = 0, 50, 100$  and  $150$  MeV.

We observe that at temperature  $T = 0$  if we consider the contribution of scalar quark condensates only then the values of mass shift for the scalar  $D_0$  mesons are found to be 73.30 and 150.68 MeV at nuclear matter density  $\rho_B = \rho_0$  and  $4\rho_0$  respectively. When we consider the individual contributions of  $\langle q^\dagger i D_0 q \rangle$ ,  $\langle \bar{q} i D_0 i D_0 q \rangle$  and  $\langle \frac{\alpha_s G G}{\pi} \rangle$  condensates then the values of mass-shift at  $\rho_0$  ( $4\rho_0$ ) and  $T = 0$  are observed to be 4.01(16.07), 12.70(37.87) and 6.68(25.96) MeV respectively. From above discussion we observe that the maximum




FIG. 6: (Color online) Figure shows the contribution of individual condensates to the mass shift of scalar mesons  $D_0$  as a function of the squared Borel mass parameter,  $M^2$ . The subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ .

contribution to the in-medium modification of scalar  $D_0$  mesons is from light quark condensate  $\langle \bar{q}q \rangle$ . Note that leaving the quark condensate,  $\langle q^\dagger i D_0 q \rangle$ , the all other condensates have been evaluated within chiral SU(3) model in the present investigation. So in figure (6) we also show the variation of the mass shift of scalar  $D_0$  mesons as a function of squared Borel mass parameter when we neglect the contribution of  $\langle q^\dagger i D_0 q \rangle$  condensate. We observe that if we neglect the contribution of  $\langle q^\dagger i D_0 q \rangle$  condensate then the values of mass shift are found






FIG. 7: (Color online) Figure shows the contribution of individual condensates to the mass shift of vector mesons  $D^*$  as a function of the squared Borel mass parameter,  $M^2$ . The subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ .

to be 78.70 and 158.99 MeV at densities  $\rho_0$  and  $4\rho_0$  respectively. Note that considering the contribution of all condensates, at temperature  $T = 0$ , the values of mass shift were 75.88 and 148.05 MeV at densities  $\rho_0$  and  $4\rho_0$  respectively. So we observe that if we neglect the condensate  $\langle q^\dagger i D_0 q \rangle$  then there is a percentage change of 4 % and 7 % in the mass shift of  $D_0$  mesons at densities  $\rho_0$  and  $4\rho_0$  respectively. From figure (7), we observe that for vector mesons,  $D^*$  at temperature  $T = 0$  and baryon density  $\rho_B = \rho_0$  ( $4\rho_0$ ), the values of mass




FIG. 8: (Color online) Figure shows the contribution of individual condensates to the mass shift of axial vector mesons  $D_1$  as a function of the squared Borel mass parameter,  $M^2$ . The subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ .

shift due to condensates  $\langle \bar{q}q \rangle$ ,  $\langle q^\dagger i D_0 q \rangle$ ,  $\langle \bar{q} g_s \sigma G q \rangle$ ,  $\langle \bar{q} i D_0 i D_0 q \rangle$  and  $\langle \frac{\alpha_s G G}{\pi} \rangle$  are observed to be -82.15(-151.15), 1.689(6.749), 18.95(57.16), -5.49(-6.16) and 3.54 (14.93) MeV respectively. For temperature  $T = 100$  MeV the above values of mass shift at baryon density  $\rho_B = \rho_0$  ( $4\rho_0$ ) changes to -73.01 (-142.35), 1.689 (6.75), 17.85 (56.34), -4.53 (-5.43) and 3.43 (14.64) MeV respectively. For the axial vector mesons,  $D_1$  the values of mass shift due to individual

	T = 0		T = 100 MeV	
	$\rho_B = \rho_0$	$\rho_B = 4\rho_0$	$\rho_B = \rho_0$	$\rho_B = 4\rho_0$
Total	224	420	196	399
$\langle \bar{q}q \rangle$	223	422	195	402
$\langle q^\dagger i D_0 q \rangle$	2.26	9.04	2.26	9.04
$\langle \bar{q} i D_0 i D_0 q \rangle$	6.60	20.04	5.99	18.69
$\langle \frac{\alpha_s GG}{\pi} \rangle$	3.49	13.35	3.58	13.57
$\langle q^\dagger i D_0 q \rangle = 0$	226	426	197	405

TABLE I: The above table gives the values of mass shift of scalar mesons  $B_0$  (in MeV units) due to individual condensates at temperatures  $T = 0$  and 100 MeV. For each value of temperature the values are tabulated for baryon densities  $\rho_B = \rho_0$  and  $4\rho_0$ .

condensates  $\langle \bar{q}q \rangle$ ,  $\langle q^\dagger i D_0 q \rangle$ ,  $\langle \bar{q} g_s \sigma G q \rangle$ ,  $\langle \bar{q} i D_0 i D_0 q \rangle$  and  $\langle \frac{\alpha_s GG}{\pi} \rangle$  are observed to be 79.31(152), 0.1516(0.607), -8.77(-22.63), 8.27(21.76) and 2.34 (10.08) MeV respectively. For temperature  $T = 100$  MeV the above values of mass shift at baryon density  $\rho_B = \rho_0$  ( $4\rho_0$ ) changes to 70 (145), 0.151 (0.607), -7.99(-22.04), 7.58 (21.26) and 2.23 (9.81) respectively. In figures (9), (10) and (11) we have shown the contributions of individual condensates to the mass shift of scalar mesons,  $B_0$ , vector mesons,  $B^*$  and axial vector mesons,  $B_1$  respectively. The subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ . In table (I), (II) and (III) we have tabulated the values of mass shift for scalar mesons,  $B_0$ , vector mesons,  $B^*$  and axial vector mesons,  $B_1$  respectively. The values of mass shift have been given at baryon densities  $\rho_0$  and  $4\rho_0$  and temperatures  $T = 0$  and 100 MeV.

In ref. [22] the properties of scalar mesons  $D_0$  and  $B_0$  had been studied using QCD sum rules and the observed values of mass shift at nuclear saturation density were 69 and 217 MeV respectively. The properties of  $D_0$  mesons had also been studied in ref. [24] using coupled channel approach. There an extra widening from the already large width of the resonance in free space was observed for the  $D_0$  meson. The properties of vector mesons ( $D^*$  and  $B^*$ ) and axial-vector ( $D_1$  and  $B_1$ ) had been studied using QCD sum rules in ref. [23]. For vector mesons  $D^*$  and  $B^*$  the values of mass shift were  $-71$  and  $-380$  MeV respectively whereas




FIG. 9: (Color online) Figure shows the contribution of individual condensates to the mass shift of scalar mesons  $B_0$  as a function of the squared Borel mass parameter,  $M^2$ . The subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ .

for axial vector mesons  $D_1$  and  $B_1$  these values changes to 72 and 264 MeV respectively. The observed positive values of mass shift for  $D_0$  and  $D_1$  mesons in our present investigation and also in earlier investigations indicate that these mesons feels repulsive interactions in the nuclear medium and their in-medium mass increases as function of baryonic density. This means the chances of decay of higher charmonium states to these heavy charmed mesons pairs are suppressed and hence these mesons may not cause a decrease in the production




FIG. 10: Figure shows the contribution of individual condensates to the mass shift of vector mesons  $B_*$  as a function of the squared Borel mass parameter,  $M^2$ . The subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ .

of  $J/\psi$  mesons in heavy-ion collisions. However for the vector mesons  $D^*$  we observe the negative values of mass shift and hence we conclude that they feel attractive interactions in the nuclear medium. The decrease in the mass of vector mesons  $D^*$  may cause the decay of higher charmonium states to  $D^*\bar{D}^*$  pairs and hence it may be a cause of  $J/\psi$  suppression in heavy-ion collision experiments.

	T = 0		T = 100 MeV	
	$\rho_B = \rho_0$	$\rho_B = 4\rho_0$	$\rho_B = \rho_0$	$\rho_B = 4\rho_0$
Total	-367	-705.69	-318.08	-665.88
$\langle \bar{q}q \rangle$	-373.11	-728.58	-323.61	-687.98
$\langle q^\dagger i D_0 q \rangle$	1.477	5.9057	1.477	5.9057
$\langle \bar{q} g_s \sigma G q \rangle$	12.43	36.77	11.67	36.18
$\langle \bar{q} i D_0 i D_0 q \rangle$	-3.98	-6.42	-3.40	-5.98
$\langle \frac{\alpha_s G G}{\pi} \rangle$	1.64	7.56	1.51	7.21
$\langle q^\dagger i D_0 q \rangle = 0$	-366.34	-702.26	-317.28	-662.48

TABLE II: The above table gives the values of mass shift of vector mesons  $B^*$  (in MeV units) due to individual condensates at temperatures  $T = 0$  and 100 MeV. For each value of temperature the values are tabulated for baryon densities  $\rho_B = \rho_0$  and  $4\rho_0$ .

	T = 0		T = 100 MeV	
	$\rho_B = \rho_0$	$\rho_B = 4\rho_0$	$\rho_B = \rho_0$	$\rho_B = 4\rho_0$
Total	266.51	492.22	232.75	467.29
$\langle \bar{q}q \rangle$	270.11	503.57	236.35	478.86
$\langle q^\dagger i D_0 q \rangle$	.88	3.53	0.88	3.53
$\langle \bar{q} g_s \sigma G q \rangle$	-6.34	-16.79	-5.83	-16.40
$\langle \bar{q} i D_0 i D_0 q \rangle$	4.79	12.53	4.39	12.23
$\langle \frac{\alpha_s G G}{\pi} \rangle$	1.41	6.39	1.31	6.12
$\langle q^\dagger i D_0 q \rangle = 0$	266.75	493.11	232.99	468.20

TABLE III: (Color online) The above table gives the values of mass shift of axial vector mesons  $B_1$  (in MeV units) due to individual condensates at temperatures  $T = 0$  and 100 MeV. For each value of temperature the values are tabulated for baryon densities  $\rho_B = \rho_0$  and  $4\rho_0$ .

## V. SUMMARY

In the present paper we investigated the mass modifications of scalar mesons ( $D_0, B_0$ ), vector mesons ( $D^*, B^*$ ) and axial vector mesons ( $D_1, B_1$ ) at finite density and temperature of the nuclear medium. We used QCD sum rules along with chiral SU(3) model to investigate




FIG. 11: (Color online) Figure shows the contribution of individual condensates to the mass shift of axial vector mesons  $B_1$  as a function of the squared Borel mass parameter,  $M^2$ . The subplots (a), (c) and (e) show the results at temperature  $T = 0$ , whereas the subplots (b), (d) and (f) are plotted for temperature  $T = 100$  MeV. We have shown the results at baryon densities  $\rho_0$ ,  $2\rho_0$  and  $4\rho_0$ .

the properties of above mentioned mesons. Using chiral  $SU(3)$  model we found the values of quark and gluon condensates at finite density and temperature of the medium which were further used within QCD sum rules to find the in-medium masses of scalar, vector and axial vector mesons. We observed a positive value of mass shift for the scalar mesons ( $D_0$  and  $B_0$ ) and axial vector mesons ( $D_1$  and  $B_1$ ) i.e. their in-medium mass found to be more as

compared to the vacuum mass. For a constant value of temperature the values of mass shift for these scalar and axial-vector mesons are found to increase as we increase the density of the nuclear medium. On the other hand for a constant value of density the temperature of the medium causes a decrease in the values of mass shift of scalar and axial-vector mesons. The vector mesons  $D^*$  and  $B^*$  are found to have negative values of mass shift. It means their in-medium masses are small as compared to vacuum masses. For a constant value of density, as a function of temperature, the magnitude of mass shift of  $D^*$  and  $B^*$  mesons decreases. However, for a constant value of temperature, as a function of density the magnitude of mass shift of vector mesons  $D^*$  and  $B^*$  are found to increase. We have also investigated the effects of individual terms on the mass shift of mesons. It was found that the scalar quark condensates,  $\bar{q}q$  has maximum contribution to the in-medium modification of scalar, vector and axial vector mesons. The effects of density and temperature of the medium on the scattering lengths of scalar, vector and axial vector mesons were also investigated. We observed the positive value of scattering lengths for scalar and axial vector mesons whereas for the vector mesons a negative value of scattering lengths were observed. Also it was found that the magnitude of the scattering lengths of scalar, vector and axial-vector mesons decreases as we move from low to high value of density or temperature of the nuclear medium. The present investigation of medium modification of scalar, vector and axial vector mesons will be helpful for understanding their production rate and also the phenomenon of  $J/\psi$  suppression in the Compressed Baryonic matter experiment at FAIR, GSI.

### Acknowledgments

Author is thankful to Amruta Mishra for fruitful discussions on chiral SU(3) model and QCD sum rules. The financial support from the Department of Science and Technology (DST), India for research project under fast track scheme for young scientists (SR/FTP/PS-209/2012) is gratefully acknowledged.

- 
- [1] W.S. Chung, C.M. Ko and G.Q. Li, Nucl.Phys. A **641**, 357 (1998).
  - [2] D.B. Kaplan and A.E. Nelson, Phys. Lett. B **175**, 57 (1986).
  - [3] A.E. Nelson and D.B. Kaplan, Phys. Lett. B **192**, 193 (1987).



- [4] G.E. Brown and M. Rho, Phys. Rep. **269**, 333 (1996).
- [5] P. Papazoglou, D. Zschesche, S. Schramm, J. Schaffner-Bielich, H. Stöcker, and W. Greiner, Phys. Rev. C **59**, 411 (1999).
- [6] A. Mishra, E. L. Bratkovskaya, J. Schaffner-Bielich, S. Schramm and H. Stöcker, Phys. Rev. C **70**, 044904 (2004).
- [7] A. Mishra and S. Schramm, Phys. Rev. C **74**, 064904 (2006), A. Mishra, S. Schramm and W. Greiner, Phys. Rev. C **78**, 024901 (2008).
- [8] Amruta Mishra, Arvind Kumar, Sambuddha Sanyal, S. Schramm, Eur. Phys. J. A **41**, 205 (2009).
- [9] Amruta Mishra and Arindam Mazumdar, Phys. Rev. C **79**, 024908 (2009).
- [10] A. Mishra, E. L. Bratkovskaya, J. Schaffner-Bielich, S.Schramm and H. Stöcker, Phys. Rev. C **69**, 015202 (2004).
- [11] Arvind Kumar and Amruta Mishra, Phys, Rev. C **81**, 065204 (2010).
- [12] Arvind Kumar and Amruta Mishra , Eur. Phys, J. A **47**, 164 (2011).
- [13] S.H. Lee and K. Morita, Pramana Journal of Physics **72**, 97 (2009).
- [14] K. Morita and S.H. Lee, Phys. Rev. C **77**, 064904 (2008).
- [15] F. Klingl *et.al.*, Phys. Rev. Lett. **82**, 3396 (1999).
- [16] Yuji Koike and Arata Hayashigaki, Prog. Theo. Phys. **98**, 631 (1997).
- [17] T. Hilger,R. Thomas, B. Kampfer, Phys. Rev. C **79**, 025202 (2009).
- [18] Arata Hayashigaki, Phys. Lett. B **487**, 96 (2000).
- [19] E. Oset, A. Ramos, Nucl. Phys. A **635**, 99 (1998).
- [20] Volker Koch, Phys. Lett. B **337**, 7 (1994).
- [21] L. Tolós, D. Cabrera and A. Ramos, Phys. Rev. C **78**, 045205 (2008).
- [22] Zhi-Gang Wang, Tao Huang, Phys. Rev. C **84**, 048201 (2011).
- [23] Zhi-Gang Wang, Int. J. Mod. Phys. A **28**, 1350049 (2013) .
- [24] Laura Tolos et al, Nucl. Phys. A **827**, 249(2009)
- [25] T. Hilger and B. Kampfer, Nucl. Phys. A **207**, 277(2010)
- [26] N. Masera for the HELIOS-3 collaboration, Nucl. Phys. A **590**, 93c (1995).
- [27] G. Agakichiev et al (CERES collaboration), Phys. Rev. Lett. **75**, 1272 (1995); G. Agakichiev et al (CERES collaboration), Phys. Lett. B **422**, 405 (1998); G. Agakichiev et al (CERES collaboration), Nucl. Phys. A **661**, 23c (1999).

- [28] R. J. Porter et al (DLS collaboration), Phys. Rev. Lett. **79**, 1229 (1997); W. K. Wilson et al (DLS collaboration), Phys. Rev. C **57**, 1865 (1998).
- [29] A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel, Phys. Lett. B **659**, 149 (2008).
- [30] Arvind Kumar and Amruta Mishra, Phys. Rev. C **82**, 045207 (2010).
- [31] D. Zschesche, A. Mishra, S. Schramm, H. Stöcker and W. Greiner, Phys. Rev. C **70**, 045202 (2004).
- [32] A. Mishra, K. Balazs, D. Zschesche, S. Schramm, H. Stöcker, and W. Greiner, Phys. Rev. C **69**, 024903 (2004).
- [33] S. Weinberg, Phys. Rev. **166**, 1568 (1968).
- [34] S. Coleman, J. Wess, B. Zumino, Phys. Rev. **177**, 2239 (1969); C.G. Callan, S. Coleman, J. Wess, B. Zumino, Phys. Rev. **177**, 2247 (1969).
- [35] W. A. Bardeen and B. W. Lee, Phys. Rev. **177**, 2389 (1969).
- [36] J. Schechter, Phys. Rev. D **21**, 3393 (1980).
- [37] Youngshin Kwon, Chihiro Sasaki, Wolfram Weise, Phys. Rev. C **81**, 065203 (2010).
- [38] S. Zschocke, O.P. Pavlenko, B. Kampfer, Eur. Phys. J. A **15**, 529 (2002).
- [39] R. Thomas, T. Hilger, B. Kampfer, Nucl. Phys. A **795**, 19 (2007).
- [40] R. J. Furnstahl and B.D. Serot, Phys. Rev. C **41**, 262 (1992); J Theis, G. Graebner, G. Buchwald, J. Maruhn, W. Greiner and J. Polonyi, Phys. Rev. D **28**, 2286 (1983).
- [41] Josef Sollfrank, Ulrich Heinz, Z. Phys. C **65**, 111 (1995).
- [42] Zhou Li-Juan, Zheng Bo, Zhong Hong-wei, Ma Wei-xing, arXiv:1403.6965 [hep-ph].
- [43] David E. Miller, Acta Phys. Polon. B **28**, 2937 (1997).
- [44] TAN Yu-Hong, ZHANG Xiao-Bing, LI Lei and NING Ping-Zhi, Commun. Theor. Phys. (Beijing, China) **37**, 215 (2002).
- [45] M. Lutz, Phys. Lett. B **426**, 12 (1998).
- [46] A. Ramos and E. Oset, Nucl. Phys. A **671**, 481 (2000).
- [47] M. Lutz and C. L. Korpa, Nucl. Phys. A **700**, 309 (2002).
- [48] L. Tolós, A. Ramos, and A. Polls, Phys. Rev. C **65**, 054907 (2002).