

Next-to-leading order $pp \rightarrow W' \rightarrow tb$ production at 14 TeV and 33 TeV

Zack Sullivan*

Department of Physics, Illinois Institute of Technology, Chicago, Illinois 60616-3793, USA

(Dated: August 15, 2013)

I update the predicted leading order and next-to-leading order cross sections and total widths for W' bosons that decay to top and bottom quarks at 14 TeV and 33 TeV pp colliders ($pp \rightarrow W' \rightarrow tb$). Separate tables are included for right- and left-handed bosons. Theoretical uncertainties are completely dominated by parton distribution function uncertainties, and are computed for $W'_{\pm}$ production at a 14 TeV pp collider.

New charged vector currents, generically called W' bosons, appear in many models with extended gauge explanations of electroweak symmetry breaking. As long as these new bosons couple to fermions, a model-independent search can probe the parameter space of all such models by looking for the decay of the W' boson into a final state involving a top quark and bottom quark [1]. While the general case of mixed right- and left-handed couplings can be probed, nearly all models assume a term enters the Lagrangian of the form

$$\mathcal{L} = \frac{g'}{2\sqrt{2}} V'_{ij} W'_\mu \bar{f}^i \gamma^\mu (1 \pm \gamma_5) f^j + \text{H.c.}, \quad (1)$$

where g' plays the role of g_{SM} from the Standard Model (SM), and V'_{ij} is a generalized Cabibbo-Kobayashi-Maskawa (CKM) matrix that also allows for lepton generation mixing.

Searches for W' bosons decaying to a top and bottom quark final state [1] have been performed since Run I of the Fermilab Tevatron [2]. The now standard analysis involves searching for a $lvbj$ invariant mass peak, where each object is isolated, and applies loose W and top-mass constraints to reduce backgrounds [1]. Last year we updated the standard analysis to optimize the search for the 7 TeV and 8 TeV runs of the Large Hadron Collider [3]. The ATLAS [4] and CMS [5] Collaborations used the cross sections of Ref. [3] to publish strong lower bounds on the mass of a W' boson. Current mass limits are ~ 2000 GeV if the W' boson has Standard Model-like (SM) couplings.

A previous full simulation study of the reach at a 14 TeV pp collider was performed in 2003 [6]. The conclusions of that study showed the mass reach with SM-like couplings should be ~ 5 – 5.5 TeV, or about $1/3$ of the collider energy. This limit is directly attributable to the loss of quark-antiquark luminosity in a pp collider at large- x proton momentum fraction. More important, Ref. [6] presented the model-independent reach in the coupling ratio g'/g_{SM} versus W' mass, and compared it to a few classes of models. One of the key results of Ref. [1] was that all models with narrow-width W' bosons fall somewhere in this space. Hence, all experimental measurements should produce limits, or a discovery, as a function of g'/g_{SM} versus $M_{W'}$.

Just prior to the final meeting of the Community Summer Study, in Ref. [7] we proposed a new analysis that expands the reach in the tb final state by searching for a “boosted-top” tag, where the Wj decay products of the top quark decay can not be isolated. In addition, we introduced a new “boosted-bottom” tag to identify bottom jets in the TeV range in order to overcome dijet backgrounds. This new search is optimized for TeV dijets with one boosted-top and one boosted-bottom tag. Using this search method it should be possible to extend the existing mass reach for W' bosons with SM-like couplings to about 2600 GeV, and the model-independent reach in coupling ratio g'/g_{SM} by nearly a factor of two if $M_{W'} > 1500$ GeV with existing 8 TeV LHC data. This analysis should be done for the upcoming 14 TeV run of the LHC as well.

This white paper summarizes the leading order (LO) and next-to-leading order (NLO) cross sections and widths for $pp \rightarrow W' \rightarrow tb$ production for the upcoming 14 TeV run of the Large Hadron Collider, and the proposed 33 TeV run examined by the Community Summer Study (CSS). Detailed study of the kinematics has shown that in both the standard [3] and boosted [7] analyses, the NLO correction can be treated as a K -factor times LO. Hence, the tables below list both LO and NLO results for a wide range of possible W' masses.

All cross sections are calculated using the code from Ref. [1], updated for modern parton distribution function (PDF) sets and CKM matrix values. Leading order cross sections are calculated with CTEQ6L1 parton distribution functions (PDFs) [8], while NLO cross sections use CT10 [9] PDFs. The top-quark mass is taken to be 173.2 ± 1 GeV [10], and $\alpha_s = 0.118 \pm 0.001$. Uncertainties due to the top-quark mass and α_s are negligible ($< 1\%$) for the W' masses shown. Estimates of the NLO theoretical uncertainty due to choice of scale is also comparatively small (1–2%) in all cases. Hence, the central values of the predictions are shown to three decimal places. Using the standard CTEQ

*Electronic address: Zack.Sullivan@IIT.edu

Modified Tolerance Method for error determination (first published for this cross section [1]) and CT10, the current PDF uncertainty ranges from 10–30% depending on the W' mass, and whether it is right- or left-handed. These uncertainties should come down with improved fitting of large- x PDFs using LHC data.

Cross sections are calculated for $t\bar{b}$ and $\bar{t}b$ separately, for both 14 TeV and 33 TeV pp colliders. Cross sections at 14 TeV for right-handed W'_R bosons are listed by mass at LO and NLO in femtobarns in Table I. Current PDF uncertainties are shown for the NLO cross sections. Cross sections for left-handed W'_L bosons are listed by mass at LO and NLO in femtobarns in Table II. Note, the left-handed cross sections assume *no interference* with the standard model production process. See Sec. II of Ref. [3] for a description of how to use the left-handed normalizations. Cross sections at 33 TeV are listed for W'_R and W'_L in Tables III and IV, respectively. LO and NLO total widths for the W' bosons are also provided in all tables for use in simulation studies.

Many models that attempt to explain electroweak symmetry breaking predict the existence of W' bosons. This white paper provides the LO and NLO cross sections and total widths necessary to determine the model-independent reach at 14 TeV and 33 TeV pp colliders. Since resonant production is expected up to $\sim 1/3$ of the collider energy, a 33 TeV pp collider should be able to observe or exclude most perturbative models with W' bosons with masses below 10 TeV.

Acknowledgments

This work is supported by the U.S. Department of Energy under Contract No. DE-SC0008347.

-
- [1] Zack Sullivan, Phys. Rev. D **66**, 075011 (2002), [hep-ph/0207290].
 - [2] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **90**, 081802 (2003), [hep-ex/0209030].
 - [3] Daniel Duffy and Zack Sullivan, Phys. Rev. D **86**, 075018 (2012), [arXiv:1208.4858 [hep-ph]].
 - [4] G. Aad *et al.* (ATLAS Collaboration), Phys. Rev. Lett. **109**, 081801 (2012), [arXiv:1205.1016 [hep-ex]].
 - [5] S. Chatrchyan *et al.* (CMS Collaboration), Phys. Lett. B **718**, 1229 (2013), [arXiv:1208.0956 [hep-ex]].
 - [6] Zack Sullivan, “How to rule out Little Higgs (and constrain many other models) at the LHC,” in *Proceedings of the XXXVIIIth Rencontres de Moriond: QCD and High Energy Hadronic Interactions*, Les Arcs, Savoie, France, March 22–29, 2003, edited by Étienne Augé and Jean Trân Thanh Vân (Thê Giói Publishers, Hanoi, 2003), p. 379, [hep-ph/0306266].
 - [7] Daniel Duffy and Zack Sullivan, arXiv:1307.1820 [hep-ph], submitted to Phys. Rev. Lett..
 - [8] J. Pumplin *et al.*, J. High Energy Phys. **0207**, 012 (2002), [hep-ph/0201195].
 - [9] H. -L. Lai *et al.*, Phys. Rev. D **82**, 074024 (2010), [arXiv:1007.2241 [hep-ph]].
 - [10] T. Aaltonen *et al.* (CDF and D0 Collaborations), Phys. Rev. D **86**, 092003 (2012), [arXiv:1207.1069 [hep-ex]].

TABLE I: LO and NLO total width in (GeV) and cross section with PDF errors in (fb) vs. W'_R mass for $pp \rightarrow W'_R \rightarrow t\bar{b}(\bar{t}b)$ at $\sqrt{S} = 14$ TeV, where the decay to leptons is not allowed.

Mass (GeV)	Γ_{LO} (GeV)	$\sigma_{LO}^{t\bar{b}}$ (fb)	$\sigma_{LO}^{\bar{t}b}$ (fb)	Γ_{NLO} (GeV)	$\sigma_{NLO}^{t\bar{b}}$ (fb)	$\sigma_{NLO}^{\bar{t}b}$ (fb)
500	11.97	87300	48400	12.47	112000 $^{+7600}_{-5200}$	62700 $^{+5800}_{-1600}$
750	18.58	22700	11300	19.24	29000 $^{+1400}_{-1800}$	14600 $^{+700}_{-700}$
1000	25.07	7960	3580	25.88	9970 $^{+960}_{-350}$	4680 $^{+270}_{-300}$
1250	31.51	3350	1380	32.46	4210 $^{+200}_{-420}$	1780 $^{+170}_{-100}$
1500	37.93	1580	600	39.03	1950 $^{+120}_{-180}$	779 $^{+67}_{-64}$
1750	44.33	805	284	45.58	984 $^{+58}_{-116}$	369 $^{+33}_{-40}$
2000	50.72	432	142	52.11	514 $^{+60}_{-42}$	186 $^{+20}_{-17}$
2250	57.11	242	75.5	58.65	286 $^{+27}_{-33}$	98.7 $^{+11.0}_{-11.1}$
2500	63.49	139	41.4	65.17	163 $^{+14.0}_{-26.3}$	54.3 $^{+6.8}_{-6.8}$
2750	69.86	81.6	23.4	71.70	93.9 $^{+13.0}_{-13.7}$	30.9 $^{+4.2}_{-4.4}$
3000	76.24	48.8	13.6	78.22	55.5 $^{+8.1}_{-9.1}$	18.1 $^{+3.0}_{-2.8}$
3250	82.61	29.6	8.06	84.73	33.3 $^{+5.3}_{-6.5}$	10.8 $^{+1.8}_{-1.9}$
3500	88.99	18.1	4.88	91.23	20.2 $^{+3.5}_{-4.3}$	6.57 $^{+1.43}_{-1.02}$
3750	95.36	11.2	3.02	97.76	12.4 $^{+2.5}_{-2.6}$	4.13 $^{+0.84}_{-0.82}$
4000	101.7	7.05	1.90	104.3	7.72 $^{+1.70}_{-1.82}$	2.62 $^{+0.63}_{-0.49}$
4250	108.1	4.47	1.226	110.8	4.86 $^{+1.15}_{-1.17}$	1.71 $^{+0.41}_{-0.34}$
4500	114.5	2.87	0.806	117.3	3.13 $^{+0.74}_{-0.80}$	1.13 $^{+0.29}_{-0.21}$
4750	120.8	1.87	0.543	123.8	2.05 $^{+0.52}_{-0.51}$	0.765 $^{+0.205}_{-0.137}$
5000	127.2	1.24	0.375	130.3	1.37 $^{+0.35}_{-0.33}$	0.535 $^{+0.126}_{-0.104}$
5250	133.6	0.841	0.266	136.8	0.945 $^{+0.239}_{-0.211}$	0.380 $^{+0.091}_{-0.065}$
5500	139.9	0.586	0.194	143.3	0.671 $^{+0.161}_{-0.136}$	0.278 $^{+0.062}_{-0.044}$
5750	146.3	0.419	0.145	149.8	0.493 $^{+0.103}_{-0.095}$	0.209 $^{+0.042}_{-0.030}$
6000	152.7	0.309	0.112	156.3	0.371 $^{+0.080}_{-0.054}$	0.161 $^{+0.028}_{-0.022}$
6250	159.0	0.235	0.088	162.8	0.290 $^{+0.050}_{-0.040}$	0.127 $^{+0.020}_{-0.014}$
6500	165.4	0.184	0.071	169.3	0.231 $^{+0.035}_{-0.027}$	0.102 $^{+0.014}_{-0.010}$
6750	171.8	0.148	0.059	175.8	0.188 $^{+0.026}_{-0.019}$	0.084 $^{+0.009}_{-0.008}$
7000	178.1	0.121	0.049	182.3	0.157 $^{+0.018}_{-0.014}$	0.070 $^{+0.008}_{-0.006}$

TABLE II: LO and NLO total width in (GeV) and cross section with PDF errors in (fb) vs. W'_L mass for $pp \rightarrow W'_L \rightarrow t\bar{b}(\bar{t}b)$ at $\sqrt{S} = 14$ TeV, where the decay to leptons is allowed, but no interference is included.

Mass (GeV)	Γ_{LO} (GeV)	$\sigma_{LO}^{t\bar{b}}$ (fb)	$\sigma_{LO}^{\bar{t}b}$ (fb)	Γ_{NLO} (GeV)	$\sigma_{NLO}^{t\bar{b}}$ (fb)	$\sigma_{NLO}^{\bar{t}b}$ (fb)
500	16.21	64100	35600	16.71	84300 ⁺²⁰⁰⁰ ₋₆₉₀₀	47200 ⁺²⁰⁰⁰ ₋₃₂₀₀
750	24.95	16800	8370	25.60	21700 ⁺¹²⁰⁰ ₋₁₀₀₀	11000 ⁺⁶⁰⁰ ₋₅₀₀
1000	33.56	5950	2680	34.36	7610 ⁺³³⁰ ₋₅₂₀	3510 ⁺²⁷⁰ ₋₂₀₀
1250	42.12	2510	1040	43.07	3190 ⁺¹²⁰ ₋₃₅₀	1350 ⁺¹¹⁰ ₋₉₀
1500	50.66	1190	452	51.76	1480 ⁺⁹⁰ ₋₁₄₀	594 ⁺⁴² ₋₅₁
1750	59.18	606	215	60.43	742 ⁺⁴⁸ ₋₆₉	282 ⁺²¹ ₋₂₈
2000	67.69	326	108	69.08	395 ⁺²⁸ ₋₄₄	143 ⁺¹³ ₋₁₆
2250	76.20	183	57.5	77.74	219 ⁺¹⁷ ₋₃₁	76.1 ^{+7.2} _{-9.2}
2500	84.70	105	31.7	86.39	124 ⁺¹⁶ ₋₁₄	41.9 ^{+5.3} _{-5.0}
2750	93.20	62.2	18.1	95.03	72.7 ^{+8.0} _{-12.0}	24.0 ^{+3.4} _{-3.2}
3000	101.7	37.4	10.6	103.7	42.9 ^{+6.0} _{-6.7}	14.2 ^{+2.1} _{-2.2}
3250	110.2	22.8	6.33	112.3	25.9 ^{+4.1} _{-4.6}	8.52 ^{+1.52} _{-1.25}
3500	118.7	14.1	3.88	120.9	15.9 ^{+2.6} _{-3.3}	5.28 ^{+0.98} _{-0.90}
3750	127.2	8.82	2.43	129.6	9.86 ^{+1.87} _{-2.07}	3.32 ^{+0.73} _{-0.54}
4000	135.7	5.59	1.55	138.2	6.23 ^{+1.24} _{-1.38}	2.16 ^{+0.44} _{-0.41}
4250	144.2	3.59	1.02	146.8	3.98 ^{+0.87} _{-0.90}	1.42 ^{+0.30} _{-0.27}
4500	152.6	2.34	0.683	155.5	2.61 ^{+0.58} _{-0.62}	0.958 ^{+0.225} _{-0.168}
4750	161.1	1.55	0.469	164.1	1.74 ^{+0.40} _{-0.39}	0.662 ^{+0.155} _{-0.110}
5000	169.6	1.05	0.331	172.7	1.19 ^{+0.28} _{-0.26}	0.471 ^{+0.103} _{-0.077}
5250	178.1	0.731	0.240	181.4	0.842 ^{+0.186} _{-0.167}	0.343 ^{+0.068} _{-0.056}
5500	186.6	0.521	0.179	190.0	0.612 ^{+0.126} _{-0.113}	0.256 ^{+0.048} _{-0.036}
5750	195.1	0.382	0.136	198.6	0.459 ^{+0.082} _{-0.077}	0.195 ^{+0.035} _{-0.024}
6000	203.6	0.288	0.107	207.2	0.353 ^{+0.056} _{-0.054}	0.153 ^{+0.023} _{-0.018}
6250	212.1	0.223	0.085	215.9	0.277 ^{+0.043} _{-0.031}	0.122 ^{+0.017} _{-0.012}
6500	220.6	0.177	0.070	224.5	0.225 ^{+0.029} _{-0.024}	0.100 ^{+0.011} _{-0.010}
6750	229.0	0.144	0.058	233.1	0.185 ^{+0.020} _{-0.019}	0.082 ^{+0.008} _{-0.007}
7000	237.5	0.119	0.048	241.7	0.155 ^{+0.014} _{-0.015}	0.069 ^{+0.007} _{-0.005}

TABLE III: LO and NLO total width in (GeV) and cross section in (fb) vs. W'_R mass for $pp \rightarrow W'_R \rightarrow t\bar{b}(\bar{t}b)$ at $\sqrt{S} = 33$ TeV, where the decay to leptons is not allowed.

Mass (GeV)	Γ_{LO} (GeV)	$\sigma_{LO}^{t\bar{b}}$ (fb)	$\sigma_{LO}^{\bar{t}b}$ (fb)	Γ_{NLO} (GeV)	$\sigma_{NLO}^{t\bar{b}}$ (fb)	$\sigma_{NLO}^{\bar{t}b}$ (fb)
500	11.97	281000	183000	12.47	345000	229000
1000	25.07	31100	17900	25.88	38400	22900
1500	37.93	7610	3960	39.03	9560	5030
2000	50.72	2660	1270	52.11	3290	1620
2500	63.49	1130	496	65.17	1410	638
3000	76.24	543	221	78.22	665	281
4000	101.7	157	56.3	104.3	188	72.1
5000	127.2	54.3	17.5	130.3	63.7	22.6
6000	152.7	20.8	6.23	156.3	24.0	8.09
7000	178.1	8.51	2.43	182.3	9.53	3.18
8000	203.6	3.64	1.02	208.4	4.00	1.35
9000	229.1	1.62	0.457	234.4	1.76	0.609
10000	254.5	0.750	0.220	260.4	0.812	0.297
11000	280.0	0.363	0.114	286.3	0.395	0.155
12000	305.4	0.187	0.064	312.3	0.208	0.087
13000	330.9	0.104	0.039	338.3	0.119	0.053
14000	356.4	0.063	0.025	364.3	0.074	0.035
15000	381.8	0.041	0.018	390.3	0.051	0.024

TABLE IV: LO and NLO total width in (GeV) and cross section with PDF errors in (fb) vs. W'_L mass for $pp \rightarrow W'_L \rightarrow t\bar{b}(\bar{t}b)$ at $\sqrt{S} = 33$ TeV, where the decay to leptons is allowed, but no interference is included.

Mass (GeV)	Γ_{LO} (GeV)	$\sigma_{LO}^{t\bar{b}}$ (fb)	$\sigma_{LO}^{\bar{t}b}$ (fb)	Γ_{NLO} (GeV)	$\sigma_{NLO}^{t\bar{b}}$ (fb)	$\sigma_{NLO}^{\bar{t}b}$ (fb)
500	16.21	206000	134000	16.71	254000	172000
1000	33.56	23200	13400	34.36	29600	17100
1500	50.66	5700	2970	51.76	7210	3850
2000	67.69	1990	952	69.08	2490	1220
2500	84.70	848	374	86.39	1050	488
3000	101.7	408	167	103.7	504	216
4000	135.7	119	42.8	138.2	143	55.7
5000	169.6	41.2	13.5	172.7	48.5	17.4
6000	203.6	15.9	4.83	207.2	18.4	6.32
7000	237.5	6.56	1.91	241.7	7.43	2.51
8000	271.5	2.85	0.818	276.2	3.18	1.09
9000	305.4	1.29	0.377	310.7	1.43	0.506
10000	339.4	0.612	0.187	345.2	0.676	0.253
11000	373.3	0.306	0.100	379.7	0.340	0.137
12000	407.3	0.164	0.058	414.1	0.186	0.079
13000	441.2	0.095	0.036	448.6	0.111	0.050
14000	475.1	0.059	0.024	483.1	0.071	0.033
15000	509.1	0.040	0.017	517.5	0.049	0.023