

# Meson dominance implications for the bottomness preserving decays of the $B_c^+$ meson

Peter Lichard

*Institute of Physics, Silesian University in Opava,  
746 01 Opava, Czech Republic*  
and

*Institute of Experimental and Applied Physics,  
Czech Technical University in Prague,  
128 00 Prague, Czech Republic*

Encouraged by a recent observation of the  $B_c^+ \rightarrow B_s^0 + \pi^+$  decay by the LHCb collaboration we present the meson dominance predictions for other weak decays of the  $B_c^+$  into  $B_s^0$  or  $B^0$  in the form of branching ratios to the observed decay.

PACS numbers: 13.25.Hw, 13.20.He, 14.40.Nd

The LHCb collaboration at the CERN LHC collider has recently announced [1] the observation of the bottomness-preserving (BP) decay  $B_c^+ \rightarrow B_s^0 + \pi^+$  with significance in excess of five standard deviations independently in two decay channels of  $B_s^0$ . This sets a hope that similar BP decays of the  $B_c^+$  meson may be found either in the already accumulated data or after the LHC reopens with higher energy and luminosity in 2015.

In this note we present the estimates of the branching ratios of other BP decays of the  $B_c^+$  relative to the already observed one. We use the meson dominance (MD) model [2], which describes well [3] the meson decays that fall into the “external W-emission” category according to the quark-diagram nomenclature [4].

The decay of the  $B_c^+$  (or, to be general,  $P_1$ ) into the neutral meson  $B_s^0$  ( $P_2$ ) and the  $\pi^+$  ( $P_3$ ) is described by a diagram depicted in Fig. 1. The diagram contains first a strong interaction vertex entered by  $P_1$ ,  $P_2$ , and a positive vector meson  $V$  (here  $D_s^{*+}$ ) with flavor quantum numbers required by the conservation laws. The vector meson then couples to the gauge boson  $W^+$ , which in turn couples to the outgoing pseudoscalar meson  $P_3$  (here  $\pi^+$ ). The corresponding partial decay width is given by

$$\Gamma(P_1 \rightarrow P_2 + P_3) = \frac{G_F^2 X_{P_1 P_2 V} Z_{P_3}}{16\pi m_1^3} (x-y)^2 \lambda^{1/2}(x, y, m_3^2), \quad (1)$$

where  $x = m_1^2$ ,  $y = m_2^2$ , and function  $\lambda$  is defined by  $\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz$ . The definition of the dimensionless parameter  $X_{P_1 P_2 V}$

$$X_{P_1 P_2 V} = \left| w_V V_V \frac{g_{V P_1 P_2}}{g_\rho} \right|^2$$

includes the strong coupling constant  $g_{V P_1 P_2}$ , the  $\rho\pi\pi$




FIG. 1. Decay  $B_c^+ \rightarrow B_s^0 + \pi^+$  in the MD model.

coupling constant  $g_\rho$ , the element  $V_V$  of the CKM matrix pertinent to valence quark and antiquark of the meson  $V$ , and a dimensionless parameter  $w_V \approx 1$  characterizing the deviation of the  $V - W^+$  coupling from the  $\rho^+ - W^+$  one. At present, there is no chance of getting the value of  $X_{B_c^+ B_s^0 D_s^{*+}}$  from its components. But as we are going to relate the branching fractions of the various  $B_c^+ \rightarrow B_s^0$  transitions, this unknown quantity will cancel.

The parameter  $Z_P$ , which is another important ingredient of Eq. (1), is defined in terms of the pseudoscalar decay constant  $f_P$  and the CKM matrix element  $V_P$  corresponding to the valence quark composition of a particular pseudoscalar meson  $P$  by

$$Z_P = |f_P V_P|^2. \quad (2)$$

Its value is determined from the muonic decay width given by the formula [2]

$$\Gamma(P \rightarrow \mu^+ + \nu_\mu) = \frac{G_F^2}{8\pi} Z_P m_\mu^2 m_P \left( 1 - \frac{m_\mu^2}{m_P^2} \right)^2 \times [1 + \mathcal{O}(\alpha)].$$

We have included the radiative corrections  $\mathcal{O}(\alpha)$  following [5]. The results obtained by using the experimental muonic decay widths [6] are shown in Table I.

Now we are ready to calculate the branching ratio

$$\frac{\mathcal{B}(B_c^+ \rightarrow B_s^0 + K^+)}{\mathcal{B}(B_c^+ \rightarrow B_s^0 + \pi^+)} = \frac{Z_{K^+}}{Z_{\pi^+}} \sqrt{\frac{\lambda(m_{B_c^+}^2, m_{B_s^0}^2, m_{K^+}^2)}{\lambda(m_{B_c^+}^2, m_{B_s^0}^2, m_{\pi^+}^2)}} = (6.470 \pm 0.025) \times 10^{-2},$$

TABLE I. Parameters  $Z_P$  characterizing the coupling of pseudoscalar mesons to the charged gauge boson and their sources. For definition, see Eq. (2).

P	$Z_P$ (MeV <sup>2</sup> )	Source
$\pi^+$	$(1.6158 \pm 0.0019) \times 10^4$	$\pi^+ \rightarrow \mu^+ \nu_\mu$
$K^+$	$(1.2307 \pm 0.0033) \times 10^3$	$K^+ \rightarrow \mu^+ \nu_\mu$




FIG. 2. The semileptonic decay of the  $B_c^+$  to  $B_s^0$  meson in the MD model.

which shows that the decay  $B_c^+ \rightarrow B_s^0 + K^+$  will not be probably observed soon.

Let us turn now to the semileptonic decays. The decay  $B_c^+ (P_1) \rightarrow B_s^0 (P_2) + \ell^+ + \nu_\ell$  is in the MD model described by diagram in Fig. 2. The differential decay width in  $t$ , which is the square of the four-momentum transfer from  $P_1$  to  $P_2$ , is given by

$$\frac{d\Gamma}{dt} = \frac{G_F^2 X_{P_1 P_2 V}}{3(4\pi m_1)^3} \frac{(t-z)^2}{t^3} \lambda^{1/2}(x, y, t) \left( \frac{r}{r-t} \right)^2 \times \left[ (2t+z)\lambda(x, y, t) + \frac{3z}{r^2}(x-y)^2(r-t)^2 \right]. \quad (3)$$

Newly defined parameters are  $r = m_V^2$  and  $z = m_\ell^2$ . Formula (3) is a simplified but equivalent version of Eq. (4.3) from Ref. [2]. After factorizing out the unknown multiplication constant we can perform the numerical integration of (3) and get the total semileptonic decay width (up to that multiplication constant). After dividing it by (1), the unknown  $X$ -factor cancels and we are getting

$$\frac{\mathcal{B}(B_c^+ \rightarrow B_s^0 + e^+ + \nu_e)}{\mathcal{B}(B_c^+ \rightarrow B_s^0 + \pi^+)} = 0.392 \pm 0.006$$

and

$$\frac{\mathcal{B}(B_c^+ \rightarrow B_s^0 + \mu^+ + \nu_\mu)}{\mathcal{B}(B_c^+ \rightarrow B_s^0 + \pi^+)} = 0.367 \pm 0.006.$$

We can also compare the branching fractions of the modes with different  $B$ -mesons in the final states, *e.g.*,

$$\frac{\mathcal{B}(B_c^+ \rightarrow B^0 + \pi^+)}{\mathcal{B}(B_c^+ \rightarrow B_s^0 + \pi^+)} = R_X \frac{m_{B_c^+}^2 - m_{B^0}^2}{m_{B_c^+}^2 - m_{B_s^0}^2} \times \sqrt{\frac{\lambda(m_{B_c^+}^2, m_{B^0}^2, m_{\pi^+}^2)}{\lambda(m_{B_c^+}^2, m_{B_s^0}^2, m_{\pi^+}^2)}},$$

where

$$R_X = \frac{X_{B_c^+ B^0 D^{*+}}}{X_{B_c^+ B_s^0 D^{*+}}}$$

is an unknown quantity. We can get a crude estimate of its value if assuming that the light flavor SU(3) symmetry is not badly broken. Then we can write

$$R_X \approx \left| \frac{V_{cd}}{V_{cs}} \right|^2 \approx 5.3 \times 10^{-2}$$

and

$$\frac{\mathcal{B}(B_c^+ \rightarrow B^0 + \pi^+)}{\mathcal{B}(B_c^+ \rightarrow B_s^0 + \pi^+)} \approx 7 \times 10^{-2}.$$

The semileptonic transitions of  $B_c^+$  to  $B^0$  will be suppressed relative to those to  $B_s^0$  by the same factor.

To conclude: The semileptonic decays of the  $B_c^+$  to  $B_s^0$  meson are, based on the magnitude of their branching fractions, the best candidates for the experimental observation. The branching fraction of  $B_c^+ \rightarrow B_s^0 + e^+ + \nu_e$  ( $B_c^+ \rightarrow B_s^0 + \mu^+ + \nu_\mu$ ) is about thirty-nine (thirty-seven) per cent of that of the already observed decay  $B_c^+ \rightarrow B_s^0 + \pi^+$ .

I am indebted to Dr. Tim Gershon for useful correspondence and to Dr. Josef Jurán for discussions. This work was supported by the Research Program MSM6840770029 of the Czech Ministry of Education, Youth and Sports.

- 
- [1] R. Aaij *et al.* (LHCb Collaboration), arXiv: 1308.4544 [hep-ex].  
 [2] P. Lichard, Phys. Rev. D **55**, 5385 (1997).  
 [3] P. Lichard, in preparation.

- [4] L.-L. Chau, Phys. Rep. **95**, 1 (1983).  
 [5] V. Cirigliano and I. Rosell, JHEP **10**, 5 (2007).  
 [6] J. Beringer *et al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).