Hamiltonian systems with an infinite number of localized travelling waves

Georgy L. Alfimov[a], Elina V. Medvedeva[a], and Dmitry E. Pelinovsky[b,c]
[a] *Moscow Institute of Electronic Engineering, Zelenograd, Moscow, 124498, Russia*
[b] *Department of Mathematics, McMaster University, Hamilton, Ontario, Canada and*
[c] *Department of Applied Mathematics, Nizhny Novgorod State Technical University, Nizhny Novgorod, Russia*

In many Hamiltonian systems, propagation of steadily travelling solitons or kinks is prohibited because of resonances with linear excitations. We show that Hamiltonian systems with resonances may admit an infinite number of travelling solitons or kinks if the closest to the real axis singularities in the complex upper half-plane of limiting asymptotic solution are of the form $z_\pm = \pm\alpha + i\beta$, $\alpha \neq 0$. This quite a general statement is illustrated by examples of the fifth-order Korteweg–de Vries-type equation, the discrete cubic-quintic Klein–Gordon equation, and the nonlocal double sine–Gordon equations.

arXiv:1309.0183v1 [nlin.PS] 1 Sep 2013

Introduction. Nonlinear localized travelling waves such as bright or dark solitons are key concepts for many branches of modern physics, including nonlinear optics, theory of magnets, theory of Josephson junctions, etc. It is known that in many dispersive systems the presence of these nonlinear entities is strongly restricted due to *resonances* with linear excitations. These resonances take place in Hamiltonian systems of various origin, such as the fifth-order Korteweg–de Vries equation [1], nonlinear lattices [2–4] and models with complex dispersion and nonlocal interactions [5, 6]. As a result, it is quite typical that in such Hamiltonian systems the localized excitations either do not exist at all or they only exist for specific values of some external parameters. In the last case the nonlinear excitations are called *embedded solitons* (i.e. solitons "embedded" into the spectrum of linear waves). These embedded solitons have been discovered in hydrodynamics, nonlinear optics and other fields of modern physics [7].

To give an example, consider an operator equation

$$L_\varepsilon u = F(u), \tag{1}$$

for a function $u(\xi)$ where L_ε is a Fourier multiplier operator in ξ space with even symbol $\widehat{L}(k)$ in k space, $F(u)$ is a nonlinear function and ε is a parameter. The prototypical examples of problems leading to Eq.(1) are the generalized Korteweg–de Vries equation,

$$u_t + (F(u))_x + M_\varepsilon u_x = 0, \tag{2}$$

or discrete or nonlocal Klein-Gordon equations,

$$u_{tt} - M_\varepsilon u + F(u) = 0, \tag{3}$$

for $u(x,t)$, where M_ε is a Fourier multiplier operator in x space. Above $\xi = x - vt$ is the travelling wave coordinate and the operator L_ε in Eq.(1) includes both M_ε and v. We assume that in the both cases $\varepsilon = 0$ implies a degeneration of the problem with $L_0 = \partial_\xi^2$.

Consider a solitary wave $u(\xi)$ which is asymptotic to the equilibrium state $u \equiv 0$ as $\xi \to \pm\infty$ (the case of kink wave which is asymptotic to a pair of equilibrium states $u \equiv u_\pm$ as $\xi \to \pm\infty$ can be analyzed in a similar way). Then the resonances correspond to the real roots of the dispersion equation near $u \equiv 0$

$$\widehat{L}_\varepsilon(k) = F'(0). \tag{4}$$

If for some value of ε, there exist a *single* pair of real roots $k = \pm k_0$ in Eq.(4), we are in situation when the resonance prohibits propagation of regular solitons in Eq.(2) and Eq.(3) and the embedded solitons may appear. In this case, the velocity v of the soliton, typically, *is not arbitrary* but should be "adjusted" to avoid "gluing" with linear modes. In general, v belongs to some discrete set. This set may be empty (i.e, no localized waves propagate), or include finite or infinite number of values. The case when Eq.(4) has more then one pair of real roots is more complex and the presence of localized excitations in this case is highly doubtful.

In this paper we address the following question: *Are there some conditions which would indicate to existence of infinitely many embedded solitons described by Eq.(1)? If this is possible, can we describe this infinite set asymptotically?* As a result, we present conditions for existence of countable infinite sequence of embedded solitons. The main assumption is that the limiting solution of Eq.(1) as $\varepsilon \to 0$, being extended in a complex plane, should have a pair of symmetric singularities in the upper half-plane. We give an asymptotic formula for values $\{\varepsilon_n\}$ as $n \to \infty$, for which embedded solitons exist. In terms of Eq.(2) and Eq.(3) this means a presence of an infinite number of velocities v for nonlinear localized excitations.

Up to the moment no rigorous proof of this asymptotic theory has been found. We give some heuristic explanation of the mechanism behind the asymptotic formula and illustrate this result with three numerical examples. Surprisingly, it has been observed that in some cases the asymptotic formula predicts the parameters even of lowest embedded solitons from this sequence with reasonable accuracy.

We note that the idea that two symmetric singularities in the upper half-plane can be related to the countable infinite sequence of tangential intersections of stable and unstable manifolds can be found in [8] for the primary intersection point of the two-dimensional symplectic maps. In this paper we generalize this principle to more general

class of physically relevant systems.

Main result. Consider Eq. (1), where $u(\xi)$ is real-valued function defined on $\mathbb{R}$. Assume that L_ε is a real operator which depends continuously on real parameter ε and satisfies $L_0 = \partial_\xi^2$. The Fourier symbol $\hat{L}_\varepsilon(k)$ is supposed to be an even function of k.

Assume now that (a) the equation $F(u) = 0$ has zero solution $u = 0$ with $F'(0) > 0$; (b) the dispersion equation (4) has only one pair of real roots $k = \pm k(\varepsilon)$ such that $k(\varepsilon) \to \infty$ as $\varepsilon \to 0$; and (c) the equation

$$u'' = F(u) \tag{5}$$

has an even localized solution $\tilde{u}(\xi)$ such that $\tilde{u}(\xi) \to 0$ as $\xi \to \pm\infty$. In addition, the key assumption of our asymptotic theory is that the solution $\tilde{u}(\xi)$ can be continued into the complex plane and the closest to the real axis singularities of $\tilde{u}(\xi)$ in the upper half-plane are given by the pair $z_\pm = \pm\alpha + i\beta$, with $\alpha, \beta > 0$, which is symmetric with respect to the imaginary axis.

Then, we expect the existence of an infinite sequence of values $\{\varepsilon_n\}$ such that for each $\varepsilon = \varepsilon_n$, Eq.(1) has a soliton solution $u(\xi)$ with $u(\xi) \to 0$ as $\xi \to \pm\infty$, and this sequence obeys the following asymptotic law

$$k(\varepsilon_n) \sim (n\pi + \varphi_0)/\alpha \tag{6}$$

where φ_0 is a phase constant that depends on L_ε and $\tilde{u}$.

This result can be extended naturally to the case of kink solutions of Eq.(1) connecting a pair of equilibrium states $u = u_\pm$, such that $F(u_\pm) = 0$ and $F'(u_-) = F'(u_+) > 0$. In this case, $F'(u_\pm)$ appears instead of $F'(0)$ in the dispersion relation (4), whereas the differential equation (5) is assumed to have a kink solution $\tilde{u}(\xi)$ such that $\tilde{u}(\xi) \to u_\pm$ as $\xi \to \pm\infty$.

Justification. Let us give some heuristic arguments for justification of the main result. Introduce $v(\xi) = u(\xi) - \tilde{u}(\xi)$, $N(v) = F(\tilde{u} + v) - F(\tilde{u}) - F'(\tilde{u})v$ and $H_\varepsilon = L_0 - L_\varepsilon$. Then, we have

$$(L_\varepsilon - F'(\tilde{u}))v = H_\varepsilon\tilde{u} + N(v). \tag{7}$$

Based on our assumptions, we take as granted that (a) the function $H_\varepsilon\tilde{u}(\xi)$ can be continued into the upper complex half-plane and its closest to the real axis singularities are $z_\pm = \pm\alpha + i\beta$, whereas (b) the homogeneous linearized equation

$$(L_\varepsilon - F'(\tilde{u}))v = 0 \tag{8}$$

has a pair of solutions $\varphi_\varepsilon^\pm(\xi) = e^{\pm ik(\varepsilon)\xi}\psi_\varepsilon^\pm(\xi)$, where $k(\varepsilon)$ is the only positive root of Eq.(4) and $\psi_\varepsilon^\pm(\xi) \to 1$ as $\varepsilon \to 0$. The latter hypothesis is quite natural, since $v(\xi) = e^{\pm ik(\varepsilon)\xi}$ are solutions of $(L_\varepsilon - F'(0))v = 0$.

Then for small ε, the term $H_\varepsilon\tilde{u}$ in right-hand side of the inhomogeneous equation (7) dominates and the solvability condition for this inhomogeneous equation [9] can be written approximately as the orthogonality condition

$$0 = J_\pm(\varepsilon) \;\approx\; \int_{-\infty}^{\infty} e^{\pm ik(\varepsilon)\xi} H_\varepsilon\tilde{u}(\xi)\, d\xi. \tag{9}$$

The asymptotic value of the integral in (9) as $\varepsilon \to 0$ is determined by the closest to the real axis singularities of integrand in the complex plane (the *Darboux principle*, see [10]). Since $H_\varepsilon\tilde{u}(\xi)$ is even, bounded, and real-valued for real ξ, the main contribution comes from $z_\pm = \pm\alpha + i\beta$ and $J_+ = J_- \equiv J$.

In the simplest case, when the integrand has poles of order n in the points $z_\pm$, the result is simply a sum of the residues in these poles multiplied by $2\pi i$. In a more complicated case, the singularities $z_\pm$ of $H_\varepsilon\tilde{u}$ can be rational or transcendental branch points. For both cases, since $H_\varepsilon\tilde{u}(x)$ is even and real for real x, we can write

$$H_\varepsilon\tilde{u}(\xi) \sim C(\varepsilon)e^{i\pi\kappa/2}(\xi - z_+)^\kappa, \quad \xi \to z_+ \tag{10}$$

$$H_\varepsilon\tilde{u}(\xi) \sim \overline{C(\varepsilon)}e^{i\pi\kappa/2}(\xi - z_-)^\kappa, \quad \xi \to z_-, \tag{11}$$

where κ is a real number, $\kappa \neq 0, 1, 2, \ldots$. It is naturally to assume that $C(\varepsilon) \sim C_0\varepsilon^q$ as $\varepsilon \to 0$ for some values of C_0 and q. Then applying standard formulas [13] in the asymptotic limit $k(\varepsilon) \to \infty$ as $\varepsilon \to 0$, we conclude that

$$J(\varepsilon) \;\sim\; \frac{4\pi\varepsilon^q|C_0|e^{-\beta k(\varepsilon)}}{(k(\varepsilon))^{\kappa+1}\Gamma(-\kappa)}\cos\left(\alpha k(\varepsilon) + \phi_0\right), \tag{12}$$

where $\phi_0 = \arg(C_0)$. Consequently, zeros of $J(\varepsilon)$ obey the asymptotic formula (6) with $\varphi_0 = \pi/2 - \phi_0$.

If $\tilde{u}(\xi)$ is a symmetric kink solution of Eq.(5), the reasoning remains the same up to the point that $H_\varepsilon\tilde{u}(\xi)$ is now odd in ξ.

Examples. The validity of the main result has been confirmed by many numerical studies. Below we give three illustrative examples that concern problems of different physical origins.

Example 1. Consider the equation

$$\varepsilon^2 u'''' + u'' - u + ru^2 - u^3 = 0, \tag{13}$$

where ε is a parameter. Eq.(13) arises in hydrodynamics where it describes travelling waves for the fifth-order KdV equation [1]. If $\varepsilon = 0$ and $r > 3/\sqrt{2}$, Eq.(13) has an exact soliton solution

$$\tilde{u}(\xi) = \frac{3}{\sqrt{r^2 - \frac{9}{2}}\cosh\xi + r}. \tag{14}$$

Closest to the real axis singularities in the upper complex half-plane are the simple poles

$$z_\pm(r) = \pm\operatorname{arctanh}\frac{3}{\sqrt{2}r} + i\pi. \tag{15}$$

Since $H_\varepsilon = -\varepsilon^2\partial_\xi^4$, we note that the singularities of $H_\varepsilon\tilde{u}$ are situated in (15) and are poles of order $n = 5 = -\kappa$.

The expansion (10) holds with $C(\varepsilon) = \varepsilon^2 C_0$, where C_0 is purely imaginary. Therefore, $\phi_0 = \pi/2$ in Eq. (12) and $\varphi_0 = 0$ in Eq. (6).

The dispersion relation (4) reads as $\varepsilon^2 k^4 - k^2 - 1 = 0$ and it has one positive root $k_0(\varepsilon)$ such that $k_0(\varepsilon) \sim 1/\varepsilon$ as $\varepsilon \to 0$. According to the main result, we expect that there exists an infinite sequence of values $\{\varepsilon_n\}$ such that Eq.(13) has soliton solutions for $\varepsilon = \varepsilon_n$ with the asymptotic formula

$$\pi n \varepsilon_n \sim \alpha = \operatorname{arctanh} \frac{3}{\sqrt{2} r} \quad \text{as} \quad n \to \infty. \tag{16}$$

Numerical computations strongly support this prediction. Fig.1 shows the values $\alpha/(\pi\varepsilon_n)$ which approach to integers for larger values of n. The profiles of the three lowest solitons corresponding to points A, B, and C are shown on the inserts by solid line, together with the limiting soliton (14) by dotted line. The discrepancy reduces quickly for larger values of n.




FIG. 1: Soliton solutions of Eq.(13) with $r = 2.3$. Values of $\alpha/(\pi\varepsilon_n)$ are shown by asterisks. Profiles of the first three solitons are shown on the inserts by solid line, the dotted line shows the limiting soliton (14).

Example 2. Consider the nonlocal double sine-Gordon equation

$$u_{tt} = \int_{\mathbb{R}} K_\varepsilon(|x-y|) u_{yy} dy + \sin u + 2a \sin 2u, \tag{17}$$

where $a > 0$ is a parameter. In particular, this equation arises in nonlocal Josephson electrodynamics where it describes layered structures [6] (the second sine harmonic is important if they include, for instance, ferromagnetic layers, [11]). A list of possible kernels K_ε which arises in Josephson models can be found in [12]. We assume that K_0 is the Dirac distribution such that Eq. (17) with $\varepsilon = 0$ reduces to the classical double sine-Gordon equation. If we denote the Fourier transform of K_ε by $\hat{K}_\varepsilon(k)$, then $\hat{K}_\varepsilon(k) \to \hat{K}_0 = 1$ as $\varepsilon \to 0$.

Travelling wave solutions $u(\xi) = u(x - vt)$ of Eq. (17) satisfy the equation

$$v^2 u_{\xi\xi} = \int_{\mathbb{R}} K_\varepsilon(|\xi - \xi'|) u_{\xi'\xi'} d\xi' + \sin u + 2a \sin 2u. \tag{18}$$

If $\varepsilon = 0$, Eq.(18) reads

$$(1 - v^2) u'' = \sin u + 2a \sin 2u, \tag{19}$$

where $v^2 < 1$ is assumed. We consider 2π-kink solutions with boundary conditions at infinity

$$\lim_{\xi \to -\infty} u(\xi) = 0, \quad \lim_{\xi \to +\infty} u(\xi) = 2\pi.$$

For $a > 0$, Eq. (19) has exact 2π-kink solution

$$\tilde{u}(\xi) = \pi + 2 \arctan\left(\frac{1}{\sqrt{1+4a}} \sinh\left(\frac{\sqrt{1+4a}}{\sqrt{1-v^2}} \xi \right) \right) \tag{20}$$

and the closest singularities to the real axis are the two logarithmic branching points $z_\pm = \pm\alpha + i\beta$, where

$$\alpha = \frac{\sqrt{1-v^2}}{2\sqrt{1+4a}} \operatorname{arccosh}(1+8a), \quad \beta = \frac{\pi\sqrt{1-v^2}}{2\sqrt{1+4a}}.$$

The dispersion relation

$$-v^2 k^2 + k^2 \hat{K}_\varepsilon(k) = 1 + 4a$$

is assumed to have a single pair of real roots $k = \pm k(\varepsilon)$ for all $v^2 < 1$. In particular, if K_ε is the Kac-Baker kernel

$$K_\varepsilon(|\zeta|) = \frac{1}{2\varepsilon} \exp\left(-\frac{|\zeta|}{\varepsilon} \right), \quad \hat{K}_\varepsilon(k) = \frac{1}{1 + \varepsilon^2 k^2}, \tag{21}$$

then this assumption is satisfied and

$$k_0(\varepsilon) \sim \frac{\sqrt{1-v^2}}{\varepsilon v}, \quad \text{as} \quad \varepsilon \to 0.$$

Let us present arguments that Eq.(18) with the kernel (21) admits an infinite sequence of the 2π-kink solutions. We note that this equation can be reduced to the system of differential equations

$$\begin{cases} v^2 u_{\xi\xi} = q + \sin u + 2a \sin 2u, \\ -\varepsilon^2 q_{\xi\xi} + q = u_{\xi\xi}, \end{cases} \tag{22}$$

where an additional variable q is introduced. For the limiting kink $\tilde{u}$, we denote a solution of the second equation of the system (22) by $\tilde{q}$. Now since $H_\varepsilon \tilde{u} = \tilde{u}'' - \tilde{q}$, we understand that $H_\varepsilon \tilde{u}$ has a double pole in $\tilde{u}''$ at $z_\pm = \pm\alpha + i\beta$ in addition to the logarithmic branching points in $\tilde{q}$. Hence $n = 2 = -\kappa$ and the expansion (10) holds with $C(\varepsilon) \to C_0$ as $\varepsilon \to 0$, where C_0 is real. Therefore, $\phi_0 = 0$ in Eq. (12) and $\varphi_0 = \pi/2$ in Eq. (6).

According to the main result, we expect that there exists an infinite sequence of values $\{\varepsilon_n\}$ such that Eq.(18) with the kernel (21) has a 2π-kink solution for $\varepsilon = \varepsilon_n$ with the asymptotic formula as $n \to \infty$,

$$\pi(1+2n)\varepsilon_n \sim \delta = \frac{(1-v^2)\operatorname{arccosh}(1+8a)}{v\sqrt{1+4a}}. \tag{23}$$

Using an appropriate shooting method [14], we compute numerically the values of ε, for which there exist

2π-kink solutions of Eq. (18) with the kernel (21). Numerical calculations strongly confirm the existence of the sequence $\{\varepsilon_n\}$ as well as its asymptotic properties (23). The values of $\delta/(\pi\varepsilon_n)$ for $a = 1/8$ and $v = 0.1$ are given in Table I and approach closer to odd integers for larger values of n.

Evidently, each value ε_n depends on the parameter v; however from the physical viewpoint, the inverse functions $v_n(\varepsilon)$ are more important. Fig.2 represents the dependence of the velocities v_n versus ε for the first three 2π-kink solutions. The corresponding profiles of the 2π-kinks (solid line) at the points A, B and C are shown in the inserts together with the limiting kink (20) (dotted line). The difference between the actual kink and the limiting kink (20) is not visible already for kinks at the points B and C.

$1+2n$	1	3	5	7	9	11
$\delta/(\pi\varepsilon_n)$	3.7168	4.9763	6.3699	7.8595	9.4541	11.1396

TABLE I: The values of $\delta/(\pi\varepsilon_n)$ for which Eq.(18) with the kernel (21) admits 2π-kink solution for $a = 1/8$ and $v = 0.1$.




FIG. 2: Kink solutions of Eq.(18) with the kernel (21). Values of v_n versus ε are shown for the first three solutions. Profiles of the first three kinks are shown in the inserts by solid line, the dotted line shows the limiting kink (20).

Example 3. Discrete Klein-Gordon equation is one of the basic equations describing lattice dynamics in various contexts, from solid state physics to biophysics [2]. Travelling waves of the discrete Klein–Gordon equation satisfy the equation

$$v^2 u_{\xi\xi} = \varepsilon^{-2}(u(\xi+\varepsilon) - 2u(\xi) + u(\xi-\varepsilon)) + F(u), \quad (24)$$

where ε is the spacing between lattice sites and F is a nonlinear function. Bistable nonlinearity

$$F(u) = u(1-u^2)(1+\gamma u^2), \quad \gamma > 0, \quad (25)$$

may support kinks which satisfy the boundary conditions

$$\lim_{\xi\to\pm\infty} u(\xi) = \pm 1.$$

If $\gamma = 0$, Eq.(24) corresponds to the classical ϕ^4 model, where no travelling kinks were previously found [4]. We anticipate that for $\gamma > 0$ and v fixed there exists an infinite sequence of travelling kinks for discrete values of parameter ε.

If $\varepsilon = 0$, Eq.(24) reads

$$(1-v^2)u'' + u(1-u^2)(1+\gamma u^2) = 0, \quad (26)$$

where $v^2 < 1$ is assumed. For $\gamma > 0$, Eq. (26) has exact kink solution

$$\tilde{u}(\xi) = \frac{\sqrt{3+\gamma}\tanh(\eta\xi)}{\sqrt{3(1+\gamma) - 2\gamma\tanh^2(\eta\xi)}}, \ \eta = \frac{\sqrt{1+\gamma}}{\sqrt{2(1-v^2)}} \quad (27)$$

and the closest singularities to the real axis are the two square root branching points $z_\pm = \pm\alpha + i\beta$, where

$$\alpha = \frac{\sqrt{1-v^2}}{\sqrt{2(1+\gamma)}}\operatorname{arccosh}\left(\frac{3+5\gamma}{3+\gamma}\right), \quad \beta = \frac{\pi\sqrt{1-v^2}}{\sqrt{2(1+\gamma)}}.$$

If $\gamma > 0$ the dispersion relation

$$-v^2k^2 + 4\varepsilon^{-2}\sin^2(k\varepsilon/2) + 2(1+\gamma) = 0$$

has a single pair of real roots $k = \pm k(\varepsilon)$ for $v \in (v_0, 1)$ for some $v_0 \equiv v_0(\gamma) > 0$. More than one pair of roots exist (always, an odd number) for $v \in (0, v_0)$. We note that $k(\varepsilon) \sim p_0/\varepsilon$ as $\varepsilon \to 0$, where p_0 is a positive root of the transcendental equation $-v^2p_0^2 + 4\sin^2(p_0/2) = 0$. This equation has a single pair of real roots for $v \in (\tilde{v}_0, 1)$ where $\tilde{v}_0 \approx 0.22$ [4].

For the limiting kink $\tilde{u}$, we have

$$H_\varepsilon\tilde{u} = \tilde{u}'' - \varepsilon^{-2}(\tilde{u}(\xi+\varepsilon) - 2\tilde{u}(\xi) + \tilde{u}(\xi-\varepsilon)).$$

The first term yields the expansion (10) with $\kappa = -5/2$ and $C(\varepsilon) \sim C_0 e^{5i\pi/4}$ as $\varepsilon \to 0$ near the singular points $z_\pm = \pm\alpha + i\beta$, where C_0 is purely imaginary. The second term yields the expansion (10) with $\kappa = -1/2$ and $C(\varepsilon) \sim \varepsilon^{-2}C_0 e^{i\pi/4}$ as $\varepsilon \to 0$, where C_0 is purely imaginary. Both terms yield equal contribution in ε that depends on v. Nevertheless, for both terms, $\phi_0 = -\pi/4$ in Eq. (12) and $\varphi_0 = 3\pi/4$ in Eq. (6).

According to the main result, we expect that there exists an infinite sequence of values $\{\varepsilon_n\}$ such that Eq.(24) has a kink solution for ε_n with the asymptotic formula

$$\pi(4n+3)\varepsilon_n \sim \chi = 4p_0\alpha \quad \text{as} \quad n \to \infty. \quad (28)$$

Using Newton's method with the fourth-order finite-difference approximation of the second derivative, we compute numerically the values of ε, for which kink solutions of Eq.(24) exist. Numerical calculations strongly confirm the existence of the sequence $\{\varepsilon_n\}$ as well as its asymptotic properties (28). The values of $\chi/(\pi\varepsilon_n)$ for $\gamma = 5$ and $v = 0.6$ are given in Table II with satisfactory agreement.

Fig.3 shows the first two kink solutions of Eq.(24) with $\gamma = 5$ and $v = 0.6$ (thick and thin solid lines) together

$3+4n$	3	7	11	15
$\chi/(\pi\varepsilon_n)$	3.5303	7.3547	11.1520	15.0329

TABLE II: The values of ε for which Eq.(24) admits kink solutions for $\gamma = 5$ and $v = 0.6$.




FIG. 3: Profiles of the first (thick solid line) and second (thin solid line) kink solutions and the limiting solution (26) (dotted line) for Eq.(24) with $\gamma = 5$ and $v = 0.6$.

with the limiting solution of Eq.(27) (dotted line). The difference between the third kink solution and the limiting kink is not visible in the figure.

Conclusion. We have shown on three prototypical examples that Hamiltonian systems with resonances may admit an infinite sequence of travelling solitons or kinks if the leading-order asymptotic solution has a pair of symmetric singularities in the upper half-plane. This simple but universal observation reveals the way why travelling solitons or kinks have increased mobility in some nonlinear systems with resonances but not in the others.

To emphasize the universality of this prediction, we mention two additional examples earlier studied in literature, where the mechanism for existence of the infinite sequence of travelling solitons or kinks also should take place. First, three families of radiationless travelling solitons were captured numerically in the saturable discrete nonlinear Schrödinger equation [15]. Although the leading-order asymptotic solution is not available in the closed form, one can show that it has a pair of symmetric singularities in the upper half-plane, hence one can anticipate that there exist infinitely many families of travelling solitons in this example. Second, two moving kinks were reported in the modified Peyrard–Remoissenet potential of the discrete Klein–Gordon equation [3]. Since this model is a modification of the discrete double sine–Gordon equation, one can anticipate the existence of an infinite sequence of families of the moving kinks in this example. Our study of some other models of this kind (e.g. various nonlocal generalizations of double and triple sine-Gordon model) also confirms the asymptotical formula (6).

Acknowledgments. Authors are grateful to A.S.Malishevskii for useful discussions. The work of GLA and EVM was supported by Russian federal program "Scientific and educational personnel of the innovative Russia", grant 14.B37.21.1273. The work of DP is supported by the ministry of education and science of Russian Federation (Project 14.B37.21.0868).

[1] T. Kawahara, J. Phys. Soc. Japan **33**, 260 (1972); J.K. Hunter and J. Scheule, Physica D **32**, 253 (1988); A.R. Champneys and M.D. Groves, J. Fluid Mech. **342**, 199 (1997); A.R. Champneys, Nonlinearity **14**, 87 (2001); A. Tovbis and D. Pelinovsky, Nonlinearity **19**, 2277 (2006)

[2] Y. Zoloratyuk, J.C. Eilbeck, and A.V. Savin, Physica D **108**, 81 (1997); P.G. Kevrekidis, Physica D **183**, 68 (2003); A.A. Aigner, A.R. Champneys, and V.M. Rothos, Physica D **186**, 148 (2003)

[3] A.V. Savin, Y. Zolotaryuk, and J.C. Eilbeck, Physica D **138**, 267 (2000);

[4] G. Iooss and D.E. Pelinovsky, Physica D **216**, 327 (2006); O.F. Oxtoby, D.E. Pelinovsky, and I.V. Barashenkov, Nonlinearity **19**, 217 (2006)

[5] G.L. Alfimov, V.M. Eleonsky, and N.E. Kulagin, Chaos **2**, 565 (1992); G.L. Alfimov, V.M. Eleonsky, N.E. Kulagin, and N.V.Mitzkevich, Chaos **3**, 405 (1993); G.L. Alfimov, V.M. Eleonsky, and L.M. Lerman, Chaos **8**, 257 (1998)

[6] Yu. M. Aliev, K.N. Ovchinnikov, V.P. Silin, S.A. Uryupin, JETP **80**, 551 (1995); S. Savel'ev, V. Yampol'skii, A. Rakhmanov, F. Nori, Rep. Prog. Phys. **73**, 026501 (2010)

[7] A.R. Champneys, B.A. Malomed, and M.J. Friedman, Phys. Rev. Lett. **80**, 4168 (1998); A.R. Champneys, B.A. Malomed, J. Yang, and D.J. Kaup, Physica D **152-153**, 340 (2001); T. Wagenknecht and A.R. Champneys, Physica D **177**, 50 (2003)

[8] V.G. Gelfreich, V.F. Lazutkin, and M.B. Tabanov, Chaos **1**, 137 (1991); V. Gelfreich and C. Simo, Discr. Cont. Dynam. Syst. **B 10**, 511 (2008).

[9] V. Vougalter and V. Volpert, Proc. Edinb. Math. Soc. **54**, 249 (2011); Anal. Math. Phys. **2012**, 473 (2012).

[10] J. Boyd, *Chebyshev and Fourier Spectral Methods* (Dover Publishers, 2001)

[11] A. A. Golubov, M. Yu. Kupriyanov, and E. Il'ichev, Review of Modern Physics **76**, 411 (2004);

[12] A.A. Abdumalikov, G.L. Alfimov, A.S. Malishevskii, Supercond. Sci. Technol. **22**, 023001 (2009)

[13] J.D. Murray, *Asymptotic Analysis* (Springer, 1984)

[14] G.L. Alfimov and E.V. Medvedeva, Phys. Rev. E. **84**, 056606 (2011)

[15] T.R.O. Melvin, A.R. Champneys, P.G. Kevrekidis, and J. Cuevas, Phys. Rev. Lett. **97**, 124101 (2006); Physica D **237**, 551 (2008).