

CP violation in bipair neutrino mixing

Teruyuki Kitabayashi*, Masaki Yasuè

Department of Physics, Tokai University, 4-1-1 Kitakaname, Hiratsuka, Kanagawa, 259-1292, Japan

Abstract

There are experimentally determined two best-fit points for the atmospheric neutrino mixing angle θ_{23} : $\sin^2 \theta_{23} = 0.413$ (case A) and $\sin^2 \theta_{23} = 0.594$ (case B). In the bipair neutrino mixing scheme, we predict $\sin^2 \theta_{23} = \sqrt{2} - 1$ (case 1) to be consistent with the case A and $\sin^2 \theta_{23} = 2 - \sqrt{2}$ (case 2) to be consistent with the case B. If the case B is realized in nature, the bipair neutrino mixing provides a unique neutrino model consistent with the observation $\sin^2 \theta_{23} = 0.594$. However, the reactor neutrino mixing angle θ_{13} is predicted to be $\sin^2 \theta_{13} = 0$, which is inconsistent with the observation. We propose a new modification scheme to yield $\sin^2 \theta_{13} \neq 0$ utilizing the charged lepton contribution and study its effect on both of CP-violating Dirac and Majorana phases, which is numerically estimated. It is found that there appear striking differences between the case 1 and the case 2 in their phase structure.

Keywords: Bipair neutrino mixing, Charged lepton contribution, CP-violating phases

1. Introduction

The results from the neutrino oscillation experiments have provided us with robust evidence that neutrinos have tiny masses and their flavor states are mixed with each other [1, 2, 3, 4]. A global analysis shows that the best-fit values of the mixing angles with 1σ allowed range are obtained as [5]

$$\begin{aligned}\sin^2 \theta_{12} &= 0.302^{+0.013}_{-0.012}, \\ \sin^2 \theta_{23} &= 0.413^{+0.037}_{-0.025} \oplus 0.594^{+0.021}_{-0.022}, \\ \sin^2 \theta_{13} &= 0.0227^{+0.0023}_{-0.0024},\end{aligned}\tag{1}$$

where θ_{12}, θ_{23} and θ_{13} stand for the solar, atmospheric and reactor neutrino mixing angle, respectively.¹ There are two best-fit points for the atmospheric neutrino mixing angle θ_{23} . We call these two best-fit points case A and case B as follows:

$$\begin{aligned}\sin^2 \theta_{23} &= 0.413 \quad (\text{case A}), \\ \sin^2 \theta_{23} &= 0.594 \quad (\text{case B}).\end{aligned}\tag{2}$$

Neutrino mixings are described by the Pontecorvo-Maki-Nakagawa-Sakata mixing matrix U_{PMNS} [8, 9], which is parameterized by three mixing angles $\theta_{12,23,13}$ as well as one CP-violating Dirac phase δ_{CP} and two CP-violating Majorana phases $\alpha_{2,3}$ [10] to be $U_{PMNS} = UK$ with

$$\begin{aligned}U &= \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{CP}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{CP}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{CP}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{CP}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{CP}} & c_{23}c_{13} \end{pmatrix}, \\ K &= \text{diag.}(1, e^{i\alpha_2}, e^{i\alpha_3}),\end{aligned}\tag{3}$$

*Corresponding author

Email addresses: teruyuki@keyaki.cc.u-tokai.ac.jp (Teruyuki Kitabayashi), yasue@keyaki.cc.u-tokai.ac.jp (Masaki Yasuè)

¹Another global analysis is also reported [6, 7]

where $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$ ($i, j = 1, 2, 3$) [9]. We will also use the notation of $t_{ij} = \tan \theta_{ij}$.

There is a theoretical prediction of these mixing angles based on the bipair neutrino mixing scheme [11]. In the bipair neutrino mixing scheme, there are two sets of solutions referred to as case 1 and case 2. We obtain $\sin^2 \theta_{23} = \sqrt{2} - 1 (= 0.414)$ in the case 1, which is close to $\sin^2 \theta_{23}$ in the case A, and $\sin^2 \theta_{23} = 2 - \sqrt{2} (= 0.586)$ in the case 2, which is close to $\sin^2 \theta_{23}$ in the case B. At the same time, $\sin^2 \theta_{12} = 1 - 1/\sqrt{2} (= 0.293)$ and $\sin^2 \theta_{13} = 0$ are predicted. Although various other possibilities realizing the observed mixing angles $\theta_{12,23}$ have been proposed [12, 13, 14, 15, 16, 17, 18], the bipair neutrino mixing can be a good candidate of neutrino mixing schemes for the both of the case A and the case B. Especially, if the case B is chosen as a right atmospheric neutrino mixing angle, the bipair neutrino mixing scheme is the only theoretical model that has a consistent prediction with the case B.

The bipair neutrino mixing predicts $\sin^2 \theta_{13} = 0$ such as in Refs.[13, 14, 15, 16, 17, 18], which is inconsistent with the observation. It is expected that additional contributions to the mixing angles are produced by the charged lepton contributions [19] if some of the nondiagonal matrix elements of charged lepton mass matrix are nonzero so that the reactor mixing angle can be shifted to lie in the allowed region. Two modification schemes of the bipair neutrino mixing to induce $\sin^2 \theta_{13} \neq 0$ to be consistent with the observation are discussed in Ref.[20] and Ref.[21]. The first modification scheme is not suitable to study effects of CP-violating Majorana phases while the second modification scheme is not suitable to study those of CP-violating Dirac phase.

In this letter, we discuss third modification scheme based on the parameterization by Pascoli, Petcov and Rodejohann [22] and evaluate charged lepton contribution to both of CP-violating Dirac and Majorana phases in the bipair neutrino mixing scheme. We estimate sizes of these CP-violating phases, whose correlations are numerically obtained and discuss expected effects of CP violation in the case 1 and the case 2.

2. Modified bipair neutrino mixing

The bipair neutrino mixing is described by a mixing matrix U with $\theta_{13} = 0$, which is equipped with two pairs of identical magnitudes of matrix elements to be denoted by U_{ij}^0 ($i, j=1,2,3$). There are two cases of the bipair texture:

- The case 1 of the bipair neutrino mixing ($|U_{12}^0| = |U_{32}^0|$ and $|U_{22}^0| = |U_{23}^0|$) is parameterized by U_{BP1}^0 containing only one mixing angle θ_{12} :

$$U_{BP1}^0 = \begin{pmatrix} c & s & 0 \\ -t^2 & t & t \\ st & -s & t/c \end{pmatrix}, \quad (4)$$

where $c = c_{12}$, $s = s_{12}$ and $t = t_{12}$. The mixing angles are predicted to be

$$\begin{aligned} \sin^2 \theta_{12}^{BP1} &= 1 - 1/\sqrt{2} = 0.293, \\ \sin^2 \theta_{23}^{BP1} &= \sqrt{2} - 1 = 0.414, \end{aligned} \quad (5)$$

and $\sin^2 \theta_{13}^{BP1} = 0$. The case 1 of the bipair neutrino mixing well describes the 1σ data of the solar neutrino mixing in Eq.(1) and the case A of the atmospheric neutrino mixing in Eq.(2).

- The case 2 of the bipair neutrino mixing ($|U_{12}^0| = |U_{22}^0|$ and $|U_{32}^0| = |U_{33}^0|$) is parameterized by U_{BP2}^0 :

$$U_{BP2}^0 = \begin{pmatrix} c & s & 0 \\ -st & s & t/c \\ t^2 & -t & t \end{pmatrix}. \quad (6)$$

The mixing angles are predicted to be

$$\begin{aligned} \sin^2 \theta_{12}^{BP2} &= 1 - 1/\sqrt{2} = 0.293, \\ \sin^2 \theta_{23}^{BP2} &= 2 - \sqrt{2} = 0.586, \end{aligned} \quad (7)$$

and $\sin^2 \theta_{13}^{BP2} = 0$. The case 2 of the bipair neutrino mixing well describes the 1σ data of the solar neutrino mixing in Eq.(1) and the case B of the atmospheric neutrino mixing in Eq.(2).

Our modified bipair neutrino mixing is described by a specific type of U_{PMNS} , which will be constructed by the parameterization proposed by Pascoli, Petcov and Rodejohann [22]. Namely, U_{PMNS} is given by

$$U_{PMNS} = \tilde{U}^0 K, \quad \tilde{U}^0 \equiv U_\ell^\dagger P U_\nu, \quad (8)$$

where U_ℓ (as well as U_R to be used later) and U_ν , respectively, arise from the diagonalization of the charged lepton mass matrix M_ℓ and of the neutrino mass matrix M_ν and P is defined by

$$P = \text{diag.}(1, e^{i\phi_2}, e^{i\phi_3}). \quad (9)$$

The lepton mass matrices satisfy the relations of $M_\ell = U_\ell M_\ell^{diag} U_R^\dagger$ and $M_\nu = U_\nu M_\nu^{diag} U_\nu^T$ where M_ℓ^{diag} and M_ν^{diag} are, respectively, the diagonal mass matrix of the charged lepton and of the neutrino.

The charged lepton mixing matrix U_ℓ can be parameterized by three mixing angles $\theta_{12,23,13}^\ell$ and one CP-violating Dirac phase δ as follows:

$$U_\ell = \begin{pmatrix} c_{12}^\ell c_{13}^\ell & s_{12}^\ell c_{13}^\ell & s_{13}^\ell e^{-i\delta} \\ -s_{12}^\ell c_{23}^\ell - c_{12}^\ell s_{23}^\ell s_{13}^\ell e^{i\delta} & c_{12}^\ell c_{23}^\ell - s_{12}^\ell s_{23}^\ell s_{13}^\ell e^{i\delta} & s_{23}^\ell c_{13}^\ell \\ s_{12}^\ell s_{23}^\ell - c_{12}^\ell c_{23}^\ell s_{13}^\ell e^{i\delta} & -c_{12}^\ell s_{23}^\ell - s_{12}^\ell c_{23}^\ell s_{13}^\ell e^{i\delta} & c_{23}^\ell c_{13}^\ell \end{pmatrix}, \quad (10)$$

where $c_{ij}^\ell = \cos \theta_{ij}^\ell$ and $s_{ij}^\ell = \sin \theta_{ij}^\ell$ ($i, j = 1, 2, 3$). To be more specific, we adapt the Cabibbo-Kobayashi-Maskawa like parameterization of the matrix U_ℓ [22, 23], which is represented by small parameters having magnitude of the order of Wolfenstein parameter $\lambda \sim 0.227$ or less. In the small angle approximation, the charged lepton mixing matrix can be approximated to be:

$$U_\ell = \begin{pmatrix} 1 - \frac{\epsilon_{12}^2 + \epsilon_{13}^2}{2} & \epsilon_{12} & e^{-i\delta} \epsilon_{13} \\ -\epsilon_{12} - \epsilon_{23} \epsilon_{13} e^{i\delta} & 1 - \frac{\epsilon_{12}^2 + \epsilon_{23}^2}{2} & \epsilon_{23} \\ \epsilon_{12} \epsilon_{23} - e^{i\delta} \epsilon_{13} & -\epsilon_{23} - \epsilon_{12} \epsilon_{13} e^{i\delta} & 1 - \frac{\epsilon_{23}^2 + \epsilon_{13}^2}{2} \end{pmatrix}, \quad (11)$$

where $\epsilon_{ij} \equiv s_{ij}^\ell$. Since Eq.(11) is non-unitary, the resulting neutrino mixing matrix is not unitary. Current data do not allow a big unitarity violation effect [24]. We demand that the charged lepton mixing matrix is unitary within 1%, which is roughly λ^3 . Therefore, we have retained the terms of the second order in ϵ_{ij} in Eq.(11). Although numerical calculations are performed with approximate unitarity kept, results of our estimations will be shown within the first order in ϵ_{ij} to avoid apparent complexity of derived expressions. We mention that there are other types of approximation for charged lepton mixing matrix [19].

If M_ℓ is non-diagonal, U_ℓ is associated with M_ℓ . We construct such a mass matrix that yields Eq.(11). One possible form of U_ℓ arises from the wishful thinking of the similarity between the quark and lepton sectors [23]. Moreover, with SU(5) GUT relation, we can make simple mass matrix of charged leptons [25]. In minimal SU(5) GUT, the charged lepton mass matrix M_ℓ and the down quark mass matrices M_d are related each other to satisfy the relation $M_\ell = M_d^T$; therefore, we expect $U_R = U_d$. The Cabibbo-Kobayashi-Maskawa matrix U_{CKM} for quark sector is given by $U_{CKM} = U_u^\dagger U_d$. Given that the quark mixing angles are small, either both U_u and U_d are nearly diagonal or they are nearly equal. One possible choice is that U_d is nearly diagonal and U_R is also nearly diagonal. In this case, the mass matrix M_ℓ for with $M_\ell^{diag} = \text{diag.}(m_e, m_\mu, m_\tau)$ is approximately given by

$$M_\ell = U_\ell M_\ell^{diag} U_R^\dagger \sim \begin{pmatrix} m_e & m_\mu \epsilon_{12} & m_\tau e^{-i\delta} \epsilon_{13} \\ -m_e \epsilon_{12} & m_\mu & m_\tau \epsilon_{23} \\ -m_e e^{i\delta} \epsilon_{13} & -m_\mu \epsilon_{23} & m_\tau \end{pmatrix}, \quad (12)$$

where $U_R \sim 1$, where terms of the first order in ϵ_{ij} are retained for the sake of simplicity. This is one example of the charged lepton mass matrix, which is consistent with Eq.(11).

To estimate the charged lepton contributions to the mixing angles in the bipair neutrino mixing, we take the neutrino mixing matrix to be either $U_\nu = U_{BP1}^0$ or $U_\nu = U_{BP2}^0$. As a definition of the bipair neutrino mixing, there is no phase parameter in U_{BP1}^0 and U_{BP2}^0 . The mixing matrix U has three real parameters $\epsilon_{12}, \epsilon_{13}, \epsilon_{23}$ and five phases $\delta, \phi_2, \phi_3, \alpha_2, \alpha_3$.

In the case of $U_\nu = U_{BP1}^0$, we obtain the mixing matrix \tilde{U}_{BP1}^0 for \tilde{U}^0 as

$$\begin{aligned} \tilde{U}_{BP1}^0 &= \begin{pmatrix} c & s & 0 \\ -e^{i\phi_2}t^2 & e^{i\phi_2}t & e^{i\phi_2}t \\ e^{i\phi_3}st & -e^{i\phi_3}s & e^{i\phi_3}t/c \end{pmatrix} \\ &+ \begin{pmatrix} e^{i\phi_2}t^2\epsilon_{12} - \tilde{s}t\epsilon_{13} & -e^{i\phi_2}t\epsilon_{12} + \tilde{s}\epsilon_{13} & -e^{i\phi_2}t\epsilon_{12} - \frac{\tilde{s}}{c^2}\epsilon_{13} \\ c\epsilon_{12} - e^{i\phi_3}st\epsilon_{23} & s\epsilon_{12} + e^{i\phi_3}s\epsilon_{23} & -e^{i\phi_3}\frac{t}{c}\epsilon_{23} \\ e^{i\delta}c\epsilon_{13} - e^{i\phi_2}t^2\epsilon_{23} & e^{i\delta}s\epsilon_{13} + e^{i\phi_2}t\epsilon_{23} & e^{i\phi_2}t\epsilon_{23} \end{pmatrix}, \end{aligned} \quad (13)$$

where we define $\tilde{s} = e^{-i(\delta-\phi_3)}s$. From $U_{PMNS} = \tilde{U}_{BP1}^0 K$ and the following general relations

$$\sin^2 \theta_{12} = \frac{|U_{12}|^2}{|U_{11}|^2 + |U_{12}|^2}, \quad \sin^2 \theta_{23} = \frac{|U_{23}|^2}{|U_{23}|^2 + |U_{33}|^2}, \quad \sin^2 \theta_{13} = |U_{13}|^2, \quad (14)$$

where $U_{ij} = (U_{PMNS})_{ij}$ ($i, j=1, 2, 3$), the mixing angles are obtained as follows:

$$\begin{aligned} \sin^2 \theta_{12} &= \left(1 - 2^{5/4}\epsilon_{12}\cos\phi_2 + 2\epsilon_{13}\cos(\delta - \phi_3)\right) \sin^2 \theta_{12}^{BP1}, \\ \sin^2 \theta_{23} &= \left(1 - 2^{5/4}\epsilon_{23}\cos(\phi_2 - \phi_3)\right) \sin^2 \theta_{23}^{BP1}, \\ \sin^2 \theta_{13} &= \left(\epsilon_{12}^2 + 2^{5/4}\epsilon_{12}\epsilon_{13}\cos(\delta + \phi_2 - \phi_3)\right) \sin^2 \theta_{23}^{BP1} + 2\epsilon_{13}^2 \sin^2 \theta_{12}^{BP1}. \end{aligned} \quad (15)$$

In the case of $U_\nu = U_{BP2}^0$, we obtain the mixing matrix \tilde{U}_{BP2}^0 as

$$\begin{aligned} \tilde{U}_{BP2}^0 &= \begin{pmatrix} c & s & 0 \\ -e^{i\phi_2}st & e^{i\phi_2}s & e^{i\phi_2}t/c \\ e^{i\phi_3}t^2 & -e^{i\phi_3}t & e^{i\phi_3}t \end{pmatrix} \\ &+ \begin{pmatrix} e^{i\phi_2}st\epsilon_{12} - \tilde{t}t\epsilon_{13} & -e^{i\phi_2}s\epsilon_{12} + \tilde{t}\epsilon_{13} & -e^{i\phi_2}\frac{t}{c}\epsilon_{12} - \tilde{t}\epsilon_{13} \\ c\epsilon_{12} - e^{i\phi_3}t^2\epsilon_{23} & s\epsilon_{12} + e^{i\phi_3}t\epsilon_{23} & -e^{i\phi_3}\frac{t}{c}\epsilon_{23} \\ e^{i\delta}c\epsilon_{13} - e^{i\phi_2}st\epsilon_{23} & e^{i\delta}s\epsilon_{13} + e^{i\phi_2}t\epsilon_{23} & e^{i\phi_2}\frac{t}{c}\epsilon_{23} \end{pmatrix}, \end{aligned} \quad (16)$$

where we define $\tilde{t} = e^{-i(\delta-\phi_3)}t$. The mixing angles from $U_{PMNS} = \tilde{U}_{BP2}^0 K$ and Eq.(14) are obtained as follows:

$$\begin{aligned} \sin^2 \theta_{12} &= \left(1 - 2\epsilon_{12}\cos\phi_2 + 2^{5/4}\epsilon_{13}\cos(\delta - \phi_3)\right) \sin^2 \theta_{12}^{BP2}, \\ \sin^2 \theta_{23} &= \left(1 - 2^{3/4}\epsilon_{23}\cos(\phi_2 - \phi_3)\right) \sin^2 \theta_{23}^{BP2}, \\ \sin^2 \theta_{13} &= \sqrt{2} \left(\epsilon_{13}^2 + 2^{5/4}\epsilon_{12}\epsilon_{13}\cos(\delta + \phi_2 - \phi_3)\right) \sin^2 \theta_{12}^{BP2} + \epsilon_{12}^2 \sin^2 \theta_{23}^{BP2}. \end{aligned} \quad (17)$$

3. CP-violating phases

There are rephasing invariant quantities for CP-violating Dirac phase as well as CP-violating Majorana phases. It is known that CP-violating Dirac phase δ_{CP} is determined by the Jarlskog invariant J [26] to be:

$$\sin \delta_{CP} = \frac{\text{Im}(U_{11}U_{22}U_{12}^*U_{21}^*)}{c_{12}c_{23}c_{13}^2s_{12}s_{23}s_{13}}. \quad (18)$$

One can also find that CP-violating Majorana phases $\alpha_{2,3}$ are determined to be:

$$\alpha_2 = \arg(U_{12}U_{11}^*), \quad \alpha_3 = \arg(U_{13}U_{11}^*) + \delta_{CP}, \quad (19)$$

which can be used to predict CP-violating Majorana phases from U_{PMNS} .

To find Eq.(19), e.g., rephasing invariant expressions for Majorana phases $\alpha_{2,3}$, it is useful to employ general parameterization of U instead of Eq.(3), which is containing three Dirac phases δ , ρ and τ as well as one additional phase γ [22, 27]:

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\gamma} & 0 \\ 0 & 0 & e^{-i\gamma} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23}e^{i\tau} \\ 0 & -s_{23}e^{-i\tau} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{pmatrix} \cdot \begin{pmatrix} c_{12} & s_{12}e^{i\rho} & 0 \\ -s_{12}e^{-i\rho} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (20)$$

and three Majorana phases $\varphi_{1,2,3}$,

$$Q = \begin{pmatrix} e^{i\varphi_1} & 0 & 0 \\ 0 & e^{i\varphi_2} & 0 \\ 0 & 0 & e^{i\varphi_3} \end{pmatrix}, \quad (21)$$

giving $U_{PMNS} = UQ$. After δ_{CP} is taken to be $\delta_{CP} = \delta + \rho + \tau$, we readily reach

$$U_{PMNS} = P' \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{CP}} \\ -(c_{23}s_{12} + s_{23}c_{12}s_{13}e^{i\delta_{CP}}) & c_{23}c_{12} - s_{23}s_{12}s_{13}e^{i\delta_{CP}} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}c_{12}s_{13}e^{i\delta_{CP}} & -(s_{23}c_{12} + c_{23}s_{12}s_{13}e^{i\delta_{CP}}) & c_{23}c_{13} \end{pmatrix} Q', \quad (22)$$

where

$$P' = \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\gamma} & 0 \\ 0 & 0 & e^{-i(\gamma+\tau)} \end{pmatrix}, \quad Q' = \begin{pmatrix} e^{i\beta_1} & 0 & 0 \\ 0 & e^{i\beta_2} & 0 \\ 0 & 0 & e^{i\beta_3} \end{pmatrix}, \quad (23)$$

with $\beta_1 = \varphi_1 - \rho$, $\beta_2 = \varphi_2$ and $\beta_3 = \varphi_3 + \tau$.

CP-violating Dirac phase in the form of the rephasing invariant quantity, we can make the following product as one possible choice:

$$U_{11}U_{22}U_{12}^*U_{21}^*, \quad (24)$$

which is independent of the redundant phases as well as $\alpha_{2,3}$. The Jarlskog invariant J is expressed to be:

$$J = \text{Im}(U_{11}U_{22}U_{12}^*U_{21}^*), \quad (25)$$

giving

$$\text{Im}(U_{11}U_{22}U_{12}^*U_{21}^*) = c_{12}c_{23}c_{13}^2 s_{12}s_{23}s_{13} \sin \delta_{CP}. \quad (26)$$

In the similar way, it is possible to describe CP-violating Majorana phases in terms of rephasing invariant quantities. In fact, using Eq.(22), one can find that $U_{12}U_{11}^*$, $U_{13}U_{11}^*$ and $U_{13}U_{12}^*$ [28], which are independent of redundant phases, lead to

$$\begin{aligned} U_{12}U_{11}^* &= s_{12}c_{13}c_{12}c_{13}e^{i(\beta_2-\beta_1)}, \\ U_{13}U_{11}^* &= s_{13}c_{12}c_{13}e^{i(\beta_3-\beta_1-\delta_{CP})}, \\ U_{13}U_{12}^* &= s_{13}s_{12}c_{13}e^{i(\beta_3-\beta_2-\delta_{CP})}, \end{aligned} \quad (27)$$

from which

$$\begin{aligned}\beta_2 - \beta_1 &= \alpha_2 = \arg(U_{12}U_{11}^*), \\ \beta_3 - \beta_1 &= \alpha_3 = \arg(U_{13}U_{11}^*) + \delta_{CP}, \\ \beta_3 - \beta_2 &= \arg(U_{13}U_{12}^*) + \delta_{CP},\end{aligned}\tag{28}$$

can be derived.

4. Numerical analysis

We have performed the random search to find the parameter set of $\{\epsilon_{12,13,23}, \delta, \phi_2, \phi_3, \delta_{CP}, \alpha_2, \alpha_3\}$. To enhance the predictability, we assume that $\epsilon_{12} = \epsilon_{13} = \epsilon_{23} = \epsilon$. The unitarity of U_ℓ dictates that

$$U_\ell^\dagger U_\ell = I + \begin{pmatrix} 0 & 0 & -e^{-i\delta} \\ 0 & 2\cos\delta & 0 \\ -e^{i\delta} & 0 & 0 \end{pmatrix} \epsilon^3,\tag{29}$$

for $\epsilon^3 \leq \mathcal{O}(0.01)$. The mixing matrices $\tilde{U}_{BP1, BP2}^0$ are accordingly estimated to be:

$$\begin{aligned}\tilde{U}^0 &= \begin{pmatrix} c & s & 0 \\ -e^{i\phi_2} s c_{23} & e^{i\phi_2} c c_{23} & e^{i\phi_2} s_{23} \\ e^{i\phi_3} s s_{23} & -e^{i\phi_3} c s_{23} & e^{i\phi_3} c_{23} \end{pmatrix} \\ &+ \begin{pmatrix} s(e^{i\phi_2} c_{23} - e^{-i(\delta-\phi_3)} s_{23}) & -c(e^{i\phi_2} c_{23} - e^{-i(\delta-\phi_3)} s_{23}) & -(e^{i\phi_2} s_{23} + e^{-i(\delta-\phi_3)} c_{23}) \\ c - e^{i\phi_3} s s_{23} & s + e^{i\phi_3} c s_{23} & -e^{i\phi_3} c_{23} \\ e^{i\delta} c - e^{i\phi_2} s c_{23} & e^{i\delta} s + e^{i\phi_2} c c_{23} & e^{i\phi_2} s_{23} \end{pmatrix} \epsilon \\ &+ \begin{pmatrix} -\begin{pmatrix} c - e^{-i(\delta-\phi_2)} s c_{23} \\ -e^{i\phi_3} s s_{23} \end{pmatrix} & -\begin{pmatrix} s + e^{-i(\delta-\phi_2)} c c_{23} \\ +e^{i\phi_3} c s_{23} \end{pmatrix} & e^{-i(\delta-\phi_2)} s_{23} - e^{i\phi_3} c_{23} \\ s(e^{i\phi_2} c_{23} - e^{-i(\delta-\phi_3)} s_{23}) & -c(e^{i\phi_2} c_{23} - e^{-i(\delta-\phi_3)} s_{23}) & -(e^{i\phi_2} s_{23} + e^{-i(\delta-\phi_3)} c_{23}) \\ -e^{i\phi_3} s s_{23} & e^{i\phi_3} c s_{23} & -e^{i\phi_3} c_{23} \end{pmatrix} \epsilon^2,\end{aligned}\tag{30}$$

where $c_{23} = t/s, s_{23} = t$ for \tilde{U}_{BP1}^0 and $c_{23} = t, s_{23} = t/c$ for \tilde{U}_{BP2}^0 . The input parameters are $\epsilon, \delta, \phi_2, \phi_3$ while the output parameters are CP-violating phases $\delta_{CP}, \alpha_2, \alpha_3$. We vary the input parameters in the range of $0 \leq \epsilon \leq 0.227$ and $-\pi \leq \delta, \phi_2, \phi_3 \leq \pi$.

For the case 1, we have searched the consistent parameter set with the 1σ data of the solar and reactor neutrino mixing angles in Eq.(1) and of the case A of the atmospheric mixing angle in Eq.(2), e.g., $\sin^2 \theta_{12} = 0.302^{+0.013}_{-0.012}$, $\sin^2 \theta_{23} = 0.413^{+0.037}_{-0.025}$, and $\sin^2 \theta_{13} = 0.0227^{+0.0023}_{-0.0024}$. Fig.1 shows the result of our numerical analysis in the case 1 of the modified bipair neutrino mixing:

1. The $\delta_{CP} - \epsilon$ plot shows that²
 - δ_{CP} is constrained to be $|\delta_{CP}| \gtrsim 0.5$;
 - δ_{CP} is constrained to be $|\delta_{CP}| \gtrsim 1.0$ for the minimal value of ϵ around 0.1;
 - the smaller magnitude of δ_{CP} is allowed for the larger value of ϵ .
2. The $\alpha_2 - \epsilon$ plot shows that
 - α_2 is constrained to be $|\alpha_2| \lesssim 0.5$;
 - α_2 is constrained to be $|\alpha_2| \lesssim 0.1$ for the smaller value of ϵ satisfying $0.09 \lesssim \epsilon \lesssim 0.14$;
 - the larger magnitude of α_2 is allowed for the larger value of ϵ .

²The range of δ_{CP} is taken to be $-\pi/2 \sim \pi/2$.




Figure 1: The dependence of δ_{CP} and $\alpha_{2,3}$ on ϵ in the case 1.




Figure 2: Same as Fig.1 but in the case 2.

3. The $\alpha_3 - \epsilon$ plot shows that

- α_3 is constrained to be $|\alpha_3| \lesssim \pi/2$;
- α_3 is constrained to be $|\alpha_3| \lesssim 0.5$ for the minimal value of ϵ around 0.1;
- the larger magnitude of α_3 is allowed for the larger value of ϵ .

For the case 2, we have searched the consistent parameter set with the 1σ data of the solar and reactor neutrino mixing angles in Eq.(1) and of the case B of the atmospheric mixing angle in Eq.(2), e.g., $\sin^2 \theta_{12} = 0.302^{+0.013}_{-0.012}$, $\sin^2 \theta_{23} = 0.594^{+0.021}_{-0.022}$, and $\sin^2 \theta_{13} = 0.0227^{+0.0023}_{-0.0024}$. Fig.2 shows the result of our numerical analysis in the case 2 of the modified bipair neutrino mixing:

1. The $\delta_{CP} - \epsilon$ plot shows that

- δ_{CP} is constrained to be $|\delta_{CP}| \lesssim 0.5$ for $\epsilon \gtrsim 0.12$;
- no constraint on δ_{CP} arises for $\epsilon \lesssim 0.12$;
- the smaller magnitude of δ_{CP} is allowed in the broad range of $0.09 \lesssim \epsilon \lesssim 0.22$.

2. The $\alpha_2 - \epsilon$ plot shows that

- α_2 is constrained to be $|\alpha_2| \lesssim 0.6$;
- the smaller magnitude of α_2 is allowed for $0.09 \lesssim \epsilon \lesssim 0.12$;
- the larger magnitude of α_2 is allowed for the larger value of ϵ .

3. The $\alpha_3 - \epsilon$ plot shows that

- α_3 is constrained to be $|\alpha_3| \lesssim \pi$;
- α_3 is constrained to be either $\alpha_3 \sim 0$ around its minimal value or $|\alpha_3| \sim \pi$ around its maximal value in the broad range of $0.12 \lesssim \epsilon \lesssim 0.22$;
- α_3 takes any allowed values for $0.09 \lesssim \epsilon \lesssim 0.12$.




Figure 3: The mutual dependence of δ_{CP} and $\alpha_{2,3}$ in the case 1.




Figure 4: Same as Fig.3 but in the case 2.

It is also useful to present correlations among measurable quantities for the direct comparison with future experimental data of CP-violating phases. Shown in Fig.3 and Fig.4 are mutual dependence of CP-violating phases δ_{CP} and $\alpha_{2,3}$.

For the case 1, we see the following mutual dependence in Fig.3:

1. The $\alpha_2 - \delta_{CP}$ plot shows that $|\alpha_2| \lesssim 0.5$ and that
 - $|\alpha_2|$ approaches to its maximal value around 0.5 as $|\delta_{CP}|$ approaches its minimal value around 0.5;
 - $|\alpha_2|$ approaches to its minimal value around 0 as $|\delta_{CP}|$ approaches its maximal value around $\pi/2$;
2. The $\alpha_3 - \delta_{CP}$ plot shows that $|\alpha_3| \lesssim \pi/2$ and that
 - $|\alpha_3|$ approaches to its maximal value of $\pi/2$ as $|\delta_{CP}|$ approaches to its minimal value around 0.5;
 - $|\alpha_3|$ approaches to its minimal value of 0 as $|\delta_{CP}|$ approaches to its maximal value of $\pi/2$;
 - roughly speaking, α_3 is scattered around the straight line of $\alpha_3 = 3(2\delta_{CP} \pm \pi)/4$.
3. The $\alpha_3 - \alpha_2$ plot shows that
 - $|\alpha_2|$ approaches to its maximal value around 0.5 for $|\alpha_3| \sim 1$;
 - α_3 is constrained to be $|\alpha_3| \lesssim 1.0$ for $|\alpha_2| \lesssim 0.1$.

On the other hand, for the case 2, we see the following mutual dependence in Fig.4:

1. The $\alpha_2 - \delta_{CP}$ plot shows that $|\alpha_2| \lesssim 0.6$ and
 - δ_{CP} takes any allowed values for $\alpha_2 \sim 0$;
 - α_2 takes any allowed values for $|\delta_{CP}| \lesssim 0.5$.
2. The $\alpha_3 - \delta_{CP}$ plot shows that
 - $|\alpha_3|$ approaches to its maximal value of π as $|\delta_{CP}|$ approaches to its minimal value of 0;
 - $|\alpha_3|$ approaches to its minimal value of 0 as $|\delta_{CP}|$ approaches to its maximal value of $\pi/2$;

- roughly speaking, α_3 is scattered around the straight line of $\alpha_3 = 2\delta_{CP} \pm \pi$ as well as around $\alpha_3 = 0$;
 - δ_{CP} takes any allowed values for $\alpha_3 \sim 0$,
3. The $\alpha_3 - \alpha_2$ plot shows that
- α_3 is constrained to be either $\alpha_3 \sim 0$ around its minimal value or $|\alpha_3| \sim \pi$ around its maximal value for $0.2 \lesssim |\alpha_2| \lesssim 0.6$;
 - α_2 (α_3) takes any allowed values for $\alpha_3 \sim 0$ ($\alpha_2 \sim 0$).

It can be stated that the case 1 is excluded for $|\delta_{CP}| \lesssim 0.3$ and that the case 2 is excluded for $|\delta_{CP}| \gtrsim 0.5$ if $|\alpha_2| \gtrsim 0.2$.

5. Summary and discussions

The bipair neutrino mixing has predicted the solar neutrino mixing angle θ_{12} and the atmospheric neutrino mixing angle θ_{23} to be consistent with the currently observed experimental ones:

- For θ_{12} , our prediction is $\sin^2 \theta_{12} = 0.293$ to be consistent with the best-fit value of $\sin^2 \theta_{12} = 0.302$.
- For θ_{23} , our predictions are $\sin^2 \theta_{23} = 0.414$ in the case 1 to be consistent with the best-fit value of $\sin^2 \theta_{23} = 0.413$ (case A) and $\sin^2 \theta_{23} = 0.586$ in the case 2 to be consistent with the best-fit value of $\sin^2 \theta_{23} = 0.594$ (case B).

For the reactor neutrino mixing angle θ_{13} , our prediction of $\sin^2 \theta_{13} = 0$ is inconsistent with the best-fit value of $\sin^2 \theta_{13} = 0.0227$. We have discussed effects on neutrino mixings from the charged lepton contribution, which induce appropriate sizes of θ_{13} as well as CP-violating Dirac phase δ_{CP} as a measurable quantity. CP-violating Majorana phases α_2 and α_3 are determined by U_{PMNS} as rephasing invariant quantities to be $\alpha_2 = \arg(U_{12}U_{11}^*)$ and $\alpha_3 = \arg(U_{13}U_{11}^*) + \delta_{CP}$.

From numerical calculations based on the simplification of $\epsilon_{12} = \epsilon_{13} = \epsilon_{23} = \epsilon$, where $\epsilon_{12,13,23}$ characterizes three types of charged lepton contributions, we have found that there are striking differences between the case 1 and the case 2 in the relation of $\delta_{CP} - \epsilon$, $\alpha_2 - \epsilon$ and $\alpha_3 - \epsilon$ in Fig.1 and Fig.2 as well as in the relation of $\alpha_2 - \delta_{CP}$, $\alpha_3 - \delta_{CP}$ and $\alpha_3 - \alpha_2$ in Fig.3 and Fig.4. We can use these differences to predict CP-violating phases for comparison with the results of the future global analysis. For example, if the case A of the global analysis for the atmospheric neutrino mixing angle $\sin^2 \theta_{23} = 0.413$ to be a correct mixing angle, we can expect that the case 1 is realized and that the following predictions are obtained:

1. The larger CP-violating Dirac phase around $|\delta_{CP}| \sim \pi/2$ favors for
 - the suppressed charged lepton contribution of $\epsilon \lesssim 0.1$ as shown in Fig.1;
 - the suppressed CP-violating Majorana phase α_3 around $\alpha_3 \sim 0$ as shown in Fig.3.
2. The smaller CP-violating Dirac phase around $|\delta_{CP}| \sim 0.5$ (as the maximally allowed value) favors for
 - the larger charged lepton contribution of $\epsilon \sim 0.23$ as shown in Fig.1;
 - the larger CP-violating Majorana phase α_3 around $|\alpha_3| \sim \pi/2$ (as the maximally allowed value) as shown in Fig.3.
3. α_3 is scattered around the straight line of $\alpha_3 = 3(2\delta_{CP} \pm \pi)/4$.

On the other hand, if the case B of the global analysis for the atmospheric neutrino mixing angle $\sin^2 \theta_{23} = 0.594$ to be a correct mixing angle, we can expect that the case 2 is realized and that the following predictions are obtained:

1. The larger CP-violating Dirac phase around $|\delta_{CP}| \sim \pi/2$ favors is possible for
 - the suppressed charged lepton contribution of $\epsilon \lesssim 0.12$ as shown in Fig.2;
 - the suppressed CP-violating Majorana phase α_3 around $\alpha_3 \sim 0$ as shown in Fig.4.

2. The smaller CP-violating Dirac phase $|\delta_{CP}| \lesssim 0.5$ favors for
 - the mild charged lepton contribution of $\epsilon \gtrsim 0.12$ as shown in Fig.2;
 - the suppressed CP-violating Majorana phase α_3 around $\alpha_3 \sim 0, \pm\pi$ as shown in Fig.4.
3. α_3 is scattered around the straight line of $\alpha_3 = 2\delta_{CP} \pm \pi$ as well as staying around $\alpha_3 \sim 0$.

We can also observe that the case 1 is excluded for $|\delta_{CP}| \lesssim 0.5$ while the case 2 is excluded for $|\delta_{CP}| \gtrsim 0.5$ if $|\alpha_2| \gtrsim 0.2$.

References

- [1] Y. Fukuda, et al. (Super-Kamiokande Collaboration), Phys. Rev. Lett. **81** (1998) 1562; Phys. Rev. Lett. **82** (1999) 2430; T. Kajita, Nucl. Phys. B. Proc. Suppl. **77** (1999) 123. See also, T. Kajita and Y. Totsuka, Rev. Mod. Phys. **73** (2001) 85.
- [2] J.N. Bahcall, W.A. Fowler, I. Iben and R.L. Sears, Astrophys. J. **137** (1963) 344; J.N. Bahcall, Phys. Rev. Lett. **12** (1964) 300; R. Davis Jr., Phys. Rev. Lett. **12** (1964) 303; R. Davis Jr., D.S. Harmer and K.C. Hoffman, Phys. Rev. Lett. **20** (1968) 1205; J.N. Bahcall, N.A. Bahcall and G. Shaviv, Phys. Rev. Lett. **20** (1968) 1209; J.N. Bahcall and R. Davis Jr., Science **191** (1976) 264; Y. Fukuda, et al. (Super-Kamiokande Collaboration), Phys. Rev. Lett. **81** (1998) 1158; Phys. Rev. Lett. **81** (1998) 4279, Erratum; B.T. Cleveland, et al. (Super-Kamiokande Collaboration), Astrophys. J. **496** (1998) 505; W. Hampel, et al. (GNO Collaboration), Phys. Lett. B **447** (1999) 127; Q.A. Ahmed, et al. (SNO Collaboration), Phys. Rev. Lett. **87** (2001) 071301; Phys. Rev. Lett. **89** (2002) 011301.
- [3] See for example, K. Eguchi, et al. (KamLAND Collaboration), Phys. Rev. Lett. **90** (2003) 021802; S. Abe, et al. (KamLAND Collaboration), Phys. Rev. Lett. **100** (2008) 221803; F.P. An, et al., Phys. Rev. Lett. **108** (2012) 171803.
- [4] See for example, K. Abe, et al. (T2K Collaboration), Phys. Rev. Lett. **107** (2011) 041801; P. Adamson, et al. (MINOS Collaboration), Phys. Rev. Lett. **107** (2011) 021801.
- [5] M.C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, JHEP **12** (2012) 123.
- [6] G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo and A.M. Rotunno, Phys. Rev. D **86** (2012) 013012.
- [7] D.V. Forero, M. Tórtola and J.W.F. Valle, Phys. Rev. D **86** (2012) 073012.
- [8] B. Pontecorvo, JETP (USSR) **34** (1958) 247; Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870.
- [9] J. Beringer, et.al. (Particle Data Group), Phys. Rev. D **86** (2012) 010001.
- [10] S.M. Bilenky, J. Hosek and S.T. Petcov, Phys. Lett. **94B** (1980) 495; J. Schechter and J.W.F. Valle, Phys. Rev. D **22** (1980) 2227; M. Doi, T. Kotani, H. Nishiura, K. Okuda and E. Takasugi, Phys. Lett. **102B** (1981) 323.
- [11] T. Kitabayashi and M. Yasuè, Phys. Lett. B **696** (2011) 478.
- [12] For a recent review, G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82** (2010) 2701.
- [13] P.F. Harrison, D.H. Perkins and W.G. Scott, Phys. Lett. B **530** (2002) 167.
- [14] H. Fritzsch and Z.-Z. Xing, Phys. Lett. B **372** (1996) 265.
- [15] V.D. Barger, S. Pakvasa, T.J. Weiler and K. Whisnant, Phys. Lett. B **437** (1998) 107.
- [16] Y. Kajiyama, M. Raidal and A. Strumia, Phys. Rev. D **76** (2007) 117301.
- [17] W. Rodejohann, Phys. Lett. B **671** (2009) 267.
- [18] C. Giunti, Nucl. Phys. B. Proc. Suppl. **117** (2003) 24;
- [19] See for example, C. Giunti and M. Tanimoto, Phys. Rev. D **66** (2002) 053013; Phys. Rev. D **66** (2002) 113006; Z.Z. Xing, Phys. Lett. B **533** (2002) 85; W. Rodejohann, Phys. Rev. D **69** (2004) 033005; P.H. Frampton, S.T. Petcov and W. Rodejohann, Nucl. Phys. B **687** (2004) 31; G. Altarelli, F. Feruglio and I. Masina, Nucl. Phys. B **689** (2004) 157; A. Romanino, Phys. Rev. D **70** (2004) 013003; K. A. Hochmuth, S. T. Petcov and W. Rodejohann, Phys. Lett. B **654** (2007) 177; S. Dev, S. Gupta and R. R. Gautam, Phys. Lett. B **704** (2011) 527; J. A. Acosta, A. Aranda, M. A. Buen-Abad and A. D. Rojas, Phys. Lett. B **718** (2013) 1413.
- [20] T. Kitabayashi and M. Yasuè, Phys. Lett. B **713** (2012) 206.
- [21] T. Kitabayashi, Mod. Phys. Lett. A **28** (2013) 1350016.
- [22] S. Pascoli, S.T. Petcov and W. Rodejohann, Phys. Rev. D **68** (2003) 093007.
- [23] C. Giunti and M. Tanimoto, in Ref.[19].
- [24] See for example, S. Antusch, C. Biggio, E. F-Matinez, M. B. Gavela and J. L-Pavón, JHEP **10** (2006) 084; E. F-Matinez, M. B. Gavela, J. L-Pavón and O. Yasuda, Phys. Lett. B **649** (2007) 427; Z.-Z. Xing and S. Zhou, Phys. Lett. B **666** (2008) 166; S. Goswami and T. Ota, Phys. Rev. D **78** (2008) 033012; W. Rodejohann, Phys. Lett. B **684** (2010) 40; S. Antusch, S. Blanchet, M. Blennow and E. F-Matinez, JHEP **01** (2010) 017; D. V. Forero, S. Morisi, M. Tórtola and J. W. F. Valle, JHEP **09** (2011) 142; S. Khan, Nucl. Phys. B **864** (2012) 38.
- [25] G. Altarelli, F. Feruglio and I. Masina, in Ref.[19].
- [26] C. Jarlskog, Phys. Rev. Lett. **554** (1985) 1039.
- [27] T. Baba and M. Yasuè, Phys. Rev. D **75** (2007) 055001; Phys. Rev. D **77** (2008) 075008; Prog. Theor. Phys. **123** (2010) 659; Z.Z. Xing and Y.-L. Zhou, Phys. Lett. B **693** (2010) 584; B. Adhikary, M. Chakraborty and A. Ghosal, “Masses, mixing angles and phases of general Majorana neutrino mass matrix: application to broken cyclic symmetric mass matrix”, arXiv:1307.0988 [hep-ph]
- [28] J.F. Nieves and P.B. Pal, Phys. Rev. D **36** (1987) 315; Phys. Rev. D **64** (2001) 315.