

Penguin Pollution in $B \rightarrow J/\psi V$ Decays and Impact on the Extraction of the $B_s - \bar{B}_s$ mixing phase

Xin Liu

*School of Physics and Electronic Engineering, Jiangsu Normal University,
Xuzhou, Jiangsu 221116, People's Republic of China*

Wei Wang

*Helmholtz-Institut für Strahlen-und Kernphysik and Bethe Center for
Theoretical Physics, Universität Bonn, D-53115 Bonn, Germany*

Yuehong Xie

*Institute of Particle Physics, Huazhong Normal University,
Wuhan, Hubei 430079, People's Republic of China
and University of Edinburgh, Edinburgh EH9 3JZ, UK*

We formulate the most-general time-dependent distributions of $B_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ in which the direct CP violation is explicitly incorporated. We investigate the $B \rightarrow J/\psi V$ decays in the perturbative QCD approach where V is a light vector meson. Apart from the leading-order factorizable contributions, we also take into account QCD vertex corrections and the hard-spectator diagrams. With the inclusion of these sizeable corrections, most of our theoretical results for CP-averaged branching ratios, polarization fractions, CP-violating asymmetries, and relative phases are in good consistency with the available data. Based on the global agreement, we further explore the penguin contributions and point out that the ϕ_s extracted from $B_s \rightarrow J/\psi\phi$ can be shifted away by $\mathcal{O}(10^{-3})$.

PACS numbers: 13.25.Hw, 12.38.Bx, 14.40.Nd

I. INTRODUCTION

In the standard model (SM) of particle physics, CP violation arises from the non-vanishing complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix. Constraints stemming from the unitarity of CKM matrix can be pictorially represented as triangles, the length of whose sides are products of CKM matrix elements and the angles are relative phases between them. Due to the comparable size of the three sides, the so-called (bd) triangle from $V_{tb}V_{td}^* + V_{cb}V_{cd}^* + V_{ub}V_{ud}^* = 0$ is mostly discussed in the heavy flavor physics. In contrast, the (bs) triangle, $V_{tb}V_{ts}^* + V_{cb}V_{cs}^* + V_{ub}V_{us}^* = 0$, has a small complex phase, and provide a null test of SM. Thus it is also of great interest and deserves more theoretical and experimental efforts.

In the SM, the non-vanishing phase in (bs) triangle is related to the $B_s - \bar{B}_s$ mixing phase. States of B_s^0 or \bar{B}_s^0 at $t = 0$ can evolve in time and get mixed with each other. These states at t can be denoted as $B_s(t)$ and $\bar{B}_s(t)$. Since both the B_s^0 and \bar{B}_s^0 can decay into the same final state like $J/\psi\phi$, there is an indirect CP asymmetry (CPA) between the rates of $B_s(t) \rightarrow J/\psi\phi$ and $\bar{B}_s(t) \rightarrow J/\psi\phi$, quantified by

$$\text{Im} \left[\frac{q}{p} \frac{\bar{A}_f}{A_f} \right]. \quad (1)$$

Here the A_f and \bar{A}_f are the B_s and \bar{B}_s decay amplitudes which are dominated by the $b \rightarrow c\bar{c}s$ transition. Since the CKM factors $V_{cb}V_{cs}^*$ are real in the standard parametrization of CKM, the indirect CPA defined in Eq. (1) measures the phase in q/p defined with the form: $\phi_s = -\arg(q/p)$. This phase ϕ_s is tiny in the SM, and in particular $\phi_s = -2\beta_s = -2\arg[-V_{ts}V_{tb}^*/(V_{cs}V_{cb}^*)]$ [1]:

$$\phi_s = (-0.036 \pm 0.002) \text{ rad}. \quad (2)$$

The observation of a large non-zero value would be a signal for new physics beyond the SM.

The ϕ_s extraction has greatly benefited from measurements of time-dependent observables in $B_s/\bar{B}_s \rightarrow J/\psi\phi$, the analogue of the golden-channel $B \rightarrow J/\psi K_S$. In the experimental retrospect, many progresses have been made in the past few years. Thanks to the large amount of data sample collected on the Tevatron and LHC experiments, the result for ϕ_s is getting more and more precise [2–6]. Recently based on the data of 1.0fb^{-1} collected at 7 TeV in 2011, the LHCb collaboration gives [7]

$$\phi_s^{J/\psi\phi} = (0.07 \pm 0.09 \pm 0.01) \text{ rad}, \quad (3)$$

which is in agreement with the SM value in Eq. 2 when the errors are taken into account. Moreover new alternative channels are proposed and in particular the $B_s \rightarrow J/\psi f_0(980)$ is believed to have the supplementary power in reducing the error in ϕ_s [8, 9]. A characteristic feature of this mode is that $f_0(980)$ is a 0^{++} scalar meson, and thus the final state $J/\psi f_0$ is a CP eigenstate. In contrast with the $B_s \rightarrow J/\psi \phi$, it does not request to perform the angular decomposition, and therefore the time-dependent analysis is greatly simplified. Agreement on branching ratios (BRs) is found between theoretical calculation [10–13] and experimental measurements [14–16], while the ϕ_s is reported by the LHCb collaboration [7]

$$\phi_s^{J/\psi f_0} = (-0.14^{+0.17}_{-0.16} \pm 0.01) \text{ rad.} \quad (4)$$

However it is necessary to point out that, due to the unknown internal structure of the scalar $f_0(980)$, the extracted results for ϕ_s may be contaminated by various hadronic corrections in this process [13].

On the theoretical side, although decays of the B_s/B_s^0 meson into $J/\psi(\phi/f_0)$ are mainly governed by the $b \rightarrow c\bar{c}s$ transition at the quark level, there are indeed penguin contributions with non vanishing different weak phases. Thus the indirect CPA can be shifted away from the ϕ_s . Though intuitively penguin contribution is expected to be small in the SM, a complete and reliable estimate of its effects by some QCD-inspired approach is not yet available. Such estimate will become mandatory soon especially when confronted with the gradually-reducing experimental error. As a reference, after the upgrade of LHC the error can be diminished to $\Delta\phi_s \sim 0.008$ [6].

The main purpose of this work is to estimate the penguin contributions in the $B_{u/d/s} \rightarrow J/\psi V (V = \rho, \omega, \phi, K^*)$ decays and explore the impact to the CPA measurement. To do so, we will present the time-dependent angular distributions, in which the direct CP asymmetry is incorporated. Instead of using the flavor SU(3) symmetry to relate the effects in $B_s \rightarrow J/\psi \phi$ and the counterpart of B decay modes [17, 18], we will adopt the QCD-based factorization approach to directly compute both tree amplitudes and penguin amplitudes. In particular, the perturbative QCD factorization approach (pQCD) [19–22] will be used in this work, the same approach that has been applied to study the $B \rightarrow J/\psi P$ [23–25] and estimate the penguin contribution to ΔS in $B \rightarrow J/\psi K_S$ [26]. Recent development of this approach can be found in Refs. [27, 28]. Apart from the leading-order (LO) contributions, we will also include the next-to-leading order (NLO) corrections in α_s , which are sizeable especially to penguin contributions.

The rest of this paper is organized as follows. We derive the time-dependent angular distributions in Sec. II. Section III is devoted to the ingredients of the basic formalism in the pQCD approach. The analytic expressions for the $B_{u/d/s} \rightarrow J/\psi V$ decay amplitudes in the pQCD approach are also collected in this section. Numerical results for the CP-averaged branching ratios (BRs), polarization fractions, relative phases, and CP-violating asymmetries of the considered decays are given in Sec. IV. We summarize this work and conclude in Sec. V. Some calculation formulas are relegated to the appendix.

II. HELICITY-BASED ANGULAR DISTRIBUTIONS OF $B_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$




FIG. 1. Kinematics of $B_s/\bar{B}_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$. The moving direction of the K^+K^- pair in the B_s/\bar{B}_s rest frame is chosen as the z axis. The polar angle θ_K (θ_l) is defined as the angle between the flight direction of K^+ (l^+) and the z axis in the ϕ (J/ψ) rest frame. ϕ is the azimuthal angle between the two decay planes of ϕ and J/ψ .

The decay distributions can be expressed in terms of helicity angles, θ_K, θ_l, ϕ . The convention on the kinematics in $B_s/\bar{B}_s \rightarrow \phi(\rightarrow K^+K^-)J/\psi(\rightarrow l^+l^-)$ is illustrated in Fig. 1. The moving direction of the K^+K^- pair in the B_s/\bar{B}_s rest frame is chosen as the z axis. The polar angle θ_K (θ_l) is defined as the angle between the flight direction of K^+ (l^+) and the z axis in the ϕ (J/ψ) rest frame. ϕ is the azimuthal angle between the two decay planes of ϕ and J/ψ .

TABLE I. The time-dependent angular functions defined in Eqs. (5) and (6), as discussed in the text. In the amplitude A_i^j , the superscript j denotes the spin, while the subscript $i = 0, ||, \perp$ corresponds to the three polarization configurations of the K^+K^- system. A_0^j is also usually referred to as A_S in the literature. Some abbreviations have been used for cosine and sine functions, for instance $c_K = \cos \theta_K$, $s_K = \sin \theta_K$, $s_{2K} = \sin(2\theta_K)$

f_k	N_k	a_k	b_k	c_k	d_k
$c_K^2 s_l^2$	$\frac{ A_0^1 ^2 + \bar{A}_0^1 ^2}{2}$	1	$-\frac{2 \lambda_0^1 }{1+ \lambda_0^1 ^2} \cos(\phi_0^1)$	$\frac{1- \lambda_0^1 ^2}{1+ \lambda_0^1 ^2}$	$\frac{2 \lambda_0^1 }{1+ \lambda_0^1 ^2} \sin(\phi_0^1)$
$\frac{s_K^2(1-c_\phi^2 c_l^2)}{2}$	$\frac{ A_{ }^1 ^2 + \bar{A}_{ }^1 ^2}{2}$	1	$-\frac{2 \lambda_{ }^1 }{1+ \lambda_{ }^1 ^2} \cos(\phi_{ }^1)$	$\frac{1- \lambda_{ }^1 ^2}{1+ \lambda_{ }^1 ^2}$	$\frac{2 \lambda_{ }^1 }{1+ \lambda_{ }^1 ^2} \sin(\phi_{ }^1)$
$\frac{s_K^2(1-s_\phi^2 c_l^2)}{2}$	$\frac{ A_{\perp}^1 ^2 + \bar{A}_{\perp}^1 ^2}{2}$	1	$\frac{2 \lambda_{\perp}^1 }{1+ \lambda_{\perp}^1 ^2} \cos(\phi_{\perp}^1)$	$\frac{1- \lambda_{\perp}^1 ^2}{1+ \lambda_{\perp}^1 ^2}$	$-\frac{2 \lambda_{\perp}^1 }{1+ \lambda_{\perp}^1 ^2} \sin(\phi_{\perp}^1)$
$s_K^2 s_l^2 s_\phi c_\phi$	$ A_{\perp}^1 A_{ }^1 $	$\frac{1}{2} \begin{bmatrix} \sin(\delta_{\perp}^1 - \delta_{ }^1) - \lambda_{\perp}^1 \lambda_{ }^1 \\ \sin(\delta_{\perp}^1 - \delta_{ }^1 - \phi_{\perp}^1 + \phi_{ }^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \lambda_{\perp}^1 \sin(\delta_{\perp}^1 - \delta_{ }^1 - \phi_{\perp}^1) \\ + \lambda_{ }^1 \sin(\delta_{ }^1 - \delta_{\perp}^1 - \phi_{ }^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \sin(\delta_{\perp}^1 - \delta_{ }^1) + \lambda_{\perp}^1 \lambda_{ }^1 \\ \sin(\delta_{\perp}^1 - \delta_{ }^1 - \phi_{\perp}^1 + \phi_{ }^1) \end{bmatrix}$	$-\frac{1}{2} \begin{bmatrix} \lambda_{\perp}^1 \cos(\delta_{\perp}^1 - \delta_{ }^1 - \phi_{\perp}^1) \\ + \lambda_{ }^1 \cos(\delta_{ }^1 - \delta_{\perp}^1 - \phi_{ }^1) \end{bmatrix}$
$\frac{\sqrt{2}}{4} s_{2K} s_{2l} c_\phi$	$ A_0^1 A_{ }^1 $	$\frac{1}{2} \begin{bmatrix} \cos(\delta_0^1 - \delta_{ }^1) + \lambda_0^1 \lambda_{ }^1 \\ \cos(\delta_0^1 - \delta_{ }^1 - \phi_0^1 + \phi_{ }^1) \end{bmatrix}$	$-\frac{1}{2} \begin{bmatrix} \lambda_0^1 \cos(\delta_0^1 - \delta_{ }^1 - \phi_0^1) \\ + \lambda_{ }^1 \cos(\delta_{ }^1 - \delta_0^1 - \phi_{ }^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \cos(\delta_0^1 - \delta_{ }^1) - \lambda_0^1 \lambda_{ }^1 \\ \cos(\delta_0^1 - \delta_{ }^1 - \phi_0^1 + \phi_{ }^1) \end{bmatrix}$	$-\frac{1}{2} \begin{bmatrix} \lambda_0^1 \sin(\delta_0^1 - \delta_{ }^1 - \phi_0^1) \\ + \lambda_{ }^1 \sin(\delta_{ }^1 - \delta_0^1 - \phi_{ }^1) \end{bmatrix}$
$\frac{\sqrt{2}}{4} s_{2K} s_{2l} s_\phi$	$ A_0^1 A_{\perp}^1 $	$\frac{1}{2} \begin{bmatrix} \sin(\delta_0^1 - \delta_{\perp}^1) - \lambda_0^1 \lambda_{\perp}^1 \\ \sin(\delta_0^1 - \delta_{\perp}^1 - \phi_0^1 + \phi_{\perp}^1) \end{bmatrix}$	$-\frac{1}{2} \begin{bmatrix} \lambda_0^1 \sin(\delta_0^1 - \delta_{\perp}^1 - \phi_0^1) \\ + \lambda_{\perp}^1 \sin(\delta_{\perp}^1 - \delta_0^1 - \phi_{\perp}^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \sin(\delta_0^1 - \delta_{\perp}^1) + \lambda_0^1 \lambda_{\perp}^1 \\ \sin(\delta_0^1 - \delta_{\perp}^1 - \phi_0^1 + \phi_{\perp}^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \lambda_0^1 \cos(\delta_0^1 - \delta_{\perp}^1 - \phi_0^1) \\ + \lambda_{\perp}^1 \cos(\delta_{\perp}^1 - \delta_0^1 - \phi_{\perp}^1) \end{bmatrix}$
$\frac{1}{3} s_l^2$	$\frac{ A_0^0 ^2 + \bar{A}_0^0 ^2}{2}$	1	$\frac{2 \lambda_0^0 }{1+ \lambda_0^0 ^2} \cos(\phi_0^0)$	$\frac{1- \lambda_0^0 ^2}{1+ \lambda_0^0 ^2}$	$-\frac{2 \lambda_0^0 }{1+ \lambda_0^0 ^2} \sin(\phi_0^0)$
$\frac{2s_K s_l c_l c_\phi}{\sqrt{6}}$	$ A_0^0 A_{ }^1 $	$\frac{1}{2} \begin{bmatrix} \cos(\delta_0^0 - \delta_{ }^1) - \lambda_0^0 \lambda_{ }^1 \\ \cos(\delta_0^0 - \delta_{ }^1 - \phi_0^0 + \phi_{ }^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \lambda_0^0 \cos(\delta_0^0 - \delta_{ }^1 - \phi_0^0) \\ - \lambda_{ }^1 \cos(\delta_{ }^1 - \delta_0^0 - \phi_{ }^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \cos(\delta_0^0 - \delta_{ }^1) + \lambda_0^0 \lambda_{ }^1 \\ \cos(\delta_0^0 - \delta_{ }^1 - \phi_0^0 + \phi_{ }^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \lambda_0^0 \sin(\delta_0^0 - \delta_{ }^1 - \phi_0^0) \\ - \lambda_{ }^1 \sin(\delta_{ }^1 - \delta_0^0 - \phi_{ }^1) \end{bmatrix}$
$\frac{2s_K s_l c_l s_\phi}{\sqrt{6}}$	$ A_0^0 A_{\perp}^{1*} $	$\frac{1}{2} \begin{bmatrix} \sin(\delta_0^0 - \delta_{\perp}^1) + \lambda_0^0 \lambda_{\perp}^1 \\ \sin(\delta_0^0 - \delta_{\perp}^1 - \phi_0^0 + \phi_{\perp}^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \lambda_0^0 \sin(\delta_0^0 - \delta_{\perp}^1 - \phi_0^0) \\ - \lambda_{\perp}^1 \sin(\delta_{\perp}^1 - \delta_0^0 - \phi_{\perp}^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \sin(\delta_0^0 - \delta_{\perp}^1) - \lambda_0^0 \lambda_{\perp}^1 \\ \sin(\delta_0^0 - \delta_{\perp}^1 - \phi_0^0 + \phi_{\perp}^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} - \lambda_0^0 \cos(\delta_0^0 - \delta_{\perp}^1 - \phi_0^0) \\ + \lambda_{\perp}^1 \cos(\delta_{\perp}^1 - \delta_0^0 - \phi_{\perp}^1) \end{bmatrix}$
$\frac{2c_K s_l^2}{\sqrt{3}}$	$ A_0^0 A_0^1 $	$\frac{1}{2} \begin{bmatrix} \cos(\delta_0^0 - \delta_0^1) - \lambda_0^0 \lambda_0^1 \\ \cos(\delta_0^0 - \delta_0^1 - \phi_0^0 + \phi_0^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \lambda_0^0 \cos(\delta_0^0 - \delta_0^1 - \phi_0^0) \\ - \lambda_0^1 \cos(\delta_0^1 - \delta_0^0 - \phi_0^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \cos(\delta_0^0 - \delta_0^1) + \lambda_0^0 \lambda_0^1 \\ \cos(\delta_0^0 - \delta_0^1 - \phi_0^0 + \phi_0^1) \end{bmatrix}$	$\frac{1}{2} \begin{bmatrix} \lambda_0^0 \sin(\delta_0^0 - \delta_0^1 - \phi_0^0) \\ - \lambda_0^1 \sin(\delta_0^1 - \delta_0^0 - \phi_0^1) \end{bmatrix}$

The full decay amplitudes can be calculated using the helicity amplitudes and for a detailed discussion we also refer the reader to Refs. [29–32]. In the presence of S-wave K^+K^- the angular distribution for $B_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ at the time t of the state that was a pure B_s at $t = 0$ is given as

$$\frac{d^4\Gamma(t)}{dm_{KK}^2 d\cos\theta_K d\cos\theta_l d\phi} = \sum_{k=1}^{10} h_k(t) f_k(\theta_K, \theta_l, \phi), \quad (5)$$

where the time-dependent functions $h_k(t)$ are given as

$$h_k(t) = \frac{3}{4\pi} e^{-\Gamma t} \left\{ a_k \cosh \frac{\Delta\Gamma t}{2} + b_k \sinh \frac{\Delta\Gamma t}{2} + c_k \cos(\Delta m t) + d_k \sin(\Delta m t) \right\}. \quad (6)$$

Here $\Delta m = m_H - m_L$, $\Delta\Gamma = \Gamma_L - \Gamma_H$, and $\Gamma = (\Gamma_L + \Gamma_H)/2$. For the state that was a \bar{B}_s at $t = 0$, the signs of c_k and d_k should be reversed. The explicit results for these coefficients are collected in Table I, in which we have used

$$\lambda_i^j = \eta_i^j \frac{q}{p} \frac{\bar{A}_i^j}{A_i^j} \equiv |\lambda_i^j| e^{-i\phi_i^j}. \quad (7)$$

η_i^j is the CP-eigenvalue of the final state, j is the spin and i is the polarization/helicity of the K^+K^- system.

III. PERTURBATIVE QCD CALCULATION

Because of the large mass of the bottom quark, for convenience, we will work in the B meson rest frame, where B denotes any of the B_d, B_u, B_s mesons. Throughout this paper, we will use light-cone coordinate (P^+, P^-, \mathbf{P}_T) to describe the meson's momenta with the definitions

$$P^\pm = \frac{p_0 \pm p_3}{\sqrt{2}} \quad \text{and} \quad \mathbf{P}_T = (p_1, p_2). \quad (8)$$

Then for $B \rightarrow J/\psi V$ decays, the involved momenta can be written as

$$P_1 = \frac{m_B}{\sqrt{2}}(1, 1, \mathbf{0}_T), \quad P_2 = \frac{m_B}{\sqrt{2}}(1 - r_3^2, r_2^2, \mathbf{0}_T), \quad P_3 = \frac{m_B}{\sqrt{2}}(r_3^2, 1 - r_2^2, \mathbf{0}_T). \quad (9)$$

The J/ψ (V) meson is chosen to move on the plus (minus) z direction carrying the momentum P_2 (P_3), $r_2 = m_{J/\psi}/m_B$, and $r_3 = m_V/m_B$. In the numerical calculation, higher-order terms in r_3 can be neglected, as $r_3^2 \sim 0.04$ is numerically small. The longitudinal and transverse polarization vectors are denoted by ϵ^L and ϵ^T with the explicit forms:

$$\epsilon_2^L = \frac{m_B}{\sqrt{2}m_{J/\psi}}(1 - r_3^2, -r_2^2, \mathbf{0}_T) \quad \text{and} \quad \epsilon_3^L = \frac{m_B}{\sqrt{2}m_V}(-r_3^2, 1 - r_2^2, \mathbf{0}_T). \quad (10)$$

The transverse ones are parameterized as $\epsilon_2^T = (0, 0, \mathbf{1}_T)$ and $\epsilon_3^T = (0, 0, \mathbf{1}_T)$. Putting the (light) quark momenta in B , J/ψ and V mesons as k_1 , k_2 , and k_3 , respectively, we have

$$k_1 = (x_1 P_1^+, 0, \mathbf{k}_{1T}), \quad k_2 = x_2 P_2 + (0, 0, \mathbf{k}_{2T}), \quad k_3 = x_3 P_3 + (0, 0, \mathbf{k}_{3T}). \quad (11)$$

In the pQCD approach the decay amplitude for the $B \rightarrow J/\psi V$ can be written in a factorized form:

$$\begin{aligned} \mathcal{A}(B \rightarrow J/\psi V) &\sim \int dx_1 dx_2 dx_3 b_1 db_1 b_2 db_2 b_3 db_3 \\ &\times \text{Tr} \left[C(t) \Phi_B(x_1, b_1) \Phi_{J/\psi}(x_2, b_2) \Phi_V(x_3, b_3) H(x_i, b_i, t) S_t(x_i) e^{-S(t)} \right], \end{aligned} \quad (12)$$

where b_i is the conjugate space coordinate of k_{iT} , and t is the largest energy scale in function $H(x_i, b_i, t)$. The explicit forms will be given in the following.

A. Wave Functions and Distribution Amplitudes

The heavy B meson is usually treated as a heavy-light system and its light-cone wave function can generally be defined as

$$\begin{aligned} \Phi_{B, \alpha\beta, ij} &\equiv \langle 0 | \bar{b}_{\beta j}(0) q_{\alpha i}(z) | B(P) \rangle \\ &= \frac{i\delta_{ij}}{\sqrt{2N_c}} \int dx d^2 k_T e^{-i(xP^- z^+ - k_T z_T)} \{ (P^+ + m_B) \gamma_5 \phi_B(x, k_T) \}_{\alpha\beta}; \end{aligned} \quad (13)$$

where the indices i, j and α, β are the color indices and Lorentz indices, respectively. $P(m)$ is the momentum(mass) of the B meson, N_c is the color factor, and k_T is the intrinsic transverse momentum of the light quark in B meson. Note that there in principle are two Lorentz structures of the wave function to be considered in the numerical calculations, however, the contribution induced by the second Lorentz structure is numerically negligible [33, 34].

In Eq. (13), the $\phi_B(x, k_T)$ is the B meson light-cone distribution amplitude (LCDA). It is convenient to work in the impact coordinate space and the LCDA has the normalization

$$\int_0^1 dx \phi_B(x, b=0) = \frac{f_B}{2\sqrt{2N_c}}, \quad (14)$$

where f_B is the decay constant. The following form has been widely used in the pQCD approach [35, 36]

$$\phi_B(x, b) = N_B x^2 (1-x)^2 \exp \left[-\frac{1}{2} \left(\frac{xm_B}{\omega_B} \right)^2 - \frac{\omega_B^2 b^2}{2} \right]. \quad (15)$$

Based on the rich experimental data on the B mesons in recent years, the shape parameter ω_B has been fixed at about 0.40 GeV [35], while it is $\omega_B = 0.50$ GeV for the B_s meson [37].

The nonlocal matrix elements associated with longitudinally and transversely polarized J/ψ mesons are decomposed into

$$\Phi_{J/\psi, \alpha\beta, ij}^L \equiv \langle J/\psi(P, \epsilon_L) | \bar{c}(z)_{\beta j} c(0)_{\alpha i} | 0 \rangle = \frac{\delta_{ij}}{\sqrt{2N_c}} \int_0^1 dx e^{ixP \cdot z} \left\{ m_{J/\psi} \not{\epsilon}_L \phi_{J/\psi}(x) + \not{\epsilon}_L P \phi_{J/\psi}^t(x) \right\}_{\alpha\beta}, \quad (16)$$

$$\Phi_{J/\psi, \alpha\beta, ij}^T \equiv \langle J/\psi(P, \epsilon_T) | \bar{c}(z)_{\beta j} c(0)_{\alpha i} | 0 \rangle = \frac{\delta_{ij}}{\sqrt{2N_c}} \int_0^1 dx e^{ixP \cdot z} \left\{ m_{J/\psi} \not{\epsilon}_T \phi_{J/\psi}^v(x) + \not{\epsilon}_T P \phi_{J/\psi}^T(x) \right\}_{\alpha\beta}, \quad (17)$$

respectively. This defines the twist-2 distribution amplitudes $\phi_{J/\psi}$ and $\phi_{J/\psi}^T$, and the twist-3 distribution amplitudes $\phi_{J/\psi}^t$ and $\phi_{J/\psi}^v$ with the c quark carrying the fractional momentum xP . With the inclusion of the relativistic corrections, the distribution amplitudes for the J/ψ have been derived as [38]

$$\begin{aligned} \phi_{J/\psi}(x) &= \phi_{J/\psi}^T(x) = 9.58 \frac{f_{J/\psi}}{2\sqrt{2N_c}} x(1-x) \left[\frac{x(1-x)}{1-2.8x(1-x)} \right]^{0.7}, \\ \phi_{J/\psi}^t(x) &= 10.94 \frac{f_{J/\psi}}{2\sqrt{2N_c}} (1-2x)^2 \left[\frac{x(1-x)}{1-2.8x(1-x)} \right]^{0.7}, \\ \phi_{J/\psi}^v(x) &= 1.67 \frac{f_{J/\psi}}{2\sqrt{2N_c}} [1 + (2x-1)^2] \left[\frac{x(1-x)}{1-2.8x(1-x)} \right]^{0.7}. \end{aligned} \quad (18)$$

Here the twist-3 LCDAs $\phi^{t,v}$ vanish at the endpoint due to the additional factor $[x(1-x)]^{0.7}$.

Up to twist-3, the light-cone wave function for a light vector meson is given as

$$\begin{aligned} \Phi_{V, \alpha\beta, ij}^L &\equiv \langle V(P, \epsilon_L) | \bar{q}(z)_{\beta j} q(0)_{\alpha i} | 0 \rangle \\ &= \frac{\delta_{ij}}{\sqrt{2N_c}} \int_0^1 dx e^{ixP \cdot z} \left\{ m_V \not{\epsilon}_L \phi_V(x) + \not{\epsilon}_L P \phi_V^t(x) + m_V \phi_V^s(x) \right\}_{\alpha\beta}, \end{aligned} \quad (19)$$

$$\begin{aligned} \Phi_{V, \alpha\beta, ij}^T &\equiv \langle V(P, \epsilon_T) | \bar{q}(z)_{\beta j} q(0)_{\alpha i} | 0 \rangle \\ &= \frac{\delta_{ij}}{\sqrt{2N_c}} \int_0^1 dx e^{ixP \cdot z} \left\{ m_V \not{\epsilon}_T \phi_V^v(x) + \not{\epsilon}_T P \phi_V^T(x) + m_V i\epsilon_{\mu\nu\rho\sigma} \gamma_5 \gamma^\mu \epsilon_T^\nu n^\rho v^\sigma \phi_V^a(x) \right\}_{\alpha\beta}, \end{aligned} \quad (20)$$

for longitudinal polarization and transverse polarization, respectively. x is the momentum fraction carried by the quark in the meson, and $n = (1, 0, \mathbf{0}_T)$ and $v = (0, 1, \mathbf{0}_T)$ are dimensionless light-like unit vectors. In the above equation, we have adopted the convention $\epsilon^{0123} = 1$ for the Levi-Civita tensor $\epsilon^{\mu\nu\alpha\beta}$.

TABLE II. Decay constants for the light vector mesons (in MeV)

f_ρ	f_ρ^T	f_ω	f_ω^T	f_{K^*}	$f_{K^*}^T$	f_ϕ	f_ϕ^T
209 ± 2	165 ± 9	195 ± 3	145 ± 10	217 ± 5	185 ± 10	231 ± 4	200 ± 10

The twist-2 LCDAs can be expanded in terms of Gegenbauer polynomials:

$$\phi_V(x) = \frac{3f_V}{\sqrt{2N_c}} x(1-x) \left[1 + 3a_{1V}^{\parallel} (2x-1) + a_{2V}^{\parallel} \frac{3}{2} (5(2x-1)^2 - 1) \right], \quad (21)$$

$$\phi_V^T(x) = \frac{3f_V^T}{\sqrt{2N_c}} x(1-x) \left[1 + 3a_{1V}^{\perp} (2x-1) + a_{2V}^{\perp} \frac{3}{2} (5(2x-1)^2 - 1) \right]. \quad (22)$$

f_V and f_V^T are the longitudinal and transverse decay constants. f_V can be extracted from the data on $V^0 \rightarrow l^+ l^-$ and $\tau \rightarrow V^- \bar{\nu}$ [39], while the transverse decay constants f_V^T are taken from Ref. [40]. We collect these quantities in Tab. II. For the Gegenbauer moments of the light-vector mesons, we take the recent updates [41]:

$$a_{1K^*}^{\parallel} = 0.03 \pm 0.02, \quad a_{2K^*}^{\parallel} = 0.11 \pm 0.09, \quad a_{2\rho}^{\parallel} = a_{2\omega}^{\parallel} = 0.15 \pm 0.07, \quad a_{2\phi}^{\parallel} = 0.18 \pm 0.08; \quad (23)$$

$$a_{1K^*}^{\perp} = 0.04 \pm 0.03, \quad a_{2K^*}^{\perp} = 0.10 \pm 0.08, \quad a_{2\rho}^{\perp} = a_{2\omega}^{\perp} = 0.14 \pm 0.06, \quad a_{2\phi}^{\perp} = 0.14 \pm 0.07. \quad (24)$$

The asymptotic forms of the twist-3 distribution amplitudes $\phi_V^{t,s}$ and $\phi_V^{v,a}$ will be used in this work:

$$\phi_V^t(x) = \frac{3f_V^T}{2\sqrt{2N_c}} (2x-1)^2, \quad \phi_V^s(x) = -\frac{3f_V^T}{2\sqrt{2N_c}} (2x-1), \quad (25)$$

$$\phi_V^v(x) = \frac{3f_V}{8\sqrt{2N_c}} (1 + (2x-1)^2), \quad \phi_V^a(x) = -\frac{3f_V}{4\sqrt{2N_c}} (2x-1). \quad (26)$$

In the pQCD approach, the above choices of vector meson LCDAs can successfully explain not only the measured branching ratios but also polarization fractions for the $B \rightarrow K^* \phi$, $B \rightarrow K^* \rho$ and $B \rightarrow \rho \rho$ [42, 43].

B. Perturbative Calculations




FIG. 2. (Color online) Typical Feynman diagrams contributing to $B \rightarrow J/\psi V$ decays at leading-order in α_s in the pQCD approach.

The effective Hamiltonian H_{eff} can be written as [44]

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left\{ V_{cb}^* V_{cD} [C_1(\mu) O_1^c(\mu) + C_2(\mu) O_2^c(\mu)] - V_{tb}^* V_{tD} \left[\sum_{i=3}^{10} C_i(\mu) O_i(\mu) \right] \right\} + \text{H.c.}, \quad (27)$$

with the Fermi constant $G_F = 1.16639 \times 10^{-5} \text{GeV}^{-2}$, the light $D = d, s$ quark, and Wilson coefficients $C_i(\mu)$ at the renormalization scale μ . The local four-quark operators $O_i (i = 1, \dots, 10)$ are written as

(1) current-current(tree) operators

$$O_1^c = (\bar{D}_\alpha c_\beta)_{V-A} (\bar{c}_\beta b_\alpha)_{V-A}, \quad O_2^c = (\bar{D}_\alpha c_\alpha)_{V-A} (\bar{c}_\beta b_\beta)_{V-A}, \quad (28)$$

(2) QCD penguin operators

$$\begin{aligned} O_3 &= (\bar{D}_\alpha b_\alpha)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\beta)_{V-A}, \quad O_4 = (\bar{D}_\alpha b_\beta)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\alpha)_{V-A}, \\ O_5 &= (\bar{D}_\alpha b_\alpha)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\beta)_{V+A}, \quad O_6 = (\bar{D}_\alpha b_\beta)_{V-A} \sum_{q'} (\bar{q}'_\beta q'_\alpha)_{V+A}, \end{aligned} \quad (29)$$

(3) electroweak penguin operators

$$\begin{aligned} O_7 &= \frac{3}{2} (\bar{D}_\alpha b_\alpha)_{V-A} \sum_{q'} e_{q'} (\bar{q}'_\beta q'_\beta)_{V+A}, \quad O_8 = \frac{3}{2} (\bar{D}_\alpha b_\beta)_{V-A} \sum_{q'} e_{q'} (\bar{q}'_\beta q'_\alpha)_{V+A}, \\ O_9 &= \frac{3}{2} (\bar{D}_\alpha b_\alpha)_{V-A} \sum_{q'} e_{q'} (\bar{q}'_\beta q'_\beta)_{V-A}, \quad O_{10} = \frac{3}{2} (\bar{D}_\alpha b_\beta)_{V-A} \sum_{q'} e_{q'} (\bar{q}'_\beta q'_\alpha)_{V-A}, \end{aligned} \quad (30)$$

with the color indices α, β and the notations $(\bar{q}' q')_{V \pm A} = \bar{q}' \gamma_\mu (1 \pm \gamma_5) q'$. The index q' in the summation of the above operators runs through u, d, s, c , and b . Abbreviations of Wilson coefficients will be used

$$a_1 = C_2 + \frac{C_1}{3}, \quad a_2 = C_1 + \frac{C_2}{3}, \quad a_i = C_i + \frac{C_{i \pm 1}}{3} (i = 3 - 10), \quad (31)$$

where the upper(lower) sign applies, when i is odd(even).

In the perturbative QCD approach, the scale t in Wilson coefficients $C_i(t)$, hard-kernel $H(x_i, b_i, t)$ and Sudakov factor $e^{-S(t)}$ is chosen as the largest energy scale in the gluon and/or the quark propagators of a given Feynman diagram, in order to suppress the higher order corrections and improve the convergence of the perturbative calculation.

In the range of $t < m_b$ or $t \geq m_b$, the number of active quarks is $N_f = 4$ or $N_f = 5$, respectively. The explicit expressions of the LO and NLO C_i can be found, for instance, in Ref. [44].

In the LO calculation, we shall use the LO Wilson coefficients $C_i(m_W)$, the LO renormalization group (RG) evolution matrix $U(t, m)^{(0)}$ for the Wilson coefficient and the LO $\alpha_s(t)$:

$$\alpha_s(t) = \frac{4\pi}{\beta_0 \ln \left[t^2 / \Lambda_{QCD}^2 \right]}, \quad (32)$$

where $\beta_0 = (33 - 2N_f)/3$. In the NLO contributions, the NLO Wilson coefficients $C_i(m_W)$, the NLO RG evolution matrix $U(t, m, \alpha)$ (see Eq. (7.22) in Ref. [44]) and the $\alpha_s(t)$ at two-loop level will be used:

$$\alpha_s(t) = \frac{4\pi}{\beta_0 \ln \left[t^2 / \Lambda_{QCD}^2 \right]} \cdot \left\{ 1 - \frac{\beta_1}{\beta_0^2} \cdot \frac{\ln \left[\ln \left[t^2 / \Lambda_{QCD}^2 \right] \right]}{\ln \left[t^2 / \Lambda_{QCD}^2 \right]} \right\}, \quad (33)$$

where $\beta_1 = (306 - 38N_f)/3$. Using $\Lambda_{QCD}^{(5)} = 0.225$ GeV, we get $\Lambda_{QCD}^{(4)} = 0.287$ GeV (0.326 GeV) for LO (NLO) case. As discussed in Ref. [45, 46], it is reasonable to choose $\mu_0 = 1.0$ GeV as the lower cut-off for the hard scale t .

Decay amplitudes $\mathcal{A}^{(\sigma)}$ for the $B \rightarrow J/\psi(P_2, \epsilon_2^*)V(P_3, \epsilon_3^*)$ can be decomposed into three independent Lorentz structures

$$\begin{aligned} \mathcal{A}^{(\sigma)} &= \epsilon_{2\mu}^*(\sigma) \epsilon_{3\nu}^*(\sigma) \left[a g^{\mu\nu} + \frac{b}{m_{J/\psi} m_V} P_1^\mu P_1^\nu + i \frac{c}{m_{J/\psi} m_V} \epsilon^{\mu\nu\alpha\beta} P_{2\alpha} P_{3\beta} \right] \\ &\equiv m_B^2 \mathcal{M}_L + m_B^2 \mathcal{M}_N \epsilon_2^*(\sigma = T) \cdot \epsilon_3^*(\sigma = T) + i \mathcal{M}_T \epsilon^{\alpha\beta\gamma\rho} \epsilon_{2\alpha}^*(\sigma) \epsilon_{3\beta}^*(\sigma) P_{2\gamma} P_{3\rho}, \end{aligned} \quad (34)$$

where the superscript σ denotes the helicity of the final vector meson. The $\mathcal{M}_i (i = L, N, T)$ are written in terms of the Lorentz-invariant amplitudes a , b and c

$$\begin{aligned} m_B^2 \mathcal{M}_L &= a \epsilon_2^*(L) \cdot \epsilon_3^*(L) + \frac{b}{m_{J/\psi} m_V} \epsilon_2^*(L) \cdot P_3 \epsilon_3^*(L) \cdot P_2, \\ m_B^2 \mathcal{M}_N &= a \epsilon_2^*(T) \cdot \epsilon_3^*(T), \\ m_B^2 \mathcal{M}_T &= \frac{c}{r_2 r_3}. \end{aligned} \quad (35)$$

By taking various contributions from the relevant Feynman diagrams into consideration, one can derive the total decay amplitudes for the $B \rightarrow J/\psi V$ as

$$\begin{aligned} \mathcal{M}^h &= F^h f_{J/\psi} \left\{ V_{cb}^* V_{cd(s)} a_2 - V_{tb}^* V_{td(s)} \left(a_3 + a_5 + a_7 + a_9 \right) \right\} \\ &\quad + M^h \left\{ V_{cb}^* V_{cd(s)} C_2 - V_{tb}^* V_{td(s)} \left(C_4 - C_6 - C_8 + C_{10} \right) \right\}, \end{aligned} \quad (36)$$

where the superscript h standing for the three polarizations L, N , and T , respectively. Here F and M stands for the contributions of factorizable and non-factorizable diagrams from $(V - A)(V - A)$ operators. The LO factorization amplitudes derived from Fig. 2 for three polarizations can be read as,

$$\begin{aligned} F^L &= \zeta 8\pi C_F m_B^4 \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) (r_2^2 - 1) \\ &\quad \times \left\{ \left[(r_2^2 - 1)x_3 - 1 \right] \phi_V(x_3) + r_3(2x_3 - 1) \phi_V^s(x_3) - r_3[2(r_2^2 - 1)x_3 + r_2^2 + 1] \phi_V^t(x_3) \right\} \\ &\quad \times h_{fs}(x_1, x_3, b_1, b_3) E_{fs}(t_a) - \left[2r_3 \phi_V^s(x_3) \right] h_{fs}(x_3, x_1, b_3, b_1) E_{fs}(t_b) \Big\}, \end{aligned} \quad (37)$$

$$\begin{aligned} F^N &= \zeta 8\pi C_F m_B^4 \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) r_2 \\ &\quad \times \left\{ - \left[(r_2^2 - 1)[r_3(r_2^2 - 1)x_3 \phi_V^a(x_3) + \phi_V^T(x_3)] + r_3[(r_2^2 - 1)x_3 - 2] \phi_V^v(x_3) \right] E_{fs}(t_a) \right. \\ &\quad \left. \times h_{fs}(x_1, x_3, b_1, b_3) + r_3(r_2^2 - 1) \left[(r_2^2 - 1) \phi_V^a(x_3) - \phi_V^v(x_3) \right] h_{fs}(x_3, x_1, b_3, b_1) E_{fs}(t_b) \right\}, \end{aligned} \quad (38)$$

$$\begin{aligned}
F^T = & \zeta 16\pi C_F m_B^4 \int_0^1 dx_1 dx_3 \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1, b_1) r_2 \\
& \times \left\{ - \left[r_3 x_3 \phi_V^v(x_3) - \phi_V^T(x_3) + r_3 [(r_2^2 - 1)x_3 - 2] \phi_V^a(x_3) \right] h_{fs}(x_1, x_3, b_1, b_3) \right. \\
& \left. \times E_{fs}(t_a) - r_3 \left[(r_2^2 - 1) \phi_V^a(x_3) - \phi_V^v(x_3) \right] h_{fs}(x_3, x_1, b_3, b_1) E_{fs}(t_b) \right\}, \quad (39)
\end{aligned}$$

where the expressions of E_i and h_i can be found in the Appendix. The isospin factor is given as $\zeta = 1$ except $\zeta = -\frac{1}{\sqrt{2}}$, $\frac{1}{\sqrt{2}}$ for ρ^0 , ω .

For the nonfactorizable spectator diagrams, Fig. 2(c) and 2(d), all three meson wave functions are involved. For the $(V - A)(V - A)$ operators, the corresponding decay amplitude is

$$\begin{aligned}
M^L = & \zeta \frac{16\sqrt{6}}{3} \pi C_F m_B^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) (r_2^2 - 1) \\
& \times \left\{ [\phi_V(x_3) - 2r_3 \phi_V^t(x_3)] \left[x_3 \phi_{J/\psi}(x_2) + (2x_2 - x_3) r_2^2 \phi_{J/\psi}(x_2) \right. \right. \\
& \left. \left. - 2r_2 r_c \phi_{J/\psi}^t(x_2) \right] \right\} h_{nfs}(x_1, x_2, x_3, b_1, b_2) E_{nfs}(t_{nfs}), \quad (40)
\end{aligned}$$

$$\begin{aligned}
M^N = & \zeta \frac{32\sqrt{6}}{3} \pi C_F m_B^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \\
& \times \left\{ (r_2^2 - 1) [r_c \phi_{J/\psi}^T(x_2) - r_2 x_2 \phi_{J/\psi}^V(x_2)] \phi_V^T(x_3) + r_3 \left[r_c (1 + r_2^2) \phi_{J/\psi}^T(x_2) - r_2 [x_2 (1 + r_2^2) \right. \right. \\
& \left. \left. + x_3 (1 - r_2^2)] \phi_{J/\psi}^V(x_2) \right] \phi_V^v(x_3) \right\} h_{nfs}(x_1, x_2, x_3, b_1, b_2) E_{nfs}(t_{nfs}), \quad (41)
\end{aligned}$$

$$\begin{aligned}
M^T = & -\zeta \frac{64\sqrt{6}}{3} \pi C_F m_B^4 \int_0^1 dx_1 dx_2 dx_3 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \\
& \times \left\{ [r_c \phi_{J/\psi}^T(x_2) - r_2 x_2 \phi_{J/\psi}^V(x_2)] \phi_V^T(x_3) + r_3 \left[-r_c (1 + r_2^2) \phi_{J/\psi}^T(x_2) + r_2 [x_2 (1 + r_2^2) \right. \right. \\
& \left. \left. + x_3 (1 - r_2^2)] \phi_{J/\psi}^V(x_2) \right] \phi_V^a(x_3) \right\} h_{nfs}(x_1, x_2, x_3, b_1, b_2) E_{nfs}(t_{nfs}). \quad (42)
\end{aligned}$$

$r_c = m_c/m_B$ with m_c as the charm quark mass.




FIG. 3. (Color online) Typical Feynman diagrams contributing to $B \rightarrow J/\psi V$ decays known at Next-to-Leading Order level, in which the contributions will be combined into the Wilson coefficients associated with the factorizable contributions.

As was pointed out in Ref. [24], for these considered $B \rightarrow J/\psi V$ decays, the vertex corrections (see Fig. 3a-3d) will contribute at the current NLO level, in which their effects can be combined into the Wilson coefficients associated with the factorizable contributions [47]:

$$a_2^h = C_1 + \frac{C_2}{N_c} + \frac{\alpha_s}{4\pi} \frac{C_F}{N_c} C_2 \left(-18 + 12 \ln \frac{m_b}{\mu} + f_I^h \right), \quad (43)$$

$$a_{3,9}^h = C_{3,9} + \frac{C_{4,10}}{N_c} + \frac{\alpha_s}{4\pi} \frac{C_F}{N_c} C_{4,10} \left(-18 + 12 \ln \frac{m_b}{\mu} + f_I^h \right), \quad (44)$$

$$a_{5,7}^h = C_{5,7} + \frac{C_{6,8}}{N_c} + \frac{\alpha_s}{4\pi} \frac{C_F}{N_c} C_{6,8} \left(6 - 12 \ln \frac{m_b}{\mu} - f_I^h \right). \quad (45)$$

The function f_I^h is given as [47]

$$f_I^0 = f_I + g_I(1 - r_2^2), \quad f_I^\pm = f_I, \quad (46)$$

with

$$\begin{aligned} f_I = \frac{2\sqrt{2N_c}}{f_{J/\psi}} & \left[\int_0^1 dx_2 \phi_{J/\psi}^L(x_2) \left\{ \frac{2r_2^2 x_2}{1 - r_2^2(1 - x_2)} + (3 - 2x_2) \frac{\ln x_2}{1 - x_2} \right. \right. \\ & + \left(-\frac{3}{1 - r_2^2 x_2} + \frac{1}{1 - r_2^2(1 - x_2)} - \frac{2r_2^2 x_2}{[(1 - r_2^2(1 - x_2))^2]} \right) r_2^2 x_2 \ln(r_2^2 x_2) \\ & + \left. \left(3(1 - r_2^2) + 2r_2^2 x_2 + \frac{2r_2^4 x_2^2}{1 - r_2^2(1 - x_2)} \right) \frac{\ln(1 - r_2^2) - i\pi}{1 - r_2^2(1 - x_2)} \right\} \\ & + \int_0^1 dx_2 \phi_{J/\psi}^T(x_2) \left\{ -8x_2^2 \frac{\ln x_2}{1 - x_2} + \frac{8r_2^2 x_2^2 \ln(r_2^2 x_2)}{1 - r_2^2(1 - x_2)} - 8r_2^2 x_2^2 \frac{\ln(1 - r_2^2) - i\pi}{1 - r_2^2(1 - x_2)} \right\} \Big], \end{aligned} \quad (47)$$

$$\begin{aligned} g_I = \frac{2\sqrt{2N_c}}{f_{J/\psi}} & \left[\int_0^1 dx_2 \phi_{J/\psi}^L(x_2) \left\{ \frac{-4x_2}{(1 - r_2^2)(1 - x_2)} \ln x_2 + \frac{r_2^2 x_2}{[1 - r_2^2(1 - x_2)]^2} \ln(1 - r_2^2) \right. \right. \\ & + \left(\frac{1}{(1 - r_2^2 x_2)^2} - \frac{1}{[1 - r_2^2(1 - x_2)]^2} + \frac{2(1 + r_2^2 - 2r_2^2 x_2)}{(1 - r_2^2)(1 - r_2^2 x_2)^2} \right) r_2^2 x_2 \ln(r_2^2 x_2) - i\pi \frac{r_2^2 x_2}{[1 - r_2^2(1 - x_2)]^2} \Big\} \\ & + \int_0^1 dx_2 \phi_{J/\psi}^T(x_2) \left\{ \frac{8x_2^2}{(1 - r_2^2)(1 - x_2)} \ln x_2 - \frac{8x_2^2 r_2^2}{(1 - r_2^2)(1 - r_2^2 x_2)} \ln(r_2^2 x_2) \right\} \Big]. \end{aligned} \quad (48)$$

IV. NUMERICAL RESULTS AND DISCUSSIONS

In this section, we will present the theoretical predictions on the CP-averaged BRs, CP-averaged polarization fractions, and CP-violating asymmetries for those considered $B \rightarrow J/\psi V$ decay modes.

A. Input quantities

The masses (in units of GeV) and B meson lifetime (in ps) are taken from Particle Data Group [39]

$$\begin{aligned} m_W &= 80.41, \quad m_B = 5.28, \quad m_{B_s} = 5.37, \quad m_b = 4.8; \\ \tau_{B_u} &= 1.641, \quad \tau_{B_d} = 1.519, \quad \tau_{B_s} = 1.497, \quad m_{J/\psi} = 3.097. \end{aligned} \quad (49)$$

For the CKM matrix elements, we adopt the Wolfenstein parametrization up to $\mathcal{O}(\lambda^5)$ and the updated parameters $A = 0.811$, $\lambda = 0.22535$, $\bar{\rho} = 0.131_{-0.013}^{+0.026}$, and $\bar{\eta} = 0.345_{-0.014}^{+0.013}$ [39].

B. CP-averaged Branching Ratios, Polarization Fractions, and Relative Phases

In this subsection, we will analyze the CP-averaged BRs of the considered $B \rightarrow J/\psi V$ decays in the pQCD approach. For $B \rightarrow J/\psi V$ decays, the decay rate can be written explicitly as,

$$\Gamma = \frac{G_F^2 |\mathbf{P}_c|}{16\pi m_B^2} \sum_{\sigma=L,N,T} \mathcal{A}_\sigma^\dagger \mathcal{A}_\sigma, \quad (50)$$

where $|\mathbf{P}_c|$ is the three-momentum of the final vector meson. Based on the helicity amplitudes (35), we have defined the transversity amplitudes as

$$\mathcal{A}_L = \mathcal{M}_L, \quad \mathcal{A}_\parallel = \sqrt{2}\mathcal{M}_N, \quad \mathcal{A}_\perp = r_2 r_3 \sqrt{2(r^2 - 1)} \mathcal{M}_T, \quad (51)$$

for the longitudinal, parallel, and perpendicular polarizations, respectively, with the ratio $r = P_2 \cdot P_3 / (m_B^2 r_2 r_3)$. Using the decay amplitudes obtained in last section, it is straightforward to calculate the CP-averaged BRs with uncertainties

as displayed in Eqs. (55)-(58). The dominant errors are induced by the shape parameters $\omega_B = 0.40 \pm 0.04$ ($\omega_B = 0.50 \pm 0.05$) GeV for $B(B_s)$ meson, the uncertainties of the decay constants f_M , the Gegenbauer moments a_i for the light vector mesons, the charm quark mass $m_c = (1.50 \pm 0.15)$ GeV, and CKM matrix elements $(\bar{\rho}, \bar{\eta})$, respectively. It is worthwhile to stress that the variation of the CKM parameters has small effects on the CP-averaged BRs and polarization fractions and thus will be neglected in the numerical results as shown in Eqs. (55)-(58) and Eqs. (85)-(93).

The theoretical predictions in the pQCD approach for the CP-averaged BRs of the decays under consideration within errors are given as follows:

- for $\bar{b} \rightarrow \bar{s}$ decay channels,

$$\text{BR}(B_d \rightarrow J/\psi K^{*0}) = 1.23^{+0.30}_{-0.23}(\omega_B)^{+0.16}_{-0.16}(f_M)^{+0.10}_{-0.09}(a_i)^{+0.22}_{-0.20}(m_c) \left[1.23^{+0.42}_{-0.36} \right] \times 10^{-3}, \quad (52)$$

$$\begin{aligned} \text{BR}(B_u \rightarrow J/\psi K^{*+}) &= \frac{\tau_{B_u}}{\tau_{B_d}} \cdot \text{BR}(B_d \rightarrow J/\psi K^{*0}) \\ &= 1.33^{+0.32}_{-0.25}(\omega_B)^{+0.17}_{-0.17}(f_M)^{+0.11}_{-0.10}(a_i)^{+0.24}_{-0.22}(m_c) \left[1.33^{+0.45}_{-0.39} \right] \times 10^{-3}, \end{aligned} \quad (53)$$

$$\text{BR}(B_s \rightarrow J/\psi \phi) = 1.02^{+0.29}_{-0.22}(\omega_B)^{+0.14}_{-0.12}(f_M)^{+0.06}_{-0.06}(a_i)^{+0.16}_{-0.15}(m_c) \left[1.02^{+0.36}_{-0.30} \right] \times 10^{-3}; \quad (54)$$

- for $\bar{b} \rightarrow \bar{d}$ decay channels,

$$\text{BR}(B_d \rightarrow J/\psi \rho^0) = 2.7^{+0.7}_{-0.5}(\omega_B)^{+0.5}_{-0.3}(f_M)^{+0.2}_{-0.1}(a_i)^{+0.5}_{-0.4}(m_c) \left[2.7^{+1.0}_{-0.7} \right] \times 10^{-5}, \quad (55)$$

$$\begin{aligned} \text{BR}(B_u \rightarrow J/\psi \rho^+) &= 2 \cdot \frac{\tau_{B_u}}{\tau_{B_d}} \cdot \text{BR}(B_d \rightarrow J/\psi \rho^0) \\ &= 5.8^{+1.5}_{-1.1}(\omega_B)^{+1.1}_{-0.6}(f_M)^{+0.4}_{-0.2}(a_i)^{+1.1}_{-0.9}(m_c) \left[5.8^{+2.2}_{-1.6} \right] \times 10^{-5}, \end{aligned} \quad (56)$$

$$\text{BR}(B_d \rightarrow J/\psi \omega) = 2.3^{+0.5}_{-0.4}(\omega_B)^{+0.2}_{-0.3}(f_M)^{+0.1}_{-0.1}(a_i)^{+0.3}_{-0.4}(m_c) \left[2.3^{+0.6}_{-0.7} \right] \times 10^{-5}; \quad (57)$$

$$\text{BR}(B_s \rightarrow J/\psi \bar{K}^{*0}) = 4.1^{+1.1}_{-0.8}(\omega_B)^{+0.5}_{-0.5}(f_M)^{+0.3}_{-0.3}(a_i)^{+0.7}_{-0.7}(m_c) \left[4.1^{+1.4}_{-1.2} \right] \times 10^{-5}. \quad (58)$$

The values given in the square parentheses are obtained by adding various errors in quadrature.

Experimentally, the available measurements of the branching ratios for the considered decay modes are as follows [39, 48, 49],

$$\text{BR}^{\text{ex.}}(B_d \rightarrow J/\psi K^{*0}) = (1.34 \pm 0.06) \times 10^{-3}, \quad (59)$$

$$\text{BR}^{\text{ex.}}(B_u \rightarrow J/\psi K^{*+}) = (1.43 \pm 0.08) \times 10^{-3}, \quad (60)$$

$$\text{BR}^{\text{ex.}}(B_s \rightarrow J/\psi \phi) = (1.09^{+0.28}_{-0.23}) \times 10^{-3}, \quad (61)$$

$$\text{BR}^{\text{ex.}}(B_d \rightarrow J/\psi \rho^0) = (2.7 \pm 0.4) \times 10^{-5}, \quad (62)$$

$$\text{BR}^{\text{ex.}}(B_u \rightarrow J/\psi \rho^+) = (5.0 \pm 0.8) \times 10^{-5}, \quad (63)$$

$$\text{BR}^{\text{ex.}}(B_d \rightarrow J/\psi \omega) = (2.4 \pm 0.7) \times 10^{-5}, \quad (64)$$

$$\text{BR}^{\text{ex.}}(B_s \rightarrow J/\psi \bar{K}^{*0}) = (4.4 \pm 0.9) \times 10^{-5}. \quad (65)$$

It is necessary to point out that the LHCb results for B_s decays correspond to the time-integrated quantities, while theory predictions made in the above refer to the branching fractions at $t = 0$ [50]. These two quantities can differ by 10% by definition. The precision of the measurements for the $\bar{b} \rightarrow \bar{d}$ modes can be improved once more data samples are collected in future.

From the above results, one can see that most of our NLO pQCD predictions on BRs agree well with the existing experimental measurements within the uncertainties. Meanwhile, one can observe that the decay rates for the $\bar{b} \rightarrow \bar{s}$ transition processes, i.e., $B_{u/d} \rightarrow J/\psi K^*$ and $B_s \rightarrow J/\psi \phi$, are generally much larger than those for the $\bar{b} \rightarrow \bar{d}$ transition ones, i.e., $B_{u/d} \rightarrow J/\psi \rho$, $B_d \rightarrow J/\psi \omega$, and $B_s \rightarrow J/\psi \bar{K}^{*0}$. This is due to the CKM hierarchy for two kinds of process: the CKM factors $V_{cb}V_{cs}$ in $b \rightarrow s$ are about 4 times larger than the $V_{cb}V_{cd}$ for $b \rightarrow d$ process. The remanent differences arise from the SU(3) symmetry breaking effects in the hadronic parameters, such as decay constants, mesonic masses, distribution amplitudes, etc.. This analysis may provide theoretical ground for the use of SU(3) symmetry in B and B_s decays to hunt for a scalar glueball in Ref. [51, 52]

Here, we will also explore some interesting relations among those considered decay channels,

- The ratio $R_{\omega/\rho}$ between the branching ratios of $B_d \rightarrow J/\psi\omega$ and $B_d \rightarrow J/\psi\rho^0$ decays can be given as,

$$R_{\omega/\rho}^{\text{th.}} \equiv \frac{\text{BR}(B_d \rightarrow J/\psi\omega)}{\text{BR}(B_d \rightarrow J/\psi\rho^0)} \approx 0.85_{-0.03}^{+0.01}(\omega_B)_{-0.04}^{+0.00}(f_M)_{-0.02}^{+0.00}(a_i)_{-0.04}^{+0.00}(m_c)[0.85_{-0.07}^{+0.01}], \quad (66)$$

where most theoretical errors have been cancelled out in the ratio. This prediction is in good consistency with the LHCb measurement [49] within errors,

$$\frac{\text{BR}(B_d \rightarrow J/\psi\omega)}{\text{BR}(B_d \rightarrow J/\psi\rho^0)} = 0.89_{-0.23}^{+0.20}. \quad (67)$$

Theoretically, both decay modes embrace the same transition at the quark level, which means the involved QCD behavior is similar. The differences between their CP-averaged branching ratios come from their different decay constants and masses.

- The ratio of BRs of $B_s \rightarrow J/\psi\bar{K}^{*0}$ and $B_d \rightarrow J/\psi K^{*0}$ decays is predicted as

$$R_{s/d}^{\text{th.}} \equiv \frac{\text{BR}(B_s \rightarrow J/\psi\bar{K}^{*0})}{\text{BR}(B_d \rightarrow J/\psi K^{*0})} \approx 0.0333_{-0.0007}^{+0.0011}(\omega_B)_{-0.0004}^{+0.0001}(f_M)_{-0.0021}^{+0.0021}(a_i)_{-0.0002}^{+0.0001}(m_c)[0.0333_{-0.0022}^{+0.0024}], \quad (68)$$

which agrees well with that shown in Ref. [48]

$$\frac{\text{BR}(B_s \rightarrow J/\psi\bar{K}^{*0})}{\text{BR}(B_d \rightarrow J/\psi K^{*0})} = 0.0343_{-0.006}^{+0.006}, \quad (69)$$

and also with the CDF results [53]

$$\frac{\text{BR}(B_s \rightarrow J/\psi\bar{K}^{*0})}{\text{BR}(B_d \rightarrow J/\psi K^{*0})} = 0.062 \pm 0.028, \quad (70)$$

where the $\mathcal{BR}(B_s \rightarrow J/\psi K^{*0})$ measured by CDF Collaboration is $[8.3 \pm 3.8] \times 10^{-5}$ [53].

- The ratio of the branching ratios of two B_s decay channels can be predicted as,

$$R_{K^*/\phi}^{\text{th.}} \equiv \frac{\text{BR}(B_s \rightarrow J/\psi\bar{K}^{*0})}{\text{BR}(B_s \rightarrow J/\psi\phi)} \approx 0.040_{-0.000}^{+0.001}(\omega_B)_{-0.000}^{+0.001}(f_M)_{-0.001}^{+0.001}(a_i)_{-0.000}^{+0.001}(m_c)[0.040_{-0.001}^{+0.002}], \quad (71)$$

which is also in good agreement with the entry derived from the available data [39, 48],

$$\frac{\text{BR}(B_s \rightarrow J/\psi\bar{K}^{*0})}{\text{BR}(B_s \rightarrow J/\psi\phi)} \approx 0.040_{-0.0119}^{+0.0133}. \quad (72)$$

- In those two $\bar{b} \rightarrow \bar{s}$ transition modes, the theoretical ratio of $\text{BR}(B_d \rightarrow J/\psi K^{*0})$ to $\text{BR}(B_s \rightarrow J/\psi\phi)$ is

$$\begin{aligned} R_{d/s}^{\text{th.}} &\equiv \frac{\text{BR}(B_d \rightarrow J/\psi K^{*0})}{\text{BR}(B_s \rightarrow J/\psi\phi)} \approx 1.21_{-0.04}^{+0.03}(\omega_B)_{-0.02}^{+0.00}(f_M)_{-0.02}^{+0.02}(a_i)_{-0.03}^{+0.01}(m_c)[1.21_{-0.06}^{+0.04}], \\ R_{s/d}^{\text{th.}} &\equiv \frac{\text{BR}(B_s \rightarrow J/\psi\phi)}{\text{BR}(B_d \rightarrow J/\psi K^{*0})} \approx 0.83_{-0.02}^{+0.03}(\omega_B)_{-0.00}^{+0.01}(f_M)_{-0.02}^{+0.01}(a_i)_{-0.02}^{+0.01}(m_c)[0.83_{-0.03}^{+0.03}]. \end{aligned} \quad (73)$$

which is consistent well with the existing data [39],

$$\frac{\text{BR}(B_d \rightarrow J/\psi K^{*0})}{\text{BR}(B_s \rightarrow J/\psi\phi)} \approx 1.22_{-0.27}^{+0.32}. \quad (74)$$

We will also study the polarization fractions for $B \rightarrow J/\psi V$ decay modes:

$$f_{L(\parallel, \perp)} \equiv \frac{|\mathcal{A}_{L(\parallel, \perp)}|^2}{|\mathcal{A}_L|^2 + |\mathcal{A}_\parallel|^2 + |\mathcal{A}_\perp|^2}. \quad (75)$$

By definition these fractions satisfy the relation:

$$f_L + f_\parallel + f_\perp = 1. \quad (76)$$

The CP-averaged polarization fractions are predicted as follows

- for $B_{u/d} \rightarrow J/\psi K^*$ decays,

$$f_L(B_{u/d} \rightarrow J/\psi K^*) = 48.7^{+0.8}_{-0.7}(\omega_B)^{+1.6}_{-1.5}(f_M)^{+3.8}_{-3.8}(a_i)^{+0.7}_{-1.2}(m_c) \left[48.7^{+4.3}_{-4.3} \right] \%, \quad (77)$$

$$f_\parallel(B_{u/d} \rightarrow J/\psi K^*) = 30.3^{+0.4}_{-0.5}(\omega_B)^{+0.9}_{-0.9}(f_M)^{+2.2}_{-2.1}(a_i)^{+1.2}_{-0.9}(m_c) \left[30.3^{+2.7}_{-2.5} \right] \%, \quad (78)$$

$$f_\perp(B_{u/d} \rightarrow J/\psi K^*) = 21.0^{+0.2}_{-0.4}(\omega_B)^{+0.8}_{-0.9}(f_M)^{+1.6}_{-1.8}(a_i)^{+0.2}_{-0.0}(m_c) \left[21.0^{+1.8}_{-2.1} \right] \%. \quad (79)$$

Compared to the LHCb measurement [54]:

$$f_\parallel(B_{u/d} \rightarrow J/\psi K^*) = (22.7 \pm 1.2)\%, \quad f_\perp(B_{u/d} \rightarrow J/\psi K^*) = (20.1 \pm 0.9)\%, \quad (80)$$

our result for f_\perp is in agreement while the one for f_\parallel is a bit higher than the data by about 2σ .

- for $B_s \rightarrow J/\psi \phi$ mode,

$$f_L(B_s \rightarrow J/\psi \phi) = 50.7^{+0.8}_{-0.8}(\omega_B)^{+1.5}_{-1.4}(f_M)^{+3.1}_{-3.0}(a_i)^{+0.5}_{-1.1}(m_c) \left[50.7^{+3.6}_{-3.6} \right] \%, \quad (81)$$

$$f_\parallel(B_s \rightarrow J/\psi \phi) = 29.8^{+0.6}_{-0.4}(\omega_B)^{+0.8}_{-0.7}(f_M)^{+1.7}_{-1.7}(a_i)^{+1.2}_{-0.8}(m_c) \left[29.8^{+2.3}_{-2.0} \right] \%, \quad (82)$$

$$f_\perp(B_s \rightarrow J/\psi \phi) = 19.4^{+0.4}_{-0.3}(\omega_B)^{+0.7}_{-0.6}(f_M)^{+1.4}_{-1.3}(a_i)^{+0.4}_{-0.0}(m_c) \left[19.4^{+1.7}_{-1.5} \right] \%, \quad (83)$$

which are also in agreement with the recent measurement by the LHCb Collaboration [2]

$$f_L = [49.7 \pm 3.3]\%, \quad f_\perp = [23.7 \pm 1.9]\%. \quad (84)$$

- for $B_{u/d} \rightarrow J/\psi \rho$ decays,

$$f_L(B_{u/d} \rightarrow J/\psi \rho) = 51.8^{+0.9}_{-0.7}(\omega_B)^{+1.7}_{-1.6}(f_M)^{+2.8}_{-2.6}(a_i)^{+0.0}_{-0.4}(m_c) \left[51.8^{+3.4}_{-3.2} \right] \%, \quad (85)$$

$$f_\parallel(B_{u/d} \rightarrow J/\psi \rho) = 27.7^{+0.5}_{-0.4}(\omega_B)^{+0.9}_{-0.9}(f_M)^{+1.6}_{-1.4}(a_i)^{+0.8}_{-0.3}(m_c) \left[27.7^{+2.1}_{-1.7} \right] \%, \quad (86)$$

$$f_\perp(B_{u/d} \rightarrow J/\psi \rho) = 20.4^{+0.3}_{-0.3}(\omega_B)^{+0.7}_{-0.8}(f_M)^{+1.3}_{-1.2}(a_i)^{+0.5}_{-0.2}(m_c) \left[20.4^{+1.6}_{-1.5} \right] \%. \quad (87)$$

- for $B_d \rightarrow J/\psi \omega$ mode,

$$f_L(B_d \rightarrow J/\psi \omega) = 53.5^{+0.9}_{-0.8}(\omega_B)^{+2.2}_{-2.0}(f_M)^{+2.6}_{-2.5}(a_i)^{+0.0}_{-0.5}(m_c) \left[53.5^{+3.5}_{-3.3} \right] \%, \quad (88)$$

$$f_\parallel(B_d \rightarrow J/\psi \omega) = 26.9^{+0.6}_{-0.5}(\omega_B)^{+1.1}_{-1.0}(f_M)^{+1.5}_{-1.4}(a_i)^{+0.8}_{-0.4}(m_c) \left[26.9^{+2.1}_{-1.8} \right] \%, \quad (89)$$

$$f_\perp(B_d \rightarrow J/\psi \omega) = 19.5^{+0.3}_{-0.3}(\omega_B)^{+1.0}_{-1.0}(f_M)^{+1.3}_{-1.1}(a_i)^{+0.5}_{-0.2}(m_c) \left[19.5^{+1.7}_{-1.5} \right] \%. \quad (90)$$

- for $B_s \rightarrow J/\psi \bar{K}^*$ mode,

$$f_L(B_s \rightarrow J/\psi \bar{K}^{*0}) = 50.9_{-1.0}^{+0.9}(\omega_B)_{-1.6}^{+1.8}(f_M)_{-3.9}^{+3.7}(a_i)_{-1.2}^{+0.4}(m_c) \left[50.9_{-4.5}^{+4.2} \right] \% , \quad (91)$$

$$f_{\parallel}(B_s \rightarrow J/\psi \bar{K}^{*0}) = 29.1_{-0.6}^{+0.5}(\omega_B)_{-1.0}^{+0.7}(f_M)_{-2.2}^{+2.0}(a_i)_{-0.9}^{+1.1}(m_c) \left[29.1_{-2.6}^{+2.4} \right] \% , \quad (92)$$

$$f_{\perp}(B_s \rightarrow J/\psi \bar{K}^{*0}) = 20.1_{-0.3}^{+0.4}(\omega_B)_{-0.9}^{+0.8}(f_M)_{-1.7}^{+1.7}(a_i)_{-0.0}^{+0.4}(m_c) \left[20.1_{-1.9}^{+2.0} \right] \% . \quad (93)$$

The measurement of polarization fractions for $B_s \rightarrow J/\psi \bar{K}^{*0}$ decay by the LHCb Collaboration [48] is

$$f_L = [50 \pm 8]\% , \quad f_{\parallel} = [19_{-8}^{+10}]\% . \quad (94)$$

In terms of the transversity amplitudes, we can define their relative phases ϕ_{\parallel} and ϕ_{\perp}

$$\phi_{\parallel} \equiv \arg \frac{\mathcal{A}_{\parallel}}{\mathcal{A}_L} \quad \text{and} \quad \phi_{\perp} \equiv \arg \frac{\mathcal{A}_{\perp}}{\mathcal{A}_L} . \quad (95)$$

We then predict the CP-averaged relative phases for the considered $B \rightarrow J/\psi V$ decays as

- for $B_{u/d} \rightarrow J/\psi K^*$ decays,

$$\phi_{\parallel} = 2.65_{-0.08}^{+0.10} \quad \text{rad} , \quad \phi_{\perp} = 2.59_{-0.11}^{+0.08} \quad \text{rad} ; \quad (96)$$

- for $B_s \rightarrow J/\psi \phi$ mode,

$$\phi_{\parallel} = 2.74_{-0.07}^{+0.07} \quad \text{rad} , \quad \phi_{\perp} = 2.65_{-0.08}^{+0.08} \quad \text{rad} ; \quad (97)$$

- for $B_{u/d} \rightarrow J/\psi \rho$ decays,

$$\phi_{\parallel} = 2.58_{-0.08}^{+0.10} \quad \text{rad} , \quad \phi_{\perp} = 2.52_{-0.11}^{+0.09} \quad \text{rad} ; \quad (98)$$

- for $B_d \rightarrow J/\psi \omega$ mode,

$$\phi_{\parallel} = 2.58_{-0.11}^{+0.09} \quad \text{rad} , \quad \phi_{\perp} = 2.51_{-0.12}^{+0.10} \quad \text{rad} ; \quad (99)$$

- for $B_s \rightarrow J/\psi \bar{K}^{*0}$ mode

$$\phi_{\parallel} = 2.67_{-0.08}^{+0.09} \quad \text{rad} , \quad \phi_{\perp} = 2.59_{-0.11}^{+0.10} \quad \text{rad} ; \quad (100)$$

where various errors from the input parameters have been added in quadrature.

C. CP Asymmetries

As for the direct CP-violating asymmetry in these considered modes, considering the involved three polarizations, whose definitions are as follows,

$$A_{CP}^{\text{dir}} \equiv \frac{\bar{\Gamma} - \Gamma}{\bar{\Gamma} + \Gamma} = \frac{|\bar{\mathcal{A}}(\bar{B} \rightarrow \bar{f}_{\text{final}})|^2 - |\mathcal{A}(B \rightarrow f_{\text{final}})|^2}{|\bar{\mathcal{A}}(\bar{B} \rightarrow \bar{f}_{\text{final}})|^2 + |\mathcal{A}(B \rightarrow f_{\text{final}})|^2} , \quad (101)$$

where Γ and \mathcal{A} denote the decay rate and decay amplitude of $B \rightarrow J/\psi V$ decays, respectively, and $\bar{\Gamma}$ and $\bar{\mathcal{A}}$ are the charge conjugation one correspondingly. It is conventional to combine the three polarization fractions in Eq. (75) with those of its CP-conjugate \bar{B} decay, and to quote the six resulting observables corresponding to transversity amplitudes as direct induced CP asymmetries [55]. The direct CP asymmetries in transversity basis can be defined as

$$A_{CP}^{\text{dir}, \alpha} = \frac{\bar{f}_{\alpha} - f_{\alpha}}{\bar{f}_{\alpha} + f_{\alpha}} , \quad (\alpha = L, \parallel, \perp) , \quad (102)$$

where the definition of \bar{f} is same as that in Eq.(75) but for the corresponding \bar{B} decay.

The direct CP-violating asymmetries for those $B \rightarrow J/\psi V$ decays are

$$A_{CP}^{\text{dir}}(B \rightarrow J/\psi K^*) = 3.19_{-0.32}^{+0.42}(\omega_B)_{-0.09}^{+0.12}(f_M)_{-0.12}^{+0.14}(a_i)_{-0.12}^{+0.25}(m_c)_{-0.12}^{+0.13}(\text{CKM}) \times 10^{-4}, \quad (103)$$

$$A_{CP}^{\text{dir}}(B_s \rightarrow J/\psi \phi) = 2.66_{-0.37}^{+0.45}(\omega_B)_{-0.07}^{+0.10}(f_M)_{-0.08}^{+0.11}(a_i)_{-0.12}^{+0.14}(m_c)_{-0.10}^{+0.10}(\text{CKM}) \times 10^{-4}, \quad (104)$$

$$A_{CP}^{\text{dir}}(B \rightarrow J/\psi \rho) = -5.15_{-0.70}^{+0.57}(\omega_B)_{-0.16}^{+0.18}(f_M)_{-0.24}^{+0.22}(a_i)_{-0.30}^{+0.05}(m_c)_{-0.19}^{+0.21}(\text{CKM}) \times 10^{-3}, \quad (105)$$

$$A_{CP}^{\text{dir}}(B \rightarrow J/\psi \omega) = -5.08_{-0.68}^{+0.55}(\omega_B)_{-0.21}^{+0.19}(f_M)_{-0.25}^{+0.22}(a_i)_{-0.27}^{+0.00}(m_c)_{-0.18}^{+0.20}(\text{CKM}) \times 10^{-3}, \quad (106)$$

$$A_{CP}^{\text{dir}}(B_s \rightarrow J/\psi \bar{K}^{*0}) = -4.15_{-0.70}^{+0.74}(\omega_B)_{-0.14}^{+0.18}(f_M)_{-0.22}^{+0.27}(a_i)_{-0.20}^{+0.27}(m_c)_{-0.16}^{+0.16}(\text{CKM}) \times 10^{-3}. \quad (107)$$

We also give the results for direct CP asymmetries corresponding to three polarizations,

- for $B_{u/d} \rightarrow J/\psi K^*$ decays,

$$A_{CP}^{\text{dir},L} = 3.04_{-0.54}^{+0.66} \times 10^{-4}, \quad A_{CP}^{\text{dir},||} = 3.39_{-0.37}^{+0.47} \times 10^{-4}, \quad A_{CP}^{\text{dir},\perp} = 3.26_{-0.44}^{+0.64} \times 10^{-4}; \quad (108)$$

- for $B_s \rightarrow J/\psi \phi$ mode,

$$A_{CP}^{\text{dir},L} = 2.69_{-0.49}^{+0.62} \times 10^{-4}, \quad A_{CP}^{\text{dir},||} = 2.71_{-0.40}^{+0.39} \times 10^{-4}, \quad A_{CP}^{\text{dir},\perp} = 2.52_{-0.40}^{+0.55} \times 10^{-4}; \quad (109)$$

- for $B_{u/d} \rightarrow J/\psi \rho$ decays,

$$A_{CP}^{\text{dir},L} = -4.15_{-0.96}^{+0.78} \times 10^{-3}, \quad A_{CP}^{\text{dir},||} = -6.32_{-0.81}^{+0.72} \times 10^{-3}, \quad A_{CP}^{\text{dir},\perp} = -6.09_{-1.09}^{+0.84} \times 10^{-3}; \quad (110)$$

- for $B_d \rightarrow J/\psi \omega$ mode,

$$A_{CP}^{\text{dir},L} = -3.84_{-0.96}^{+0.81} \times 10^{-3}, \quad A_{CP}^{\text{dir},||} = -6.63_{-0.90}^{+0.74} \times 10^{-3}, \quad A_{CP}^{\text{dir},\perp} = -6.36_{-1.16}^{+0.82} \times 10^{-3}; \quad (111)$$

- for $B_s \rightarrow J/\psi \bar{K}^{*0}$ mode

$$A_{CP}^{\text{dir},L} = -3.83_{-0.90}^{+1.14} \times 10^{-3}, \quad A_{CP}^{\text{dir},||} = -4.63_{-0.71}^{+0.80} \times 10^{-3}, \quad A_{CP}^{\text{dir},\perp} = -4.27_{-1.05}^{+0.76} \times 10^{-3}; \quad (112)$$

in which various errors have been again added in quadrature.

D. Impact of Penguin Contamination

Based on the encouraging agreements of our theoretical calculations with the available data on branching ratios and polarisations, we now study the penguin impacts on the mixing-phase in $B_s \rightarrow J/\psi \phi$ decay:

$$\phi_s^{\text{eff}} = -\arg \left[\frac{q}{p} \frac{\bar{\mathcal{A}}_f^\alpha}{\mathcal{A}_f^\alpha} \right] = \phi_s + \Delta\phi_s, \quad (113)$$

where α denotes three polarization configurations L , $||$, and \perp , and $\mathcal{A}_f^\alpha(\bar{\mathcal{A}}_f^\alpha)$ stands for the decay amplitude of $B_s \rightarrow J/\psi \phi(\bar{B}_s \rightarrow J/\psi \phi)$, which can be decomposed into [18]

$$\mathcal{A}_f^\alpha(B_s \rightarrow J/\psi \phi) = V_{cb}^* V_{cs}(T_c^\alpha + P_c^\alpha + P_t^\alpha) + V_{ub}^* V_{us}(P_u^\alpha + P_t^\alpha), \quad (114)$$

Here, the unitarity relation $V_{tb}^* V_{ts} = -V_{cb}^* V_{cs} - V_{ub}^* V_{us}$ for the CKM matrix elements has been used. The dominant tree amplitude T_c^α contributes to the branching ratio of $B_s \rightarrow J/\psi \phi$ decay, and P_c^α , P_u^α , and P_t^α are the penguin pollution in the decay. In this work, we have calculated the T_c^α and P_t^α while the u-quark and c-quark penguin are not included. Then the charge conjugation amplitude for $B_s \rightarrow J/\psi \phi$ decay is

$$\bar{\mathcal{A}}_f^\alpha(\bar{B}_s \rightarrow J/\psi \phi) = V_{cb} V_{cs}^*(T_c^\alpha + P_c^\alpha + P_t^\alpha) + V_{ub} V_{us}^*(P_u^\alpha + P_t^\alpha). \quad (115)$$

TABLE III. Factorization amplitudes \mathcal{A}_L , $\mathcal{A}_{||}$ and \mathcal{A}_\perp (without CKM factors) for the hadronic $B \rightarrow J/\psi V$ decays in the pQCD approach, where only the central values are quoted. The results are given in units of GeV^3 .

Decay modes		Tree Operators	Penguin Operators ($\times 10^{-2}$)
$B_{u/d} \rightarrow J/\psi K^*$	L	$-0.260 + i0.959$	$-1.608 + i2.571$
	$ $	$-0.175 - i0.775$	$0.325 - i2.020$
	\perp	$-0.188 - i0.635$	$0.157 - i1.617$
$B_s \rightarrow J/\psi \phi$	L	$-0.158 + i0.933$	$-1.147 + i2.376$
	$ $	$-0.167 - i0.713$	$0.158 - i1.784$
	\perp	$-0.185 - i0.563$	$0.010 - i1.378$
$B_{u/d} \rightarrow J/\psi \rho$	L	$0.210 - i0.620$	$1.005 - i1.665$
	$ $	$0.113 + i0.472$	$-0.179 + i1.231$
	\perp	$0.123 + i0.398$	$-0.085 + i1.014$
$B_d \rightarrow J/\psi \omega$	L	$-0.203 + i0.573$	$-0.928 + i1.539$
	$ $	$-0.100 - i0.426$	$0.186 - i1.112$
	\perp	$-0.109 - i0.356$	$0.093 - i0.910$
$B_s \rightarrow J/\psi \bar{K}^*$	L	$-0.211 + i0.783$	$-1.026 + i1.975$
	$ $	$-0.125 - i0.606$	$0.130 - i1.506$
	\perp	$-0.144 - i0.495$	$0.010 - i1.208$

For simplicity, one can introduce the ratio

$$a_f e^{i\delta_f + i\gamma} \equiv \frac{V_{ub}^* V_{us} (P_u^\alpha + P_t^\alpha)}{V_{cb}^* V_{cs} (T_c^\alpha + P_c^\alpha + P_t^\alpha)}, \quad (116)$$

where γ is the weak phase of V_{ub}^* . This leads to the ratio of amplitudes

$$\frac{\bar{\mathcal{A}}_f^\alpha}{\mathcal{A}_f^\alpha} = \frac{1 + a_f e^{i\delta_f - i\gamma}}{1 + a_f e^{i\delta_f + i\gamma}} \simeq 1 - 2ia_f \cos \delta_f \sin \gamma, \quad (117)$$

and the phase shift:

$$\Delta\phi_s \simeq \arcsin(2a_f \cos \delta_f \sin \gamma). \quad (118)$$

To proceed, we present the factorisation amplitudes (factoring out the CKM matrix elements) in Table. III. Using the $B_s \rightarrow J/\psi \phi$ channel as the example, if we include the LO and vertex corrections, we find the results:

$$\begin{aligned} \Delta\phi_s(L) &\approx 0.72 \times 10^{-3}; \\ \Delta\phi_s(||) &\approx 0.71 \times 10^{-3}; \\ \Delta\phi_s(\perp) &\approx 0.69 \times 10^{-3}. \end{aligned} \quad (119)$$

The vertex corrections give the dominant contribution. The addition of hard-scattering diagrams gives

$$\begin{aligned} \Delta\phi_s(L) &\approx 0.96 \times 10^{-3}; \\ \Delta\phi_s(||) &\approx 0.84 \times 10^{-3}; \\ \Delta\phi_s(\perp) &\approx 0.80 \times 10^{-3}. \end{aligned} \quad (120)$$

By including various parametric errors, we therefore give the quantity $\Delta\phi_s$ from our pQCD calculation as follows,

$$\begin{aligned} \Delta\phi_s(L) &\approx 0.96_{-0.03}^{+0.04}(\omega_B)_{-0.00}^{+0.02}(f_M)_{-0.01}^{+0.01}(a_i)_{-0.02}^{+0.03}(m_c)_{-0.03}^{+0.04}(\text{CKM}) [0.96_{-0.05}^{+0.07(+0.05)}] \times 10^{-3}; \\ \Delta\phi_s(||) &\approx 0.84_{-0.02}^{+0.02}(\omega_B)_{-0.00}^{+0.00}(f_M)_{-0.01}^{+0.01}(a_i)_{-0.01}^{+0.00}(m_c)_{-0.04}^{+0.03}(\text{CKM}) [0.84_{-0.05}^{+0.04(+0.02)}] \times 10^{-3}; \\ \Delta\phi_s(\perp) &\approx 0.80_{-0.01}^{+0.01}(\omega_B)_{-0.00}^{+0.00}(f_M)_{-0.01}^{+0.01}(a_i)_{-0.02}^{+0.00}(m_c)_{-0.03}^{+0.03}(\text{CKM}) [0.80_{-0.04}^{+0.03(+0.01)}] \times 10^{-3}. \end{aligned} \quad (121)$$

The values as given in the parentheses are only from the variation of hadronic parameters and have been added in quadrature. The deviation $\Delta\phi_s$ is found to be of $\mathcal{O}(10^{-3})$ in the standard model with the pQCD approach by taking into account the known NLO contributions, specifically, vertex corrections. This finding can be examined in the ongoing LHCb experiment and under-designed Super B factory and may provide an important standard model reference for verifying the existing new physics from the $B_s \rightarrow J/\psi\phi$ data.

Due to the large amount of data sample, the LHCb experiment is able to perform an analysis of the angular distribution of $B_s \rightarrow J/\psi\phi$. So the coefficients given in Table I could be viewed as experimental observables. Our predictions for the P-wave coefficients (in units of 10^{-3}) are as follows:

f_k	Δa_k	Δb_k	Δc_k	Δd_k
$c_K^2 s_l^2$	—	0.6	-0.3	1.1
$\frac{s_K^2(1-c_\phi^2 c_l^2)}{2}$	—	0.7	0.9	-1.1
$\frac{s_K^2(1-s_\phi^2 c_l^2)}{2}$	—	-0.7	1.2	1.3
$s_K^2 s_l^2 s_\phi c_\phi$	-0.1	1.1	6.4	1.1
$\sqrt{2} s_K c_K s_l c_l c_\phi$	-1.0	1.0	0.3	0.03
$\sqrt{2} s_K c_K s_l c_l s_\phi$	1.1	-0.02	-44	-1.4

The first three coefficients a_k have been chosen as 1, and thus it is not meaningful to discuss penguin effects to them. We find most of the results for the other coefficients are of order 10^{-3} . The other four coefficients given in Table I are the S-wave and the interference terms. The study of them requests the calculation of $B_s \rightarrow J/\psi(K^+K^-)_S$, presumably dominated by the $f_0(980)$, and thus is left out in this work.

In our calculation, only perturbative expansions at NLO in α_s and at leading power in $1/m_b$ are taken into account. The robustness of this calculation may be challenged by higher order QCD and power corrections, and also by long distance effects. The former can be in principle improved when higher order calculation is available. These include various sources. According to the discussions in Ref. [26], the penguin correction from the u -quark loop P_u^α is of $\mathcal{O}(\alpha_s^2)$, which needs a two-loop calculation for the corresponding amplitude. While, the correction from c -quark loop has the same phase with leading-order diagram and thus will not contribute to the $\Delta\phi_s$.

The long distance effects are often parametrized as the rescattering mechanism, and for instance see Ref. [56–58]. To have a sizeable contribution, the intermediate states shall have a large production rates and preferable overlap with $J/\psi\phi$ final states. The states that satisfy these constraints include $D_s^\pm \bar{D}_s^\mp$ (with $\mathcal{BR}(B_s \rightarrow D_s^\pm \bar{D}_s^\mp) \sim 0.5\%$) and their spin counterparts. The rescattering contribution can be in principle included when the CP violation for $B_s \rightarrow D_s \bar{D}_s$ is measured in future. However from theoretical viewpoint, it is likely that the rescattering mechanism will not include large $\Delta\phi_s$. Taking the $B_s \rightarrow D_s \bar{D}_s$ as the example, whose factorisation amplitude is given as

$$\mathcal{M}(B_s \rightarrow D_s^+ \bar{D}_s^-) = \frac{G_F}{\sqrt{2}} f_{D_s} (m_{B_s}^2 - m_{D_s}^2) F_0^{B_s \rightarrow D_s}(m_{D_s}^2) \{V_{cb} V_{cs}^* a_1 - V_{tb} V_{ts}^* [a_4 + a_{10} + (a_6 + a_8) R]\}, \quad (122)$$

with $R = 2m_{D_s}^2/(m_b - m_c)/(m_c + m_s)$, we find that the penguin effects are much smaller than the tree operator with the $a_1 \sim 1$. This should be in contrast with the case in $B_s \rightarrow J/\psi\phi$, where the $a_2 \sim 0.1$.

It is also suggested that the penguin contributions can be constrained by the SU(3)-symmetry related decay mode. To validate this proposal, one may explore the SU(3) symmetry breaking effects and thus we also calculate the modules of normalised amplitudes (in units of GeV^3) for $B_s \rightarrow J/\psi \bar{K}^{*0}$ and $B_s \rightarrow J/\psi\phi$ decays,

$$|\mathcal{A}_L(B_s \rightarrow J/\psi \bar{K}^{*0})| \approx 0.814_{-0.133}^{+0.140}, \quad |\mathcal{A}_L(B_s \rightarrow J/\psi\phi)| \approx 0.949_{-0.152}^{+0.159}; \quad (123)$$

$$|\mathcal{A}_\parallel(B_s \rightarrow J/\psi \bar{K}^{*0})| \approx 0.617_{-0.099}^{+0.110}, \quad |\mathcal{A}_\parallel(B_s \rightarrow J/\psi\phi)| \approx 0.730_{-0.112}^{+0.126}; \quad (124)$$

$$|\mathcal{A}_\perp(B_s \rightarrow J/\psi \bar{K}^{*0})| \approx 0.513_{-0.087}^{+0.100}, \quad |\mathcal{A}_\perp(B_s \rightarrow J/\psi\phi)| \approx 0.590_{-0.096}^{+0.109}. \quad (125)$$

which result in the ratios:

$$\begin{aligned} \left| \frac{\mathcal{A}_L(B_s \rightarrow J/\psi \bar{K}^{*0})}{\mathcal{A}_L(B_s \rightarrow J/\psi\phi)} \right| &= 0.858_{-0.196}^{+0.206}, \\ \left| \frac{\mathcal{A}_\parallel(B_s \rightarrow J/\psi \bar{K}^{*0})}{\mathcal{A}_\parallel(B_s \rightarrow J/\psi\phi)} \right| &= 0.845_{-0.188}^{+0.210}, \\ \left| \frac{\mathcal{A}_\perp(B_s \rightarrow J/\psi \bar{K}^{*0})}{\mathcal{A}_\perp(B_s \rightarrow J/\psi\phi)} \right| &= 0.869_{-0.204}^{+0.234}. \end{aligned} \quad (126)$$

These values indicate that the SU(3) symmetry breaking effects may reach 20% in the decay amplitudes and thus can be examined by future experiments.

V. SUMMARY

Up to this date, the CKM mechanism has successfully described almost all available data on flavor physics and CP violation, which has continued to motivate more precise tests of CP violation in the heavy flavor sector. The fact that the $B_s - \bar{B}_s$ mixing angle ϕ_s is tiny provides an ideal test of the SM and offers an opportunity to probe the new physics. To achieve this goal renders the precise theoretical predictions and experimental measurements important. The reduction of experimental uncertainties seems to have a promising prospect in near future, due to the large amount of data sample (to be) collected at LHC and the forthcoming Super KEKB factory.

On the other side, it is in necessity to learn about theoretical contamination. What has been explored in this work is an attempt to fill this gap. We have computed both tree and penguin amplitudes in the perturbative QCD approach, in which the leading-order contributions and NLO QCD corrections are taken into account. With the inclusion of these sizeable corrections, our theoretical results for CP-averaged branching ratios, polarization fractions, CP-violating asymmetries, and relative phases are in good consistency with the available data. Based on the global agreement, we have explored the penguin contributions and discussed the impact on ϕ_s extracted from $B_s \rightarrow J/\psi\phi$. Adopting the k_T factorization approach, we found that the results for ϕ_s can be shifted by about 10^{-3} while some angular coefficients can receive corrections about 10^{-2} . The future experiments can examine these predictions.

ACKNOWLEDGMENTS

W.Wang is grateful Yu Jia for valuable discussions on the phase convention. The work of X.Liu is supported by the National Natural Science Foundation of China under Grants No. 11205072, by a project funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), and by the Research Fund of Jiangsu Normal University under Grant No. 11XLR38. The work of W.Wang is supported by the DFG and the NSFC through funds provided to the Sino-German CRC 110 ‘‘Symmetries and the Emergence of Structure in QCD’’.

Appendix A: Related Functions in the perturbative QCD factorization

We show here the function h_i 's, coming from the Fourier transformations of the function $H^{(0)}$,

$$h_{fs}(x_1, x_3, b_1, b_3) = K_0 \left(\sqrt{x_1 x_3 (1 - r_2^2)} m_B b_1 \right) \left[\theta(b_1 - b_3) K_0 \left(\sqrt{x_3 (1 - r_2^2)} m_B b_1 \right) I_0 \left(\sqrt{x_3 (1 - r_2^2)} m_B b_3 \right) \right. \\ \left. + \theta(b_3 - b_1) K_0 \left(\sqrt{x_3 (1 - r_2^2)} m_B b_3 \right) I_0 \left(\sqrt{x_3 (1 - r_2^2)} m_B b_1 \right) \right] S_t(x_3), \quad (\text{A1})$$

$$h_{nfs}(x_1, x_2, x_3, b_1, b_2) = \left\{ \theta(b_2 - b_1) I_0(m_B \sqrt{x_1 x_3 (1 - r_2^2)} b_1) K_0(m_B \sqrt{x_1 x_3 (1 - r_2^2)} b_2) \right. \\ \left. + (b_1 \leftrightarrow b_2) \right\} \cdot \begin{cases} K_0(m_B F_{(1)} b_2), & \text{for } F_{(1)}^2 > 0 \\ \frac{\pi i}{2} H_0^{(1)}(m_B \sqrt{|F_{(1)}^2|} b_2), & \text{for } F_{(1)}^2 < 0 \end{cases}, \quad (\text{A2})$$

where J_0 is the Bessel function, K_0 and I_0 are the modified Bessel functions with $K_0(-ix) = -(\pi/2)Y_0(x) + i(\pi/2)J_0(x)$. The $F_{(j)}$'s are defined by

$$F_{(1,2)}^2 = (x_1 - x_2)[x_3 + (x_2 - x_3)r_2^2] + r_c^2. \quad (\text{A3})$$

The threshold resummation form factor $S_t(x_i)$ is adopted from Ref. [35]

$$S_t(x) = \frac{2^{1+2c} \Gamma(3/2 + c)}{\sqrt{\pi} \Gamma(1 + c)} [x(1 - x)]^c, \quad (\text{A4})$$

where the parameter $c = 0.3$. This function is normalized to unity.

The Sudakov factors are used as

$$S_{ab}(t) = s(x_1 P_1^+, b_1) + s(x_3 P_3^-, b_3) + s((1-x_3)P_3^-, b_3) - \frac{1}{\beta_1} \left[\ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_3\Lambda)} \right], \quad (\text{A5})$$

$$S_{cd}(t) = s(x_1 P_1^+, b_1) + s(x_2 P_2^+, b_2) + s((1-x_2)P_2^+, b_2) + s(x_3 P_3^-, b_1) + s((1-x_3)P_3^-, b_1) - \frac{1}{\beta_1} \left[2 \ln \frac{\ln(t/\Lambda)}{-\ln(b_1\Lambda)} + \ln \frac{\ln(t/\Lambda)}{-\ln(b_2\Lambda)} \right]. \quad (\text{A6})$$

The scale t_i 's in the above equations are chosen as

$$\begin{aligned} t_a &= \max(\sqrt{x_3(1-r_2^2)}m_B, 1/b_1, 1/b_3), \\ t_b &= \max(\sqrt{x_1(1-r_2^2)}m_B, 1/b_1, 1/b_3), \\ t_{nfs} &= \max(\sqrt{x_1 x_3(1-r_2^2)}m_B, \sqrt{(x_1-x_2)[x_3+(x_2-x_3)r_2^2]} + r_c^2 m_B, 1/b_1, 1/b_2). \end{aligned} \quad (\text{A7})$$

The scale t_i 's are chosen as the maximum energy scale appearing in each diagram to kill the large logarithmic radiative corrections.

-
- [1] J. Charles, O. Deschamps, S. Descotes-Genon, R. Itoh, H. Lacker, A. Menzel, S. Monteil and V. Niess *et al.*, Phys. Rev. D **84**, 033005 (2011) [[arXiv:1106.4041](#)] [hep-ph].
- [2] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **108**, 101803 (2012) [[arXiv:1112.3183](#)] [hep-ex].
- [3] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **109**, 171802 (2012) [[arXiv:1208.2967](#)] [hep-ex].
- [4] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. D **85**, 032006 (2012) [[arXiv:1109.3166](#)] [hep-ex].
- [5] G. Aad *et al.* [ATLAS Collaboration], JHEP **1212**, 072 (2012) [[arXiv:1208.0572](#)] [hep-ex].
- [6] R. Aaij *et al.* [LHCb Collaboration], Eur. Phys. J. C **73**, 2373 (2013) [[arXiv:1208.3355](#)] [hep-ex].
- [7] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. D **87**, 112010 (2013) [[arXiv:1304.2600](#)] [hep-ex].
- [8] S. Stone and L. Zhang, Phys. Rev. D **79**, 074024 (2009) [[arXiv:0812.2832](#)] [hep-ph].
- [9] S. Stone and L. Zhang, [arXiv:0909.5442](#) [hep-ex].
- [10] P. Colangelo, F. De Fazio and W. Wang, Phys. Rev. D **81**, 074001 (2010) [[arXiv:1002.2880](#)] [hep-ph].
- [11] P. Colangelo, F. De Fazio and W. Wang, Phys. Rev. D **83**, 094027 (2011) [[arXiv:1009.4612](#)] [hep-ph].
- [12] O. Leitner, J. -P. Dedonder, B. Loiseau and B. El-Bennich, Phys. Rev. D **82**, 076006 (2010) [[arXiv:1003.5980](#)] [hep-ph].
- [13] R. Fleischer, R. Kneijens and G. Ricciardi, Eur. Phys. J. C **71**, 1832 (2011) [[arXiv:1109.1112](#)] [hep-ph].
- [14] R. Aaij *et al.* [LHCb Collaboration], Phys. Lett. B **698**, 115 (2011) [[arXiv:1102.0206](#)] [hep-ex].
- [15] J. Li *et al.* [Belle Collaboration], Phys. Rev. Lett. **106**, 121802 (2011) [[arXiv:1102.2759](#)] [hep-ex].
- [16] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. D **84**, 052012 (2011) [[arXiv:1106.3682](#)] [hep-ex].
- [17] B. Bhattacharya, A. Datta and D. London, Int. J. Mod. Phys. A **28**, 1350063 (2013) [[arXiv:1209.1413](#)] [hep-ph].
- [18] S. Faller, R. Fleischer and T. Mannel, Phys. Rev. D **79**, 014005 (2009) [[arXiv:0810.4248](#)] [hep-ph].
- [19] Y. -Y. Keum, H. -n. Li and A. I. Sanda, Phys. Lett. B **504**, 6 (2001) [[hep-ph/0004004](#)].
- [20] Y. Y. Keum, H. -N. Li and A. I. Sanda, Phys. Rev. D **63**, 054008 (2001) [[hep-ph/0004173](#)].
- [21] C. -D. Lu, K. Ukai and M. -Z. Yang, Phys. Rev. D **63**, 074009 (2001) [[hep-ph/0004213](#)].
- [22] C. -D. Lu and M. -Z. Yang, Eur. Phys. J. C **23**, 275 (2002) [[hep-ph/0011238](#)].
- [23] C. -H. Chen and H. -N. Li, Phys. Rev. D **71**, 114008 (2005) [[hep-ph/0504020](#)].
- [24] X. Liu, Z. -Q. Zhang and Z. -J. Xiao, Chin. Phys. C **34**, 937 (2010).
- [25] X. Liu, H. -n. Li and Z. -J. Xiao, Phys. Rev. D **86**, 011501 (2012) [[arXiv:1205.1214](#)] [hep-ph].
- [26] H. -n. Li and S. Mishima, JHEP **0703**, 009 (2007) [[hep-ph/0610120](#)].
- [27] H. -n. Li, Y. -L. Shen, Y. -M. Wang and H. Zou, Phys. Rev. D **83**, 054029 (2011) [[arXiv:1012.4098](#)] [hep-ph].
- [28] H. -n. Li, Y. -L. Shen and Y. -M. Wang, Phys. Rev. D **85**, 074004 (2012) [[arXiv:1201.5066](#)] [hep-ph].
- [29] Y. Xie, P. Clarke, G. Cowan and F. Muheim, JHEP **0909**, 074 (2009) [[arXiv:0908.3627](#)] [hep-ph].
- [30] L. Zhang and S. Stone, Phys. Lett. B **719**, 383 (2013) [[arXiv:1212.6434](#)] [hep-ph].
- [31] A. S. Dighe, I. Dunietz and R. Fleischer, Eur. Phys. J. C **6**, 647 (1999) [[hep-ph/9804253](#)].
- [32] R. Fleischer, Phys. Rev. D **60**, 073008 (1999) [[hep-ph/9903540](#)].
- [33] C. -D. Lu and M. -Z. Yang, Eur. Phys. J. C **28**, 515 (2003) [[hep-ph/0212373](#)].
- [34] Z. -T. Wei and M. -Z. Yang, Nucl. Phys. B **642**, 263 (2002) [[hep-ph/0202018](#)].
- [35] T. Kurimoto, H. -n. Li and A. I. Sanda, Phys. Rev. D **65**, 014007 (2002) [[hep-ph/0105003](#)].
- [36] T. Kurimoto, H. -n. Li and A. I. Sanda, Phys. Rev. D **67**, 054028 (2003) [[hep-ph/0210289](#)].

- [37] A. Ali, G. Kramer, Y. Li, C. -D. Lu, Y. -L. Shen, W. Wang and Y. -M. Wang, Phys. Rev. D **76**, 074018 (2007) [[hep-ph/0703162](#) [HEP-PH]].
- [38] A. E. Bondar and V. L. Chernyak, Phys. Lett. B **612**, 215 (2005) [[hep-ph/0412335](#)].
- [39] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).
- [40] H. -n. Li and S. Mishima, Phys. Rev. D **74**, 094020 (2006) [[hep-ph/0608277](#)].
- [41] P. Ball and G. W. Jones, JHEP **0703**, 069 (2007) [[hep-ph/0702100](#) [HEP-PH]].
- [42] H. -n. Li and S. Mishima, Phys. Rev. D **71**, 054025 (2005) [[hep-ph/0411146](#)].
- [43] H. -n. Li, Phys. Lett. B **622**, 63 (2005) [[hep-ph/0411305](#)].
- [44] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996) [[hep-ph/9512380](#)].
- [45] C. -H. Chen, [hep-ph/0601019](#).
- [46] Z. -J. Xiao, Z. -Q. Zhang, X. Liu and L. -B. Guo, Phys. Rev. D **78**, 114001 (2008) [[arXiv:0807.4265](#) [hep-ph]].
- [47] H. -Y. Cheng, Y. -Y. Keum and K. -C. Yang, Phys. Rev. D **65**, 094023 (2002) [[hep-ph/0111094](#)].
- [48] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. D **86**, 071102 (2012) [[arXiv:1208.0738](#) [hep-ex]].
- [49] R. Aaij *et al.* [LHCb Collaboration], Nucl. Phys. B **867**, 547 (2013) [[arXiv:1210.2631](#) [hep-ex]].
- [50] K. De Bruyn, R. Fleischer, R. Knegjens, P. Koppenburg, M. Merk and N. Tuning, Phys. Rev. D **86**, 014027 (2012) [[arXiv:1204.1735](#) [hep-ph]].
- [51] W. Wang, Y. -L. Shen and C. -D. Lu, J. Phys. G **37**, 085006 (2010) [[arXiv:0908.2216](#) [hep-ph]].
- [52] C. -D. Lu, U. -G. Meissner, W. Wang and Q. Zhao, Eur. Phys. J. A **49**, 58 (2013) [[arXiv:1301.0225](#) [hep-ph]].
- [53] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. D **83**, 052012 (2011) [[arXiv:1102.1961](#) [hep-ex]].
- [54] R. Aaij *et al.* [LHCb Collaboration], [arXiv:1307.2782](#) [hep-ex].
- [55] M. Beneke, J. Rohrer and D. Yang, Nucl. Phys. B **774**, 64 (2007) [[hep-ph/0612290](#)].
- [56] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Rev. D **69**, 054023 (2004) [[hep-ph/0310084](#)].
- [57] P. Colangelo, F. De Fazio and T. N. Pham, Phys. Lett. B **542**, 71 (2002) [[hep-ph/0207061](#)].
- [58] H. -Y. Cheng, C. -K. Chua and A. Soni, Phys. Rev. D **71**, 014030 (2005) [[hep-ph/0409317](#)].