

Mainly axion cold dark matter from natural supersymmetry

Kyu Jung Bae,¹ Howard Baer,¹ and Eung Jin Chun²

¹*Dept. of Physics and Astronomy, University of Oklahoma, Norman, OK, 73019, USA*

²*Korea Institute for Advanced Study, Seoul 130-722, Korea*

By eschewing fine-tuning from the electroweak and QCD sectors of supersymmetry (natural supersymmetry or SUSY), and by invoking the Kim-Nilles solution to the SUSY μ problem, one is led to models wherein the dark matter is comprised of a mixture of axions and Higgsino-like WIMPs. Over a large range of Peccei-Quinn breaking scale $f_a \sim 10^9 - 10^{12}$ GeV, one then expects about 90-95% axion dark matter. In such a scenario, both axion and WIMP direct detection may be expected.

PACS numbers: 12.60.-i, 95.35.+d, 14.80.Ly, 11.30.Pb

The recent discovery of a Higgs-like boson with mass $m_h \simeq 125$ GeV at the CERN LHC is a triumph of modern particle physics [1]. But it brings with it a conundrum: why is the Higgs mass so small? In the Standard Model (SM), one may calculate

$$m_h^2(\text{phys}) = m_h^2(\text{tree}) + \delta m_h^2 \quad (1)$$

where the radiative correction $\delta m_h^2 \sim -\frac{3f_t^2}{8\pi^2}\Lambda^2 + \dots$ where $f_t \sim 1$ is the top quark Yukawa coupling and Λ is a high energy cutoff/regulator which denotes the limit of validity of the effective theory. If the SM is to be valid at energy scales Λ far beyond $m_{\text{weak}} \sim 100$ GeV, then an enormous fine-tuning will be required to maintain $m_h \sim 125$ GeV.

Supersymmetric (SUSY) theories of particle physics provide all-orders cancellations of the quadratic divergences thus stabilizing the Higgs mass. In SUSY, δm_h^2 is instead logarithmically divergent, and includes terms such as

$$\delta m_h^2(\text{SUSY}) \sim -\frac{3f_t^2}{8\pi^2}(m_{Q_3}^2 + m_{U_3}^2 + A_t^2) \ln(\Lambda^2/m_{\text{SUSY}}^2) \quad (2)$$

where $m_{Q_3}^2$, $m_{U_3}^2$ and A_t^2 are soft SUSY breaking terms related to the top- and bottom- squark masses and $m_{\text{SUSY}} \sim \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$. Requiring $\delta m_h^2 \lesssim 10m_h^2$ for Λ as high as m_{GUT} implies light 3rd generation squarks $m_{\tilde{t}_{1,2}}, m_{\tilde{b}_1} \lesssim 200$ GeV [2, 3], although this rather severe fine-tuning measure ignores non-independent contributions to m_h^2 which lead to large cancellations [4]. A model-independent, more conservative measure which allows for cancellations within m_h^2 requires that the magnitude of all *weak scale* contributions to m_h^2 (or m_Z^2) be comparable to m_h^2 (or m_Z^2). For instance, the Z mass is given by

$$\frac{m_Z^2}{2} \simeq -m_{H_u}^2 - \Sigma_u^u - \mu^2 \quad (3)$$

which is valid for the ratio of Higgs field vacuum expectation values (vevs) $\tan\beta \equiv v_u/v_d$ over its typical range of 3 – 60. Here, $m_{H_u}^2$ is a soft SUSY breaking Higgs mass, μ is the superpotential Higgs/Higgsino mass and

Σ_u^u collects various radiative corrections (expressions are provided in the Appendix of Ref. [5]). For many models, $m_{H_u}^2$ is driven radiatively to large negative values $|m_{H_u}^2| \gg m_Z^2$ signaling the breakdown of electroweak symmetry. A large positive value of μ^2 must be imposed (fine-tuned) to obtain the measured Z -mass, $m_Z \simeq 91.2$ GeV. (Alternatively, if $|m_{H_u}^2|$ is small but the Σ_u^u terms become large positive, $\Sigma_u^u \gg m_Z^2$, again a large value of μ^2 must be imposed, leading again to fine-tuning.) To avoid large uncorrelated cancellations (fine-tuning) in the Z mass, then one expects $|\mu|$ and $|m_{H_u}| \sim m_Z$, or of order 100 – 200 GeV [5, 6]. In addition, requiring the dominant radiative corrections

$$\left[\Sigma_u^u(\tilde{t}_{1,2}) = \frac{3}{16\pi^2} F(m_{\tilde{t}_{1,2}}^2) \times \left[f_t^2 A_t^2 - 8g_Z^2 \left(\frac{1}{4} - \frac{2}{3}x_W \right) \Delta_t \right] \right] \quad (4)$$

(where $\Delta_t = (m_{\tilde{t}_L}^2 - m_{\tilde{t}_R}^2)/2 + m_Z^2 \cos 2\beta(\frac{1}{4} - \frac{2}{3}x_W)$, $g_Z^2 = (g^2 + g'^2)/8$, $x_W \equiv \sin^2\theta_W$ and where $F(m^2) = m^2(\log \frac{m^2}{Q^2} - 1)$ with $Q^2 \simeq m_{\tilde{t}_1} m_{\tilde{t}_2}$ the optimized scale choice for minimization of the scalar potential) to be $\lesssim 100 - 200$ GeV then requires highly mixed top squarks with mass $m_{\tilde{t}_1} \sim 1 - 2$ TeV and $m_{\tilde{t}_2} \sim 3 - 4$ TeV [5, 6].¹ Such highly mixed top squark masses lift m_h into the 125 GeV range (even with stops as light as a few TeV) since m_h is maximized for large mixing [7]. The TeV-scale top squark masses are also heavy enough to suppress anomalous contributions to $b \rightarrow s\gamma$ decay and to avoid recent LHC null results for top squark searches [5].

The above features have enormous implications for SUSY phenomenology: in this case, the lightest SUSY

¹ Using the full radiative corrections, then large values of weak scale A_t suppress $\Sigma_u^u(\tilde{t}_1)$ via the square bracket in Eq. (4) and via the F function for $\Sigma_u^u(\tilde{t}_2)$. The suppression due to mixing then allows for much larger stop masses (around the few TeV scale) than are found in generic natural SUSY models, where it is often claimed that $m_{\tilde{t}_{1,2}}$ need be $\lesssim 200 - 500$ GeV [3].

particle, often touted as a WIMP dark matter candidate, is expected to be largely Higgsino-like since the Higgsino mass is $\simeq |\mu|$ and $|\mu| \sim m_Z$. Since Higgsino-like WIMPs couple directly to the Goldstone components of the gauge bosons, they can annihilate with large rates into WW and ZZ final states. Since the WIMP thermal relic abundance is inversely proportional to its annihilation rate, then Higgsinos with mass $m_{\tilde{Z}_1} \sim 100\text{--}200$ GeV develop a relic density of $\Omega_{\tilde{Z}_1} h^2 \sim 0.005\text{--}0.01$, *i.e.* typically a factor 10–15 below the WMAP/Planck measured value [8]. Supersymmetric models with low fine-tuning (not too heavy 3rd generation squarks and low μ) are referred to as *natural SUSY* [2, 3, 5, 6] since they are devoid of large electroweak fine-tuning.

A further possibility for fine-tuning occurs in the QCD sector. To implement 't Hooft's solution to the $U(1)_A$ problem [9] (*i.e.* why there are three and not four light pions), the term

$$\frac{\bar{\theta}}{32\pi^2} F_{A\mu\nu} \tilde{F}_A^{\mu\nu} \quad (5)$$

should occur in the QCD Lagrangian, where $\bar{\theta} = \theta + \arg(\det\mathcal{M})$, \mathcal{M} is the quark mass matrix, $F_{A\mu\nu}$ is the gluon field strength and $\tilde{F}_A^{\mu\nu}$ is its dual. Measurements of the neutron electric dipole moment (EDM) require $\bar{\theta} \lesssim 10^{-10}$ so that one might require an enormous cancellation within $\bar{\theta}$ [10]. Alternatively, the PQWW solution [11] introduces an *axion* field a ; the additional axion contributions to Eq. (5) allow for $\bar{\theta}$ to dynamically settle to zero, thus solving the so-called strong CP problem.

In SUSY theories, the axion enters as one element of an axion *superfield* which necessarily contains also a spin-0 R -parity even saxion s and a spin-1/2 R -parity-odd axino \tilde{a} . Calculations of the saxion and axino masses within the context of supergravity [12] imply $m_s \sim m_{\tilde{a}} \sim m_{3/2}$ where the gravitino mass $m_{3/2}$ is expected to be of order the TeV scale. If the lightest neutralino (*e.g.* the Higgsino \tilde{Z}_1) is the lightest SUSY particle (LSP) in R -parity conserving theories, then one would expect dark matter to be comprised of *two particles*: the axion as well as the Higgsino-like SUSY WIMP. The saxion and axino couplings to matter are suppressed by the PQ breaking scale f_a which may range from $f_a \sim 10^9\text{--}10^{16}$ GeV [10]. While the saxion and axino are expected to play little or no role in terrestrial experiments, they can have an enormous impact on dark matter production in the early universe.

The PQ symmetry required to solve the strong CP problem can be implemented in two ways. In the SUSY KSVZ model [13, 14], the axion superfield couples to exotic heavy quark/squark superfields Q and \bar{Q} which carry PQ charges. The loop-induced axino-gluino-gluon coupling leads to a thermal axino production rate proportional to the re-heat temperature T_R at the end of inflation. It also allows for axino decays $\tilde{a} \rightarrow g\tilde{g}$, $\tilde{a} \rightarrow \gamma\tilde{Z}_i$

or $\tilde{a} \rightarrow Z\tilde{Z}_i$ (with $i = 1\text{--}4$). In SUSY KSVZ, axinos are sufficiently long-lived that they almost always decay after neutralino freeze-out $T_{\text{fr}} \sim m_{\tilde{Z}_1}/25$ [15]. Axinos, produced at a sufficient rate, may induce *neutralino re-annihilation* at the axino decay temperature $T_D^{\tilde{a}} = \sqrt{\Gamma_{\tilde{a}} M_P}/(\pi^2 g_*(T_D^{\tilde{a}})/90)^{1/4}$ (here, M_P is the reduced Planck mass $\simeq 2 \times 10^{18}$ GeV). Since the re-annihilation Yield $Y_{\tilde{Z}_1}^{re-ann} \equiv n_{\tilde{Z}_1}/s$ (where $n_{\tilde{Z}_1}$ is number density and s is entropy density) is inversely proportional to T_D , then when $T_D < T_{fr}$, re-annihilation also augments the neutralino abundance [15, 16]

Saxions may be produced thermally [17] (again proportional to T_R/f_a^2) or via coherent oscillations [18] (proportional to f_a^2 , so important at very large $f_a \sim 10^{14}\text{--}10^{16}$ GeV). They may decay via $s \rightarrow gg$, which leads to entropy dilution of all relics present at the time of decay, or to $\tilde{g}\tilde{g}$ or other SUSY modes, which can augment the neutralino abundance. Depending on the combinations of PQ charges q_i and PQ vevs v_i , saxions should also decay via $s \rightarrow aa$ or $s \rightarrow \tilde{a}\tilde{a}$ for $\xi \equiv \sum_i q_i^3 v_i^2/f_a^2 \sim 1$. The first of these leads to production of dark radiation which is stringently limited by WMAP/Planck [19] parametrized in terms of the number of additional neutrinos present in the universe: $\Delta N_{eff} < 1.6$ at 95% CL. If the saxion or axino decays occur much after $T_{\text{BBN}} \sim 1$ MeV, then light elements produced during Big Bang Nucleosynthesis may become dis-associated leading also to severe constraints [20].

Alternatively to SUSY KSVZ, in the SUSY DFSZ model [21–24] the PQ superfield couples directly to the Higgs superfields carrying non-trivial PQ charges:

$$W_{\text{DFSZ}} \ni \lambda \frac{S^2}{M_P} H_u H_d. \quad (6)$$

Here, S is a Minimal Supersymmetric Standard Model (MSSM) singlet but carries a PQ charge and contains the axion field. An advantage of this approach is that it provides a solution to the SUSY μ problem [22]: since the μ term is supersymmetric, one expects $\mu \sim M_P$ in contrast to phenomenology (Eq. (3)) which requires $\mu \sim m_{\text{weak}}$. In this Kim-Nilles solution, PQ charge assignments to the Higgs fields imply that the usual superpotential μ term is forbidden. Upon breaking of PQ symmetry, the field S receives a vev $\langle S \rangle \sim f_a$, so that an effective μ term is generated with $\mu \sim \lambda f_a^2/M_P \sim \lambda m_{3/2}$. For $\lambda f_a^2/M_P \sim 100$ GeV, then one may generate $\mu \sim 100$ GeV in accord with naturalness whilst $m_{\tilde{a}} \sim m_{3/2} \sim 10$ TeV in accord with LHC constraints and in accord with at least a partial decoupling solution to the SUSY flavor, CP and gravitino problems [25].

In SUSY DFSZ, the direct coupling of the axion supermultiplet to the Higgs superfields leads to thermal production rates which are independent of T_R [23, 24].

The saxion and axino thermal yields are then given by

$$Y_s^{\text{TP}} \simeq 10^{-7} \zeta_s \left(\frac{\mu}{\text{TeV}} \right)^2 \left(\frac{10^{12} \text{ GeV}}{f_a} \right)^2 \quad (7)$$

$$Y_{\tilde{a}}^{\text{TP}} \simeq 10^{-7} \zeta_{\tilde{a}} \left(\frac{\mu}{\text{TeV}} \right)^2 \left(\frac{10^{12} \text{ GeV}}{f_a} \right)^2 \quad (8)$$

where the ζ_i are model-dependent constants of order unity. Saxions can also be produced via coherent oscillations with a yield given by

$$Y_s^{\text{CO}} = 1.9 \times 10^{-6} \left(\frac{\text{GeV}}{m_s} \right) \left(\frac{\min[T_R, T_s]}{10^7 \text{ GeV}} \right) \left(\frac{f_a}{10^{12} \text{ GeV}} \right)^2 \quad (9)$$

assuming an initial saxion field amplitude of $s_0 \sim f_a$. Along with these, neutralinos will be produced via thermal freeze-out as usual given by

$$Y_{\tilde{Z}_1}^{\text{fr}} \simeq \frac{(90/\pi^2 g_*(T_{\text{fr}}))^{1/2}}{4 \langle \sigma v \rangle M_P T_{\text{fr}}} \quad (10)$$

where T_{fr} is the freeze-out temperature ($\sim m_{\tilde{Z}_1}/25$) and $\langle \sigma v \rangle$ is the thermally averaged neutralino annihilation cross section times relative velocity. Axions will also be produced at the QCD phase transition via coherent oscillations [26] given by

$$\Omega_a^{\text{std}} h^2 \simeq 0.23 f(\theta_i) \theta_i^2 \left(\frac{f_a/N_{\text{DW}}}{10^{12} \text{ GeV}} \right)^{7/6} \quad (11)$$

where the misalignment angle $0 < \theta_i < \pi$, N_{DW} is the domain wall number, and $f(\theta_i)$ is the anharmonicity factor parametrized as $f(\theta_i) = \left[\ln \left(\frac{e}{1 - \theta_i^2/\pi^2} \right) \right]^{7/6}$.

Along with the above processes, neutralinos can be produced via axino decays. In SUSY DFSZ, the dominant modes include: $\tilde{a} \rightarrow \tilde{Z}_i \phi$ (where $\phi = h, H, A$), $\tilde{Z}_i Z$ ($i = 1 - 4$), $\tilde{W}_j^\pm H^\mp$ and $\tilde{W}_j^\pm W^\mp$ ($j = 1 - 2$). Summing over decay modes and neglecting phase space factors, the axino width is

$$\Gamma_{\tilde{a}} \sim \frac{c_H^2}{4\pi} \left(\frac{\mu}{v_{PQ}} \right)^2 m_{\tilde{a}}, \quad (12)$$

where c_H is an order one parameter for the axino (saxion) coupling arising from Eq. (6) and $v_{PQ} \equiv \sqrt{\sum_i q_i^2 v_i^2} \sim f_a$ in terms of PQ charges and vevs. This tends to greatly exceed the value obtained in SUSY KSVZ.

For illustration, we adopt the Standard Underabundance SUSY benchmark (SUA) scenario from Ref. [5, 27] which assumes a Higgsino mass scale $\mu \sim 150 \text{ GeV}$ while sparticles which do not contribute to naturalness are at high masses $\gg 1 \text{ TeV}$, safely beyond the LHC reach. Our results are hardly sensitive to the selected benchmark so long as $|\mu| \sim m_Z$ while the remaining sparticles are quite heavy. The axino decay temperature for the SUA point using the complete decay widths is shown in Fig. 1. The




FIG. 1: Decay temperature of DFSZ axinos vs. PQ scale f_a for the SUA benchmark.

spike is due to a resonant axino-neutralino mixing effect when $|m_{\tilde{a}} - m_{\tilde{Z}_i}| \lesssim \mu v / f_a$ and can be ignored in our analyses. Comparing against the neutralino freeze-out temperature for a 135 GeV Higgsino-like LSP ($T_{fr} \sim 5 \text{ GeV}$), we see that $T_D^{\tilde{a}}$ tends to *exceed* T_{fr} for $f_a \lesssim 10^{12} \text{ GeV}$. In this case, the axino-produced neutralinos will thermalize and their abundance is determined by the usual thermal freeze-out. For higher $f_a \gtrsim 10^{12}$, the axinos decay after neutralino freeze-out. The injection of a huge population of decay-produced neutralinos into the cosmic plasma at temperatures well below freeze-out leads to neutralino re-annihilation at $T_D^{\tilde{a}}$. Solving the Boltzmann equation for the neutralino re-annihilation abundance leads to a Yield value given by [15, 16]²

$$Y_{\tilde{Z}_1}^{\text{re-ann}} \simeq \frac{(90/\pi^2 g_*(T_D^{\tilde{a}}))^{1/2}}{4 \langle \sigma v \rangle M_P T_D^{\tilde{a}}} \quad (13)$$

which essentially replaces T_{fr} by $T_D^{\tilde{a}}$ in Eq. (10). Since $T_D^{\tilde{a}} < T_{\text{fr}}$, this gives an *increased* yield over the thermal expectation. If the temperature at which the axino density equals the radiation density, $T_e^{\tilde{a}} = 4m_{\tilde{a}} Y_{\tilde{a}}/3$, is smaller than the decay temperature, axinos temporarily dominate the energy density of the universe. Axino domination rarely happens in SUSY DFSZ since then a huge value of $\mu \gtrsim 10^{-4} f_a$ is required.

Saxions, produced thermally and non-thermally, can decay via $s \rightarrow hh, HH, hH, AA, H^+H^-, ZZ, W^+W^-, ZA, W^\pm H^\mp, \tilde{Z}_i \tilde{Z}_{i'}, \tilde{W}_j^\pm \tilde{W}_{j'}^\mp$, and also to fermions and sfermions (complete decay formulae are given in [28]). For large m_s , the width is dominated by

$$\Gamma(s \rightarrow \text{Higgsinos}) \simeq \frac{c_H^2}{32\pi} \left(\frac{\mu}{v_{PQ}} \right)^2 m_s. \quad (14)$$

In addition, for $\xi \sim 1$, the decay $s \rightarrow aa$ may be sizable, leading to dark radiation [27], or $s \rightarrow \tilde{a}\tilde{a}$ may occur

² For an explicit derivation, see Eq's 7.8-7.10 of Ref. [15].




FIG. 2: Decay temperature of DFSZ saxions vs. PQ scale f_a for the SUA benchmark with $m_{\bar{a}} = 2$ TeV. Solid curves have $\xi = 1$ while dashed curves have $\xi = 0$.

as well, further augmenting the LSP abundance. The saxion decay temperature T_D^s is shown in Fig. 2. The spike is due to a resonant saxion-Higgs mixing effect when $|m_s^2 - m_{h,H}^2| \lesssim B\mu v/f_a$ and can be ignored in our analyses. For $f_a \lesssim 10^{12}$ GeV, saxions tend to decay before the neutralino freeze-out. A comparison of the saxion radiation equality temperature T_e^s against the decay temperature T_D^s shows that saxions dominate the energy density of the universe only when $f_a \gtrsim 10^{14}$ GeV for which Y_s^{CO} is large enough.

Let us now examine the contributions of neutralinos and axions to the observed dark matter density expected in the SUSY DFSZ model. Our main result is shown in Fig. 3 assuming $m_{\bar{a}} = m_s = 5$ TeV. We adopt the range of f_a starting at 10^9 GeV (stellar cooling arguments and supernovae analyses require $f_a \gtrsim 10^9$ GeV), and proceed to values $f_a \sim 10^{16}$ GeV (far above the naive closure bound $f_a \lesssim 10^{12}$ GeV gained from Eq. (11) using $\theta_i = 1$). The neutralino abundance $\Omega_{\tilde{Z}_1} h^2 \approx 0.01$ is given by the standard thermal freeze-out over a *large range* of f_a extending all the way up to $f_a \sim 10^{12}$ GeV. In this regime, the axion abundance can always be found by adjusting θ_i such that the summed abundance meets the measured value: $\Omega_{\tilde{Z}_1} h^2 + \Omega_a h^2 = 0.12$. The required value of θ_i is shown in Fig. 4. For very low $f_a \sim 10^9$ GeV, a large value of $\theta_i \sim \pi$ is required, and $\Omega_a h^2$ is dominated by the anharmonicity term. As f_a increases, the assumed initial axion field value $\theta_i f_a$ increases, so the required misalignment angle θ_i decreases. Values of $\theta_i \sim 1$ are found around $f_a \sim 10^{12}$ GeV for both $\xi = 0$ and 1. In this entire region with $f_a \sim 10^9 - 10^{12}$ GeV, we expect from natural SUSY that the relic Higgsino abundance lies at the standard freezeout value, comprising about 5-10% of the total dark matter density, while axions would comprise 90-95% of the abundance. Thus, over the commonly considered range of f_a , we expect *mainly axion cold dark matter from natural SUSY*, along with a non-negligible fraction of Higgsino-like WIMPs.

As f_a increases beyond 10^{12} GeV, axinos and saxions




FIG. 3: Neutralino and axion relic abundance from the SUSY DFSZ axion model versus PQ scale f_a for the SUA benchmark point.




FIG. 4: Axion field misalignment angle vs. f_a which is required to saturate mixed axion-neutralino abundance for $\xi = 0$ (dashed) and $\xi = 1$ (solid).

decay later than the neutralino freeze-out so that axino and saxion decays to SUSY particles add to the WIMP abundance, and $\Omega_{\tilde{Z}_1} h^2$ begins to increase. The exact value of $\Omega_{\tilde{Z}_1} h^2$ is given by the re-annihilation abundance (Eq. (13)) at temperatures $T_D < T_{\text{fr}}$ and so the neutralino abundance can be seen as the rising curves of $\Omega_{\tilde{Z}_1} h^2$ in Fig. 3 for both $\xi = 0$ and $\xi = 1$. The rapid rise of the neutralino abundance as f_a increases results in excluding the region of $f_a \gtrsim 5 \times 10^{12}$ GeV (5×10^{13} GeV) by dark matter overproduction for all values of $\xi = 0 - 1$. If we increase $m_{\bar{a}} = m_s$ to 10 (20) TeV, then the upper bound on f_a moves to 6×10^{12} (9×10^{12}) GeV for $\xi = 0$. For $\xi = 0$, the $\Omega_{\tilde{Z}_1} h^2$ curve rises steadily with f_a due to increasing production of saxions from coherent oscillations and their dominant decays to SUSY particles. This leads to subsequent neutralino re-annihilation at decreasing temperatures T_D^s . For $\xi = 1$, the dominant saxion decay mode is $s \rightarrow aa$, and decay-produced neutralinos come mainly from thermal axino production which decreases as f_a increases. One sees that $\Omega_{\tilde{Z}_1} h^2$ turns over and briefly reaches $\Omega_{\tilde{Z}_1} h^2 \simeq 0.1$ at $f_a \sim 3 \times 10^{13}$ GeV before beginning again a rise due to increasing non-thermal saxion production. It is important to note that for $\xi \sim 1$ and $f_a \gtrsim 10^{14}$ GeV, too much dark radiation is produced ($\Delta N_{\text{eff}} > 1.6$) (not shown here) and thus large

f_a is excluded by overproduction of both dark radiation and WIMPs.

Summary: Supersymmetry with not too heavy top squarks, low Higgsino mass $\mu \sim 100 - 200$ GeV and PQWW solution to the strong CP problem successfully avoids high fine-tuning in both the electroweak and QCD sectors of the theory while evading LHC constraints. The SUSY DFSZ model, wherein Higgs superfields carry PQ charge, also provides a solution to the SUSY μ problem. In such models, over a large range of PQ breaking scale $f_a \sim 10^9 - 10^{12}$ GeV, saxions and axinos typically decay before neutralino freeze-out so that the Higgsino portion of dark matter is expected to lie in the 5-10% range while axions would comprise the remainder: 90-95%. The relic Higgsinos ought to be detectable at ton scale noble liquid detectors, even with a depleted local abundance, while indirect detection should be more limited since expected rates go as the depleted abundance squared [8]. Prospects are bright for microwave cavity detection of axions since the range of f_a where mainly axion dark matter is expected should be accessible to experimental searches [29]. While corroborative searches for natural SUSY with light Higgsinos is limited at the LHC [30], a definitive Higgsino search should be possible at e^+e^- colliders with \sqrt{s} up to 500 – 600 GeV.

Acknowledgements: We thank V. Barger for comments on the manuscript and A. Lessa for collaboration on the early phase of this work. This work was supported in part by the US Department of Energy, Office of High Energy Physics.

-
- [1] G. Aad *et al.* (ATLAS Collaboration), *Phys. Lett.* **B710** (2012) 49; S. Chatrchyan *et al.* (CMS Collaboration), *Phys. Lett.* **B710** (2012) 26.
- [2] R. Kitano and Y. Nomura, *Phys. Lett.* **B631** (2005) 631 and *Phys. Rev.* **D73** (2006) 095004.
- [3] M. Papucci, J. T. Ruderman and A. Weiler, *JHEP* **1209** (2012) 035; C. Brust, A. Katz, S. Lawrence and R. Sundrum, *JHEP* **1203** (2012) 103; R. Essig, E. Izaguirre, J. Kaplan and J. G. Wacker, *JHEP* **1201** (2012) 074; H. Baer, V. Barger, P. Huang and X. Tata, *JHEP* **1205** (2012) 109.
- [4] H. Baer, V. Barger and D. Mickelson, arXiv:1309.2984 [hep-ph], *Phys. Rev.* **D**, in press.
- [5] H. Baer, V. Barger, P. Huang, A. Mustafayev and X. Tata, *Phys. Rev.* **D87** (2013) 115028.
- [6] H. Baer, V. Barger, P. Huang, A. Mustafayev and X. Tata, *Phys. Rev. Lett.* **109** (2012) 161802.
- [7] M. S. Carena and H. E. Haber, *Prog. Part. Nucl. Phys.* **50** (2003) 63 [hep-ph/0208209].
- [8] H. Baer, V. Barger and D. Mickelson, *Phys. Lett. B* **726** (2013) 330.
- [9] R. D. Peccei, *Lect. Notes Phys.* **741** (2008) 3 [hep-ph/0607268].
- [10] For a recent review, see J. E. Kim and G. Carosi, *Rev. Mod. Phys.* **82** (2010) 557.
- [11] R. Peccei and H. Quinn, *Phys. Rev. Lett.* **38** (1977) 1440 and *Phys. Rev.* **D16** (1977) 1791; S. Weinberg, *Phys. Rev. Lett.* **40** (1978) 223; F. Wilczek, *Phys. Rev. Lett.* **40** (1978) 279.
- [12] P. Moxhay and K. Yamamoto, *Phys. Lett.* **B151** (1985) 363; E. J. Chun, J. E. Kim and H. P. Nilles, *Phys. Lett.* **B287** (1992) 123; E. J. Chun and A. Lukas, *Phys. Lett.* **B357** (1995) 43; J. E. Kim, M.-S. Seo, *Nucl. Phys.* **B864** (2012) 296.
- [13] J. E. Kim, *Phys. Rev. Lett.* **43** (1979) 103; M. A. Shifman, A. Vainshtein and V. I. Zakharov, *Nucl. Phys.* **B166** (1980) 493.
- [14] J. E. Kim, *Phys. Rev. Lett.* **67** (1991) 3465
- [15] H. Baer, A. Lessa, S. Rajagopalan and W. Sreethawong, *JCAP* **1106** (2011) 031.
- [16] K.-Y. Choi, J. E. Kim, H. M. Lee and O. Seto, *Phys. Rev.* **D77** (2008) 123501.
- [17] P. Graf and F. Steffen, *JCAP* **1302** (2013) 018.
- [18] J. E. Kim, *Phys. Lett.* **B136** (1984) 378; M. Kawasaki, K. Nakayama and M. Senami, *JCAP* **0803** (2008) 009.
- [19] E. Komatsu, *et. al.*, *Astrophys. J. Suppl.* **192** (2011) 18; J. Dunkley, *et. al.*, *Astrophys. J.* **739** (2011) 52; R. Keisler, *et. al.*, *Astrophys. J.* **743** (2011) 28 .
- [20] R. H. Cyburt, *et. al.*, *JCAP* **0910** (2009) 021; M. Kawasaki, K. Kohri, T. Moroi and A. Yotsuyanagi, *Phys. Rev.* **D78** (2008) 065011; K. Jedamzik, *Phys. Rev.* **D74** (2006) 103509.
- [21] M. Dine, W. Fischler and M. Srednicki, *Phys. Lett.* **B104** (1981) 199; A. P. Zhitnitskii, *Sov. J. Nucl.* **31** (1980) 260.
- [22] J. E. Kim and H. P. Nilles, *Phys. Lett.* **B138** (1984) 150.
- [23] E. J. Chun, *Phys. Rev.* **D84** (2011) 043509.
- [24] K. J. Bae, K. Choi and S. H. Im, *JHEP* **1108**, 065 (2011); K. J. Bae, E. J. Chun and S. H. Im , *JCAP* **1203** (2012) 013.
- [25] M. Dine, A. Kagan and S. Samuel, *Phys. Lett.* **B243** (1990) 250; N. Arkani-Hamed and H. Murayama, *Phys. Rev.* **D 56** (1997) 6733; K. Agashe and M. Graesser, *Phys. Rev.* **D 59** (1998) 015007; A. Cohen, D. B. Kaplan and A. Nelson, *Phys. Lett.* **B388** (1996) 588.
- [26] L. F. Abbott and P. Sikivie, *Phys. Lett.* **B120** (1983) 133; J. Preskill, M. Wise and F. Wilczek, *Phys. Lett.* **B120** (1983) 127; M. Dine and W. Fischler, *Phys. Lett.* **B120** (1983) 137; M. Turner, *Phys. Rev.* **D33** (1986) 889; for an update, see L. Visinelli and P. Gondolo, *Phys. Rev.* **D80** (2009) 035024.
- [27] K. J. Bae, H. Baer and A. Lessa, *JCAP* **1304** (2013) 041.
- [28] K. J. Bae, H. Baer and E. J. Chun, *JCAP* **1312** (2013) 028.
- [29] L. Duffy, *et. al.*, *Phys. Rev. Lett.* **95** (2005) 091304 and *Phys. Rev.* **D74** (2006) 012006; for a review, see S. J. Asztalos, L. Rosenberg, K. van Bibber, P. Sikivie and K. Zioutas, *Ann. Rev. Nucl. Part. Sci.* **56** (2006) 293.
- [30] H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev, W. Sreethawong and X. Tata, *JHEP* **1312** (2013) 013 and arXiv:1306.3148 [hep-ph].