

Observational rotation curves and density profiles vs. the Thomas-Fermi galaxy structure theory

H. J. de Vega^{1,2*}, P. Salucci^{3†}, N. G. Sanchez^{2‡}

¹*Sorbonne Universités, UPMC (Univ. Paris VI), CNRS, Laboratoire Associé au CNRS UMR 7589, Tour 13-14, 4ème. et 5ème. étage, Boîte 126, 4, Place Jussieu, 75252 Paris, France.*

²*Observatoire de Paris, LERMA. Laboratoire Associé au CNRS UMR 8112, 61, Avenue de l'Observatoire, 75014 Paris, France.*

³*SISSA/ISAS and INFN, Trieste, Iniziativa Specifica QSKY, via Bonomea 265, I-34136, Trieste, Italia.*

26 February 2022

ABSTRACT

The Thomas-Fermi approach to galaxy structure determines selfconsistently the gravitational potential of the fermionic warm dark matter (WDM) given its distribution function $f(E)$. This framework is appropriate for macroscopic quantum systems as neutron stars, white dwarfs and WDM galaxies. Compact dwarf galaxies are near the quantum degenerate regime, while large galaxies are in the classical Boltzmann regime. We derive analytic **scaling relations** for the main galaxy magnitudes: halo radius r_h , mass M_h and phase-space density which are all well reproduced by the observational data for a large variety of galaxies. Small deviations from the exact scaling show up for compact dwarfs due to quantum macroscopic effects. We contrast the theoretical curves for the circular galaxy velocities $v_c(r)$ and density profiles $\rho(r)$ with those obtained from observations using the empirical Burkert profile. Results are **independent** of any WDM particle physics model, they only follow from the gravitational interaction of the WDM particles and their fermionic nature. The theoretical rotation curves and density profiles reproduce very well the observational curves for $r \lesssim r_h$ obtained from ten different and independent sets of data for galaxy masses from $5 \times 10^9 M_\odot$ till $5 \times 10^{11} M_\odot$. Our normalized theoretical circular velocities and normalized density profiles turn to be **universal** functions of r/r_h for **all** galaxies. In addition, they agree extremely well with the observational curves described by the Burkert profile for $r \lesssim 2 r_h$. These results show that the Thomas-Fermi approach correctly describes the galaxy structures.

Key words: Dark Matter, Galaxy structure, Galaxy Rotation Curves, Galaxy Density Profiles

1 INTRODUCTION

Dark matter (DM) is the main component of galaxies: the fraction of DM over the total galaxy mass goes from 95% for large dilute galaxies (Memola et al. 2011; Oh et al. 2008; Persic et al. 1996) till 99.99% for dwarf compact galaxies (Brodie et al. 2011; Martin et al. 2008; Walker 2012; Willman & Strader 2012; Woo et al. 2008). Therefore, the study of galaxy properties is an excellent way to disentangle the nature of DM.

Warm Dark Matter (WDM), that is dark matter

formed by particles with masses in the keV scale receives increasing attention today (Chalange Colloquium (2012); Meudon Colloquium (2012); Cosmic Frontier (2013) and references therein).

At intermediate scales ~ 100 kpc, WDM gives the **correct abundance** of substructures and therefore WDM solves the CDM overabundance of structures at small scales. (Colín et al. 2000; Sommer-Larsen & Dolgov 2001; Gao & Theuns 2007; Tikhonov et al. 2009; Zavala et al. 2009; Papastergis et al. 2011; Lovell et al. 2012, 2013; Anderhalden et al. 2013). For scales larger than 100 kpc, WDM yields the same results than CDM. Hence, WDM agrees with all the observations: small scale as well as large scale structure observations and CMB anisotropy observations.

* de Vega@lpthe.jussieu.fr

† salucci@sissa.it

‡ Norma.Sanchez@obspm.fr

WDM simulations (as Avila-Reese et al. (2001); Colín et al. (2008); Viñas et al. (2012); Gao & Theuns (2007); Lovell et al. (2012, 2013); Macciò et al. (2012); Zavala et al. (2009) and many others) are purely classical (i.e. WDM quantum dynamics is not used in those simulations). The dynamics of DM in the simulations is worked out classically from the classical Newton's equations or self-gravitating hydrodynamics (also classical, without WDM quantum effects). Quantum effects, as the DM quantum pressure, are absent in such frameworks. Inside galaxy cores, below ~ 100 pc, WDM N -body classical physics simulations do not provide the correct structures because quantum effects are important in WDM at these scales. Classical physics N -body WDM simulations without the WDM quantum pressure exhibit cusps or small cores with sizes smaller than the observed cores (Avila-Reese et al. 2001; Colín et al. 2008; Viñas et al. 2012; Macciò et al. 2012). The relevant WDM quantum effect, as discussed in our previous articles (Destri, de Vega, Sanchez 2013a, in what follows (DdVS 2013a), Destri, de Vega, Sanchez 2013b, in what follows DdVS (2013b)), is the fermionic quantum pressure.

In WDM simulations the fact that the DM is warm appears in the primordial power spectrum which is suppressed at small scales below the free streaming length. This can be implemented by a simple cutoff or through precise formulas fitting the WDM primordial power spectrum (see (DdVS 2013c) for a recent article), and through the non-zero particle velocity dispersion. In addition, in fermionic WDM, the phase space density is bounded from above by the Pauli principle (DdVS 2013a), (Tremaine & Gunn 1979). This is the only quantum (fermionic) aspect of WDM implemented in the WDM simulations. Such bound is clearly not enough to account for the quantum pressure of the WDM fermions, because the quantum pressure requires a combination of both the Pauli principle and the Heisenberg principle. And therefore, the power spectrum cutoff and the Pauli bound on the phase-space density are not enough in the simulations to enlarge the size of the WDM halos against the gravitation attraction. That is the reason why in WDM simulations the core size problem persists. The presence of a repulsive quantum fermionic pressure is crucial to enlarge enough the halos against gravitation to account for the macroscopic core sizes. The cutoff in the primordial power spectrum in the WDM simulations is enough to account for the right number of substructures and solve the CDM overabundance problem.

Besides of the cutoff in the primordial power spectrum and the Pauli bound in the phase-space density implemented in the WDM simulations, the crucial point is that in WDM simulations the N -body self-gravitating classical evolution follows the classical Newton's equations. This dynamics in the inner dense regions is far away from the quantum evolution according to the N -body Schrödinger equation and does not contain the quantum fermionic pressure. Instead, the Thomas-Fermi approach corresponds to the Schrödinger equation in the large N regime and contains from the start the quantum pressure. This is the reason why the quantum pressure is naturally contained in the Thomas-Fermi approach and it is not included so far in the classical N -body WDM simulations.

The quantum pressure is well captured in the Thomas-Fermi approach (DdVS 2013a,b). The lack of quantum pres-

sure in the WDM simulations explains why they exhibit cusps or small cores with sizes smaller than the observed cores (Macciò et al. 2012). WDM predicts correct structures and cores with the right sizes for small scales (below the kpc scale) when its **quantum** nature is taken into account (DdVS 2013a,b).

The quantum effects of matter in the inner halo regions arise because of the quantum fermionic nature and of the quantum uncertainty principle, the combined action of both translates into a non-zero quantum pressure. For macroscopic systems with a large number of particles as galaxies, this translates into macroscopic quantum effects. (Other examples in nature are He^3 , white dwarf stars and neutron stars). The quantum pressure goes as the mass of the particle to the power $-8/3$ and it is therefore much smaller for baryons than for WDM (DdVS 2013a).

The Thomas-Fermi DdVS approach applies irrespective of the WDM particle physics model. The lower bound for the WDM particle mass $m > 1.91$ keV is derived in the Thomas-Fermi approach (DdVS 2013b) from the lightest known dwarf galaxies. This value is independent of the WDM particle physics model.

The main fermionic WDM particle candidate is a sterile neutrino in the keV scale. Many models of sterile neutrinos are available by now [see for a recent review Merle (2013)]. Another fermionic WDM particle candidate is a gravitino in the keV scale.

In summary, all the small structure formation constraints on the WDM particle mass (Lovell et al. 2012, 2013; Menci et al. 2013; Nierenberg et al. 2013; Pacucci et al. 2013; Papastergis et al. 2011; Zavala et al. 2009), as well as the bounds from sterile neutrino decay into X-rays (Watson et al. 2012) favours a WDM particle mass approximately in the 2 - 3 keV range.

Bounds on the WDM particle mass from Lyman- α forest data (Viel et al. 2013) may be not so reliable since they are affected by the difficult-to-characterize non-linear growth of baryonic and DM structures (Watson et al. 2012). Besides these systematic effects from the Lyman- α data, there are uncertainties in the WDM simulations themselves mainly originated from the uncertainty on the chosen initial velocity dispersion for the particles in the simulations whose effective mass is about $10^5 M_\odot = 10^{68}$ keV each, as discussed by several authors (Lovell et al. 2012, 2013; Viel et al. 2013; Macciò et al. 2012). Namely, the effective particles in the WDM simulations are about 10^{68} times heavier than the real WDM particles. This makes difficult to infer the initial velocity distribution of the effective particles from the known initial velocity distribution of the real WDM particles.

The Lyman- α mass bounds are usually given for the thermal relic mass. This is the mass of the WDM particle if it decouples in thermal equilibrium, which is normally not the case for sterile neutrinos. The relation between the physical particle mass and the thermal mass has to be worked out explicitly for each specific particle physics model. About hundred sterile neutrino models are available today (Merle 2013) for which the Lyman- α bounds are not known. Therefore, it is not possible so far to provide precise generic Lyman- α bounds on the WDM sterile neutrino mass. At present, Lyman- α bounds on the WDM

particle mass are only available for a few specific particle physics models. For discussions on small structure formation in WDM and Lyman- α bounds on the WDM particle mass see Chalonge Colloquium (2012); Meudon Colloquium (2012).

The results presented in this paper do not depend on the precise value of the WDM particle mass m but only on the fact that m is in the keV scale.

One can determine the keV scale of the DM particle mass m but not its precise value within the keV scale just from the core radius value of dilute galaxies, those with $M_h > 10^6 M_\odot$ used in this paper.

The aim of this paper is to show that the obtained rotation curves and density profiles in the Thomas-Fermi galaxy structure theory are in well agreement with the galaxy data parametrized with Burkert profiles.

We follow here the Thomas-Fermi approach to galaxy structure for self-gravitating fermionic WDM (DdVS 2013a,b). This approach is especially appropriate to take into account quantum properties of systems with large number of particles. That is, macroscopic quantum systems as neutron stars and white dwarfs (Landau & Lifshits 1980). In this approach, the central quantity to derive is the DM chemical potential $\mu(\mathbf{r})$, which is the free energy per particle. For self-gravitating systems, the potential $\mu(\mathbf{r})$ is proportional to the gravitational potential $\phi(\mathbf{r})$,

$$\mu(\mathbf{r}) = \mu_0 - m\phi(\mathbf{r}), \quad (1)$$

μ_0 being a constant, and obeys the **self-consistent** and **nonlinear** Poisson equation

$$\nabla^2 \mu(\mathbf{r}) = -4\pi g G m^2 \int \frac{d^3 p}{(2\pi \hbar)^3} f\left(\frac{p^2}{2m} - \mu(\mathbf{r})\right). \quad (2)$$

Here G is Newton's gravitational constant, g is the number of internal degrees of freedom of the DM particle, p is the DM particle momentum and $f(E)$ is the energy distribution function. This is a semiclassical gravitational approach to determine selfconsistently the gravitational potential of the fermionic WDM given its distribution function $f(E)$.

In the Thomas-Fermi approach, DM dominated galaxies are considered in a stationary state. This is a realistic situation for the late stages of structure formation since the free-fall (Jeans) time t_{ff} for galaxies is much shorter than the age of galaxies. t_{ff} is at least one or two orders of magnitude smaller than the age of the galaxy.

We consider spherical symmetric configurations where eq.(2) becomes an ordinary nonlinear differential equation that determines self-consistently the chemical potential $\mu(r)$ and constitutes the Thomas-Fermi approach (DdVS 2013a,b). We choose for the energy distribution function a Fermi-Dirac distribution

$$f(E) = \frac{1}{e^{E/E_0} + 1},$$

where E_0 is the characteristic one-particle energy scale. E_0 plays the role of an effective temperature scale and depends on the galaxy mass. The Fermi-Dirac distribution function is justified in the inner regions of the galaxy, inside the halo radius where we find that the Thomas-Fermi density profiles perfectly agree with the observational data modeled with the empirical Burkert profile.

Observations show that the DM angular momentum is small. In spirals we have a direct proof of this fact from their bottom up general scenario of formation. In these objects we can compute from observations the disk angular momentum, if the angular momentum per unit mass is conserved during the process of disk formation, the values found imply that DM halos are not dominated by rotation (Tonini et al. 2006). Therefore, the spherical symmetric approximation makes sense. Indeed, our results confirm the consistency of such assumption.

In this paper spherical symmetry is considered for simplicity to determine the essential physical galaxy properties as the classical or quantum nature of galaxies, compact or dilute galaxies, the phase space density values, the cored nature of the mass density profiles, the galaxy masses and sizes. It is clear that DM halos are not perfectly spherical but describing them as spherically symmetric is a first approximation to which other effects can be added. In DdVS (2013a) we estimated the angular momentum effect and this yields small corrections.

Our spherically symmetric treatment captures the essential features of the gravitational dynamics and agree with the observations. Notice that we are treating the DM particles quantum mechanically through the Thomas-Fermi approach, so that expectation values are independent of the angles (spherical symmetry) but the particles move and fluctuate in all directions in totally non-spherically symmetric ways. Namely, this is more than treating purely classical orbits for particles in which only radial motion is present. The Thomas-Fermi approach to galaxies can be generalized to describe non-spherically symmetric and non-isotropic situations, by considering distribution functions which include other particle parameters like the angular momentum.

The solutions of the Thomas-Fermi equations (2) are characterized by the value of the chemical potential at the origin $\mu(0)$. Large positive values of $\mu(0)$ correspond to dwarf compact galaxies (fermions near the quantum degenerate limit), while large negative values of $\mu(0)$ yield large and dilute galaxies (classical Boltzmann regime).

Approaching the classical diluted limit yields larger and larger halo radii, galaxy masses and velocity dispersions. On the contrary, in the quantum degenerate limit we get solutions of the Thomas-Fermi equations corresponding to the **minimal** halo radii, galaxy masses and velocity dispersions.

The surface density

$$\Sigma_0 \equiv r_h \rho_0 \simeq 120 M_\odot/\text{pc}^2 \quad \text{up to } 10\% - 20\%, \quad (3)$$

has the remarkable property of being nearly **constant** and independent of luminosity in different galactic systems (spirals, dwarf irregular and spheroidals, elliptics) spanning over 14 magnitudes in luminosity and over different Hubble types (Donato et al. 2009; Spano et al. 2008). It is therefore a useful characteristic scale to express galaxy magnitudes.

To reproduce the smaller observed structures the WDM particle mass should be in the keV scale. We choose the value 2 keV as references scale to express physical magnitudes.

In this paper, we compute the circular velocity in the Thomas-Fermi approach using its expression in terms of the

chemical potential:

$$v_c(r) = \sqrt{\frac{G M(r)}{r}} = \sqrt{-\frac{r}{m} \frac{d\mu}{dr}},$$

On the other hand, the circular velocities of galaxies are known with precision from the observational data from the kinematics of thousands disk galaxies and from the information arising from other tracers of the gravitational field of galaxies as the dispersion velocities of spheroidals and weak lensing measurements (Salucci et al. (2007) and references therein). All this evidence shows that an empirical Burkert profile

$$\rho_B(r) = \frac{\rho_0}{\left(1 + \frac{r}{r_h}\right) \left[1 + \left(\frac{r}{r_h}\right)^2\right]}, \quad (4)$$

correctly reproduces the observations out to the galaxy virial radius.

In this paper, we contrast the observational curves for the circular velocities of galaxies $V_{URC,h}(r)$ and the density profiles obtained from observations using the empirical Burkert profile eq.(4) with the theoretical results $v_c(r)$ and $\rho(r)$ arising from the resolution of the Thomas-Fermi equations.

Our theoretical results follow solving the self-consistent and nonlinear Poisson equation eq.(2) which is **solely** derived from the purely **gravitational** interaction of the WDM particles and their **fermionic** nature. All results are valid for self-gravitating fermionic WDM particles which are assumed stable (or with a lifetime of the order or longer than the Hubble time). The non-gravitational interactions of the WDM particles are assumed weak enough to satisfy the particle accelerator bounds and beta decay bounds. Except for these general WDM particle properties, the framework described here does not require any particular particle physics model of WDM production. All the results reported here are independent of the details of the WDM particle physics model as the symmetry group and the values of the weak enough particle couplings.

The theoretical rotation curves and density profiles well reproduce inside the halo radius the observational curves described by the empirical Burkert profile, obtained from ten different and independent sets of data for galaxy masses from $5 \times 10^9 M_\odot$ till $5 \times 10^{11} M_\odot$.

Our theoretical circular velocities and density profiles exhibit the **universal** property as the observational curves do, and in addition, they **coincide** with the observational curves described by the empirical Burkert profile for $r \lesssim 2 r_h$.

In summary, the results presented in this paper show the ability of the Thomas-Fermi approach to correctly describe the galaxy structures.

This paper is organized as follows. In Section 2 we present the Thomas-Fermi approach to galaxy structure and we express the main galaxy magnitudes in terms of the solution of the Thomas-Fermi equation and the value of the surface density Σ_0 . We discuss the theoretical circular velocity curves, the theoretical density profiles and the remarkable universality of them. In Section 3 we present the contrast between the observational and theoretical curves for the galaxy

circular velocities and the density profiles and we discuss the universal property of these profiles. Section 4 is devoted to our conclusions.

2 GALAXY PROPERTIES IN THE THOMAS-FERMI WDM APPROACH

We consider DM dominated galaxies in their late stages of structure formation when they are relaxing to a stationary situation, at least not too far from the galaxy center.

This is a realistic situation since the free-fall (Jeans) time t_{ff} for galaxies is much shorter than the age of galaxies:

$$t_{ff} = \frac{1}{\sqrt{G \rho_0}} = 1.49 \cdot 10^7 \sqrt{\frac{M_\odot}{\rho_0 \text{ pc}^3}} \text{ yr}.$$

The observed central densities of galaxies yield free-fall times in the range from 15 million years for ultracompact galaxies till 330 million years for large dilute spiral galaxies. These free-fall (or collapse) times are small compared with the age of galaxies running in billions of years.

Hence, we can consider the DM described by a time independent and non-relativistic energy distribution function $f(E)$, where $E = p^2/(2m) - \mu$ is the single-particle energy, m is the mass of the DM particle, μ is the chemical potential (DdVS 2013a,b), related to the gravitational potential $\phi(\mathbf{r})$ by eq.(1).

In the Thomas-Fermi approach, $\rho(\mathbf{r})$ is expressed as a function of $\mu(\mathbf{r})$ through the standard integral of the DM phase-space distribution function over the momentum

$$\rho(\mathbf{r}) = \frac{g m}{2 \pi^2 \hbar^3} \int_0^\infty dp p^2 f\left(\frac{p^2}{2m} - \mu(\mathbf{r})\right), \quad (5)$$

where g is the number of internal degrees of freedom of the DM particle, with $g = 1$ for Majorana fermions and $g = 2$ for Dirac fermions. For definiteness, we will take $g = 2$ in the sequel.

We will consider spherical symmetric configurations. Then, the Poisson equation for $\phi(r)$ takes the self-consistent form

$$\begin{aligned} \frac{d^2 \mu}{dr^2} + \frac{2}{r} \frac{d\mu}{dr} &= -4\pi G m \rho(r) = \\ &= -\frac{4 G m^2}{\pi \hbar^3} \int_0^\infty dp p^2 f\left(\frac{p^2}{2m} - \mu(r)\right), \end{aligned} \quad (6)$$

where G is Newton's constant and $\rho(r)$ is the DM mass density.

Eq. (6) provides an ordinary **nonlinear** differential equation that determines **self-consistently** the chemical potential $\mu(r)$ and constitutes the Thomas-Fermi approach (DdVS 2013a,b). This is a semi-classical approach to galaxy structure in which the quantum nature of the DM particles is taken into account through the quantum statistical distribution function $f(E)$.

The DM pressure and the velocity dispersion can also be expressed as integrals over the DM phase-space distribution function as

$$P(r) = \frac{1}{3 \pi^2 m \hbar^3} \int_0^\infty dp p^4 f\left(\frac{p^2}{2m} - \mu(r)\right), \quad (7)$$

$$\langle v^2 \rangle(r) = \frac{1}{m^2} \frac{\int_0^\infty dp p^4 f\left(\frac{p^2}{2m} - \mu(r)\right)}{\int_0^\infty dp p^2 f\left(\frac{p^2}{2m} - \mu(r)\right)} = 3 \frac{P(r)}{\rho(r)}.$$

The fermionic DM mass density ρ is bounded at the origin due to the Pauli principle (DdVS 2013a) which implies the bounded boundary condition at the origin

$$\frac{d\mu}{dr}(0) = 0. \quad (8)$$

We see that $\mu(r)$ fully characterizes the DM halo structure in this Thomas-Fermi framework. The chemical potential is monotonically decreasing in r since eq.(6) implies

$$\frac{d\mu}{dr} = -\frac{GmM(r)}{r^2}, \quad M(r) = 4\pi \int_0^r dr' r'^2 \rho(r'). \quad (9)$$

In this semi-classical framework the stationary energy distribution function $f(E)$ must be given. We consider the Fermi-Dirac distribution

$$f(E) = \Psi_{FD}(E/E_0) = \frac{1}{e^{E/E_0} + 1}, \quad (10)$$

where the characteristic one-particle energy scale E_0 in the DM halo plays the role of an effective temperature. The value of E_0 depends on the galaxy mass. In neutron stars, where the neutron mass is about six orders of magnitude larger than the WDM particle mass, the temperature can be approximated by zero. In galaxies, $E_0 \sim m \langle v^2 \rangle$ turns to be non-zero but small in the range: $10^{-3} \text{ K} \lesssim E_0 \lesssim 50 \text{ K}$ which reproduce the observed velocity dispersions for $m \sim 2 \text{ keV}$. The smaller values of E_0 correspond to compact dwarf galaxies and the larger values of E_0 are for large and dilute galaxies.

Notice that for the relevant galaxy physical magnitudes, the Fermi-Dirac distribution function gives similar results to out of equilibrium distribution functions (DdVS 2013a).

The choice of Ψ_{FD} is justified in the inner regions, where relaxation to thermal equilibrium is possible. Far from the origin however, the Fermi-Dirac distribution as its classical counterpart, the isothermal sphere, produces a mass density tail $1/r^2$ that overestimates the observed tails of the galaxy mass densities. Indeed, the classical regime $\mu/E_0 \rightarrow -\infty$ is attained for large distances r since eq.(9) indicates that $\mu(r)$ is always monotonically decreasing with r .

More precisely, large positive values of the chemical potential at the origin correspond to the degenerate fermions limit which is the extreme quantum case and oppositely, large negative values of the chemical potential at the origin gives the diluted case which is the classical regime. The quantum degenerate regime describes dwarf and compact galaxies while the classical and diluted regime describes large and diluted galaxies. In the classical regime, the Thomas-Fermi equation (6)-(8) become the equations for a self-gravitating Boltzmann gas.

It is useful to introduce dimensionless variables ξ , $\nu(\xi)$

$$r = l_0 \xi, \quad \mu(r) = E_0 \nu(\xi), \quad (11)$$

where l_0 is the characteristic length that emerges from the dynamical equation (6):

$$l_0 \equiv \frac{\hbar}{\sqrt{8G}} \left[\frac{9\pi I_2(\nu_0)}{m^8 \rho_0} \right]^{\frac{1}{6}} = R_0 \left(\frac{2 \text{ keV}}{m} \right)^{\frac{4}{3}} \left[\frac{I_2(\nu_0)}{\rho_0} \frac{M_\odot}{\text{pc}^3} \right]^{\frac{1}{6}},$$

$$R_0 = 7.425 \text{ pc}, \quad (12)$$

and

$$I_n(\nu) \equiv (n+1) \int_0^\infty y^n dy \Psi_{FD}(y^2 - \nu), \quad n = 1, 2, \dots, \quad \nu_0 \equiv \nu(0), \quad \rho_0 = \rho(0), \quad (13)$$

where we use the integration variable $y \equiv p/\sqrt{2m E_0}$.

Then, in dimensionless variables, the self-consistent Thomas-Fermi equation (6) for the chemical potential $\nu(\xi)$ takes the form

$$\frac{d^2 \nu}{d\xi^2} + \frac{2}{\xi} \frac{d\nu}{d\xi} = -I_2(\nu), \quad \nu'(0) = 0. \quad (14)$$

We solve eq.(14) numerically by using as independent variable $u \equiv \ln \xi$ and then applying the fourth-order Runge-Kutta method. We solve eq.(14) for a broad range of values $\nu_0 \equiv \nu(0)$, from negative values $\nu_0 \lesssim -5$ describing galaxies in dilute regimes to positive values $\nu_0 \gtrsim 1$ corresponding to compact dwarf galaxies.

We find the main physical galaxy magnitudes, such as the mass density $\rho(r)$, the velocity dispersion $\sigma^2(r) = v^2(r)/3$ and the pressure $P(r)$, which are all r -dependent as:

$$\rho(r) = \rho_0 \frac{I_2(\nu(\xi))}{I_2(\nu_0)}, \quad \rho_0 = \frac{1}{3\pi^2} \frac{m^4}{\hbar^3} \left(\frac{2 E_0}{m} \right)^{3/2} I_2(\nu_0),$$

$$\sigma(r) = \sqrt{\frac{2 E_0}{5 m} \frac{I_4(\nu(\xi))}{I_2(\nu(\xi))}} = \quad (15)$$

$$= 1.092 \sqrt{\frac{I_4(\nu(\xi))}{I_2(\nu(\xi))}} \left(\frac{2 \text{ keV}}{m} \right)^{\frac{4}{3}} \left[\frac{\rho_0}{I_2(\nu_0)} \frac{\text{pc}^3}{M_\odot} \right]^{\frac{1}{3}} \frac{\text{km}}{\text{s}}, \quad (16)$$

$$P(r) = \frac{2 E_0}{5 m} \rho_0 \frac{I_4(\nu(\xi))}{I_2(\nu_0)} = \frac{1}{5} \left(\frac{3\pi^2 \hbar^3}{m^4} \right)^{\frac{2}{3}} \left[\frac{\rho_0}{I_2(\nu_0)} \right]^{5/3} I_4(\nu(\xi)), \quad (17)$$

As a consequence, from eqs.(9), (11), (12), (14) and (15) the total mass $M(r)$ enclosed in a sphere of radius r and the phase space density $Q(r)$ turn to be

$$\begin{aligned} M(r) &= 4\pi \frac{\rho_0 l_0^3}{I_2(\nu_0)} \int_0^\xi dx x^2 I_2(\nu(x)) = \\ &= 4\pi \frac{\rho_0 l_0^3}{I_2(\nu_0)} \xi^2 |\nu'(\xi)| = \\ &= M_0 \xi^2 |\nu'(\xi)| \left(\frac{\text{keV}}{m} \right)^4 \sqrt{\frac{\rho_0}{I_2(\nu_0)} \frac{\text{pc}^3}{M_\odot}}, \\ M_0 &= 4\pi M_\odot \left(\frac{R_0}{\text{pc}} \right)^3 = 0.8230 \cdot 10^5 M_\odot, \end{aligned} \quad (18)$$

$$Q(r) \equiv \frac{\rho(r)}{\sigma^3(r)} = 3\sqrt{3} \frac{\rho(r)}{\langle v^2 \rangle^{\frac{3}{2}}(r)} =$$

$$= \frac{\sqrt{125}}{3 \pi^2} \frac{m^4}{\hbar^3} \frac{I_2^{\frac{5}{2}}(\nu(\xi))}{I_4^{\frac{3}{2}}(\nu(\xi))}. \quad (19)$$

We have systematically eliminated the energy scale E_0 in terms of the central density ρ_0 through eq.(15). Notice that $Q(r)$ turns to be independent of E_0 and therefore from ρ_0 .

We define the core size r_h of the halo by analogy with the empirical Burkert density profile as

$$\frac{\rho(r_h)}{\rho_0} = \frac{1}{4} \quad , \quad r_h = l_0 \xi_h. \quad (20)$$

It must be noticed that the surface density

$$\Sigma_0 \equiv r_h \rho_0, \quad (21)$$

is found nearly **constant** and independent of luminosity in different galactic systems (spirals, dwarf irregular and spheroidals, elliptics) spanning over 14 magnitudes in luminosity and over different Hubble types. More precisely, all galaxies seem to have the same value for Σ_0 , namely $\Sigma_0 \simeq 120 M_\odot/\text{pc}^2$ up to 10% – 20% (Donato et al. 2009; Spano et al. 2008; Kormendy & Freeman 2004). It is remarkable that at the same time other important structural quantities as r_h , ρ_0 , the baryon-fraction and the galaxy mass vary orders of magnitude from one galaxy to another.

The constancy of Σ_0 seems unlikely to be a mere coincidence and probably reflects a physical scaling relation between the mass and halo size of galaxies. It must be stressed that Σ_0 is the only dimensionful quantity which is constant among the different galaxies.

We use here the dimensionful quantity Σ_0 to set the energy scale in the Thomas-Fermi approach. That is, we replace the central density ρ_0 in eqs.(11), (12), (15) and (18) in terms of Σ_0 eq.(21) with the following results

$$\begin{aligned} l_0 &= \frac{\hbar^{\frac{6}{5}}}{G^{\frac{3}{5}}} \left(\frac{9 \pi}{512} \right)^{\frac{1}{5}} \left[\frac{\xi_h I_2(\nu_0)}{m^8 \Sigma_0} \right]^{\frac{1}{5}} = \\ &= 4.2557 [\xi_h I_2(\nu_0)]^{\frac{1}{5}} \left(\frac{2 \text{ keV}}{m} \right)^{\frac{8}{5}} \left(\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2} \right)^{\frac{1}{5}} \text{ pc}, \\ E_0 &= \hbar^{\frac{6}{5}} \frac{G^{\frac{2}{5}}}{m^{\frac{3}{5}}} (18 \pi^6)^{\frac{1}{5}} \left[\frac{\Sigma_0}{\xi_h I_2(\nu_0)} \right]^{\frac{4}{5}} = \\ &= \frac{7.12757 \cdot 10^{-3}}{[\xi_h I_2(\nu_0)]^{\frac{4}{5}}} \left(\frac{2 \text{ keV}}{m} \right)^{\frac{3}{5}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{4}{5}} \text{ K}, \end{aligned} \quad (22)$$

and

$$r = 4.2557 \xi [\xi_h I_2(\nu_0)]^{\frac{1}{5}} \left(\frac{2 \text{ keV}}{m} \right)^{\frac{8}{5}} \left(\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2} \right)^{\frac{1}{5}} \text{ pc} \quad (23)$$

$$\rho(r) = 18.1967 \frac{I_2(\nu(\xi))}{[\xi_h I_2(\nu_0)]^{\frac{6}{5}}} \left(\frac{m}{2 \text{ keV}} \right)^{\frac{8}{5}} \left(\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2} \right)^{\frac{6}{5}} \frac{M_\odot}{\text{pc}^3},$$

$$M(r) = \frac{27312 \xi^2}{[\xi_h I_2(\nu_0)]^{\frac{3}{5}} |\nu'(\xi)|} \left(\frac{2 \text{ keV}}{m} \right)^{\frac{16}{5}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{3}{5}} M_\odot. \quad (24)$$

For a fixed value of the surface density Σ_0 , the solutions of

the Thomas-Fermi eqs.(14) are parametrized by a single parameter: the dimensionless chemical potential at the center ν_0 . That is, ν_0 is determined by the value of the halo galaxy mass

$$M_h \equiv M(r_h) \quad (25)$$

In the classical dilute limit, $\nu_0 \lesssim -5$, the analytic expressions for the main galaxies magnitudes are given by:

$$\xi_h = \frac{3.147473}{e^{\nu_0/2}} \quad , \quad \left. \frac{d\nu}{d \ln \xi} \right|_{\xi_h} = -1.839957 \quad , \quad (26)$$

$$M_h = 1.75572 \Sigma_0 r_h^2 \quad ,$$

$$M_h = \frac{67011.4}{e^{\frac{4}{5} \nu_0}} \left(\frac{2 \text{ keV}}{m} \right)^{\frac{16}{5}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{3}{5}} M_\odot,$$

$$r_h = 68.894 \sqrt{\frac{M_h}{10^6 M_\odot}} \sqrt{\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2}} \text{ pc}, \quad (27)$$

$$r = 22.728 \xi \left(\frac{M_\odot}{M_h} \right)^{\frac{1}{8}} \left(\frac{2 \text{ keV}}{m} \right)^2 \left(\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2} \right)^{\frac{1}{8}} \text{ pc}, \quad (28)$$

$$\rho(r) = 5.195045 \left(\frac{M_h}{10^4 M_\odot} \right)^{\frac{3}{4}} \left(\frac{m}{2 \text{ keV}} \right)^4 \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{3}{4}} \times e^{\nu(\xi)} \frac{M_\odot}{\text{pc}^3}, \quad (29)$$

$$\begin{aligned} M(r) &= 179.30 \left(\frac{M_h}{M_\odot} \right)^{\frac{3}{4}} \left(\frac{2 \text{ keV}}{m} \right)^2 \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{3}{8}} \times \\ &\xi \left| \frac{d\nu(\xi)}{d \ln \xi} \right| M_\odot, \end{aligned} \quad (30)$$

$$Q(0) = 1.2319 \left(\frac{10^5 M_\odot}{M_h} \right)^{\frac{5}{4}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{3}{4}} \text{ keV}^4. \quad (31)$$

These equations are accurate for $M_h \gtrsim 10^6 M_\odot$. We see that they exhibit a **scaling** behaviour for r_h vs. M_h , Q_h and $Q(0)$ vs. M_h and M_h vs. the fugacity at the center $z_0 = e^{\nu_0}$. These scaling behaviours are **very accurate** except near the degenerate limit as shown by fig. 1.

It must be stressed that (i) the scaling relations eqs.(26)-(31) are a consequence solely of the self-gravitating interaction of the fermionic WDM and (ii) the value of the WDM particle mass $m \simeq 2 \text{ keV}$ appears in the proportionality factors which is therefore confirmed by the galaxy data (see figs. 7 and 8).

In eqs.(23)-(31) we use the surface density Σ_0 as energy scale to express the theoretical results [we used the central density ρ_0 in refs. (DdVS 2013a,b)]. It is highly remarkable that our theoretical results **reproduce** the observed DM halo properties with **good precision**.

The limit opposite limit, $\nu_0 \gtrsim 1$ is the extreme quantum limit corresponding to degenerate WDM fermions. The galaxy mass and halo radius take in the degenerate limit




Figure 1. The ordinary logarithm of the theoretical halo radius \hat{r}_h vs. the ordinary logarithm of the halo mass \hat{M}_h for small galaxy masses in the Thomas-Fermi approach from eqs.(20), (24), (25), (46) and (47) (dashed green line) and the dilute regime eq.(26)-(27) (red continuous line). \hat{r}_h and \hat{M}_h are defined by eqs.(46)-(47). The dilute regime approximates very well the exact Thomas-Fermi results, r_h follows the square-root of M_h scaling behaviour of the classical regime eq.(26)-(27). This is so even near the fermion degenerate quantum limit.

their **minimum** values

$$r_h^{min} = 11.3794 \left(\frac{2 \text{ keV}}{m} \right)^{\frac{8}{5}} \left(\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2} \right)^{\frac{1}{5}} \text{ pc},$$

$$M_h^{min} = 30998.7 \left(\frac{2 \text{ keV}}{m} \right)^{\frac{16}{5}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{3}{5}} M_\odot, \quad (32)$$

while the phase-space density $Q(r)$ takes its **maximum** value

$$Q_h^{max} = 16 \frac{\sqrt{125}}{3 \pi^2} \left(\frac{m}{2 \text{ keV}} \right)^4 \text{ keV}^4 = 6.041628 \left(\frac{m}{2 \text{ keV}} \right)^4 \text{ keV}^4. \quad (33)$$

But the Thomas-Fermi equations provide a whole continuous range of galaxy solutions above the degenerate limit as discussed above and in DdVS (2013a,b).

The degenerate limit corresponds to $E_0 = 0$. In the classical dilute limit $M_h \gtrsim 10^6 M_\odot$, E_0 runs approximately from 0.02 K to 20 K.

2.1 The galaxy circular velocities

We consider now the circular velocity $v_c(r)$ defined through the virial theorem as

$$v_c(r) \equiv \sqrt{\frac{G M(r)}{r}}. \quad (34)$$




Figure 2. Normalized circular velocities $U(x) = v_c(r)/v_c(r_h)$ vs. $x = r/r_h$ from the observational data eq.(43) and from the theoretical Thomas-Fermi formula eq.(35). The theoretical curves from the Thomas-Fermi approach for ten different galaxy masses all **fall** one into each other providing an **universal** rotation curve which practically **coincides** with the observational universal curve URC for $x = r/r_h \lesssim 2$.

The circular velocity is directly related by eq.(9) to the derivative of the chemical potential

$$v_c(r) = \sqrt{-\frac{r}{m} \frac{d\mu}{dr}},$$

which in dimensionless variables takes the form

$$v_c(r) = \sqrt{-\frac{E_0}{m} \frac{d\nu}{d \ln \xi}}.$$

Expressing the energy scale E_0 in terms of the surface density using eq.(22) we have for the circular velocity the explicit expression

$$v_c(r) = 5.2537 \frac{\sqrt{-\xi \nu'(\xi)}}{[\xi_h I_2(\nu_0)]^{\frac{2}{5}}} \left(\frac{2 \text{ keV}}{m} \right)^{\frac{4}{5}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{2}{5}} \frac{\text{km}}{\text{s}}. \quad (35)$$

In the dilute Boltzmann regime the circular velocity at the core radius r_h scales as the power 1/4 of the galaxy halo mass M_h :

$$v_c(r_h) = 7.901 \left(\frac{M_h}{10^6 M_\odot} \right)^{\frac{1}{4}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{1}{4}} \frac{\text{km}}{\text{s}}. \quad (36)$$

It is important to consider the circular velocity normalized to unit at the core radius r_h

$$U(x) \equiv \frac{v_c(r)}{v_c(r_h)} = \sqrt{x \frac{\nu'(\xi)}{\nu'(\xi_h)}}, \quad x = \frac{r}{r_h}. \quad (37)$$

Explicitly solving the Thomas-Fermi eqs.(14) we find that $v_c(r)/v_c(r_h)$ is **only** function of $x = r/r_h$ and takes the

same values for all galaxy masses in the range going from $5.13 \cdot 10^9 M_\odot$ till $5.15 \cdot 10^{11} M_\odot$ as shown in fig. 2. Namely, $U(x)$ turns to be an **universal** function.

This is a remarkable result since a priori, $v_c(r)/v_c(r_h)$ could be a function of r and r_h and could be different for different galaxies.

This important result shows the ability of the Thomas-Fermi approach to correctly describe the galaxy structures.

The phase-space density $Q(r)$ can be also obtained from the circular velocity at the radius r as

$$Q_c(r) = 3 \sqrt{3} \frac{\rho(r)}{v_c^3(r)} \quad , \quad Q_{ch} = 3 \sqrt{3} \frac{\rho(r_h)}{v_c^3(r_h)} \quad (38)$$

Expressing $\rho(r_h)$ in terms of r_h eq.(26), Σ_0 eq.(21) and $v_c(r_h)$ from eq.(35) yields for the phase-space density at the halo radius:

$$Q_{ch} = 7.96204 \frac{I_2(\nu_0)}{[-\xi_h \nu'(\xi_h)]^{\frac{3}{2}}} \left(\frac{m}{2 \text{ keV}} \right)^4 \text{ keV}^4 \quad (39)$$

The numerical values of Q_{ch} turn to be larger than $Q(0)$ eq.(19) approximately by a factor of two. In the dilute regime $M_h \gtrsim 10^6 M_\odot$ they are related by

$$Q_{ch} = 2.0873 Q(0) \quad (40)$$

3 CIRCULAR VELOCITIES CONTRASTED WITH OBSERVATIONS

The circular velocities values $v_c(r)$ eq.(34) are known with precision from galaxy observational data.

3.1 The galaxy data

The kinematics of about several thousands disk galaxies, described by the Rotation Curves of Spirals, and the information obtained from other tracers of the gravitational field of galaxies, including the dispersion velocities of spheroidals and the weak lensing measurements (Salucci et al. 2007 and references therein) show that the density of the dark matter halos around galaxies of different kinds, different luminosity and Hubble types is well represented, out to the galaxy virial radius, by an empirical Burkert profile

$$\rho_B(r) = \rho_{0B} F_B \left(\frac{r}{r_{hB}} \right) \quad , \quad F_B(x) = \frac{1}{(1+x)(1+x^2)} \quad , \quad x \equiv \frac{r}{r_{hB}} \quad (41)$$

where ρ_0 stands for the central core density and r_{hB} for the core radius. The empirical Burkert profile satisfactorily fits the astronomical observations and we use the observed data of ρ_{0B} and r_{hB} for DM dominated spiral galaxies given in (Salucci et al. 2007).

Kinematical data and properties of other galaxy gravitational potential tracers are all reproduced, within their observational uncertainty by a mass model including a DM halo with a Burkert profile (see Donato et al. (2009); Salucci et al. (2007)). While some other cored DM distributions [but not the pseudo-isothermal one $\rho_0/(r^2 + a^2)$] may successfully reproduce these data, every cuspy distribution fails to do so (see e. g. Gentile et al. (2004)).

The Burkert and the Peñarrubia et al. (2012) profiles are indistinguishable with the data available at present. Any cored density profile with two free parameters (central density, core radius) that decreases faster than $1/r^2$ can be mapped into each other, just by a transformation of the parameters. Only when the determination of the DM density profile will be available at the few per cent error level (today it is 10-30 percent) we would be able to discriminate between different cored the profiles.

The circular velocities $V_{URC,h}(r)$ for the empirical Burkert density profile follow from eq.(34) and eq.(41) (Salucci et al. 2007)

$$V_{URC,h}^2(r) = 2\pi G \frac{\rho_{0B} r_{hB}^3}{r} \times \left[\ln(1+x) - \arctan x + \frac{1}{2} \ln(1+x^2) \right] \quad , \quad x = \frac{r}{r_{hB}} \quad (42)$$

Notice that normalizing $V_{URC,h}(r)$ to its value at the core radius r_{hB} yields

$$U_{URC}^2(x) \equiv \frac{V_{URC,h}^2(r)}{V_{URC,h}^2(r_h)} = \frac{3.93201}{x} \left[\ln(1+x) - \arctan x + \frac{1}{2} \ln(1+x^2) \right] \quad (43)$$

Namely, the function $U(x)_{URC}$ only depends on $x = r/r_{hB}$ and complies with the concept of Universal Rotation Curve (URC, Salucci et al. 2007): $U(x)_{URC}$ is an **universal** function.

Notice that the URC concept is valid not only for the Burkert representation of the density profile but also for other density profiles that correctly reproduce the density data.

In the Burkert profile case, the halo galaxy mass follows integrating eq.(41) from zero to r_{hB}

$$M_h = 1.59796 \rho_{0B} r_{hB}^3 = 1.59796 \Sigma_0 r_{hB}^2 \quad (44)$$

This empirical equation can be recasted in a similar form to eq.(26) of the theoretical Thomas-Fermi approach,

$$r_{hB} = 72.215 \sqrt{\frac{120}{\Sigma_0 \text{ pc}^2} \frac{M_h}{10^6}} \text{ pc} \quad (45)$$

r_h in eqs.(26) and r_{hB} (45) refer to the point where $\rho(r_h)/\rho(0) = 1/4$ and $\rho_B(r_{hB})/\rho(0) = 1/4$ both according to eq.(20), for two different density profiles: the theoretical Thomas-Fermi profile $\rho(r)$ in eq.(26) and the empirical Burkert profile $\rho(r)_B$ for the observational data.

The halo radius r_{hB} and r_h for given galaxy mass M_h and surface density Σ_0 are related by the universal relation

$$r_h = 0.95401 r_{hB} \quad ,$$

that follows from eqs.(26) and (44). Namely, due to the slight shape difference between the theoretical Thomas-Fermi and empirical Burkert profiles (see fig. 3), the Thomas-Fermi halo radius turns to be about 5% smaller than the Burkert halo radius.

It follows from eq.(14) that the theoretical Thomas-Fermi profile poses an expansion in even powers of r^2 (this is also the case for the density profiles obtained in the linear approximation from the cosmological density fluctuations in

Table 1. Observed values r_h , $\rho(0)$ and M_h covering from ultracompact galaxies to large spiral galaxies from the references: (Walker & Peñarrubia 2012; Salucci et al. 2007; dVSS 2010; Gilmore et al. 2007; Simon & Geha 2007; Simon et al. 2011; Wolf et al. 2010; Brodie et al. 2011; Willman & Strader 2012; Martinez et al. 2011; Salucci et al. 2012). The errors are generally about 20%-40% and they are explicitly given when available.

Galaxy	$\frac{r_h}{\text{pc}}$	$\rho(0)/\frac{M_\odot}{(\text{pc})^3}$	$\frac{M_h}{10^6 M_\odot}$
Willman 1	33^{+7}_{-8}	6.8 ± 3	$0.39^{+2.5}_{-1.6}$
Segue 1	52 ± 10	$2.5^{+4.1}_{-1.9}$	$0.58^{+8.2}_{-5.1}$
Leo IV	166 ± 160	$.19 \pm 0.2$	1.4 ± 1.5
Canis Venatici II	145 ± 70	0.49 ± 0.25	2.4 ± 1.1
Coma-Berenices	109 ± 40	2.09 ± 0.86	1.2 ± 0.4
Leo II	57.5^{+166}_{-51}	$4.07^{+530}_{-3.5}$	14.1 ± 0.3
Leo T	170 ± 23	0.79 ± 0.36	12.9 ± 7
Hercules	354 ± 140	0.1 ± 0.04	7.1 ± 2.6
Carina	603^{+545}_{-287}	$0.065^{+0.07}_{-0.03}$	11.7 ± 0.3
Ursa Major I	458 ± 150	0.25 ± 0.08	15 ± 4
Draco	646^{+401}_{-266}	$0.18^{+0.12}_{-0.06}$	20.4 ± 0.5
Leo I	282^{+255}_{-150}	$0.41^{+0.94}_{-0.23}$	38 ± 0.6
Sculptor	355^{+124}_{-80}	$0.25^{+0.1}_{-0.08}$	35.5 ± 0.3
Boötes I	406 ± 36	$0.22^{+0.32}_{-0.12}$	23.6^{+20}_{-10}
Canis Venatici I	596 ± 150	0.08 ± 0.02	27 ± 4
Sextans	46.8^{+115}_{-34}	$5.5^{+130}_{-5.2}$	32.4 ± 0.7
Ursa Minor	245^{+317}_{-156}	$0.41^{+1.9}_{-0.3}$	36.3 ± 0.5
Fornax	372^{+85}_{-90}	$0.19^{+0.12}_{-0.06}$	126 ± 0.06
NGC 185	450	4.09	975 ± 300
NGC 855	1063	2.64	8340 ± 1600
Small Spiral URC	5100 ± 1550	0.029 ± 0.001	6900 ± 2300
NGC 4478	1890	3.7	$6.55 \times 10^4 \pm 7 \times 10^3$
Medium Spiral URC	$1.9 \times 10^4 \pm 6 \times 10^3$	0.0076 ± 0.002	$1.01 \times 10^5 \pm 3 \times 10^4$
NGC 731	6160	0.47	$2.87 \times 10^5 \pm 3 \times 10^4$
NGC 3853	5220	0.77	$2.87 \times 10^5 \pm 4.5 \times 10^4$
NGC 499	7700	0.91	$1.09 \times 10^6 \pm 2 \times 10^5$
Large Spiral URC	$5.9 \times 10^4 \pm 1.8 \times 10^3$	$2.3 \times 10^{-3} \pm 7 \times 10^{-4}$	$1. \times 10^6 \pm 3 \times 10^5$

dVSS (2010)). On the contrary, the empirical Burkert profile eq.(41) is not an even function of r and exhibits a linear behaviour in r near the origin. This is the source of the small deviation near $r = 0$ between the theoretical Thomas-Fermi profile and the empirical Burkert profile exhibited in fig. 3.

The circular velocity at the halo radius for the empirical Burkert profile follows setting $x = 1$ in eq.(42) with the

result

$$V_{URC,h}(r_h) = 7.717 \left(\frac{M_h}{10^6 M_\odot} \right)^{\frac{1}{4}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{1}{4}} \frac{\text{km}}{\text{s}}.$$

This value is to be compared with the theoretical Thomas-Fermi result eq.(36). We see that they differ from each other by only 2.4 %, confirming again the success of the Thomas-Fermi approach to describe the galaxy structures.




Figure 3. Normalized density profiles $\rho(r)/\rho(0)$ as functions of r/r_h . We display the theoretical profiles for galaxy masses in the dilute regime $1.4 \cdot 10^5 < \hat{M}_h < 7.5 \cdot 10^{11}$, $-1.5 > \nu_0 > -20.78$. **All fall into the same and universal density profile.** We plot the empirical Burkert profile as function of r/r_h .

3.2 Comparison to observations

Our results are **independent** of the details of the WDM particle physics model. They follow from the gravitational self-interaction of WDM particles and their fermionic nature. The same remarks apply to all the Thomas-Fermi results including the lower bound $m > 1.91$ keV (DdVS 2013b).

We depict in fig. 2 the normalized circular velocities $U(x) = v_c(r)/v_c(r_h)$ vs. $x = r/r_h$ obtained on one hand from the observational data $U(x)_{URC}$ described with the empirical Burkert profile eq.(43), and on the other hand, $U(x)$ obtained from the theoretical Thomas-Fermi formula eq.(35). In general, the normalized circular velocities $v_c(r)/v_c(r_h)$ can be functions of r and r_h , which just reflects the fact that in a spherically symmetric approach only one parameter r_h shows up in the circular velocities. [The parameter r_h depends on the characteristic energy E_0 in the Fermi-Dirac distribution function at fixed surface density Σ_0 . Generalizing the Thomas-Fermi approach to non-spherically symmetric and non-isotropic situations by including other particle parameters like the angular momentum in the distribution functions would lead to density profiles depending on other parameters besides r_h .]

Here we remarkably find that the normalized circular velocities $v_c(r)/v_c(r_h)$ turn out to be functions of **only** one variable: the ratio $x = r/r_h$. [On the contrary, $v_c(r)$ is a function of r **and** r_h separately]. We see that the theoretical curves from the Thomas-Fermi approach for ten different galaxy masses **all fall one into each other**. Therefore, we find the result that the Thomas-Fermi approach provides **universal** rotation curves. Moreover, the theoretical

Thomas-Fermi curves $U(x)$ and the observational universal curve $U(x)_{URC}$ described by the empirical Burkert profile **coincide** for $r < r_h$.

We depict in fig. 3 the normalized density profiles $F(x) = \rho(r)/\rho(0)$ as functions of $x = r/r_h$ obtained from the theoretical Thomas-Fermi profiles for galaxy masses in the dilute regime $1.4 \cdot 10^5 < \hat{M}_h < 7.5 \cdot 10^{11}$, $-1.5 > \nu_0 > -20.78$. **All fall into the same and universal density profile.** The empirical Burkert profile $F_B(x)$ in fig. 3 turns to be very close to the theoretical Thomas-Fermi profile $F(x)$ except near the origin as discussed above.

We display in figs. 4 $v_c(r)$ in km/s vs. r in kpc obtained on one hand from the observational data described with the empirical Burkert profile eq.(42) and on the other hand from the theoretical Thomas-Fermi formula eq.(35). We plot in figs. 4 $v_c(r)$ for $0 < r < r_{vir}$, r_{vir} being the virial radius of the galaxy.

The corresponding halo galaxy masses M_h are indicated in figs. 4 and run from $5.13 \cdot 10^9 M_\odot$ till $5.15 \cdot 10^{11} M_\odot$.

The theoretical rotation curves reproduce the observational curves modeled with the empirical Burkert profile for $r \lesssim r_h$ justifying the use of the Fermi-Dirac distribution function eq.(10) in the Thomas-Fermi eqs.(13)-(14).

We display in figs. 5 and 6 the theoretical density profiles computed from the Thomas-Fermi equations and the observational profiles described by the empirical Burkert expression. We plot the ordinary logarithm of the density in M_\odot/pc^3 vs. r in kpc in the interval $0 < r < 4 r_h$. We see a very good agreement of the theoretical density profiles with the observations modeled with the empirical Burkert profile in all the range $0 < r < 4 r_h$.

In fig. 7 we plot the ordinary logarithm of

$$\hat{r}_h \equiv \frac{r_h}{\text{pc}} \left(\frac{m}{2 \text{ keV}} \right)^{\frac{8}{5}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{1}{5}} \quad (46)$$

versus the ordinary logarithm of

$$\hat{M}_h \equiv \frac{M_h}{M_\odot} \left(\frac{m}{2 \text{ keV}} \right)^{\frac{16}{5}} \left(\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2} \right)^{\frac{3}{5}}. \quad (47)$$

From eqs.(26), (46) and (47), we find the **scaling** relation

$$\log_{10} \hat{r}_h = \frac{1}{2} \log_{10} \hat{M}_h - 1.16182, \quad (48)$$

which is accurate for $M_h \gtrsim 10^6 M_\odot$. We see from fig. 7 that \hat{r}_h follows with precision the square-root law eq. (48) obtained from the dilute regime eq.(26) of the Thomas-Fermi equations.

For m in the keV scale, namely $m \sim 2$ keV, we obtain galaxies as nondegenerate solutions of the Thomas-Fermi equations with core radius from 10 pc to 1 Mpc as shown in fig. 7. Therefore, fermionic WDM remarkably well explains the observations of the Fornax and Sculptor dSphs (Walker & Peñarrubia 2012; Peñarrubia et al. 2012)

Actually, the dilute regime formulas (26)-(31) apply even near the fermion degenerate limit as shown by fig. 1. In fig. 1 we depicted $\log_{10} \hat{r}_h$ vs. $\log_{10} \hat{M}_h$ for the smaller galaxies $\hat{M}_{h,min} \leq \hat{M}_h < 10^5$ where $\hat{M}_{h,min} = 30999$. We see that the dilute regime eqs.(26)-(31) reproduce the Thomas-Fermi results for practically **all** galaxy masses even near the degenerate limit.

We plot in fig. 8 the ordinary logarithm of the theoretical phase-space density Q_{ch}/keV^4 vs. the ordinary logarithm of \hat{M}_h and the observational values of $\log_{10} Q_{chB}/\text{keV}^4$. We see that the theoretical phase-space density Q_{ch} reproduces very well the observational data parametrized with the empirical Burkert profile.

The errors of the data can be estimated to be about 10-20 %. The bibliographical sources are Walker & Peñarrubia (2012); Salucci et al. (2007); Salucci (2009); dVSS (2010); Gilmore et al. (2007); Simon & Geha (2007); Simon et al. (2011); Wolf et al. (2010); Brodie et al. (2011); Willman & Strader (2012); Martinez et al. (2011).

Baryons represent less than 5% of the galaxy mass (Memola et al. 2011; Oh et al. 2008; Persic et al. 1996). For dwarf galaxies baryons count for less than 0.01% of the galaxy mass (Brodie et al. 2011; Martin et al. 2008; Walker 2012; Willman & Strader 2012; Woo et al. 2008).

The self-gravity of the baryonic material is negligible while baryons are immersed in a DM halo potential well. Baryons trace the DM potential well playing the role of test particles to measure the local DM density. In fig. 8 we contrast theory against DM observational data gathered from baryons and described with the empirical Burkert profile.

4 CONCLUSIONS

The more appropriate way to decipher the nature of the dark matter is to study the properties of the physical objects formed by it: galaxies are formed overwhelmingly by dark matter since 95% to 99.99 % of their mass is dark. This is the task we pursue in the present paper.

Fermionic WDM by itself produce galaxies and structures in agreement with observations modeled with the empirical Burkert profile showing that baryonic corrections to WDM are not very important. Therefore, the effect of including baryons is expected to be a correction to the pure WDM results, consistent with the fact that dark matter is in average six times more abundant than baryons.

The theoretical curves from the Thomas-Fermi approach to galaxy structure for self-gravitating fermionic WDM (DdVS 2013a,b) practically coincide with the observed galaxy rotation curves and density profiles described with the empirical Burkert profile for $r < 2 r_h$. In addition, our approach provides scaling relations for the main galaxy magnitudes eqs.(26)-(31) as the halo radius r_h , mass M_h and phase space density well in agreement with the observational data. The set of data is displayed in Table 2.1.

Therefore, the Fermi-Dirac distribution applies in the region $r \lesssim 2 r_h$ for the whole range of galaxy masses.

Notice that (i) the scaling relations eqs.(26)-(31) are a consequence solely of the self-gravitating interaction of the fermionic WDM and (ii) the proportionality factors in these scaling relations are confirmed by the galaxy data (see figs. 7 and 8).

The galaxy relations derived in eqs.(26)-(31) are accurate for $M_h \gtrsim 10^6 M_\odot$. We see that they exhibit a scaling behaviour for r_h vs. M_h , $Q(0)$ vs. M_h and M_h vs. the fugacity at the center $z_0 = e^{\mu(0)/E_0}$. These scaling behaviours of the dilute classical regime are **very accurate** even near the

degenerate limit as shown by fig. 1. Interestingly enough, the small deviation of these scaling laws near the degenerate limit is a manifestation of the quantum effects present in compact dwarf galaxies.

The dimensionless halo radius \hat{r}_h vs. the dimensionless halo mass \hat{M}_h plotted in fig. 7 follows with precision the square-root scaling eq.(48) obtained from the Thomas-Fermi equations (6) in the dilute regime. Moreover, the observational data modeled with the empirical Burkert profile for r_h vs. M_h for a large variety of galaxies (Table 2.1, galaxy data in McConnachie 2012, Salucci et al. 2007 and Salucci 2009) are **all** satisfactorily reproduced by the theoretical Thomas-Fermi curve.

Also, as shown by fig. 8 the observational values of the phase-space density modeled with the empirical Burkert profile are well reproduced by the theoretical Thomas-Fermi results.

The theoretical circular velocities $v_c(r)$ and the theoretical density profiles $\rho(r)$ computed from the Thomas-Fermi equations (6) reproduce very well the observational curves modeled with the empirical Burkert profile for $r \lesssim r_h$ as shown in figs. 3, 4, 5 and 6. These results fully justify the use of the Fermi-Dirac distribution function in the Thomas-Fermi equations (6).

Remarkably enough, solving the Thomas-Fermi eqs.(6) we find that the theoretical circular velocities $U(x) = v_c(r)/v_c(r_h)$ as well as the normalized density profiles $F(x) = \rho(r)/\rho(0)$ are **only** functions of $x = r/r_h$ and take respectively the same value for **all** galaxy masses in the range going from $5.13 \times 10^9 M_\odot$ till $5.15 \times 10^{11} M_\odot$ as shown in figs. 2 and 3. Namely, the Thomas-Fermi approach provides **universal** functions $U(x)$ and $F(x)$ for the normalized circular velocities and normalized density profiles, respectively. Moreover, figs. 2 and 3 show that the observational universal curves and the theoretical Thomas-Fermi curves **coincide** for $r \lesssim 2 r_h$.

These important results show the ability of the Thomas-Fermi approach to correctly describe the galaxy structures.

ACKNOWLEDGMENTS

P. S. thanks the Observatoire de Paris LERMA and the CIAS for their kind invitation and hospitality.

REFERENCES

- Anderhalden D. et al. arXiv:1212.2967, JCAP, 03, 014 (2013).
- V. Avila-Reese et al., Ap J, 559, 516 (2001).
- J. P. Brodie et al., AJ, 142, 199 (2011).
- P. Colín, O. Valenzuela, V. Avila-Reese, Ap J, 542, 622 (2000).
- P. Colín, O. Valenzuela, V. Avila-Reese, Ap J, 673, 203 (2008).
- Cosmic Frontier, H. J. de Vega, N. G. Sanchez arXiv:1304.0759.
- C. Destri, H. J. de Vega, N. G. Sanchez, New Astronomy 22, 39 (2013).

- C. Destri, H. J. de Vega, N. G. Sanchez, *Astrop. Phys.*, **46**, 14 (2013).
- C. Destri, H. J. de Vega, N. G. Sanchez, *Phys. Rev. D* **88**, 083512 (2013).
- H. J. de Vega, N. G. Sánchez, *Mon. Not. R. Astron. Soc.* **404**, 885 (2010).
- H. J. de Vega, P. Salucci, N. G. Sanchez, *New Astronomy* **17**, 653 (2012).
- F. Donato et al., *MNRAS* **397**, 1169 (2009).
- L. Gao and T. Theuns, *Science*, 317, 1527 (2007).
- G. Gentile et al., *MNRAS* **351**, 903 (2004).
- G. Gilmore et al., *Ap J*, 663, 948 (2007).
- Highlights and Conclusions of the Chalonge 16th Paris Cosmology Colloquium 2012, arXiv:1307.1847.
- Highlights and Conclusions of the Chalonge Meudon Workshop 2012 arXiv:1305.7452.
- C. J. Hogan, J. J. Dalcanton, *Phys. Rev. D* **62**, 063511 (2000).
- J Kormendy, K C Freeman, *IAU Symposium*, Sydney, 220, 377 (2004), arXiv:astro-ph/0407321.
- L D Landau and E M Lifshits, *Statistical Mechanics*, Elsevier, Oxford, 1980.
- A. Lapi, A. Cavaliere, *ApJ* 692, 174 (2009).
- M. R. Lovell et al., *MNRAS*, 420, 2318 (2012).
- M. R. Lovell et al. arXiv:1308.1399.
- D. Lynden-Bell, *Mon. Not. Roy. Astron. Soc.* **136**, 101 (1967).
- A. Macciò, S. Paduroiu, D. Anderhalden, A. Schneider, B. Moore, *MNRAS* 424, 1105 (2012).
- A. W. McConnachie, *AJ*, 144, 4 (2012).
- J. Madsen, *Phys. Rev. Lett.* **64**, 2744 (1990) and *Phys. Rev. D* **44**, 999 (1991).
- N. F. Martin et al. *ApJ* 684, 1075 (2008).
- G. D. Martinez et al., *Ap J*, 738, 55 (2011).
- E. Memola et al., *A&A*, 534, A50 (2011).
- N. Menci, F. Fiore, A. Lamastra, *ApJ*, 766, 110 (2013).
- A. Merle, arXiv:1302.2625, *Int. J. Mod. Phys. D* **22**, 1330020 (2013).
- A. M. Nierenberg et al. arXiv:1302.3243, to appear in *ApJ*.
- S. H. Oh et al., *AJ*, 136, 2761 (2008).
- F. Pacucci et al. arXiv:1306.0009, to appear in *MNRAS Letters*.
- E. Papastergis et al., *Ap J*, 739, 38 (2011).
- J. Peñarrubia, A. Pontzen, M. Walker, S. Koposov, *ApJ*, 759, L42 (2012).
- M. Persic, P. Salucci, F. Stel, *MNRAS*, 281, 27 (1996).
- P. Salucci et al. *MNRAS* **378**, 41 (2007).
- P. Salucci, unpublished (2009).
- P. Salucci et al. *MNRAS* **420**, 2034 (2012).
- S. Shao et al. *MNRAS*, 430, 2346 (2013).
- J. D. Simon, M. Geha, *Ap J*, 670, 313 (2007) and references therein.
- J. D. Simon et al., *Ap. J.* 733, 46 (2011) and references therein.
- J. Sommer-Larsen, A. Dolgov, *Ap J*, 551, 608 (2001).
- M. Spano et al., *MNRAS*, 383, 297 (2008).
- S. Tremaine, M. Henon, D. Lynden-Bell, *Mon. Not. Roy. Astron. Soc.* **219**, 285 (1986).
- A. V. Tikhonov et al., *MNRAS*, 399, 1611 (2009).
- Tonini, C. Lapi, A. Shankar, F. Salucci, P. 2006 *ApJ*, 638, 13
- S. Tremaine, J. E. Gunn, *Phys. Rev. Lett.* **42**, 407 (1979).
- M. Viel et al., *Phys. Rev. D* **88**, 043502 (2013).
- J. Viñas, E. Salvador-Solé, A. Manrique, *MNRAS* 424, L6 (2012).
- Matthew Walker, private communication, 2012.
- M. Walker, J. Peñarrubia, *Ap. J.* 742, 20 (2011).
- C. R. Watson et al. *JCAP*, 03, 018 (2012).
- B. Willman and J. Strader, *AJ*, 144, 76 (2012).
- J. Wolf et al., *MNRAS*, 406, 1220 (2010) and references therein.
- J. Woo et al. *MNRAS*, 390, 1453 (2008).
- J. Zavala et al., *Ap J*, 700, 1779 (2009).




Figure 4. The velocity rotation curves $v_c(r)$ in km/s versus r in kpc for ten different independent galaxy masses M_h going from $5.13 \times 10^9 M_\odot$ till $5.15 \times 10^{11} M_\odot$. For each galaxy mass M_h , we show the two curves: the theoretical Thomas-Fermi curve and the observational curve described by the empirical Burkert profile. The Thomas-Fermi curves reproduce remarkably well the observational curves for $r \lesssim r_h$. We plot $v_c(r)$ for $0 < r < r_{vir}$, r_{vir} being the virial radius of the galaxy.




Figure 5. The theoretical Thomas-Fermi density profiles and the observational profiles described by the Burkert expression for the first five galaxy masses. We plot the ordinary logarithm of the density in M_\odot/pc^3 vs. r in kpc in the interval $0 < r < 4 r_h$. For each galaxy mass M_h , we show the two curves: the theoretical Thomas-Fermi curve and the observational Burkert curve. The agreement of the Thomas-Fermi curves to the observational curves is remarkable.




Figure 6. The theoretical Thomas-Fermi density profiles and the observational profiles described by the Burkert expression for further five galaxy masses. We plot the ordinary logarithm of the density in M_\odot/pc^3 vs. r in kpc in the interval $0 < r < 4 r_h$. For each galaxy mass M_h , we show the two curves: the theoretical Thomas-Fermi curve and the observational Burkert curve. The agreement of the Thomas-Fermi curves to the observational curves is remarkable.




Figure 7. The ordinary logarithm of $\hat{r}_h = \frac{r_h}{\text{pc}} \left(\frac{\Sigma_0 \text{ pc}^2}{120 M_\odot} \right)^{\frac{1}{5}}$ vs. the ordinary logarithm of $\hat{M}_h = \frac{M_h}{M_\odot} \left(\frac{120 M_\odot}{\Sigma_0 \text{ pc}^2} \right)^{\frac{3}{5}}$. We see that r_h follows with precision the square-root of M_h as in the dilute regime eq.(26) of the Thomas-Fermi equations. The data for M_h and r_h are taken from Table 2.1, from (McConnachie 2012) and from (Salucci et al. 2007; Salucci 2009) and they are extremely well reproduced by the theoretical Thomas-Fermi curve.




Figure 8. The theoretical $\log_{10} Q_{ch} / \text{keV}^4$ vs. the halo mass $\log_{10} \hat{M}_h$. The theoretical curve Q_{ch} is obtained from the Thomas-Fermi expression (39). The data for $Q_{ch,B}$ have been obtained from circular velocities. Galaxy Data 1 refers to data from Table 2.1, Galaxy Data 2 refers to data from (McConnachie 2012) and Galaxy Data 3 refers to spiral galaxy data from (Salucci et al. 2007; Salucci 2009) .