

PHENOMENOLOGY OF TRANSVERSE MOMENTUM DEPENDENT PARTON DENSITIES

MARIAELENA BOGLIONE

*Physics Department, University of Torino,
 Via P. Giuria 1 Torino, 10125, Italy
 boglione@to.infn.it*

Received Day Month Year
 Revised Day Month Year

This talk provides a short overview of the phenomenology of transverse momentum dependent distribution and fragmentation functions, focussing on the most recent phenomenological developments in the study of their Q^2 evolution and energy dependence.

Keywords: Transverse momentum; QCD evolution; parton densities

PACS numbers: 13.88.+e, 12.38.Bx, 13.85.Ni

The exploration of the three-dimensional structure of the nucleon, both in momentum and in configuration space, is one of the major issues in high energy hadron physics. Transverse momentum dependent parton densities (TMDs) have recently attracted huge interest, not only for their remarkable spin correlation properties, but mostly because they represent a crucial tool for the investigation of the three-dimensional structure of the nucleon.

Huge amount of experimental data on spin asymmetries in several different processes show that TMD distribution and fragmentation functions exist and are non zero. Much progress has been achieved, for instance, in the phenomenological studies of the Sivers distribution function, which represents the number density of unpolarized partons inside a transversely polarized hadron, and of the Collins fragmentation function, which is related to the probability of a polarized quark fragmenting into an unpolarized hadron.

TMD studies are readily performed in the framework of QCD factorization, within a generalized parton model that incorporates the partonic intrinsic transverse motion in the kinematics of the examined scattering processes. In this simple phenomenological approach, cross sections and spin asymmetries are generated as convolutions of distribution and (or) fragmentation TMDs with elementary scattering cross sections; for instance for semi-inclusive deep inelastic scattering (SIDIS) processes, we have:

$$d\sigma^{\ell p \rightarrow \ell' h X} = \sum_q f_{q/p}(x, \mathbf{k}_\perp; Q^2) \otimes d\sigma^{\ell q \rightarrow \ell' q} \otimes D_{h/q}(z, \mathbf{p}_\perp; Q^2). \quad (1)$$




Fig. 1. A schematic illustration of SIDIS kinematics

In the $\gamma^* - p$ c.m. frame, see Fig. 1, the measured transverse momentum, \mathbf{P}_T , of the final hadron is generated by the transverse momentum of the quark in the target proton, \mathbf{k}_\perp , and of the final hadron with respect to the fragmenting quark, \mathbf{p}_\perp . At order k_\perp/Q it is simply given by

$$\mathbf{P}_T = z \mathbf{k}_\perp + \mathbf{p}_\perp. \quad (2)$$

There is a general consensus that such a scheme holds in the kinematical region defined by

$$P_T \simeq \Lambda_{\text{QCD}} \ll Q. \quad (3)$$

The presence of the two scales, small P_T and large Q , allows to identify the contribution from the unintegrated partonic distribution ($P_T \simeq k_\perp$), while remaining in the region of validity of the QCD parton model.

Within this simple scheme we can successfully describe a wide range of unpolarized and polarized experimental data, provided we are able to model and phenomenologically determine the appropriate TMDs, including their scale evolution. Historically, in the Torino-Cagliari standard approach, TMDs are parametrized in a form in which their dependence on the lightcone momentum fraction and on the partonic intrinsic transverse momentum are factorized,

$$f_{q/p}(x, k_\perp) = f_{q/p}(x, Q^2) \frac{e^{-k_\perp^2 / \langle k_\perp^2 \rangle}}{\pi \langle k_\perp^2 \rangle}, \quad (4)$$

$$D_{h/q}(z, p_\perp) = D_{h/q}(z, Q^2) \frac{e^{-p_\perp^2 / \langle p_\perp^2 \rangle}}{\pi \langle p_\perp^2 \rangle}, \quad (5)$$

with a Q^2 -independent, normalized Gaussian factor giving the intrinsic transverse momentum distribution, multiplied by a collinear unpolarized parton distribution function (PDF) evolving with Q^2 according to DGLAP equations; $\langle k_\perp^2 \rangle$ and $\langle p_\perp^2 \rangle$ are free parameters which can be extracted from experiments. A similar parameterization is devised for polarized TMDs, like the Sivers function

$$\Delta^N f_{q/p^\uparrow}(x, k_\perp) = 2 \mathcal{N}_q(x) h(k_\perp) f_{q/p}(x, k_\perp, Q^2) \quad (6)$$

with

$$f_{q/p}(x, k_\perp) = f(x, Q^2) \frac{e^{-k_\perp^2 / \langle k_\perp^2 \rangle}}{\pi \langle k_\perp^2 \rangle}, \quad (7)$$

$$\mathcal{N}_q(x) = N_q x^{\alpha_q} (1-x)^{\beta_q} \frac{(\alpha_q + \beta_q)^{(\alpha_q + \beta_q)}}{\alpha_q^{\alpha_q} \beta_q^{\beta_q}}, \quad (8)$$

and

$$h(k_\perp) = \sqrt{2e} \frac{k_\perp}{M_1} e^{-k_\perp^2 / M_1^2}. \quad (9)$$

N_q , α_q , β_q and M_1 are free parameters which can be extracted from experimental data.

A completely analogous parameterization holds for the Collins function.

In Ref. [1] we performed a study of the unpolarized and Sivers TMDs based on EMC² measurements of the azimuthal dependence and the P_T^2 distribution of the SIDIS cross sections, where we were able to extract the Gaussian widths, $\langle k_\perp^2 \rangle$ and $\langle p_\perp^2 \rangle$, appearing in Eqs. (4) and (5), which represented the two free parameters of our fit. We found:

$$\langle k_\perp^2 \rangle = 0.25 \text{ GeV}^2, \quad \langle p_\perp^2 \rangle = 0.20 \text{ GeV}^2. \quad (10)$$

Afterwards, in Refs. [3, 4, 5] by exploiting the above parameters, we extracted the Sivers function by fitting polarized SIDIS azimuthal Sivers asymmetries from the HERMES^{6,7} and COMPASS⁸ experiments. Similarly, in a combined fit^{9,10,11} of SIDIS^{6,12,13,14} and e^+e^- ^{15,16,17} experimental data, we extracted the transversity and Collins functions simultaneously.

As new, more precise and higher statistics data are rapidly becoming available, it is due time to start wondering whether we can find any sign of TMD scale evolution in the experimental measurements we can now access. Interesting studies¹⁹ have been performed by comparing the Sivers asymmetries measured by the HERMES^{7,12} and COMPASS^{14,18} Collaborations, over similar x , z and P_T ranges, but different average values of Q^2 :

$$\langle Q^2 \rangle = 2.4 \text{ GeV}^2 \text{ for HERMES and } \langle Q^2 \rangle = 3.2 \text{ GeV}^2 \text{ for COMPASS.} \quad (11)$$

As shown in Fig. 2 it is quite evident that there is a systematic difference between the two sets of data, which can possibly be interpreted as a sign of scale evolution. A similar comparison for the analogous Collins asymmetry shows that in this case there is no evident inconsistency between the two measurements, and no indication of a possible TMD evolution for the Collins function can be deduced.

From the theory point of view, much progress has recently been achieved in the framework of TMD evolution. After the 1985 first pioneering work on QCD resummation for transverse momentum dependent processes by Collins, Soper and Sterman²⁰, the issue was revived by Idilbi, Ji, Ma and Yuan in several papers^{21,22,23} in 2004–2005 and, later on, revisited by J. Collins in his 2011 book²⁴.

This scheme of TMD evolution was finalized by Aybat and Rogers²⁵ for the unpolarized TMDs, who also proposed, together with J. Collins and J. Qiu, a model for the TMD evolution for polarized parton densities, in particular for the Sivers function²⁶.

Most of the people mentioned above are participants of this workshop, therefore the interested reader can find more details and better explanations in their contributions to these proceedings.

After the publication of these theory works, the first studies of what can more properly be defined TMD-phenomenology started to be performed: Aybat, Prokudin, Rogers²⁷ proposed a very elementary phenomenological exercise in which they compared the HERMES and COMPASS Sivers single spin asymmetry A_{UT} ,




Fig. 2. Comparison between the Sivers $A_{UT}^{\sin(\phi_h - \phi_S)}$ asymmetry, upper panel, and Collins $A_{UT}^{\sin(\phi_h + \phi_S)}$ asymmetry, lower panel, as measured by the HERMES^{7,12} and COMPASS^{14,18} Collaborations.




Fig. 3. Comparison between HERMES⁷ and COMPASS¹⁴ data and the analogous asymmetry obtained from the phenomenological model of Ref. [4] by applying the TMD evolution of Ref. [26] (dashed line) or simply using Eq. (6) (solid line).

calculated at their two fixed values of $\langle Q^2 \rangle$: 2.4 GeV² for HERMES and 3.8 for COMPASS; evolution effects were then compared, as illustrated in Fig. 3. Notice that no x dependence was taken into account, somehow washing out the most sensitive and interesting information this kind of data can provide on TMD scale evolution.

At the same time, M. Anselmino, S. Melis and myself²⁸ performed a new, more refined phenomenological analysis of the Sivers TMD evolution, again based on the comparison between HERMES and COMPASS sets of measurements on the Sivers effect, but focussing mostly on the actual Q^2 and x dependence. We have shown that TMD-evolution^{25,26} implies a strong variation with Q^2 of the functional form of the unpolarized and Sivers TMDs, as functions of the intrinsic momentum k_\perp ; moreover, our fit of all SIDIS data on the Sivers asymmetry using TMD-evolution, when compared with the same analysis performed with the simplified DGLAP-evolution, exhibits a smaller value of the total χ^2 , a reduction which mostly originates from the large Q^2 COMPASS data, which are greatly affected by the TMD evolution, as shown in Fig. 4. We then considered this as a strong indication in favor of the TMD evolution.

Without going into the details of such a complex evolution scheme, there is one particular point that I would like to address. The TMD evolution equation of the unpolarized TMD PDFs, in configuration space, is the following

$$\tilde{f}_{q/p}(x, b_T; Q) = f_{q/p}(x, Q_0) \tilde{R}(Q, Q_0, b_T) \exp \left\{ -b_T^2 \left(\langle k_\perp^2 \rangle / 4 + \frac{g_2}{2} \ln \frac{Q}{Q_0} \right) \right\}, \quad (12)$$

$$\text{where } g_K(b_T) = \frac{1}{2} g_2 b_T^2 \text{ with } g_2 = 0.68 \text{ GeV}^2 \text{ corresponding to } b_{\text{max}} = 0.5 \text{ GeV}^{-1}. \quad (13)$$

A similar equation regulates the evolution of the first derivative of the Sivers function. Eq. (12) shows that the Q^2 evolution is controlled by the logarithmic Q dependence of the b_T Gaussian width, together with the factor $\tilde{R}(Q, Q_0, b_T)$: for increasing




Fig. 4. The results obtained for the SIDIS $A_{UT}^{\sin(\phi_h - \phi_S)}$ asymmetries applying TMD evolution (red, solid lines) are compared with the analogous results found by using DGLAP evolution equations (blue, dashed lines). The green, dash-dotted lines correspond to the results obtained by using the approximated analytical TMD evolution (see text for further details). The experimental data are from HERMES ⁷ (left panel) and COMPASS ¹⁴ (right panel) Collaborations.

values of Q^2 , they are responsible for the typical broadening effect already observed in Refs. [25] and [26]. Notice that the parameter g_2 , that controls the b_T Gaussian width and its spreading, is not extracted from our fit, but taken as a fixed values from elsewhere ²⁹. We could have determined its value in our fit, and probably got a smaller value, but it is important to remember that SIDIS asymmetries are very little sensitive to the precise value of g_2 ; therefore, our choice to keep it fixed was motivated by the balance one always has to keep between number of data and number of free parameters, to obtain a reliable fit. Drell Yan processes, instead, where Q^2 s are much larger and perturbative corrections become important, are extremely sensitive to it. Therefore, one should not simply take the parameters used in this application of TMD evolution to SIDIS processes and apply them blindly to Drell Yan data or other processes. This would require a new, careful, global analysis on all SIDIS and Drell Yan data, re-starting from unpolarized cross sections.

Sun and Yuan³² have recently applied some CSS²⁰-like evolution scheme at one loop, with strong approximations which hold for moderate Q and Q_0 , to account for the TMD evolution of the unpolarized TMD PDFs, and extended this formalism to the Sivvers function as well. In this approximated scheme the evolution does not produce such a strong suppression of the Drell-Yan asymmetries. They then perform a phenomenological study on a rather limited selection of Drell-Yan and SIDIS data, showing that the evolution scheme they propose can satisfactorily describe most of them.

Remaining on Drell Yan processes, a recent preliminary study was performed by S. Melis, fitting the E288³⁰ and E605³¹ Drell-Yan data and assuming a Gaussian




Fig. 5. Gaussian fit of Drell Yan data from E288³⁰ and E605³¹ experiments.

k_{\perp} dependence. The free parameter $\langle k_{\perp}^2 \rangle$ was extracted separately for each data sets corresponding to different \sqrt{s} values. The results showed that those Drell Yan data indicate a \sqrt{s} (roughly linear) dependence in addition to the logarithmic Q^2 dependence typical of scale evolution. This is shown in Figs. 5 and 6.

Most recently D. Boer³³ has performed a study of the energy scale dependence of the Sivers asymmetry in SIDIS, although on a larger range of Q values (3 - 100 GeV): he finds that the peak of the Sivers asymmetry falls off with Q roughly like $(1/Q)^{0.7}$, quite faster than found within the CSS²⁰ evolution schemes. Moreover, the peak of the asymmetry is located around the initial scale Q_0 and moves rather slowly towards higher transverse momentum values as Q increases, which may be due to the absence of perturbative tails of the TMDs.

Important work on TMD evolution has also recently been done by Echevarria, Idilbi, Scimemi³⁴ in the framework of effective field theory which, however, has not yet reached the stage of feasible phenomenological applications.

As this is one of the opening talks of this workshop, there are no proper conclusions. Rather, I will close with a few remarks on future perspectives.

As far as TMD evolution is concerned, we have recently come a long way. We now have evolution schemes and some first attempts towards a full phenomenological study of the unpolarized distribution and fragmentation TMDs, and of the Sivers




Fig. 6. Average k_{\perp} extracted from Drell Yan data as a function of \sqrt{s} : notice that $\langle k_{\perp}^2 \rangle$ shows a linear growth with \sqrt{s} .

and Collins functions. These are very preliminary studies, which need to be refined and re-thought in a more consistent and appropriate way, especially as far as the parametrization of unknown phenomenological quantities are concerned.

From the experimental side, we certainly need more SIDIS (polarized and unpolarized) data at larger values of x (JLab 12) and spanning a larger Q^2 range (EIC) as well as more (and more precise) Drell-Yan data, for which many beautiful experiments are being planned (COMPASS, RHIC, Fermilab, NICA, JPARK). Inclusive hadron production in hadron-hadron scattering processes, as well, represent a very interesting and infinitely challenging field where to sharpen our tools.

With the new experimental data on SIDIS multiplicities coming in, we have to go back one step, re-think and re-perform a solid, global analysis of Drell-Yan as well as SIDIS unpolarized cross sections, to determine the basic parameters for the phenomenological quantities needed for the implementation of the TMD evolution schemes. Afterwards, we can proceed on a firm footing to perform the same analysis for the Sivvers, transversity and Collins TMD functions, keeping in mind the importance of finding phenomenological frameworks suitable for all processes.

Acknowledgments

I thank the organizers for their warm hospitality; in particular, Alexei Prokudin who, beside being a great collaborator, always manages to turn a conference into a very special event. I am most grateful to S. Melis and M. Anselmino for their relentless work and constant support.

References

1. M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia and A. Prokudin, *Phys. Rev.* **D71**, p. 074006 (2005).
2. J. Aubert *et al.*, *Phys.Lett.* **B130**, p. 118 (1983).

3. M. Anselmino, M. Boglione, D'Alesio, A. Kotzinian, F. Murgia and A. Prokudin, *Phys. Rev.* **D72**, p. 094007 (2005).
4. M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, S. Melis, F. Murgia, A. Prokudin, C. Türk, *Eur.Phys.J.* **A39**, 89 (2009).
5. M. Anselmino, M. Boglione, U. D'Alesio, S. Melis, F. Murgia, A. Prokudin, *J.Phys.Conf.Ser.* **295**, p. 012062 (2011).
6. A. Airapetian *et al.*, *Phys. Rev. Lett.* **94**, p. 012002 (2005).
7. A. Airapetian *et al.*, *Phys.Rev.Lett.* **103**, p. 152002 (2009).
8. M. Alekseev *et al.*, *Phys.Lett.* **B673**, 127 (2009).
9. M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, A. Prokudin, *Phys. Rev.* **D75**, p. 054032 (2007).
10. M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, A. Prokudin, S. Melis, *Nucl. Phys. Proc. Suppl.* **191**, 98 (2009).
11. M. Anselmino, M. Boglione, U. D'Alesio, S. Melis, F. Murgia, A. Prokudin, *Phys.Rev.* **D87**, p. 094019 (2013).
12. A. Airapetian *et al.*, *Phys. Lett.* **B693**, 11 (2010).
13. E. S. Ageev *et al.*, *Nucl. Phys.* **B765**, p. 31 (2007).
14. C. Adolph *et al.*, *Phys.Lett.* **B717**, 376 (2012).
15. R. Seidl *et al.*, *Phys. Rev. Lett.* **96**, p. 232002 (2006).
16. R. Seidl *et al.*, *Phys. Rev.* **D78**, p. 032011 (2008).
17. R. Seidl *et al.*, *Phys. Rev.* **D86**, p. 032011(E) (2012).
18. C. Adolph *et al.*, *Phys.Lett.* **B717**, 383 (2012).
19. A. Martin, talk presented at the XXI International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS 2013), April 22-26 2013, Marseilles, France.
20. J. C. Collins, D. E. Soper and G. F. Sterman, *Nucl.Phys.* **B250**, p. 199 (1985).
21. X.-d. Ji, J.-P. Ma and F. Yuan, *Phys. Lett.* **B597**, 299 (2004).
22. A. Idilbi, X.-d. Ji, J.-P. Ma and F. Yuan, *Phys.Rev.* **D70**, p. 074021 (2004).
23. X.-d. Ji, J.-p. Ma and F. Yuan, *Phys. Rev.* **D71**, p. 034005 (2005).
24. J. Collins, *Foundations of perturbative QCD*, Cambridge monographs on particle physics, nuclear physics and cosmology, n. 32, Cambridge University Press, Cambridge (2011).
25. S. M. Aybat and T. C. Rogers, *Phys.Rev.* **D83**, p. 114042 (2011).
26. S. M. Aybat, J. C. Collins, J.-W. Qiu and T. C. Rogers, *Phys.Rev.* **D85**, p. 034043 (2012).
27. S. M. Aybat, A. Prokudin and T. C. Rogers, *Phys.Rev.Lett.* **108**, p. 242003 (2012).
28. M. Anselmino, M. Boglione and S. Melis, *Phys.Rev.* **D86**, p. 014028 (2012).
29. F. Landry, R. Brock, P. M. Nadolsky and C. Yuan, *Phys.Rev.* **D67**, p. 073016 (2003).
30. A. Ito, R. Fisk, H. Jostlein, D. Kaplan, S. Herb *et al.*, *Phys.Rev.* **D23**, p. 604 (1981).
31. G. Moreno, C. Brown, W. Cooper, D. Finley, Y. Hsiung *et al.*, *Phys.Rev.* **D43**, 2815 (1991).
32. P. Sun and F. Yuan, *Phys.Rev.* **D88**, p. 034016 (2013).
33. D. Boer, *Nucl.Phys.* **B874**, 217 (2013).
34. M. G. Echevarria, A. Idilbi and I. Scimemi, *JHEP* **1207**, p. 002 (2012).