

Residual Symmetry Reductions and Interaction Solutions of (2+1)-Dimensional Burgers Equation

Xi-zhong Liu, Jun Yu, Bo Ren

Institute of Nonlinear Science, Shaoxing University, Shaoxing 312000, China

Abstract

The (2+1)-dimensional Burgers equation has been investigated first from prospective of symmetry by localizing the nonlocal residual symmetries and then studied by a simple generalized tanh expansion method. New symmetry reduction solutions has been obtained by using the standard Lie point symmetry group approach. A new Bäcklund transformation for Burgers equation has been given with the generalized tanh expansion method . From this BT, interactive solutions among different nonlinear excitations which is hard to obtain by other methods has also been obtained easily.

PACS numbers: 02.30.Jr, 02.30.Ik, 05.45.Yv, 47.35.Fg

Keywords: Burgers equation, residual symmetry, Bäcklund transformation, symmetry reduction solution, generalized tanh expansion method

I. INTRODUCTION

Due to the important role that nonlinear theory plays in mathematics, physics, chemistry, biology, communications, astrophysics and geophysics, much endeavor have been devoted to find important and significant properties of different type of nonlinear equations. Among which, the study on integrable systems occupies the most important place. The integrable systems are introduced by scientist with some ideal conditions to describe complex nonlinear phenomenon, such as the Korteweg-de Vries (KdV) equation [1], the nonlinear Schrödinger equation (NLS) [2], the sine-Gordon (SG) equation [3], Davey-Stewartson (DS) equation [4], the Kadomtzev-Pedviashvili (KP) [5] equation and so on. Various effective methods like the inverse scattering transformation (IST), Darboux transformation, Bäcklund transformation (BT)[6], Hirota's direct method [7], mapping and deformation approach [8], standard and extended truncated Painlevé analysis [9] etc. have been well developed to find exact solutions for integrable models.

Since the Lie group theory was introduced by Sophus Lie to study differential equations [10], the symmetry theory has been developed a lot for studying nonlinear equations. It is well known that using Lie point symmetries, the dimensions of nonlinear equations can be reduced through classical or non-classical Lie group approaches and the exact solutions of these equations can be constructed further. However, for integrable system there exist infinitely many non-local symmetries which are linked with potential symmetries [11], inverse recursion operators [12], negative hierarchies, conformal invariance, Darboux transformations, Bäcklund transformations, and so on. It would be very interesting to use these non-local symmetries for studying important properties of the corresponding equations. An direct thinking is to localize these symmetries by prolong the original system then use the traditional Lie point symmetry theory to reduce the prolonged system [13] and find new exact reduction solutions which is usually hard to be found by traditional DT or BT method.

The Painlevé analysis is usually used to study the integrability of nonlinear systems. Painlevé integrable demands that there exist Laurent series on open sets of the complex time variable solutions and these solutions are consistent, or match on the overlapping pieces of the sets on which they are defined. In other words this notion of integrability is existence of meromorphic solution. When the equations has Painlevé property, the truncated Painlevé expansion method can be used to find exact solutions of the equations. The recent

studies on Painlevé analysis reveals that the truncated Painlevé expansion is the Bäcklund transformation and the residue of the expansion is just the non-local symmetry related this Bäcklund transformation, which is called residual symmetry. In Ref. [14], the truncated Painlevé expansion method is further developed to the general the hyperbolic tangent (tanh) function expansion approach by allowing the expansion coefficients being arbitrary functions of space-time and interaction solutions between solitons and any Schrödinger waves has been obtained. Unlike the usual tanh expansion method which lose many essential information of the original equation the generalized tanh expansion method can be used to find much more general solutions retrieving the missing essential properties.

In this paper, we concentrate on the (2+1)-dimensional Burgers equation with constants a, b in the form

$$u_t = uu_y + avu_x + bu_{yy} + abu_{xx}, \quad (1a)$$

$$u_x = v_y \quad (1b)$$

via residual related symmetry reduction and generalized tanh function expansion approach to obtain new BT transformations and interaction solutions between different solitons. An equivalent form of the Burgers Eq. (2) has been derived from the generalized Painlevé integrability classification in Ref. [15].

Burgers equation is a fundamental partial differential equation in the field of fluid mechanics. It also occurs in various areas of applied physics such as the modeling of gas dynamics and traffic flow. The exact solutions of Burgers equation have been studied by many authors. In Ref. [16], the infinite many symmetries and exact solutions of Burgers equation are obtained by the repeated symmetry reduction approach.

In Sec. 2 of this paper, the residual symmetry of Burgers system is localized and then Levi transformation (the second DT) is re-obtained via Lie's first theorem. In Sec. 3, the localized symmetry group is analyzed and classified according to different transformation properties. The reduction solutions related with space-time and Möbius transformation invariance symmetries has also been obtained subsequently. In Sec. 4, the generalized tanh expansion approach is used to Burgers equation to obtain new Bäcklund transformation. Using the BT, some special explicit novel exact solutions and the interactions between them has also been investigated. It is found that completely non-elastic interactions and completely elastic interactions will occur when specific parameters be fixed. The last section

is devoted to a short summary and discussion.

II. LOCALIZATION OF THE NON-LOCAL RESIDUAL SYMMETRIES OF THE BURGERS EQUATION

By means of the Painleve analysis, various integrable properties such as the Backlund transformations, Lax pairs, infinitely many symmetries, bilinear forms, and so on, can be easily found if the studied model possesses Painleve property, i.e., it is Painleve integrable.

For the Burgers Eq. (1), the truncated Painlevé expansion takes the general form

$$u = \sum_{i=0}^{\alpha} u_i f^{i-\alpha}, \quad v = \sum_{j=0}^{\beta} v_j f^{j-\beta}, \quad (2)$$

where u_α and v_β are arbitrary solutions of the discussed system while $u_{\alpha-1}, u_{\alpha-2}, \dots, u_0$ and $v_{\beta-1}, v_{\beta-2}, \dots, v_0$ are all related to derivatives of f . In order to determine α and β , we first substitute $u \sim u_0 f^\alpha$ and $v \sim v_0 f^\beta$ into Eq. (1) and balance the nonlinear and dispersion terms to give $\alpha = 1, \beta = 1$. Hence the truncated Painlevé expansion reads

$$u = \frac{u_0}{f} + u_1, \quad v = \frac{v_0}{f} + v_1. \quad (3)$$

It can be easily proved that the residuals u_0, v_0 are the symmetries corresponding to the BT[].

With the property of Painlevé integrable the BT (3) transforms the original Burgers system (1) into its Schwarzian form

$$S(f) = 0. \quad (4)$$

That mens (4) is form invariant under the Mobious transformation

$$f \rightarrow \frac{a_1 f + b_1}{a_2 f + b_2}, \quad a_1 a_2 \neq b_1 b_2 \quad (5)$$

which means (4) possess three symmetries $\sigma^f = d_1, \sigma^f = d_2 f$ and

$$\sigma^f = d_3 f^2$$

with arbitrary constants d_1, d_2 and d_3 .

Substituting Eq. (3) into Eq. (1) yields

$$\begin{aligned}
& -abu_{1,xx} - av_1u_{1,x} - u_1u_{1,y} + u_{1,t} - bu_{1,yy} + f^{-1}(-av_1u_{0,x} - av_0u_{1,x} - abu_{0,xx} \\
& - u_1u_{0,y} + u_{0,t} - u_0u_{1,y} - bu_{0,yy}) + f^{-2}(av_1u_0f_x + 2abu_{0,x}f_x + abu_0f_{xx} + bu_0f_{yy} - av_0u_{0,x} \\
& + 2bu_{0,y}f_y - u_0u_{0,y} + u_1u_0f_y - u_0f_t) - f^{-3}u_0(2abf_x^2 - av_0f_x + 2bf_y^2 - u_0f_y) = 0, \quad (6)
\end{aligned}$$

and

$$u_{1,x} - v_{1,y} + f^{-1}(-v_{0,y} + u_{0,x}) + f^{-2}(v_0f_y - u_0f_x) = 0 \quad (7)$$

Vanishing the coefficients f^{-3} yields

$$u_0 = 2bf_y, \quad v_0 = 2bf_x \quad (8)$$

Substituting Eq. (8) into Eq. (7) yields the Burgers equation (1b) with u_1 and v_1 as solutions

$$u_{1,x} - v_{1,y} = 0. \quad (9)$$

Substituting Eq. (8) into Eq. (6) and vanishing the coefficient of f^{-2} yields

$$av_1f_x + abf_{xx} + bf_{yy} + u_1f_y - f_t = 0, \quad (10)$$

while the coefficient of f^{-1} equals Eq. (9) by using Eq. (10).

Vanishing the coefficient of f^0 in Eq. (6), we have

$$u_{1,t} - abu_{1,xx} - av_1u_{1,x} - u_1u_{1,y} - bu_{1,yy} = 0 \quad (11)$$

which is just the Burgers equation (1a) with the solution u_1 and v_1 .

Now, using the standard truncated Painleve expansion

$$\begin{aligned}
u &= \frac{2bf_y}{f} + u_1, \\
v &= \frac{2bf_x}{f} + v_1,
\end{aligned} \quad (12)$$

we have the following Bäcklund transformation theorem

Theorem 1. (12) is a BT between the solutions $\{u, v\}$ and $\{u_1, v_1\}$ if the latter is related to f by Eq. (10).

Since u_0 and v_0 are non-local symmetries corresponding to BT, one can naturally believe that it can be localized to a Lie point symmetry such that we can use the Lie's first theorem

to recover the original BT. To this end, we introduce new variables to eliminate the space derivatives of f

$$g \equiv f_x, \quad (13)$$

$$h \equiv f_y \quad (14)$$

and investigate the symmetries relations between different variables. That is to say, we have to solve the linearized system of the prolonged equations of (1), (10),(13) and (14)

$$\sigma_{u_1,x} - \sigma_{v_1,y} = 0, \quad (15a)$$

$$-ab\sigma_{v_1,xy} + \sigma_{u_1,t} - b\sigma_{u_1,yy} - \sigma_{u_1}u_{1,y} - u_1\sigma_{u_1,y} - a\sigma_{v_1}v_{1,y} - av_1\sigma_{v_1,y} = 0, \quad (15b)$$

$$\sigma_{f,x} - \sigma_g = 0, \quad (15c)$$

$$\sigma_{f,y} - \sigma_h = 0, \quad (15d)$$

$$\sigma_{f,t} - b\sigma_{f,yy} - av_1\sigma_{f,x} - a\sigma_{v_1}f_x - ab\sigma_{f,xx} - f_y\sigma_{u_1} - u_1\sigma_{f,y} = 0 \quad (15e)$$

It can be easily verified that the solution of (15) has the form

$$\sigma^{u_1} = 2bh, \quad (16a)$$

$$\sigma^{v_1} = 2bg, \quad (16b)$$

$$\sigma^g = -\frac{fg}{b}, \quad (16c)$$

$$\sigma^h = -\frac{fh}{b}, \quad (16d)$$

$$\sigma^f = -\frac{f^2}{2b}, \quad (16e)$$

if d_3 is fixed as $-\frac{1}{2b}$.

The result (16) indicates that the residual symmetries (8) is localized in the properly prolonged system (1), (10), (13) and (14) with the Lie point symmetry vector

$$V = 2bh\partial_{u_1} + 2bg\partial_{v_1} - \frac{fg}{b}\partial_g - \frac{fh}{b}\partial_h - \frac{f^2}{2b}\partial_f. \quad (17)$$

In other words, symmetries related to the truncated Painlevé expansion is just a special Lie point symmetry of the prolonged system (1), (10), (13) and (14).

Now has we obtained the localized residual symmetries, the interesting question is to ask what kind of finite transformation would be corresponding the Lie point symmetry (17). Then we has the following theorem.

Theorem 2. If $\{u_1, v_1, g, h, f\}$ is a solution of the prolonged system (1), (10), (13) and (14), then so is $\{\hat{u}_1, \hat{v}_1, \hat{g}, \hat{h}, \hat{f}\}$ with

$$\hat{u}_1 = u_1 + \frac{4hb^2\epsilon}{\epsilon f + 2b}, \quad (18a)$$

$$\hat{v}_1 = v_1 + \frac{4gb^2\epsilon}{\epsilon f + 2b} \quad (18b)$$

$$\hat{g} = \frac{4gb^2}{(\epsilon f + 2b)^2}, \quad (18c)$$

$$\hat{h} = \frac{4hb^2}{(\epsilon f + 2b)^2}, \quad (18d)$$

$$\hat{f} = \frac{2bf}{\epsilon f + 2b}, \quad (18e)$$

with arbitrary group parameter ϵ .

Proof. Using Lie's first theorem on vector (17) with the corresponding initial condition as follows

$$\frac{d\hat{u}_1(\epsilon)}{d\epsilon} = 2b\hat{h}(\epsilon), \quad \hat{u}_1(0) = u_1, \quad (19)$$

$$\frac{d\hat{v}_1(\epsilon)}{d\epsilon} = 2b\hat{g}(\epsilon), \quad \hat{v}_1(0) = v_1, \quad (20)$$

$$\frac{d\hat{g}(\epsilon)}{d\epsilon} = -\frac{\hat{f}(\epsilon)\hat{g}(\epsilon)}{b}, \quad \hat{g}(0) = g, \quad (21)$$

$$\frac{d\hat{h}(\epsilon)}{d\epsilon} = -\frac{\hat{f}(\epsilon)\hat{h}(\epsilon)}{b}, \quad \hat{h}(0) = h, \quad (22)$$

$$\frac{d\hat{f}(\epsilon)}{d\epsilon} = -\frac{\hat{f}^2(\epsilon)}{2b}, \quad \hat{f}(0) = f \quad (23)$$

one can easily obtain the solutions of the above equations stated in Theorem 1, thus the theorem is proved.

Next let us consider the Lie point symmetry of the prolonged system in the general form

$$V = X \frac{\partial}{\partial x} + Y \frac{\partial}{\partial y} + T \frac{\partial}{\partial t} + U_1 \frac{\partial}{\partial u_1} + V_1 \frac{\partial}{\partial v_1} + G \frac{\partial}{\partial g} + H \frac{\partial}{\partial h} + F \frac{\partial}{\partial f}, \quad (24)$$

which means that the prolonged system is invariant under the following transformation

$$\{x, y, t, u, v, g, h, f\} \rightarrow \{x + \epsilon X, y + \epsilon Y, t + \epsilon T, u + \epsilon U, v + \epsilon V, g + \epsilon G, h + \epsilon H, f + \epsilon F\} \quad (25)$$

with the infinitesimal parameter ϵ . Equivalently, the symmetry in the form (25) can be written as a function form as

$$\sigma_{u_1} = Xu_{1x} + Yu_{1y} + Tu_{1t} - U_1, \quad (27a)$$

$$\sigma_{v_1} = Xv_{1x} + Yv_{1y} + Tv_{1t} - V_1, \quad (27b)$$

$$\sigma_g = Xg_x + Yg_y + Tg_t - G, \quad (27c)$$

$$\sigma_h = Xh_x + Yh_y + Th_t - H, \quad (27d)$$

$$\sigma_f = Xf_x + Yf_y + Tf_t - F. \quad (27e)$$

Substituting Eq. (27) into Eq. (15) and eliminating $u_{1,t}, v_{1,y}, g_t, h_x, h_t, f_x, f_y$ and f_t in terms of the prolonged system we get more than 250 determining equations for the functions X, Y, T, U_1, V_1, G, H and F . Calculated by computer algebra, we finally get the desired result

$$\begin{aligned} X &= \frac{c_1xt}{2} + \frac{c_2x}{2} + x_0, Y = \frac{c_1ty}{2} + \frac{c_2y}{2} + c_4t + c_5, \\ T &= \frac{c_1t^2}{2} + c_2t + c_3, U_1 = -\frac{c_1y}{2} - \frac{c_1u_1t}{2} + c_6h - \frac{c_2u_1}{2} - c_4, \\ V_1 &= -\frac{2x_{0,t} + av_1c_1t + av_1c_2 - 2ac_6g + c_1x}{2a}, G = -\frac{g(bc_1t + bc_2 - 2bc_7 + 2c_6f)}{2b}, \\ H &= -\frac{h(bc_1t + bc_2 - 2bc_7 + 2c_6f)}{2b}, F = -\frac{c_6f^2 - 2bc_7f - 2bc_8}{2b}. \end{aligned} \quad (28)$$

with arbitrary constants $c_1, c_2, c_3, c_4, c_5, c_6, c_7, c_8$ and arbitrary function x_0 of t . The obtained symmetries can be categorized by different transformation invariance as follows,

(i) the residual symmetries and Mobious transformation by setting $c_1 = c_2 = c_3 = c_4 = c_5 = x_0 = 0$ and $c_6 = c_7 = c_8 = -1$

$$\sigma_{u_1} = f_y, \sigma_{v_1} = f_x, \sigma_g = -\frac{(f-b)f_x}{b}, \sigma_h = -\frac{(f-b)f_y}{b}, \sigma_f = -\frac{f^2 - 2bf - 2b}{2b} \quad (29)$$

(ii) time transformation invariant symmetries by setting $c_1 = c_2 = c_4 = c_5 = c_6 = c_7 = c_8 = x_0 = 0$ and $c_3 = 1$

$$\sigma_{u_1} = u_{1,t}, \sigma_{v_1} = v_{1,t}, \sigma_g = g_t, \sigma_h = h_t, \sigma_f = f_t \quad (30)$$

(iii) y-space transformation invariant symmetries by setting $c_1 = c_2 = c_3 = c_4 = c_6 = c_7 = c_8 = x_0 = 0$ and $c_5 = 1$

$$\sigma_{u_1} = u_{1,y}, \sigma_{v_1} = v_{1,y}, \sigma_g = g_y, \sigma_h = h_y, \sigma_f = f_y \quad (31)$$

(iv) scaling transformation invariant related symmetries by setting $c_1 = c_3 = c_4 = c_5 = c_6 = c_7 = c_8 = x_0 = 0$ and $c_2 = 2$

$$\begin{aligned}\sigma_{u_1} &= xu_{1,x} + yu_{1,y} + 2tu_{1,t} + u_1, \sigma_{v_1} = xv_{1,x} + yv_{1,y} + 2tv_{1,t} + v_1, \sigma_g = xg_x + yg_y + 2tg_t + g, \\ \sigma_h &= xh_x + yh_y + 2th_t + h, \sigma_f = xf_x + yf_y + 2tf_t\end{aligned}\quad (32)$$

(v) time relevant x-space transformation invariant symmetries by setting $c_1 = c_2 = c_3 = c_4 = c_5 = c_6 = c_7 = c_8 = 0$

$$\sigma_{u_1} = x_0u_{1,x}, \sigma_{v_1} = \frac{ax_0v_{1,x} + x_{0,t}}{a}, \sigma_g = x_0g_x, \sigma_h = x_0h_x, \sigma_f = x_0f_x \quad (33)$$

(vi) Galileo transformation invariant symmetries by setting $c_1 = c_2 = c_3 = c_5 = c_6 = c_7 = c_8 = x_0 = 0$ and $c_4 = 1$

$$\sigma_{u_1} = 1 + tu_{1,y}, \sigma_{v_1} = tv_{1,y}, \sigma_g = tg_y, \sigma_h = th_y, \sigma_f = tf_y \quad (34)$$

(vii) time-Mobious transformation invariant symmetries by setting $c_2 = c_3 = c_4 = c_5 = c_6 = c_7 = c_8 = x_0 = 0$ and $c_1 = 2$

$$\begin{aligned}\sigma_{u_1} &= xtu_{1,x} + ytu_{1,y} + t^2u_{1,t} + tu_1 + y, \sigma_{v_1} = xtv_{1,x} + ytv_{1,y} + t^2v_{1,t} + tv_1 + \frac{x}{a}, \\ \sigma_g &= xtg_x + ytg_y + t^2g_t + gt, \sigma_h = xth_x + yth_y + t^2h_t + ht, \sigma_f = xtf_x + ytf_y + t^2f_t\end{aligned}\quad (35)$$

Now let us proceed to reduce the prolonged system by standard Lie point symmetry group approach. Without loss of generality, we consider the symmetry reductions related to space-time invariant and residual symmetries by setting $c_1 = c_2 = c_4 = c_7 = 0$ and $c_3 = c_5 = c_6 = c_8 = x_0 = 1$. Substituting Eq. (28) into Eq. (27) with Eqs. (13) and (14), we get

$$\begin{aligned}\sigma_{u_1} &= u_{1,x} + u_{1,y} + u_{1,t} - f_y, \\ \sigma_{v_1} &= v_{1,x} + v_{1,y} + v_{1,t} - f_x, \\ \sigma_f &= \frac{1}{2b}(2bf_x + 2bf_y + 2bf_t + f^2 - 2b)\end{aligned}\quad (36)$$

The corresponding group invariant solutions which can be obtained by solving Eqs. (9), (10), (11) and (36) with $\sigma_{u_1} = \sigma_{v_1} = \sigma_f = 0$ has the general form

$$\begin{aligned}f &= \sqrt{2b} \tanh\left(\frac{x+F}{\sqrt{2b}}\right), u_1 = \sqrt{2b}F_Y \tanh\left(\frac{x+F}{\sqrt{2b}}\right) + U_1, \\ v_1 &= -\sqrt{2b}(F_Y + F_T - 1) \tanh\left(\frac{x+F}{\sqrt{2b}}\right) + V_1\end{aligned}\quad (37)$$

where the group invariant variables are $Y = -x + y$ and $T = -x + t$, while F , U_1 and V_1 are all group invariant functions of Y and T .

Substituting Eq. (37) into Eq. (12) we obtain the reduction solution for Burgers system (1)

$$\begin{aligned} u &= \sqrt{2b}F_Y \tanh^{-1}\left(\frac{x+F}{\sqrt{2b}}\right) + U_1, \\ v &= -\sqrt{2b}(F_Y + F_T - 1) \tanh^{-1}\left(\frac{x+F}{\sqrt{2b}}\right) + V_1 \end{aligned} \quad (38)$$

In order to get the symmetry reduction equations for the group invariant functions F , U_1 and V_1 , we first substitute Eq. (37) into Eq. (9) and find

$$U_{1,Y} + U_{1,T} + V_{1,Y} = 0. \quad (39)$$

Next, we substitute Eq. (37) into Eq. (10) and vanishing different powers of $\tanh(\frac{x+F}{\sqrt{2b}})$ we get

$$(U_1 - aV_1)F_Y + 2baF_{YT} - (1 + aV_1)F_T + b(a+1)F_{YY} + abF_{TT} + aV_1 = 0. \quad (40)$$

Finally, we get three symmetry reduction equations by substituting Eq. (37) into Eq. (11) and vanishing different powers of $\tanh(\frac{x+F}{\sqrt{2b}})$, two of them are trivial leaving only the third one

$$\begin{aligned} (aV_1 - U_1)U_{1,Y} + (aV_1 - U_1)F_Y^2 + [(1 + aV_1)F_T - baF_{TT} - 4baF_{YT} - 3b(a+1)F_{YY} \\ - aV_1]F_Y + (1 + aV_1)U_{1,T} - 2ba(F_{YY} + F_{YT})F_T + b[2aF_{YT} - (a+1)U_{1,YY} \\ - aU_{1,TT} + 2aF_{YY} - 2aU_{1,YT}] = 0 \end{aligned} \quad (41)$$

The entrance of the tanh part in Eqs. (37) indicate the intrusion of an additional soliton to the Burgers wave. In other words, the group invariant solution (37) is an interaction solution of one soliton and one general Burgers wave.

III. BÄCKLUND TRANSFORMATIONS AND INTERACTION SOLUTIONS OF THE BURGERS EQUATION

The tanh function is the simplest automorphic function which possesses the property that its derivatives can be explicitly expressed by itself. Enlightened by the form of reduction

solutions of Burgers equation in (37), we now use the general tanh function expansion method to obtain interaction solutions between solitons and Burgers waves.

The general truncated tanh expansion for (2+1)-dimensional Burgers system (1) has the form

$$u = w_1 \tanh(\lambda t + \phi) + w_0 \quad (42a)$$

$$v = p_1 \tanh(\lambda t + \phi) + p_0 \quad (42b)$$

where w_1, w_0, p_1, p_0 and ϕ are arbitrary functions of x, y and t . The λt in Eq. (42) have been added for later convenience.

Now, substituting the tanh expansion (42) into the Burgers system (1) and vanishing different powers of $\tanh(\phi)$ in both expansion equations we can prove the following Bäcklund transformation theorem.

Theorem 3. If f_1 and f_2 is the solution of the Burgers system (1), then so is u and v given by

$$u = 2b(\tanh(\phi + \lambda t) - 1)\phi_y + f_1 \quad (43a)$$

$$v = 2b(\tanh(\phi + \lambda t) - 1)\phi_x + f_2 \quad (43b)$$

with

$$\phi_t = b\phi_{yy} - 2b\phi_y^2 + f_1\phi_y + ab\phi_{xx} - 2ab\phi_x^2 + af_2\phi_x - \lambda \quad (44)$$

Proof. Substituting Eq. (42) into Burgers system (1) with $\lambda = 0$ temporarily, we get

$$\begin{aligned} & -w_1(2b\phi_y^2 + 2ab\phi_x^2 - w_1\phi_y - ap_1\phi_x) \tanh^3(\phi) + (2bw_{1,y}\phi_y + w_0w_1\phi_y + abw_1\phi_{xx} \\ & \quad - w_1w_{1,y} + 2abw_{1,x}\phi_x + ap_0w_1\phi_x - ap_1w_{1,x} + bw_1\phi_{yy} - w_1\phi_t) \tanh^2(\phi) \\ & + (-abw_{1,xx} + 2bw_1\phi_y^2 - w_1w_{0,y} - bw_{1,yy} - w_0w_{1,y} - ap_1w_1\phi_x - ap_1w_{0,x} - ap_0w_{1,x} - w_1^2\phi_y \\ & \quad + 2abw_1\phi_x^2 + w_{1,t}) \tanh(\phi) - 2abw_{1,x}\phi_x - abw_1\phi_{xx} + w_1\phi_t - 2bw_{1,y}\phi_y + w_{0,t} - abw_{0,xx} \\ & \quad - w_0w_{0,y} - ap_0w_1\phi_x - bw_{0,yy} - w_0w_1\phi_y - ap_0w_{0,x} - bw_1\phi_{yy} = 0 \end{aligned} \quad (45)$$

and

$$(-w_1\phi_x + p_1\phi_y) \tanh^2(\phi) + (w_{1,x} - p_{1,y}) \tanh(\phi) + w_1\phi_x + w_{0,x} - p_1\phi_y - p_{0,y} = 0. \quad (46)$$

Vanishing the coefficient of $\tanh^3(\phi)$ in Eq. (45) and $\tanh^2(\phi)$ in Eq. (46) we get the solutions

$$w_1 = 2b\phi_y, p_1 = 2b\phi_x \quad (47)$$

The coefficients of $\tanh^2(\phi)$ and $\tanh(\phi)$ in Eq. (45) are equivalent after using Eq. (47), which reads

$$\phi_t = b\phi_{yy} + w_0\phi_y + ab\phi_{xx} + ap_0\phi_x \quad (48)$$

After using Eq. (47), Eq. (46) is degenerated into the Burgers Eq. (1b)

$$w_{0,x} - p_{0,y} = 0 \quad (49)$$

Vanishing the coefficient of $\tanh^0(\phi)$ in Eq. (45), we get

$$f_{1,t} - f_1f_{1,y} - af_2f_{1,x} - bf_{1,yy} - abf_{1,xx} = 0 \quad (50)$$

with

$$f_1 = w_0 + 2b\phi_y \quad (51)$$

and

$$f_2 = p_0 + 2b\phi_x \quad (52)$$

It can be easily verified that f_1 and f_2 also satisfy

$$f_{1,x} - f_{2,y} = 0 \quad (53)$$

It is obvious that Eqs. (50) and (53) are just the (2+1)-dimensional Burgers system. Substituting w_0 and p_0 by solving Eqs. (51) and (52) into Eq. (48) meanwhile replacing ϕ by $\lambda t + \phi$ we get the Eq. (44).

Now substituting w_0 , p_0 , w_1 and p_1 into Eq. (42) by using Eqs. (47), (51) and (52) we finally get the Bäcklund transformation (43), then the theorem is proved.

If the seed functions are taken as constants, i.e. $f_1 = c_1$ and $f_2 = c_2$, then Eq. (44) becomes a potential Burgers equation with source λ . Even for the seed solutions taken as constants, we can still obtain rich nontrivial solutions from the Bäcklund transformation (43). For instance, the straight line solution $\phi = k_1x + k_2y + \omega t$ of Eq. (43) leads to the single kink soliton solution.

$$u = -2bk_2\{\tanh[k_2(2bk_2 - c_1)t + ak_1(2bk_1 - c_2)t - k_1x - k_2y] + 1\} + c_1 \quad (54a)$$

$$v = -2bk_1\{\tanh[k_2(2bk_2 - c_1)t + ak_1(2bk_1 - c_2)t - k_1x - k_2y] + 1\} + c_2 \quad (54b)$$

The traveling wave solution of Eq. (44) with $f_1 = c_1$ and $f_2 = c_2$ leads to a kink soliton solution

$$\phi = -\frac{(K - k_2c_1 - ak_1c_2 + \omega)(k_1x + k_2y + \omega t - e_0)}{4b(k_2^2 + ak_1^2)} - \frac{1}{2} \ln\left[1 + \exp\left(\frac{-K(k_1x + k_2y + \omega t - e_1)}{b(k_2^2 + ak_1^2)}\right)\right], \quad (55)$$

where k_1, k_2, e_0, e_1 and ω are arbitrary constants and K is related to other constants by

$$K = \sqrt{k_2^2c_1^2 - 2k_2c_1\omega + 2ak_1c_1k_2c_2 + \omega^2 - 2ac_2k_1\omega + a^2c_2^2k_1^2 - 8k_2^2\lambda b - 8\lambda abk_1^2} \quad (56)$$

Substituting the kink soliton solution (55) into Eq. (43) with $f_1 = c_1$ and $f_2 = c_2$ we obtain the two soliton fusion solution for u and v .




FIG. 1: The kink soliton solution (55).

Now we draw the corresponding pictures respectively with the solutions (55), (43a) with (55) and the interaction structure between them by plotting projectively at the $y = 1$ plane while the parameters are fixed as

$$e_0 = 0, e_1 = 0, k_1 = 1, k_2 = 1, \omega = -1, a = 1, b = 1, c_1 = 3, c_2 = 3, \lambda = 1. \quad (57)$$

Figure 1 reveals that solution (55) of Eq. (44) is a kink soliton. From figure 2 which displays the soliton structure of Eq. (43a), we see that two kink solitons fused to one at some time. Figure 3 displays three soliton interaction structure between u of (43a) with $f_1 = c_1$ and ϕ of (55). It is interesting to find from this picture that three kink solitons fused to one after interaction. Because the field v of (43b) resemble u in form, the interaction structure between the field v with ϕ of (55) has the similar form as that showing in Fig. 3.




FIG. 2: The two kink solitons fusion structure of u in Eq. (43a) with Eq. (55).




FIG. 3: Evolution of three kink solitons fusion interaction.

IV. CONCLUSION AND DISCUSSION

In summary, the (2+1)-dimensional Burgers equation is investigated by multiple ways to find interactions among different nonlinear excitations. The nonlocal residual symmetries which corresponds to the generators of truncated Painlevé expansions has been localized in an new prolonged system. The new BT 2 and symmetry reduction solutions thus has been obtained by using classical Lie point symmetry approach. The reduction solutions reveals that new soliton has been introduced to interact with background waves. Enlightened by the form of reduction solutions of Burgers equation, we further studied this system by a new generalized tanh expansion method. With this method we give a new BT theorem 3 for Burgers system. It is interesting that from BT 3 we can get rich soliton structure even

for the constant seed. Especially, two kink solitons fusion structure and evolution of three kink solitons fusing to one has been displayed by plotting projectively. This fact implies that starting from any seed solution of an integrable model, the BT 3 will introduce an additional soliton to the original seed.

Acknowledgments

The authors are in debt to Prof. S.Y. Lou for his valuable comments and suggestions. This work was supported by the National Natural Science Foundation of China under Grant No. 10875078, 11305106 and the Natural Science Foundation of Zhejiang Province of China Grant No. Y7080455, LQ13A050001.

-
- [1] Korteweg D J and de Vries F 1895 Philos. Mag. 39 422 Crighton D G 1995 Acta Appl. Math. 39 39
 - [2] C. Sulem and P. L. Sulem, The Nonlinear Schrodinger Equation C Self-Focusing and Wave Collapse, (Springer-Verlag New York, Inc. 1999), P. G. Kevrekidis, Discrete Nonlinear Schodinger Equation C Mathematical Analysis, Numerical Computations and Physical Perspectives, (Springer-Verlag Berlin Heidelberg 2009).
 - [3] Rubinstein J 1970 J. Math. Phys. 11 258 Skyrme T H R 1958 Proc. Roy. Soc. (London) A 247 260 Skyrme T H R 1961 Proc. Roy. Soc. (London) A 262 237 Perring J K and Skyrme T H R 1962 Nucl. Phys. 31 550
 - [4] Davey A and Stewartson K 1974 Proc. R. Soc. A 338 17
 - [5] Bluman G W and Kumei S 1989 *Symmetries and Differential Equations* (Berlin: Springer-Verlag)
 - [6] Matveev VB, Salle MA. Darboux transformations and solitons. Berlin, Heidelberg: Springer-Verlag; 1991.
 - [7] Hirota R. Phys Rev Lett 1971;27:1192.
 - [8] Lou SY, Ni GJ. J Math Phys 1989;30:1614; Lou SY. J Phys A: Math Gen 1999;32:4521.
 - [9] Lou SY. Z Naturforsch 1998;53a:251.

- [10] Lie S. Vorlesungen ber Differentialgleichungen mit Bekannten Infinitesimalen Transformationen. (Teuber: Leipzig; 1891) (reprinted by Chelsea: New York; 1967).
- [11] G Bluman, A Cheviakov and S Anco 2010 Appl. Math. Sci. vol 168 Symmetry Methods Partial Differential Equations (New York: Springer)
- [12] Lou S Y 1993 Phys. Lett. B 302 261
- [13] X. R. Hu, S. Y. Lou and Y. Chen, Phys. Rev. E 85 (2012) 056607.
- [14] Cheng X P, Chen C L and Lou S Y 2012 Interactions among Different Types of Nonlinear Waves Described by the KadomtsevCPetviashvili Equation arXiv:1208.3259
- [15] HONG Ke-Zhu, WU B-in, and CHEN Xian-Fen Vol. 39, No. 4, April 15, 2003
- [16] Lou S Y and Lian Z J 2005 Chin. Phys. Lett. 22 1