

# DARBOUX TRANSFORMATION AND SOLUTIONS OF THE TWO-COMPONENT HIROTA-MAXWELL-BLOCH SYSTEM\*

JIEMING YANG, CHUANZHONG LI\*\*, Tiantian LI, ZHAONENG CHENG

*Department of Mathematics, Ningbo University, Ningbo, 315211, China*

**ABSTRACT.** In this paper, we derive  $n$ -fold Darboux transformation of the two-component Hirota and the Maxwell-Bloch (TH-MB) equations and its determinant representation. Using Darboux determinant representation, we provide soliton solutions, positon solutions of the TH-MB equations.

PACS numbers: 42.65.Tg, 42.65.Sf, 05.45.Yv.

Nonlinear science as a powerful subject explains all kinds of mysteries in the challenges of science and technology today. In the nonlinear science, solitons and positons in nonlinear dynamical systems attract a lot of research in oceanography, Bose-Einstein condensate, plasma, superfluid and especially in optics([1, 2, 3, 4, 5]).

For a reduced dynamical equation, the erbium-doped fibre system was proven to allow soliton-type pulse propagation with pumping[6]. The Lax pair and the exact soliton solution for Higher-order nonlinear Schrödinger and Maxwell-Bloch (HNLS-MB) equations with pumping was derived in [6]. Kodama[7] shows that with suitable transformation and omitting the higher-order terms, higher order Nonlinear Schrödinger equation can be reduced to the Hirota equation[8]. In a similar way, after suitable choice of self steepening and self frequency effects, the HNLS-MB equations can be reduced to a coupled system of the Hirota equation and MB equation (H-MB) which is governed by femtosecond pulse propagation through doped fibre[9].

The H-MB system has been shown to be integrable with Lax pair [9]. By an efficient method which is called Darboux transformation [10], the determinant representation of  $n$ -fold Darboux transformation of AKNS system was given in [11]. In [12, 13], Darboux transformation and the rogue wave solutions of the H-MB equations were obtained using the Darboux transformation. Recently,  $n$ -fold Darboux transformation and positon solutions of the inhomogeneous Hirota-Maxwell-Bloch system is revealed in [14]. Soliton interactions in a generalized inhomogeneous coupled Hirota-Maxwell-Bloch system were also considered in [15] with symbolic computation. The two-component Hirota and Maxwell-Bloch equations [16] as a coupled version of the Hirota and Maxwell-Bloch equations includes all the important effects like the SIT in a Kerr medium with magnetisation, higher order dispersion and self-steepening. These effects play a very important role in the ultra-short pulse propagation of the order of femto seconds in nonlinear optical fibres. That is why we consider the Darboux transformation which is very useful to derive all kinds of solutions of the TH-MB equations which should be useful in explaining phenomenons of nonlinear optics.

---

\*Supported by the National Natural Science Foundation of China No. 11201251, the Natural Science Foundation of Zhejiang Province No. LY12A01007, the Natural Science Foundation of Ningbo City under Grant No 2013A610105, and the K. C. Wong Magna Fund in Ningbo University.

\*\* Corresponding author. Email: lichuanzhong@nbu.edu.cn.

The two-component Hirota-Maxwell-Bloch equation is as following[16]

$$E_{1z} = \frac{1}{2}i\beta E_{1tt} + iA\beta E_1 + \epsilon(E_{1ttt} + 3AE_{1t} + 3BE_1) - p_1, \quad (1)$$

$$E_{2z} = \frac{1}{2}i\beta E_{2tt} + iA\beta E_2 + \epsilon(E_{2ttt} + 3AE_{2t} + 3BE_2) - p_2, \quad (2)$$

$$p_{1t} = -NE_1 + M_{11}E_1 + M_{21}E_2 + \omega ip_1, \quad (3)$$

$$p_{2t} = -NE_2 + M_{12}E_1 + M_{22}E_2 + \omega ip_2, \quad (4)$$

$$M_{11t} = -E_1p_1^* - E_1^*p_1, \quad (5)$$

$$M_{12t} = -E_1^*p_2 - E_2p_1^*, \quad (6)$$

$$M_{21t} = -E_1p_2^* - E_2^*p_1, \quad (7)$$

$$M_{22t} = -E_2p_2^* - E_2^*p_2, \quad (8)$$

$$N_t = E_1p_1^* + E_1^*p_1 + E_2p_2^* + E_2^*p_2, \quad (9)$$

$$A := |E_1|^2 + |E_2|^2, \quad B := E_{1t}E_1^* + E_{2t}E_2^*. \quad (10)$$

The corresponding linear eigenvalue problem can take the form

$$\Phi_t = U\Phi, \quad (11)$$

$$\Phi_z = V\Phi, \quad (12)$$

where

$$U = \begin{pmatrix} -i\lambda & E_1 & E_2 \\ -E_1^* & i\lambda & 0 \\ -E_2^* & 0 & i\lambda \end{pmatrix} = -i\lambda\sigma_3 + U_0, \quad (13)$$

$$V = \lambda^3V_3 + \lambda^2V_2 + \lambda V_1 + V_0 + \frac{i}{\lambda + \frac{\omega}{2}}V_{-1}, \quad (14)$$

$$\begin{aligned} V_3 &:= -8i\epsilon \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad V_2 := \begin{pmatrix} 0 & E_1 & E_2 \\ -E_1^* & 0 & 0 \\ -E_2^* & 0 & 0 \end{pmatrix}; \quad V_1 := \begin{pmatrix} |E_1|^2 + |E_2|^2 & E_{1t} & E_{2t} \\ E_{1t}^* & -|E_1|^2 & -E_2E_1^* \\ E_{2t}^* & -E_1E_2^* & -|E_2|^2 \end{pmatrix}; \\ V_0 &:= \begin{pmatrix} E_{1t}E_1^* - E_{1t}^*E_1 + E_{2t}E_2^* - E_{2t}^*E_2 & E_{1tt} + 2E_1(|E_1|^2 + |E_2|^2) & E_{2tt} + 2E_2(|E_1|^2 + |E_2|^2) \\ -E_{1tt}^* - 2E_1^*(|E_1|^2 + |E_2|^2) & -(E_{1t}E_1^* - E_{1t}^*E_1) & -(E_{2t}E_1^* - E_{1t}^*E_2) \\ -E_{2tt}^* - 2E_2^*(|E_1|^2 + |E_2|^2) & -(E_{1t}E_2^* - E_{2t}^*E_1) & -(E_{2t}E_2^* - E_{2t}^*E_2) \end{pmatrix}; \\ V_{-1} &:= \frac{1}{2} \begin{pmatrix} N & p_1 & p_2 \\ p_1^* & M_{11} & M_{12} \\ p_2^* & M_{21} & M_{22} \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad U_0 = \begin{pmatrix} 0 & E_1 & E_2 \\ -E_1^* & 0 & 0 \\ -E_2^* & 0 & 0 \end{pmatrix}. \end{aligned} \quad (15)$$

Here  $E_1$  and  $E_2$  are the slowly varying amplitudes of the signal,  $p_i, i = 1, 2$  and  $M_{ij}, i, j = 1, 2$  are the electric and magnetic polarizations respectively,  $\beta$  and  $\epsilon$  are constants and  $N$  is the concentration of resonant atoms,  $*$  means complex conjugation.

For the H-MB equation and inhomogeneous H-MB equation, the Darboux transformation is constructed in [12, 14]. But for the TH-MB, because of its complexity, the Darboux transformation and reduction conditions are still not clear. Using above linear equations of TH-MB equations, one-fold Darboux transformation for TH-MB equation will be introduced in the next section.

Firstly, one consider the transformation about linear function  $\Phi$  by

$$\Phi' = T\Phi = (\lambda A - S)\Phi, \quad (16)$$

where

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}, \quad S = \begin{pmatrix} s_{11} & s_{12} & s_{13} \\ s_{21} & s_{22} & s_{23} \\ s_{31} & s_{32} & s_{33} \end{pmatrix}. \quad (17)$$

New function  $\Phi'$  is supposed to satisfy

$$\Phi'_t = U'\Phi', \quad (18)$$

$$\Phi'_z = V'\Phi'. \quad (19)$$

Then matrix  $T$  must satisfy following identities

$$T_t + TU = U'T, \quad (20)$$

$$T_z + TV = V'T. \quad (21)$$

One can choose  $T = \lambda I - S$  and then the Darboux transformation on  $E_1, E_2, p_1, p_2, N, M_{11}, M_{12}, M_{21}, M_{22}$  can be got by eq. (20) and eq. (21) as following

$$U'_0 = U_0 + i[S, \sigma_3], \quad (22)$$

$$V'_{-1} = (S + \frac{\omega}{2})V_{-1}(S + \frac{\omega}{2})^{-1}, \quad (23)$$

and  $S$  should have a condition as  $s_{21} = -s_{12}^*, s_{31} = -s_{13}^*$ .

We suppose

$$S = H\Lambda H^{-1} \quad (24)$$

where  $\Lambda = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$ ,  $H = \begin{pmatrix} \Phi_1(\lambda_1, t, z) & \Phi_1(\lambda_2, t, z) & \Phi_1(\lambda_3, t, z) \\ \Phi_2(\lambda_1, t, z) & \Phi_2(\lambda_2, t, z) & \Phi_2(\lambda_3, t, z) \\ \Phi_3(\lambda_1, t, z) & \Phi_3(\lambda_2, t, z) & \Phi_3(\lambda_3, t, z) \end{pmatrix} := \begin{pmatrix} \Phi_{1,1} & \Phi_{1,2} & \Phi_{1,3} \\ \Phi_{2,1} & \Phi_{2,2} & \Phi_{2,3} \\ \Phi_{3,1} & \Phi_{3,2} & \Phi_{3,3} \end{pmatrix}$ .

In order to satisfy the constraints of  $S$ , meanwhile make  $V'_{-1}$  having similar form as  $V_{-1}$ , i.e.  $s_{21} = -s_{12}^*, s_{31} = -s_{13}^*$ , following constraint will be considered

$$\lambda_3 = \lambda_2 = \lambda_1^*, \quad H = \begin{pmatrix} \Phi_1(\lambda_1, t, z) & \Phi_2^*(\lambda_1, t, z) & \Phi_3^*(\lambda_1, t, z) \\ \Phi_2(\lambda_1, t, z) & -\Phi_1^*(\lambda_1, t, z) & 0 \\ \Phi_3(\lambda_1, t, z) & 0 & -\Phi_1^*(\lambda_1, t, z) \end{pmatrix}.$$

As the simplest Darboux transformation, the determinant representation of one-fold Darboux transformation of the TH-MB equations will be given in the following theorem.

*Theorem 1:* The one-fold Darboux transformation of the TH-MB equations is as following

$$T_1(\lambda, \lambda_1, \lambda_2, \lambda_3) = \lambda I + t_0^{[1]} = \frac{1}{\Delta_1} \begin{pmatrix} (\mathbb{T}_1)_{11} & (\mathbb{T}_1)_{12} & (\mathbb{T}_1)_{13} \\ (\mathbb{T}_1)_{21} & (\mathbb{T}_1)_{22} & (\mathbb{T}_1)_{23} \\ (\mathbb{T}_1)_{31} & (\mathbb{T}_1)_{32} & (\mathbb{T}_1)_{33} \end{pmatrix}, \quad (25)$$

where for  $1 \leq i, j, k, m \leq 3$

$$\Delta_1 = \det(W_1)_{3 \times 3}, \quad (\mathbb{T}_1)_{km} = \det \begin{pmatrix} p_m & \lambda^{\delta_{km}} \\ W_1 & q_k^T \end{pmatrix}_{4 \times 4}.$$

The matrix  $W_1$  and row vectors  $p_m, q_k$  are defined as,

$$(W_1)_{ij} = \Phi_{j,i}, \quad (p_m)_j = \delta_{m,j}, \quad q_k = (\lambda_1 \Phi_{k,1} \quad \lambda_2 \Phi_{k,2} \quad \lambda_3 \Phi_{k,3}). \quad (26)$$

The following identities gives Darboux transformation of the TH-MB equation,

$$U^{[1]} = U + [\sigma_3, T_1], \quad V_{-1}^{[1]} = T_1|_{\lambda=-\frac{\omega}{2}} V_{-1} T_1^{-1}|_{\lambda=-\frac{\omega}{2}}. \quad (27)$$

Eqs.(27) will give one-fold Darboux transformation about  $E_1, E_2$  as

$$E_1^{[1]} = E_1 - 2is_{12} = E_1 + 2i \frac{(\mathbb{T}_1)_{12}}{\Delta_1}, \quad E_2^{[1]} = E_2 - 2is_{13} = E_2 + 2i \frac{(\mathbb{T}_1)_{13}}{\Delta_1}, \quad (28)$$

and one-fold Darboux transformation about  $p_1, p_2, N, M_{11}, M_{12}, M_{21}, M_{22}$  of the TH-MB equations in complicated forms.

Because the one-fold Darboux transformation about  $p_1, p_2, N, M_{11}, M_{12}, M_{21}, M_{22}$  of the TH-MB equations is in very complicated forms, therefore we have to omit it here for saving space. This one-fold transformation will be used to generate one-soliton solution from trivial seed solution of the TH-MB equation.

In the next part, we will give determinant representation of the n-fold Darboux transformation of the TH-MB equations. Firstly, we introduce  $3n$  eigenfunctions  $\Phi_i = \begin{pmatrix} \Phi_{1,i} \\ \Phi_{2,i} \\ \Phi_{3,i} \end{pmatrix}, i = 1, 2, \dots, 3n$  and  $\Phi_i = \Phi(\lambda = \lambda_i)$ , with constraints on eigenvalues as  $\lambda_{3n+1} = \lambda_{3n} = \lambda_{3n-1}^*$  and the reduction conditions on eigenfunctions as

$$\Phi_{3n-1,2} = \Phi_{3n,3} = -\Phi_{3n-2,1}^*, \quad \Phi_{3n-2,2} = \Phi_{3n-1,1}^*, \quad \Phi_{3n-2,3} = \Phi_{3n,1}^*, \quad \Phi_{3n-1,3} = \Phi_{3n,2} = 0. \quad (29)$$

We can also derive the n-fold Darboux transformation in form of a huge determinant in the following theorem.

*Theorem 2:* The n-fold Darboux transformation for solutions of the TH-MB equations can be represented as

$$\begin{aligned} T_n(\lambda; \lambda_1, \lambda_2, \lambda_3, \lambda_4, \dots, \lambda_{3n}) &= \lambda^n I + t_{n-1}^{[n]} \lambda^{n-1} + \dots + t_1^{[n]} \lambda + t_0^{[n]} \\ &= \frac{1}{\Delta_n} \begin{pmatrix} (\mathbb{T}_n)_{11} & (\mathbb{T}_n)_{12} & (\mathbb{T}_n)_{13} \\ (\mathbb{T}_n)_{21} & (\mathbb{T}_n)_{22} & (\mathbb{T}_n)_{23} \\ (\mathbb{T}_n)_{31} & (\mathbb{T}_n)_{32} & (\mathbb{T}_n)_{33} \end{pmatrix}, \end{aligned} \quad (30)$$

$$U_0^{[n]} = U_0 + t_{n-1}^{[n]} (-i\sigma_3) - (-i\sigma_3) t_{n-1}^{[n]}, \quad (31)$$

$$V_{-1}^{[n]} = T_n|_{\lambda=-\frac{\omega}{2}} V_{-1} T_n^{-1}|_{\lambda=-\frac{\omega}{2}}, \quad (32)$$

where for  $1 \leq i, j, k, m \leq 3n$

$$\Delta_n = \det(W_n)_{3n \times 3n}, \quad (\mathbb{T}_n)_{km} = \det \begin{pmatrix} p_m & \lambda^{n\delta_{km}} \\ W_n & q_k^T \end{pmatrix}_{(3n+1) \times (3n+1)}.$$

The matrix  $W_n$  and row vectors  $p_m, q_k$  are defined as,

$$(W_n)_{ij} = \Phi_{\bar{j},i} \lambda_i^{\lfloor \frac{j}{3} \rfloor}, \quad (p_m)_j = \delta_{m,\bar{j}} \lambda^{\lfloor \frac{j}{3} \rfloor}, \quad q_k = (\lambda_1^n \Phi_{k,1} \quad \lambda_2^n \Phi_{k,2} \quad \dots \quad \lambda_{3n}^n \Phi_{k,3n}). \quad (33)$$

$\lfloor \frac{j}{3} \rfloor$  means the floor function of  $\frac{j}{3}$ , and  $\bar{j}$  means the remainder of  $j$  modulo 3.

The  $n$ -th new solution  $E_1^{[n]}, E_2^{[n]}$  after the  $n$ -fold Darboux transformation of the TH-MB equations will be

$$E_1^{[n]} = E_1 + 2i(t_{n-1}^{[n]})_{12}, \quad E_2^{[n]} = E_2 + 2i(t_{n-1}^{[n]})_{13}, \quad (34)$$

where  $(t_{n-1}^{[n]})_{12}$  is the element at the first row and second column in the matrix of  $t_{n-1}^{[n]}$ . The formula for the new solutions  $p_1^{[n]}, p_2^{[n]}, N^{[n]}, M_{11}^{[n]}, M_{12}^{[n]}, M_{21}^{[n]}, M_{22}^{[n]}$  is just as the solutions after one fold Darboux transformation  $p_1^{[1]}, p_2^{[1]}, N^{[1]}, M_{11}^{[1]}, M_{12}^{[1]}, M_{21}^{[1]}, M_{22}^{[1]}$  in by changing  $\mathbb{T}_1$  to  $\mathbb{T}_n$ . So far, we discussed about the determinant construction of  $n$ -th Darboux transformation of the TH-MB equations. As an application of these transformations of the TH-MB equations, soliton solutions and positon solutions will be constructed in the next section. Particularly we use the 2-fold Darboux transformation to generate two-soliton solutions and positon solutions of the TH-MB equation.

We assume trivial seed solutions as  $E_1 = 0; E_2 = 0; p_1 := 0; p_2 := 0; N := 1; M_{11} := 1; M_{12} := 0; M_{21} := 0; M_{22} := 1$ , then the linear equations lead to eigenfunctions

$$\Phi_1 = \exp\left(\frac{1}{2\lambda + \omega}zi - \lambda ti\right), \quad (35)$$

$$\Phi_2 = \Phi_3 = \exp\left(\frac{(-16\epsilon\lambda^4 - 8\epsilon\lambda^3\omega + 4\beta\lambda^3 + 2\beta\lambda^2\omega + 1)zi}{2\lambda + \omega} + \lambda ti\right). \quad (36)$$

Substituting these two eigenfunctions into the one-fold Darboux transformation eq.(28) and choosing  $\lambda = \alpha_1 + \beta_1 i$ , then the following soliton solutions are obtained:

$$E_1^{[1]} = E_2^{[1]} = -4\beta_1 \frac{e^{\frac{2A}{(2\alpha_1+2i\beta_1+\omega)(-2\alpha_1+2i\beta_1-\omega)}}}{e^{\frac{16z\epsilon\alpha_1^2\beta_1(12\alpha_1^2+12\omega\alpha_1+3\omega^2+8\beta_1^2)}{(2\alpha_1+2i\beta_1+\omega)(-2\alpha_1+2i\beta_1-\omega)}} + 2e^{\frac{4\beta_1 B}{(2\alpha_1+2i\beta_1+\omega)(-2\alpha_1+2i\beta_1-\omega)}}},$$

where

$$\begin{aligned} A : &= 4t\beta_1^3 + 48z\epsilon\alpha_1^4\beta_1 + 32z\epsilon\alpha_1^2\beta_1^3 + 8z\beta\alpha_1^3\beta_1 + 8z\beta\beta_1^3\alpha_1 \\ &+ 4z\epsilon\omega^2\beta_1^3 + 4t\omega\alpha_1\beta_1 + t\alpha_1\omega^2i - 16iz\epsilon\alpha_1^5 - 4iz\beta\beta_1^4 + 4iz\beta\alpha_1^4 \\ &+ 4it\omega\alpha_1^2 + 4it\alpha_1\beta_1^2 + 48z\epsilon\omega\alpha_1^3\beta_1 + 12z\epsilon\omega^2\alpha_1^2\beta_1 + 8z\beta\omega\alpha_1^2\beta_1 \\ &+ 2z\beta\omega^2\alpha_1\beta_1 + 16z\epsilon\omega\alpha_1\beta_1^3 + 16z\epsilon\beta_1^5 + 4t\alpha_1^2\beta_1 + t\beta_1\omega^2 + 4it\alpha_1^3 \\ &+ z\beta\omega^2\alpha_1^2i + 32iz\epsilon\alpha_1^3\beta_1^2 + 48iz\epsilon\beta_1^4\alpha_1 + 4iz\beta\alpha_1^3\omega - 16iz\epsilon\alpha_1^4\omega \\ &- 4iz\epsilon\omega^2\alpha_1^3 - iz\beta\omega^2\beta_1^2 - 4iz\beta\alpha_1\beta_1^2\omega + 12iz\epsilon\omega^2\alpha_1\beta_1^2 + 48iz\epsilon\alpha_1^2\beta_1^2\omega, \\ B : &= 16z\epsilon\beta_1^4 + 4t\alpha_1^2 + 8z\beta\omega\alpha_1^2 + 2z\beta\omega^2\alpha_1 + 4t\omega\alpha_1 + 16z\epsilon\omega\alpha_1\beta_1^2 \\ &+ 4z\epsilon\omega^2\beta_1^2 + 4t\beta_1^2 + 8z\beta\beta_1^2\alpha_1 + t\omega^2 + 8z\beta\alpha_1^3. \end{aligned}$$

Similarly, substituting the eigenfunctions eq.(35)-eq.(36) into the one-fold Darboux transformation eq.(28), and taking  $\alpha = 2, \beta = -1$ , then the one-soliton solutions of the classical TH-MB equations can be obtained whose evolution is given in Fig.1, which clearly indicates that  $E_1 = E_2$  are bright solitons because their waves are above the flat non-vanishing plane. The other solutions keep initial values as following

$$N^{[1]} = 1, M_{11}^{[1]} = 1, M_{12}^{[1]} = 0, M_{21}^{[1]} = 0, M_{22}^{[1]} = 1, p_1^{[1]} = 0, p_2^{[1]} = 0. \quad (37)$$

After considering the two-fold Darboux transformation and supposing  $\alpha_1 = 1, \beta_1 = 1.5, \alpha_2 = 1.5, \beta_2 = 1, \omega = 1, \epsilon = 1, \beta = 1$ , two-soliton solutions  $E_1^{[2]}, E_2^{[2]}$  are as Fig. 2.




FIGURE 1. One solution solution  $E_1^{[1]} = E_2^{[1]}$  of the TH-MB equations when  $\epsilon = 1, \beta = 1, \omega = 1, \alpha_1 = 1, \beta_1 = 2$ .




FIGURE 2. Two solution solution  $E_1^{[2]} = E_2^{[2]}$  of the TH-MB equations when  $\alpha_1 = 1, \beta_1 = 1.5, \alpha_2 = 1.5, \beta_2 = 1, \omega = 1, \epsilon = 1, \beta = 1$ .

For the above two-soliton solutions, if the second spectral parameter  $\lambda_2$  is close to the first spectral parameter  $\lambda_1$ , doing the Taylor expansion of wave function to first order up to  $\lambda_1$  will lead to a new kind of solutions which is called positon solutions. Firstly, following four linear eigenfunctions will be got which will be used to construct the second Darboux transformation and to generate the positon solutions,

$$\Phi_{1,1} = \exp\left(\frac{1}{2\lambda_1 + \omega}zi - \lambda_1 ti\right), \quad \Phi_{4,1} = \exp\left(\frac{1}{2\lambda_2 + \omega}zi - \lambda_2 ti\right), \quad (38)$$

$$\Phi_{2,1} = \Phi_{3,1} = \exp\left(\frac{(-16\epsilon\lambda_1^4 - 8\epsilon\lambda_1^3\omega + 4\beta\lambda_1^3 + 2\beta\lambda_1^2\omega + 1)zi}{2\lambda_1 + \omega} + \lambda_1 ti\right), \quad (39)$$

$$\Phi_{5,1} = \Phi_{6,1} = \exp\left(\frac{(-16\epsilon\lambda_2^4 - 8\epsilon\lambda_2^3\omega + 4\beta\lambda_2^3 + 2\beta\lambda_2^2\omega + 1)zi}{2\lambda_2 + \omega} + \lambda_2 ti\right). \quad (40)$$

Now we take  $\lambda_2 = \lambda_1 + \epsilon(1+i)$  and using the Taylor expansion of wave function  $\Phi_{4,1}, \Phi_{5,1}, \Phi_{6,1}$  up to first order of  $\epsilon$  in terms of  $\lambda_1$ . On substitution of these manipulations into the second Darboux transformation discussed in the last section will help us to derive positon solutions. The pictorial representation of positon solutions of the TH-MB equations, i.e. the case when  $\alpha_1 = 1, \beta_1 = 1.5, \omega = 1, \epsilon = 1, \beta = 2, \omega = 1.5$  is shown in Fig.3.




FIGURE 3. One positon solution  $(E_1, E_2)$  of the TH-MB equations when  $\alpha_1 = 1, \beta_1 = 1.5, \omega = 1, \epsilon = 1, \beta = 2, \omega = 1.5$ .

When  $\alpha_1 = 1, \beta_1 = 1.5, \omega = 1, \epsilon = 1, \beta = 0, \omega = 1.5$ , i.e. the case of two-component complex modified Korteweg-de Vries(TCMKdV)-MB equations, the picture of positon solutions of the TCMKdV-MB equations is plotted in Fig.4.




FIGURE 4. One positon solution  $(E_1, E_2)$  of the TMKdV-MB equations when  $\alpha_1 = 1, \beta_1 = 1.5, \omega = 1, \epsilon = 1, \beta = 0, \omega = 1.5$ .

Comparing positons with two solitons, one can find positons are long-range analogues of solitons and slowly decreasing, oscillating solutions. Though each branch of the positon will separate finally during their propagating, the separation becomes slower and slower because  $\lambda_2$  infinitely approach to  $\lambda_1$ .

In a similar way, using the higher order Darboux transformation, one can also generate higher-order positon solutions which will not be included here.

The authors would like to thank Professor Jingsong He for helpful discussions.

## REFERENCES

- [1] Osborne A 2010 Nonlinear Ocean Waves and the Inverse Scattering Transform (New York, Elsevier)
- [2] Hasegawa A and Tappert F 1973 Appl. Phys. Lett. 23 142-144
- [3] Wu Y and Yang X 2007 Appl. Phys. Lett. 91 094104
- [4] Wu Y 2005 Phys. Rev. A 71 053820
- [5] Wu Y and Deng L 2004 Opt. Lett. 29 2064-2066
- [6] Nakkeeran K 2000 J. Phys. A: Math. Gen. 33 4377-4381
- [7] Kodama Y 1985 Phys. Lett. A 112 193-196
- [8] Hirota R 1973 J. Math. Phys. 14 805

- [9] Porsezian K and Nakkeeran K 1995 Phys. Rev. Lett. 74 2941
- [10] Matveev V B Salle M A 1991 Darboux transformations and solitons (Berlin, Springer)
- [11] He J S Zhang L Cheng Y and Li Y S 2006 Sci. China A 12 1867-78
- [12] Li C Z, He J S and Porsezian K 2013 Physical Review E 87 012913
- [13] Li C Z, He J S and Porsezian K 2013 Chin. Phys. B 22 4 044208
- [14] Li C Z, He J S 2013 SCIENCE CHINA Physics Mechanics and Astronomy(in press) arXiv:1210.2501
- [15] Xue Y S, Tian B, et al. 2011 Nonlinear Dynamics 67 2799-2806
- [16] Porsezian K, Mahalingam A and Shanmugha Sundaram P 2000 Chaos, Solitons and Fractals 11 1261-1264