Proton Structure and Tensor Gluons

George Savvidy

Institute of Nuclear and Particle Physics

Demokritos National Research Center, Ag. Paraskevi, Athens, Greece

Abstract

We consider a possibility that inside the proton and, more generally, inside the hadrons there are additional partons - tensor-gluons, which can carry a part of the proton momentum. The tensor-gluons have zero electric charge, like gluons, but have a larger spin. Inside the proton a nonzero density of the tensor-gluons can be generated by the emission of tensor-gluons by gluons. The last mechanism is typical for non-Abelian tensor gauge theories, in which there exists a gluon-tensor-tensor vertex of order g. Therefore the number of gluons changes not only because a quark may radiate a gluon or because a gluon may split into a quark-antiquark pair or into two gluons, but also because a gluon can split into two tensor-gluons. The process of gluon splitting suggests that part of the proton momentum which was carried by neutral partons is shared between vector and tensor gluons. We derive evolution equations for the parton distribution functions which take into account these new processes. The momentum sum rule allows to find the tensor-gluons contribution to the Callan-Simanzik beta function and to calculate the corresponding anomalous dimensions. This contribution changes the behavior of the structure functions, and the logarithmic correction to the Bjorken scaling becomes more mild. This also influences the unification scale at which the coupling constants of the Standard Model merge, shifting its value to lower energies of order of 40 TeV.

1 Introduction

It was predicted that the Bjorken scaling should be broken by logarithms of a transverse momentum Q^2 and that these deviations from the scaling law can be computed for the deep inelastic structure functions [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12]. In the leading logarithmic approximation the results can be formulated in the parton language [13] by assigning the well determined Q^2 dependence to the parton densities [1, 2, 14, 15, 3].

In this article we shall consider a possibility that inside the proton and, more generally, inside the hadrons there are additional partons - tensor-gluons, which can carry a part of the proton momentum. Tensor-gluons have zero electric charge, like gluons, but have a larger spin. Inside the proton a nonzero density of the tensor-gluons can be generated by the emission of tensor-gluons by gluons [16, 17, 18, 19]. The last mechanism is typical for non-Abelian tensor gauge theories, in which there exists a gluon-tensor-tensor vertex of order g [16, 17, 18, 19]. Therefore the number of gluons changes not only because a quark may radiate a gluon or because a gluon may split into a quark-antiquark pair or into two gluons [14, 15, 3], but also because a gluon can split into two tensor-gluons [16, 17, 18, 19, 20, 21]. The process of gluon splitting into tensor-gluons suggests that part of the proton momentum which was carried by neutral partons is shared between vector and tensor gluons.

The proposed model was formulated in terms of field theory Lagrangian [16, 17, 18, 19]. The Lagrangian describes the interaction of the gluons with their massless partners of higher spin [16, 17, 18, 19]. We shall call them tensor-gluons. The characteristic property of the model is that all interaction vertices between gluons and tensor-gluons have dimensionless coupling constants in four-dimensional space-time. That is, the cubic interaction vertices have only first order derivatives and the quartic vertices have no derivatives at all. These are familiar properties of the standard Yang-Mills field theory. In order to understand the physical properties of the model it was important to study the tree level scattering amplitudes. A very powerful spinor helicity technique [22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39] was used to calculate high order tree level diagrams with the participation of tensor-gluons in [20].

These tree level scattering amplitudes describe a fusion of gluons into tensor-gluons [20]. They are generalizations of the Parke-Taylor scattering amplitude to the case of two

tensor gauge bosons of spin s and (n-2) gluons. The result reads [20]:

$$M_n(1^+, ..i^-, ...k^{+s}, ..j^{-s}, ..n^+) = g^{n-2} \frac{\langle ij \rangle^4}{\prod_{r=1}^n \langle r(r+1) \rangle} \left(\frac{\langle ij \rangle}{\langle ik \rangle}\right)^{2s-2}, \tag{1.1}$$

where n is the total number of particles and the dots stand for the positive-helicity gluons. Here i is the position of the negative-helicity gluon, while k and j are the positions of the particles with helicities +s and -s respectively. For s=1 this expression reduces to the well-known result for the MHV amplitude [27]. The scattering amplitudes (1.1) were used to derive the amplitudes of splitting of gluon into tensor-gluons in [21].

Here we shall use the splitting amplitudes in order to derive the generalization of the DGLAP evolution equations for the parton distribution functions, which take into account the processes of emission of tensor-gluons by gluons. The momentum sum rule allows to find the tensor-gluons contribution to the one-loop Callan-Simanzik beta function, which takes the following form:

$$\alpha(t) = \frac{\alpha}{1 + b\alpha t}, \qquad b = \frac{\sum_{s} (12s^2 - 1)C_2(G) - 4n_f T(R)}{12\pi},$$
 (1.2)

where s is the spin of the gauge bosons. The matrix of anomalous dimensions for the twist-two operators γ_n can also be computed. The spin-dependent term in the Callan-Simanzik beta function coefficient (1.2) changes the $t = \ln(Q^2/Q_0^2)$ behavior of the structure functions demonstrating that if the tensor-gluons exist, the logarithmic correction to the Bjorken scaling changes. This also influences the unification scale at which the coupling constants of the Standard Model merge [47, 48], shifting its value to lower energies of order of $M \sim 40 \ TeV$:

$$\ln \frac{M}{\mu} = \frac{\pi}{58} \left(\frac{1}{\alpha_{el}(\mu)} - \frac{8}{3} \frac{1}{\alpha_s(\mu)} \right), \tag{1.3}$$

where $\alpha_{el}(\mu)$ and $\alpha_s(\mu)$ are the electromagnetic and strong coupling constants at scale μ .

The present paper is organized as follows. In section two the basic formulae for scattering amplitude and splitting functions are recalled, definitions and notations are specified. In section three the generalized evolution equations that describe the Q^2 dependence of parton densities is derived and the physical interpretation of the new terms is presented. In section four we obtain the one-loop Callan-Simanzik beta function for tensor-gluons and the anomalous dimensions for the singlet and nonsiglet scaling functions. In section five we discuss the unification of coupling constants of the Standard Model including one-loop contribution of the tensor-bosons. Section six contains concluding remarks and the Appendix - the details of the regularization scheme.

2 Splitting Functions

It is convenient to represent a scattering amplitude for the massless particles of momenta p_i and polarization tensors ε_i (i = 1, ..., n), which are described by irreducible massless representations of the Poincaré group and are classified by their helicities $h = \pm s$, in the following form:

$$M_n = M_n(p_1, \varepsilon_1; p_2, \varepsilon_2; ...; p_n, \varepsilon_n).$$

Further by representing the momenta p_i and polarization tensors ε_i in terms of spinors the scattering amplitude M_n can be considered as a function of spinors λ_i , $\tilde{\lambda}_i$ and helicities h_i [22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39]:

$$M_n = M_n(\lambda_1, \tilde{\lambda}_1, h_1; \dots; \lambda_n, \tilde{\lambda}_n, h_n) . \tag{2.4}$$

The advantage of the spinor representation is that introducing a complex deformation of the particles momenta one can derive a general form of the three-particle interaction vertices [34, 20]:

$$M_3(1^{h_1}, 2^{h_2}, 3^{h_3}).$$

The dimensionality of the three-point vertex $M_3(1^{h_1}, 2^{h_2}, 3^{h_3})$ is $[mass]^{D=\pm(h_1+h_2+h_3)}$.

In the generalized Yang-Mills theory [16, 17, 18, 19] all interaction vertices between highspin particles have dimensionless coupling constants, which means that the helicities of the interacting particles in the vertex are constrained by the relation $D = \pm (h_1 + h_2 + h_3) = 1$. Therefore the interaction vertex between massless tensor-gluons, the TTT-vertex, has the following form [20]:

$$M_{3} = gf^{abc} < 1, 2 >^{-2h_{1}-2h_{2}-1} < 2, 3 >^{2h_{1}+1} < 3, 1 >^{2h_{2}+1}, h_{3} = -1 - h_{1} - h_{2},$$

$$M_{3} = gf^{abc}[1, 2]^{2h_{1}+2h_{2}-1}[2, 3]^{-2h_{1}+1}[3, 1]^{-2h_{2}+1}, h_{3} = 1 - h_{1} - h_{2}, (2.5)$$

where f^{abc} are the structure constants of the internal gauge group G. In particular, considering the interaction between a gluon of helicity $h_1 = \pm 1$ and a tensor-gluon of helicity $h_2 = \pm s$, the GTT-vertex, one can find from (2.5) that

$$h_3 = \pm |s - 2|, \pm s, \pm |s + 2|$$
 (2.6)

and the corresponding gluon-tensor-tensor interaction vertices GTT have the form

$$M_3^{a_1 a_2 a_3}(1^{-s}, 2^{-1}, 3^{+s}) = g f^{a_1 a_2 a_3} \frac{\langle 1, 2 \rangle^4}{\langle 1, 2 \rangle \langle 2, 3 \rangle \langle 3, 1 \rangle} \left(\frac{\langle 1, 2 \rangle}{\langle 2, 3 \rangle} \right)^{2s-2},$$

$$M_3^{a_1 a_2 a_3}(1^{-s}, 2^{+1}, 3^{s-2}) = g f^{a_1 a_2 a_3} \frac{\langle 1, 3 \rangle^4}{\langle 1, 2 \rangle \langle 2, 3 \rangle \langle 3, 1 \rangle} \left(\frac{\langle 1, 2 \rangle}{\langle 2, 3 \rangle}\right)^{2s-2}.$$
 (2.7)

These are the vertices which reduce to the standard triple gluon vertex when s = 1. Using these vertices one can compute the scattering amplitudes of gluons and tensor-gluons. The color-ordered scattering amplitudes involving two tensor-gluons of helicities $h = \pm s$, one negative helicity gluon and (n-3) gluons of positive helicity were found in [20]:

$$\hat{M}_n(1^+, ..i^-, ...k^{+s}, ..j^{-s}, ..n^+) = ig^{n-2}(2\pi)^4 \delta^{(4)}(P^{ab}) \frac{\langle ij \rangle^4}{\prod_{l=1}^n \langle ll+1 \rangle} \left(\frac{\langle ij \rangle}{\langle ik \rangle}\right)^{2s-2}, \quad (2.8)$$

where n is the total number of particles and the dots stand for any number of positive helicity gluons, i is the position of the negative-helicity gluon, while k and j are the positions of the gluons with helicities +s and -s respectively. The expression (2.8) reduces to the famous Parke-Taylor formula [27] when s=1. In particular, the five-particle amplitude takes the following form:

$$\hat{M}_{5}(1^{+}, 2^{-}, 3^{+}, 4^{+s}, 5^{-s}) = ig^{3}(2\pi)^{4}\delta^{(4)}(P^{a\dot{b}}) \frac{\langle 25 \rangle^{4}}{\prod_{i=1}^{5} \langle ii+1 \rangle} (\frac{\langle 25 \rangle}{\langle 24 \rangle})^{2s-2}, \quad (2.9)$$

where $P^{ab} = \sum_{m=1}^{n} \lambda_m^a \tilde{\lambda}_m^b$ is the total momentum. The scattering amplitudes (2.8) and (2.9) can be used to extract splitting amplitudes of gluons and tensor-gluons [21]. The collinear behavior of the tree amplitudes has the following factorized form [26, 27, 28, 38, 39]:

$$M_n^{tree}(..., a^{\lambda_a}, b^{\lambda_b}, ...) \xrightarrow{a \parallel b} \sum_{\lambda=+1} Split_{-\lambda}^{tree}(a^{\lambda_a}, b^{\lambda_b}) \times M_{n-1}^{tree}(..., P^{\lambda}, ...),$$
 (2.10)

where $Split_{-\lambda}^{tree}(a^{\lambda_a}, b^{\lambda_b})$ denotes the splitting amplitude and the intermediate state P has momentum $k_P = k_a + k_b$ and helicity λ . Considering the amplitude (2.9) in the limit when the particles 4 and 5 become collinear, $k_4 \parallel k_5$, that is, $k_4 = zk_P$, $k_5 = (1-z)k_P$, $k_P^2 \to 0$ and z describes the longitudinal momentum sharing, one can deduce that the corresponding behavior of spinors is $\lambda_4 = \sqrt{z}\lambda_P$, $\lambda_5 = \sqrt{1-z}\lambda_P$, and that the amplitude (2.9) takes the following factorization form [21]:

$$M_5(1^+, 2^-, 3^+, 4^{+s}, 5^{-s}) = A_4(1^+, 2^-, 3^+, P^-) \times Split_+(a^{+s}, b^{-s}),$$

where

$$Split_{+}(a^{+s}, b^{-s}) = \left(\frac{1-z}{z}\right)^{s-1} \frac{(1-z)^2}{\sqrt{z(1-z)}} \frac{1}{\langle a, b \rangle}.$$
 (2.11)

In a similar way one can deduce that

$$Split_{+}(a^{-s}, b^{+s}) = \left(\frac{z}{1-z}\right)^{s-1} \frac{z^2}{\sqrt{z(1-z)}} \frac{1}{\langle a, b \rangle}.$$
 (2.12)

Considering different collinear limits $k_1 \parallel k_5$ and $k_3 \parallel k_4$ one can get [21]

$$Split_{+s}(a^{+}, b^{-s}) = \frac{(1-z)^{s+1}}{\sqrt{z(1-z)}} \frac{1}{\langle a, b \rangle}, \quad Split_{+s}(a^{-s}, b^{+}) = \frac{z^{s+1}}{\sqrt{z(1-z)}} \frac{1}{\langle a, b \rangle}$$
(2.13)

and

$$Split_{-s}(a^{+s}, b^{+}) = \frac{z^{-s+1}}{\sqrt{z(1-z)}} \frac{1}{\langle a, b \rangle}, \quad Split_{-s}(a^{+}, b^{+s}) = \frac{(1-z)^{-s+1}}{\sqrt{z(1-z)}} \frac{1}{\langle a, b \rangle}. \quad (2.14)$$

The set of splitting amplitudes (2.11)-(2.14) $G \to TT$, $T \to GT$ and $T \to TG$ reduces to the full set of gluon splitting amplitudes [26, 27, 28, 38, 39] when s = 1.

Since the collinear limits of the scattering amplitudes are responsible for parton evolution [3] we can extract from the above expressions the Altarelli-Parisi splitting probabilities for tensor-gluons. Indeed, the residue of the collinear pole in the square (of the factorized amplitude (2.10)) gives Altarelli-Parisi splitting probability P(z):

$$P(z) = C_2(G) \sum_{h_P, h_a, h_b} |Split_{-h_P}(a^{h_a}, b^{h_b})|^2 s_{ab}.$$
(2.15)

where $s_{ab} = 2k_a \cdot k_b = \langle a, b \rangle [a, b]$. The invariant operator C_2 for the representation R is defined by the equation $t^a t^a = C_2(R)$ 1 and $tr(t^a t^b) = T(R)\delta^{ab}$. Substituting the splitting amplitudes (2.11)-(2.14) into (2.15) we are getting

$$P_{TG}(z) = C_2(G) \left[\frac{z^4}{z(1-z)} \left(\frac{z}{1-z} \right)^{2s-2} + \frac{(1-z)^4}{z(1-z)} \left(\frac{1-z}{z} \right)^{2s-2} \right],$$

$$P_{GT}(z) = C_2(G) \left[\frac{1}{z(1-z)} \left(\frac{1}{1-z} \right)^{2s-2} + \frac{(1-z)^4}{z(1-z)} (1-z)^{2s-2} \right],$$

$$P_{TT}(z) = C_2(G) \left[\frac{z^4}{z(1-z)} z^{2s-2} + \frac{1}{z(1-z)} \left(\frac{1}{z} \right)^{2s-2} \right].$$
(2.16)

The momentum conservation in the vertices clearly fulfils because these functions satisfy the relations

$$P_{TG}(z) = P_{TG}(1-z), \quad P_{GT}(z) = P_{TT}(1-z), \qquad z < 1.$$
 (2.17)

In the leading order the kernel $P_{TG}(z)$ has a meaning of variation per unit t of the probability density of finding a tensor-gluon inside the gluon, $P_{GT}(z)$ - of finding gluon inside the

tensor-gluon and $P_{TT}(z)$ - of finding tensor-gluon inside the tensor-gluon. For completeness we shall present also quark and gluon splitting functions [3]:

$$P_{qq}(z) = C_2(R) \frac{1+z^2}{1-z},$$

$$P_{Gq}(z) = C_2(R) \frac{1+(1-z)^2}{z},$$

$$P_{qG}(z) = T(R)[z^2 + (1-z)^2],$$

$$P_{GG}(z) = C_2(G) \left[\frac{1}{z(1-z)} + \frac{z^4}{z(1-z)} + \frac{(1-z)^4}{z(1-z)} \right],$$
(2.18)

where $C_2(G) = N$, $C_2(R) = \frac{N^2 - 1}{2N}$, $T(R) = \frac{1}{2}$ for the SU(N) groups.

We have to notice that the limit $s \to 1/2$ in the scattering amplitudes (2.8) and (2.9) reduces them to the tree level gluon scattering amplitudes into a quark pair. Thus the formula has larger validity area than the area in which it has been initially derived [20]. This fact should be visible in the splitting probabilities as well. Indeed let us consider the tensor splitting functions (2.16) and take the limit to the half-integer spin $s \to 1/2$. One can see that tensor-gluon splitting probabilities (2.16) reduce to the quark-gluon splitting probabilities (2.18)

$$P_{TT}(z) \to P_{qq}(z), \quad P_{GT}(z) \to P_{Gq}(z), \quad P_{TG}(z) \to P_{qG}(z).$$
 (2.19)

Taking into account the remark which we made after the formula (2.14) one can see that the substitution $s \to 1$ gives gluon-gluon splitting function as well

$$\frac{1}{2}(P_{TG}(z) + P_{GT}(z) + P_{TT}(z)) \to P_{GG}(z). \tag{2.20}$$

Having in hand the new set of splitting probabilities for tensor-gluons (2.16) we can hypothesize that a possible emission of tensor-gluons should produce a tensor-gluon "shower" of neutral partons inside the proton in additional to the quark and gluon "shower". Our goal is to derive DGLAP equations [3, 5, 6, 7, 8, 9, 10, 4] which will take into account these new emission processes.

3 Generalization of DGLAP Equation

It is well known that the deep inelastic structure functions can be expressed in terms of parton quark densities. If $q^i(x)$ is the density of quarks of type i (summed over colors)

inside a proton target with fraction x of the proton longitudinal momentum in the infinite momentum frame then the scaling structure functions can be represented in the form

$$2F_1(x) = F_2(x)/x = \sum_i Q_i^2[q^i(x) + \bar{q}^i(x)].$$

The scaling behavior of the structure functions is broken and the results can be formulated by assigning a well determined Q^2 dependence to the parton densities. This can be achieved by introducing the integro-differential equations which describe the Q^2 dependence of quark $q^i(x,t)$ and gluon densities G(x,t), where $t = \ln(Q^2/Q_0^2)$ [3, 5, 6, 7, 8, 9, 10, 4].

Let us see what will happen if one supposes that there are additional partons - tensorsgluons - inside the proton. In accordance with our hypothesis there is an additional emission of tensor-gluons in the proton, therefore one should introduce the corresponding density T(x,t) of tensor-gluons (summed over colors) inside the proton in the P_{∞} frame. We can derive the integro-differential equations that describe the Q^2 dependence of parton densities in this general case. They are:

$$\frac{dq^{i}(x,t)}{dt} = \frac{\alpha(t)}{2\pi} \int_{x}^{1} \frac{dy}{y} \left[\sum_{j=1}^{2n_{f}} q^{j}(y,t) P_{q^{i}q^{j}}(\frac{x}{y}) + G(y,t) P_{q^{i}G}(\frac{x}{y}) \right], \tag{3.21}$$

$$\frac{dG(x,t)}{dt} = \frac{\alpha(t)}{2\pi} \int_{x}^{1} \frac{dy}{y} \left[\sum_{j=1}^{2n_{f}} q^{j}(y,t) P_{Gq^{j}}(\frac{x}{y}) + G(y,t) P_{GG}(\frac{x}{y}) + T(y,t) P_{GT}(\frac{x}{y}) \right],$$

$$\frac{dT(x,t)}{dt} = \frac{\alpha(t)}{2\pi} \int_{x}^{1} \frac{dy}{y} \left[G(y,t) P_{TG}(\frac{x}{y}) + T(y,t) P_{TT}(\frac{x}{y}) \right].$$

The $\alpha(t)$ is the running coupling constant ($\alpha = g^2/4\pi$). In the leading logarithmic approximation $\alpha(t)$ is of the form

$$\frac{\alpha}{\alpha(t)} = 1 + b \alpha t \quad , \tag{3.22}$$

where $\alpha = \alpha(0)$ and b is the one-loop Callan-Simanzik coefficient, which, as we shall see below, receives an additional contribution from the tensor-gluons. Here the indices i and j run over quarks and antiquarks of all flavors. The number of quarks of a given fraction of momentum changes when a quark looses momentum by radiating a gluon, or a gluon inside the proton may produce a quark-antiquark pair [3]. Similarly the number of gluons changes because a quark may radiate a gluon or because a gluon may split into a quark-antiquark pair or into two gluons or into two tensor-gluons. This last possibility is realized, because, as we have seen, in non-Abelian tensor gauge theories there is a triple vertex GTT (2.7) of a gluon and two tensor-gluons of order g [16, 17, 18, 19]. This interaction should be taken into consideration, and we added the term T(y,t) $P_{GT}(\frac{x}{y})$ in the second equation (3.21). The density of tensor-gluons T(x,t) changes when a gluon splits into two tensor-gluons or when a tensor-gluon radiates a gluon. This evolution is described by the last equation (3.21).

In order to guarantee that the total momentum of the proton, that is, of all partons is unchanged, one should impose the following constraint:

$$\frac{d}{dt} \int_0^1 dz z \left[\sum_{i=1}^{2n_f} q^i(z,t) + G(z,t) + T(z,t) \right] = 0.$$
 (3.23)

Using the evolution equations (3.21) one can express the derivatives of the densities in (3.23) in terms of kernels and to see that the following momentum sum rules should be fulfilled:

$$\int_{0}^{1} dz z [P_{qq}(z) + P_{Gq}(z)] = 0,$$

$$\int_{0}^{1} dz z [2n_{f} P_{qG}(z) + P_{GG}(z) + P_{TG}(z)] = 0,$$

$$\int_{0}^{1} dz z [P_{GT}(z) + P_{TT}(z)] = 0.$$
(3.24)

Before analyzing these momentum sum rules let us first inspect the behavior of the tensorgluon kernels (2.16) at the end points z = 0, 1. As one can see, they are singular at the boundary values similarly to the case of the standard kernels (2.18). Though there is a difference here: the singularities are of higher order compared to the standard case [3]. Therefore one should define the regularization procedure for the singular factors $(1-z)^{-2s+1}$ and z^{-2s+1} reinterpreting them as the distributions $(1-z)^{-2s+1}_+$ and z^{-2s+1}_+ , similar to the Altarelli-Parisi regularization [3]. We shall define them in the following way:

$$\int_{0}^{1} dz \frac{f(z)}{(1-z)_{+}^{2s-1}} = \int_{0}^{1} dz \frac{f(z) - \sum_{k=0}^{2s-2} \frac{(-1)^{k}}{k!} f^{(k)}(1)(1-z)^{k}}{(1-z)^{2s-1}},$$

$$\int_{0}^{1} dz \frac{f(z)}{z_{+}^{2s-1}} = \int_{0}^{1} dz \frac{f(z) - \sum_{k=0}^{2s-2} \frac{1}{k!} f^{(k)}(0) z^{k}}{z^{2s-1}},$$

$$\int_{0}^{1} dz \frac{f(z)}{z_{+}^{2s-1}} = \int_{0}^{1} dz \frac{f(z) - (1-z)f(0) - zf(1)}{z(1-z)},$$
(3.25)

where f(z) is any test function which is sufficiently regular at the end points and, as one can see, the defined substraction guarantees the convergence of the integrals. Using the

same arguments as in the standard case [3] we should add the delta function terms into the definition of the diagonal kernels so that they will completely determine the behavior of $P_{qq}(z)$, $P_{GG}(z)$ and $P_{TT}(z)$ functions. The first equation in the momentum sum rule (3.24) remains unchanged because there is no tensor-gluon contribution into the quark evolution. The second equation in the momentum sum rule (3.24) will take the following form (see Appendix for details):

$$\int_{0}^{1} dz z [2n_{f} P_{qG}(z) + P_{GG}(z) + P_{TG}(z) + b_{G}\delta(z - 1)] =$$

$$= \int_{0}^{1} dz z [2n_{f} T(R)[z^{2} + (1 - z)^{2}] + C_{2}(G) \left[\frac{1}{z(1 - z)} + \frac{z^{4}}{z(1 - z)} + \frac{(1 - z)^{4}}{z(1 - z)} \right] +$$

$$+ C_{2}(G) \left[\frac{z^{4}}{z(1 - z)} \left(\frac{z}{1 - z} \right)^{2s - 2} + \frac{(1 - z)^{4}}{z(1 - z)} \left(\frac{1 - z}{z} \right)^{2s - 2} \right] \right] + b_{G} =$$

$$= \frac{2}{3} n_{f} T(R) - \frac{11}{6} C_{2}(G) - \frac{12s^{2} - 1}{6} C_{2}(G) + b_{G} = 0.$$
(3.26)

From this result we can extract an additional contribution to the one-loop Callan-Symanzik beta function for gluons b_{GG} arising from the tensor-gluon loop. Indeed, the first beta-function coefficient enters into this expression because the momentum sum rule (3.24) implicitly comprises unitarity, thus the one-loop effects [3]. In (3.26) we have three terms which come from quark loops:

$$b_{GG}^q = -\frac{2n_f}{3}T(R), (3.27)$$

from the gluon loop:

$$b_{GG}^G = \frac{11}{6}C_2(G) \tag{3.28}$$

and from the gauge boson loop of spin s:

$$b_{GG}^{T} = \frac{12s^2 - 1}{6}C_2(G), \quad s = 1, 2, 3, 4, \dots$$
 (3.29)

It is a very interesting result because at s=1 we are rediscovering the asymptotic freedom result [1, 2]. For larger spins the tensor-gluon contribution into the Callan-Simanzik beta function has the same signature as the standard gluons, which means that tensor-gluons "accelerate" the asymptotic freedom (3.22) of the strong interaction coupling constant $\alpha(t)$. The contribution is increasing quadratically with spin of the tensor-gluons, that is, at large transfer momentum the strong coupling constant tends to zero faster compared to the standard case:

$$\alpha(t) = \frac{\alpha}{1 + b\alpha t} \,, \tag{3.30}$$

where

$$b = \frac{\sum_{s} (12s^2 - 1)C_2(G) - 4n_f T(R)}{12\pi}, \quad s = 1, 2, \dots$$
 (3.31)

In particular, the presence of the spin-two tensor-gluons in the proton will give

$$b = \frac{58C_2(G) - 4n_f T(R)}{12\pi}. (3.32)$$

Surprisingly, a similar result based on the parametrization of the charge renormalization taken in the form $b = (-1)^{2s}(A+Bs^2)$ was conjectured by Curtright [40]. Here A represents an orbital contribution and Bs^2 - the anomalous magnetic moment contribution [41, 42, 43]. The unknown coefficients A and B were found by comparing the suggested parametrization with the known results for s = 0, 1/2 and 1.

It is also possible to consider a straitforward generalization of the result obtained for the effective action in Yang-Mills theory long ago [41, 42, 43, 44] to the higher spin gauge bosons. With the spectrum of the tensor-gluons in the external chromomagnetic field $\lambda = (2n + 1 + 2s)gH + k_{\parallel}^2$ one can perform a summation of the modes and get an exact result for the one-loop effective action similar to [41, 44]:

$$\epsilon = \frac{H^2}{2} + \frac{(gH)^2}{4\pi} b \left[\ln \frac{gH}{u^2} - \frac{1}{2} \right], \tag{3.33}$$

where

$$b = -\frac{2C_2(G)}{\pi} \zeta(-1, \frac{2s+1}{2}) = \frac{12s^2 - 1}{12\pi} C_2(G), \tag{3.34}$$

and $\zeta(-1,q) = -\frac{1}{2}(q^2 - q + \frac{1}{6})$ is the generalized zeta function*. Because the coefficient in front of the logarithm defines the beta function [41, 42], one can see that (3.34) is in agreement with the result (3.29). It is also interesting to mention that the spectrum of the open strings in the background magnetic field has a similar spectrum with the gyromagnetic ration equal to two, as it was pointed out by Bachas and Fabre in [45].

The third equation in the momentum sum rule (3.24) will take the following form:

$$\int_{0}^{1} dz z [P_{TT}(z) + P_{GT}(z) + b_{T} \delta(z - 1)] =$$

$$= C_{2}(G) \int_{0}^{1} dz z \left[\frac{z^{2s+1}}{1-z} + \frac{1}{(1-z)z^{2s-1}} + \frac{1}{z(1-z)^{2s-1}} + \frac{(1-z)^{2s+1}}{z} \right] + b_{T} \Rightarrow$$

$$\Rightarrow Reg = C_{2}(G) \int_{0}^{1} dz \left[\frac{z^{2s+2} - 1}{1-z} + (1-z)^{2s+1} \right] + b_{T} =$$

^{*}The generalized zeta function is defined as $\zeta(p,q) = \sum_{k=0}^{\infty} \frac{1}{(k+q)^p} = \frac{1}{\Gamma(p)} \int_0^{\infty} dt t^{-1+p} \frac{e^{-qt}}{1-e^{-t}}$.

$$= -C_2(G) \sum_{j=1}^{2s+1} \frac{1}{j} + b_T = 0.$$
 (3.35)

And again we can extract the one-loop coefficient of the Callan-Symanzik beta function now for tensor-gluon of spin s, which has the form

$$b_{TT} = C_2(G) \sum_{j=1}^{2s+1} \frac{1}{j}, \quad s = 1, 2, 3, 4, \dots$$
 (3.36)

As one can see, at s=1 we have

$$C_2(G) (1 + \frac{1}{2} + \frac{1}{3}) = \frac{11}{6}C_2(G),$$

and it coincides with the one loop contribution of the gluons (3.28). The coefficient grows as $\ln s$. This growth is slower than s^2 in (3.31). The reason is that, as we shall discuss at the end of this section, the splitting of tensor-gluons into pair of tensor-gluons was discarded in the derivation of the evolution equation. In this article we limit ourselves by considering only emissions which always involve the standard gluons and lower-spin tensors, ignoring infinite "stair" of transitions between tensor-gluons.

In summary, we have to add $\delta(z-1)$ to the diagonal kernels $P_{qq}(z)$, $P_{GG}(z)$ and $P_{TT}(z)$ with the coefficients which have been determined by using the momentum sum rule (3.24) guaranteeing the total momentum conservation of a hadron:

$$P_{qq}(z) = C_2(R) \left[\frac{1+z^2}{(1-z)_+} + \frac{3}{2} \delta(z-1) \right],$$

$$P_{GG}(z) = 2C_2(G) \left[\frac{z}{(1-z)_+} + \frac{1-z}{z} + z(1-z) \right] + \frac{\sum_s (12s^2 - 1)C_2(G) - 4n_f T(R)}{6} \delta(z-1),$$

$$P_{TT}(z) = C_2(G) \left[\frac{z^{2s+1}}{(1-z)_+} + \frac{1}{(1-z)z_+^{2s-1}} + \sum_{j=1}^{2s+1} \frac{1}{j} \delta(z-1) \right].$$
(3.37)

Thus we completed the definition of the kernels appearing in the evolution equations (3.21).

At the end of this section we shall discuss, what type of processes could also be included into the evolution equations (3.21). In (3.21) we ignore contribution of the high-spin fermions \tilde{q}^i of spin s+1/2, which are the partners of the standard quarks [16, 17, 18, 19], supposing that they are even heavier than the top quark. That is, all kernels $P_{q\tilde{q}}$, $P_{G\tilde{q}}$, $P_{T\tilde{q}}$, $P_{\tilde{q}q}$,

(3.21) one could also include the kernels $P_{TT'}$ which describe the emission of tensor-gluons by tensor-gluons, the TT'T'' vertex (2.5) [16, 17, 18, 19]. That also can be done and the number of evolution equations in that case will tend to infinity. In this article we shall limit ourselves considering only emissions which always involve the standard gluons and lower spin tensors, ignoring infinite "stair" of transitions between tensor-gluons. We shall consider a more general case in a separate work.

It is also natural to ask what will happen if one takes into consideration the contribution of tensor-gluons of all spins into the beta function[†]. One can suggest two scenarios. In the first one the high spin gluons, let us say of $s \geq 3$, will get large mass and therefore can be ignored at a given energy scale. In the second case, when all of them remain massless, then one can suggest the Reimann zeta function regularization, similar to the Brink-Nielsen regularization [46], which gives:

$$b = C_2(G) \sum_{s=1}^{\infty} \frac{(12s^2 - 1)}{12\pi} = C_2(G) \left[\frac{1}{\pi} \zeta(-2) - \frac{1}{12\pi} \zeta(0) \right] = \frac{1}{24\pi} C_2(G), \tag{3.38}$$

where $\zeta(-2) = 0$, $\zeta(0) = -1/2$ and the theory remains asymptotically free with a smaller beta coefficient, but this summation requires further justification.

4 Moments of the Scaling Structure Functions

The moments of the scaling structure functions measure the Fourier transform of the coefficients $C^{(n)}(Q^2, g)$ of the Wilson's operator-product expansion of currents [15]:

$$\int_{0}^{1} dx x^{n-1} F_{i}(x, Q^{2}) \qquad \sim \qquad C_{i}^{(n)}(t) < n | O^{(n)} | n > .$$

$$Q^{2} \to \infty$$

Their logarithmic deviations from Bjorken scaling are obtainable from calculation of anomalous dimensions γ_n of the corresponding operators and beta function coefficient b in (3.31):

$$C_i^{(n)}(t) = C_i^{(n)}(0) \left[\frac{\alpha}{\alpha(t)}\right]^{\gamma_n/2\pi b}.$$

[†]I would like to thank John Iliopoulos and Constantin Bachas for rasing this question.

Therefore one should calculate the matrix of relevant anomalous dimensions of the twisttwo operators. They are represented in the terms of moments of the kernels in (3.21):

$$\gamma_{n} = \int_{0}^{1} dz z^{n-1} \begin{pmatrix} P_{qq}(z) & 2n_{f} P_{qG}(z) & 0 \\ P_{Gq}(z) & P_{GG}(z) & P_{GT}(z) \\ 0 & P_{TG}(z) & P_{TT}(z) \end{pmatrix} = \begin{pmatrix} \gamma_{n}^{qq} & 2n_{f} \gamma_{n}^{qG} & 0 \\ \gamma_{n}^{Gq} & \gamma_{n}^{GG} & \gamma_{n}^{GT} \\ 0 & \gamma_{n}^{TG} & \gamma_{n}^{TT} \end{pmatrix}, \quad (4.39)$$

where the corresponding integrals for quarks and gluons are well known [1]:

$$\int_{0}^{1} dz z^{n-1} P_{qq}(z) = C_{2}(R) \left[-\frac{1}{2} + \frac{1}{n(n+1)} - 2 \sum_{j=2}^{n} \frac{1}{j} \right],$$

$$\int_{0}^{1} dz z^{n-1} P_{Gq}(z) = C_{2}(R) \left[\frac{2+n+n^{2}}{n(n^{2}-1)} \right],$$

$$2n_{f} \int_{0}^{1} dz z^{n-1} P_{qG}(z) = 2n_{f} T(R) \left[\frac{2+n+n^{2}}{n(n+1)(n+2)} \right],$$

$$\int_{0}^{1} dz z^{n-1} P_{GG}(z) = C_{2}(G) \left[-\frac{1}{6} + \frac{2}{n(n-1)} + \frac{2}{(n+1)(n+2)} - \frac{2}{n(n+1)(n+2)} - \frac{2}{n(n+1)(n+2)} \right].$$

$$-2 \sum_{j=2}^{n} \frac{1}{j} - \frac{2n_{f} T(R)}{3C_{2}(G)} + \frac{\sum_{j=2}^{n} (12s^{2} - 1)}{6} \right].$$
(4.40)

We have to calculate the new terms in the matrix of anomalous dimensions which are the contribution of tensor-gluons. For tensors-gluons kernel $P_{TG}(z)$ they are:

$$\int_{0}^{1} dz z^{n-1} P_{TG}(z) = C_{2}(G) \left[\sum_{k=2s-1}^{2s+n} \frac{(-1)^{k} (2s+n)!}{k! (2s+n-k)!} \frac{1}{k-2s+2} \right] +$$

$$C_{2}(G) \left[\sum_{k=2s-n}^{2s+1} \frac{(-1)^{k} (2s+1)!}{k! (2s+1-k)!} \frac{1}{k+1+n-2s} \right], \quad n \leq 2s-1$$

$$\int_{0}^{1} dz z^{n-1} P_{TG}(z) = C_{2}(G) \left[\sum_{k=2s-1}^{2s+n} \frac{(-1)^{k} (2s+n)!}{k! (2s+n-k)!} \frac{1}{k-2s+2} \right] +$$

$$C_{2}(G) \left[\frac{\Gamma(2s+2)\Gamma(n-2s+1)}{\Gamma(n+3)} \right], \qquad n \geq 2s,$$

for the gluons-tensors kernel $P_{GT}(z)$:

$$\int_{0}^{1} dz z^{n-1} P_{GT}(z) = C_{2}(G) \left[\frac{\Gamma(2s+2)\Gamma(n-1)}{\Gamma(n+2s+1)} \right], \quad n \leq 2s$$

$$\int_{0}^{1} dz z^{n-1} P_{GT}(z) = C_{2}(G) \left[\sum_{k=2s-1}^{n-2} \frac{(-1)^{k} (n-2)!}{k! (n-k-2)!} \frac{1}{k-2s+2} + \frac{\Gamma(2s+2)\Gamma(n-1)}{\Gamma(n+2s+1)} \right], \quad n \geq 2s+1$$

$$(4.42)$$

and for the tensors-tensors kernel $P_{TT}(z)$:

$$\int_{0}^{1} dz z^{n-1} P_{TT}(z) = -C_{2}(G) \sum_{j=2+2s}^{n+2s} \frac{1}{j}, \qquad n \le 2s,$$

$$\int_{0}^{1} dz z^{n-1} P_{TT}(z) = -C_{2}(G) \left[\sum_{j=2+2s}^{n+2s} \frac{1}{j} + \sum_{j=1}^{n-2s} \frac{1}{j} \right], \qquad n \ge 2s+1 \qquad (4.43)$$

where s = 2, 3, 4, ... For the nonsinglet piece of the structure functions the relevant anomalous dimension is γ_n^{qq} , and it does not have any additional contribution from tensor-gluons, therefore for nonsinglet pieces of the structure functions one can get

$$\int_0^1 dx x^{n-2} F^{NS}(x, Q^2) \qquad \sim \qquad C_{NS}^{(n)} \left[\frac{\alpha}{\alpha(t)} \right]^{A_n^{NS}}, \tag{4.44}$$

$$Q^2 \to \infty$$

where

$$A_n^{NS} = \frac{\gamma_n^{qq}}{2\pi b} = -\frac{3C_2(R)}{\sum_s (12s^2 - 1)C_2(G) - 4n_f T(R)} \left[1 - \frac{2}{n(n+1)} + 4\sum_{j=2}^n \frac{1}{j} \right]. \tag{4.45}$$

The difference between it and the standard case comes from the value of the coefficient b in the Callan-Simanzik beta function (3.31). This result shows that if tensor-gluons exist in the strong interaction, the logarithmic correction to the Bjorken scaling becomes more mild. For the $SU(3)_c$ ($C_2(G) = 3$, $C_2(R) = 4/3$, T(R) = 1/2, $n_f = 3$) and spin-two gluons s = 2 we are getting

$$A_2^{NS} = -\frac{2}{63}, \quad A_3^{NS} = -\frac{25}{504}, \quad \dots \quad A_n^{NS} = -\frac{1}{21} \ln n$$
 (4.46)

instead of $A_2^{NS} = -\frac{16}{81}$, $A_3^{NS} = -\frac{25}{81}$, ..., $A_n^{NS} = -\frac{8}{27} \ln n$ in the standard case [15].

For the singlet pieces of the structure functions one should take into account the mixing of gluon, fermion and tensor operators, so that

$$\int_{0}^{1} dx x^{n-2} F^{S}(x, Q^{2}) \qquad \sim \qquad C_{S}^{(n)} \left[\frac{\alpha}{\alpha(t)} \right]^{A_{n}^{S}}, \tag{4.47}$$

$$Q^{2} \to \infty$$

where $A_n^S = \gamma_n/2\pi b$ are the eigenvalues of the matrix (4.39). The matrix of anomalous dimensions for arbitrary n is given by (4.40),(4.41), (4.42) and (4.43).

5 Unification of Coupling Constants of Standard Model

It is interesting to know how the contribution of tensor-gluons changes the high energy behavior of the coupling constants of the Standard Model [47, 48]. The coupling constants are evolving in accordance with the formulae

$$\frac{1}{\alpha_i(M)} = \frac{1}{\alpha_i(\mu)} + 2b_i \ln \frac{M}{\mu}, \quad i = 1, 2, 3,$$
 (5.48)

where we shall consider only the contribution of the lower s=2 tensor-bosons:

$$2b = \frac{58C_2(G) - 4n_f T(R)}{6\pi}. (5.49)$$

For the $SU(3)_c \times SU(2)_L \times U(1)$ group with its coupling constants α_3, α_2 and α_1 and six quarks $n_f = 6$ and SU(5) unification group we will get

$$2b_3 = \frac{1}{2\pi}54$$
, $2b_2 = \frac{1}{2\pi}\frac{104}{3}$, $2b_1 = -\frac{1}{2\pi}4$,

so that solving the system of equations (5.48) one can get

$$\ln \frac{M}{\mu} = \frac{\pi}{58} \left(\frac{1}{\alpha_{el}(\mu)} - \frac{8}{3} \frac{1}{\alpha_s(\mu)} \right), \tag{5.50}$$

where $\alpha_{el}(\mu)$ and $\alpha_s(\mu)$ are the electromagnetic and strong coupling constants at scale μ . If one takes $\alpha_{el}(M_Z) = 1/128$ and $\alpha_s(M_Z) = 1/10$ one can get that coupling constants have equal strength at energies of order

$$M \sim 4 \times 10^4 GeV = 40 \ TeV$$

which is much smaller than the scale $M \sim 10^{14} GeV$ in the absence of the tensor-gluons contribution. The value of the weak angle [47, 48] remains intact :

$$\sin^2 \theta_W = \frac{1}{6} + \frac{5}{9} \frac{\alpha_{el}(M_Z)}{\alpha_s(M_Z)},\tag{5.51}$$

as well as the coupling constant at the unification scale remains of the same order $\bar{\alpha}(M) = 0.01$.

6 Conclusion

Let us summarize what are the physical consequences of the tensor-gluons hypothesis. Among all parton distributions, the gluon density G(x,t) is one of the least constrained functions since it does not couple directly to the photon in deep-inelastic scattering measurements of the proton F_2 structure function. Therefore it is only indirectly constrained by scaling violations and by the momentum sum rule which resulted in the fact that only half of the proton momentum is carried by charged constituents - the quarks - and that the other part is ascribed to the neutral constituents. As it was suggested in the article the process of gluon splitting leads to the emission of tensor-gluons and therefore a part of the proton momentum which is carried by the neutral constituents here is shared between gluons and tensor-gluons. The density of neutral partons in the proton is therefore given by the sum of two functions: G(x,t) + T(x,t), where T(x,t) is the density of the tensor-gluons. To disentangle these contributions and to decide which piece of the neutral partons is the contribution of gluons and which one is of the tensor-gluons one should measure the helicities of the neutral components, which seems to be a difficult task. The other test of the proposed model will be the consistency of the mild Q^2 behavior of the moments of the structure functions with the experimental data.

The gluon density can be directly constrained also by jet production [49]. In suggested model the situation is such that the standard quarks cannot radiate tensor-gluons (such a vertex is absent in the model [16, 17, 18, 19]), therefore only gluons are radiated by quarks. A radiated gluon then can split into a pair of tensor-gluons without obscuring the structure of the observed three-jet final states. Thus it seems that there is no obvious contradiction with the existing experimental data. Our hypotheses may be wrong, but the uniqueness and simplicity of suggested extension seems to be the reasons for serious consideration.

In the last section we also observed that the unification scale at which standard coupling constants are merging is shifted to lower energies telling us that it may be that the new physics is round the corner. Whether all these phenomena are consistent with experiment is an open question.

This work was supported in part by the General Secretariat for Research and Technology of Greece and from the European Regional Development Fund MIS-448332-ORASY (NSRF 2007-13 ACTION, KRIPIS).

7 Note added

In supersymmetric extensions of the Standard Model [50, 51] the gluons and quarks have natural partners - gluinos of spin s=1/2 and squarks of spin s=0. If the gluinos appear as elementary constituents of the hadrons then the theory predicts the existence of new hadronic states, the R-hadrons [52, 53]. These new hadronic states can be produced in ordinary hadronic collisions and decay into ordinary hadrons with the radiation of massless photino - the massless partner of the photon, which takes out a conserved R-parity. Depending on the model the gluinos may be massless or massive, depending on the remaining unbroken symmetries and the representation content of the theory. The existing experimental data give evidence that most probably they have to be very heavy [54, 55].

It seems that phenomenological limitation on the existence of the R-hadrons is much stronger that the limitation on the existence of the tensor-gluons inside the ordinary hadrons. This is because gluinos change the statistics of the ordinary hadrons: the proton has to have a partner - the R-proton which is a boson and is indeed a new hadron. The existence of tensor-gluon partons inside the proton does not predict a new hadronic state, a proton remains a proton. The tensor-gluons change proton dynamical properties - its structure functions. What are the experimental limitations on that dynamics should be studied in details.

I would like to thank Pierre Fayet for illuminating discussions of the supersymmetry phenomenology and physics of R-hadrons and John Iliopoulos for helpful discussions of the experimental data.

8 Appendix

The momentum sum rule integrals (3.24), (3.26) and (3.35) can be calculated in two different ways: with direct regularization of the integrals near z = 1 and near z = 0 as it is defined in (3.25), or using the substitution w = 1 - z in the second term of the integrand and then regularizing the resulting integrand near z = 1. As one can convinced both methods give the identical results. Therefore we shell first calculate the integral over $P_{TG}(z)$ in (3.26) by direct regularization (3.25) as follows:

$$C_2(G) \int_0^1 dz \left[\frac{z^{2s+2}}{(1-z)^{2s-1}} + \frac{(1-z)^{2s+1}}{z^{2s-2}} \right] \Rightarrow Reg$$

$$= C_2(G) \int_0^1 dz \left[\frac{z^{2s+2}}{(1-z)_+^{2s-1}} + \frac{(1-z)^{2s+1}}{z_+^{2s-2}} \right] = -(-1)^{2s} \frac{12s^2 - 1}{6} C_2(G).$$
 (8.52)

The same integral we shall calculate now using the substitution w = 1 - z in the second term of the integrand and evaluating the sum

$$C_2(G) \int_0^1 dz \left[\frac{z^{2s+2}}{(1-z)^{2s-1}} + \frac{z^{2s+1}}{(1-z)^{2s-2}} \right] = C_2(G) \int_0^1 dz \frac{z^{2s+1}}{(1-z)^{2s-1}}$$

and then one should regularize the integral near z = 1 as it is defined in (3.25)

$$C_2(G) \int_0^1 dz \frac{z^{2s+1}}{(1-z)_+^{2s-1}} = -(-1)^{2s} \frac{12s^2 - 1}{6} C_2(G), \tag{8.53}$$

which is identical to the previous result.

References

- [1] D. J. Gross and F. Wilczek, *Ultraviolet Behavior Of Non-Abelian Gauge Theories*, Phys. Rev. Lett. **30**, 1343 (1973).
- [2] H. D. Politzer, *Reliable Perturbative Results For Strong Interactions?*, Phys. Rev. Lett. **30**, 1346 (1973).
- [3] G. Altarelli and G. Parisi, Asymptotic Freedom in Parton Language, Nucl. Phys. B 126 (1977) 298.
- [4] Y. L. Dokshitzer, Calculation of the Structure Functions for Deep Inelastic Scattering and e+ e- Annihilation by Perturbation Theory in Quantum Chromodynamics., Sov. Phys. JETP 46 (1977) 641 [Zh. Eksp. Teor. Fiz. 73 (1977) 1216].
- [5] V. N. Gribov and L. N. Lipatov, Deep inelastic e p scattering in perturbation theory, Sov. J. Nucl. Phys. **15** (1972) 438 [Yad. Fiz. **15** (1972) 781].
- [6] V. N. Gribov and L. N. Lipatov, e+ e- pair annihilation and deep inelastic e p scattering in perturbation theory, Sov. J. Nucl. Phys. 15 (1972) 675 [Yad. Fiz. 15 (1972) 1218].
- [7] L. N. Lipatov, The parton model and perturbation theory, Sov. J. Nucl. Phys. 20 (1975)
 94 [Yad. Fiz. 20 (1974) 181].

- [8] V. S. Fadin, E. A. Kuraev and L. N. Lipatov, On the Pomeranchuk Singularity in Asymptotically Free Theories, Phys. Lett. B **60** (1975) 50.
- [9] E. A. Kuraev, L. N. Lipatov and V. S. Fadin, The Pomeranchuk Singularity in Non-abelian Gauge Theories, Sov. Phys. JETP 45 (1977) 199 [Zh. Eksp. Teor. Fiz. 72 (1977) 377].
- [10] I. I. Balitsky and L. N. Lipatov, The Pomeranchuk Singularity in Quantum Chromodynamics, Sov. J. Nucl. Phys. 28 (1978) 822 [Yad. Fiz. 28 (1978) 1597].
- [11] N. Cabibbo and R. Petronzio, Two Stage Model Of Hadron Structure: Parton Distributions And Their Q**2 Dependence, Nucl. Phys. B 137 (1978) 395.
- [12] L. V. Gribov, E. M. Levin and M. G. Ryskin, Singlet Structure Function at Small x: Unitarization of Gluon Ladders, Nucl. Phys. B **188** (1981) 555.
- [13] J. D. Bjorken and E. A. Paschos, *Inelastic Electron Proton and gamma Proton Scattering*, and the Structure of the Nucleon, Phys. Rev. **185** (1969) 1975.
- [14] D. J. Gross and F. Wilczek, Asymptotically Free Gauge Theories. 1, Phys. Rev. D 8 (1973) 3633.
- [15] D. J. Gross and F. Wilczek, Asymptotically Free Gauge Theories. 2., Phys. Rev. D 9 (1974) 980.
- [16] G. Savvidy, Non-Abelian tensor gauge fields: Generalization of Yang-Mills theory, Phys. Lett. B 625 (2005) 341
- [17] G. Savvidy, Non-abelian tensor gauge fields. I, Int. J. Mod. Phys. A 21 (2006) 4931;
- [18] G. Savvidy, Non-abelian tensor gauge fields. II, Int. J. Mod. Phys. A 21 (2006) 4959;
- [19] G. Savvidy, Extension of the Poincaré Group and Non-Abelian Tensor Gauge Fields, Int. J. Mod. Phys. A 25 (2010) 5765 [arXiv:1006.3005 [hep-th]].
- [20] G. Georgiou and G. Savvidy, Production of non-Abelian tensor gauge bosons. Tree amplitudes and BCFW recursion relation, Int. J. Mod. Phys. A 26 (2011) 2537 [arXiv:1007.3756 [hep-th]].

- [21] I. Antoniadis and G. Savvidy, Conformal invariance of tensor boson tree amplitudes, Mod. Phys. Lett. A 27 (2012) 1250103 [arXiv:1107.4997 [hep-th]].
- [22] F. A. Berends, R. Kleiss, P. De Causmaecker, R. Gastmans and T. T. Wu, Single Bremsstrahlung Processes In Gauge Theories, Phys. Lett. B 103 (1981) 124.
- [23] R. Kleiss and W. J. Stirling, Spinor Techniques For Calculating P Anti-P \rightarrow W⁺⁻/Z⁰ + Jets, Nucl. Phys. B **262** (1985) 235.
- [24] Z. Xu, D. H. Zhang and L. Chang, Helicity Amplitudes for Multiple Bremsstrahlung in Massless Nonabelian Gauge Theories, Nucl. Phys. B 291 (1987) 392.
- [25] J. F. Gunion and Z. Kunszt, Improved Analytic Techniques For Tree Graph Calculations And The G G Q Anti-Q Lepton Anti-Lepton Subprocess, Phys. Lett. B 161 (1985) 333.
- [26] L. J. Dixon, Calculating scattering amplitudes efficiently, arXiv:hep-ph/9601359.
- [27] S. J. Parke and T. R. Taylor, An Amplitude for n Gluon Scattering, Phys. Rev. Lett. 56 (1986) 2459.
- [28] F. A. Berends and W. T. Giele, Recursive Calculations for Processes with n Gluons, Nucl. Phys. B 306 (1988) 759.
- [29] E. Witten, Perturbative gauge theory as a string theory in twistor space, Commun. Math. Phys. **252** (2004) 189 [arXiv:hep-th/0312171].
- [30] F. Cachazo, P. Svrcek and E. Witten, Gauge theory amplitudes in twistor space and holomorphic anomaly, JHEP **0410** (2004) 077 [hep-th/0409245].
- [31] F. Cachazo, Holomorphic anomaly of unitarity cuts and one-loop gauge theory amplitudes, hep-th/0410077.
- [32] R. Britto, F. Cachazo and B. Feng, New recursion relations for tree amplitudes of gluons, Nucl. Phys. B **715** (2005) 499 [arXiv:hep-th/0412308].
- [33] R. Britto, F. Cachazo, B. Feng and E. Witten, Direct proof of tree-level recursion relation in Yang-Mills theory, Phys. Rev. Lett. **94** (2005) 181602 [arXiv:hep-th/0501052].

- [34] P. Benincasa and F. Cachazo, Consistency Conditions on the S-Matrix of Massless Particles, arXiv:0705.4305 [hep-th].
- [35] F. Cachazo, P. Svrcek and E. Witten, MHV vertices and tree amplitudes in gauge theory, JHEP **0409** (2004) 006 [arXiv:hep-th/0403047].
- [36] G. Georgiou, E. W. N. Glover and V. V. Khoze, Non-MHV Tree Amplitudes in Gauge Theory, JHEP **0407** (2004) 048 [arXiv:hep-th/0407027].
- [37] N. Arkani-Hamed and J. Kaplan, On Tree Amplitudes in Gauge Theory and Gravity, JHEP 0804 (2008) 076 [arXiv:0801.2385 [hep-th]].
- [38] F. A. Berends and W. T. Giele, Multiple Soft Gluon Radiation in Parton Processes, Nucl. Phys. B 313 (1989) 595.
- [39] M. L. Mangano and S. J. Parke, Quark Gluon Amplitudes in the Dual Expansion, Nucl. Phys. B 299 (1988) 673.
- [40] T. L. Curtright, Charge Renormalization And High Spin Fields, Phys. Lett. B 102 (1981) 17.
- [41] G. K. Savvidy, Infrared Instability of the Vacuum State of Gauge Theories and Asymptotic Freedom, Phys. Lett. B **71** (1977) 133.
- [42] S. G. Matinyan and G. K. Savvidy, Vacuum Polarization Induced by the Intense Gauge Field, Nucl. Phys. B 134 (1978) 539.
- [43] I. A. Batalin, S. G. Matinyan and G. K. Savvidy, Vacuum Polarization by a Source-Free Gauge Field, Sov. J. Nucl. Phys. 26 (1977) 214 [Yad. Fiz. 26 (1977) 407].
- [44] D. Kay, Supersymmetric Savvidy State For Su(2), Su(3) And Su(4), Phys. Rev. D 28 (1983) 1562.
- [45] C. Bachas and C. Fabre, Threshold effects in open string theory, Nucl. Phys. B 476 (1996) 418 [hep-th/9605028].
- [46] L. Brink and H. B. Nielsen, A physical interpretation of the jacobi imaginary transformation and the critical dimension in dual models, Phys. Lett. B 43 (1973) 319.

- [47] H. Georgi and S. L. Glashow, Unity of All Elementary Particle Forces, Phys. Rev. Lett. 32 (1974) 438.
- [48] H. Georgi, H. R. Quinn and S. Weinberg, *Hierarchy of Interactions in Unified Gauge Theories*, Phys. Rev. Lett. **33** (1974) 451.
- [49] J. R. Ellis, M. K. Gaillard and G. G. Ross, Search for Gluons in e+ e- Annihilation, Nucl. Phys. B 111 (1976) 253 [Erratum-ibid. B 130 (1977) 516].
- [50] P. Fayet, Supersymmetry and Weak, Electromagnetic and Strong Interactions, Phys. Lett. B 64 (1976) 159.
- [51] P. Fayet, Spontaneously Broken Supersymmetric Theories of Weak, Electromagnetic and Strong Interactions, Phys. Lett. B **69** (1977) 489.
- [52] G. R. Farrar and P. Fayet, Phenomenology of the Production, Decay, and Detection of New Hadronic States Associated with Supersymmetry, Phys. Lett. B **76** (1978) 575.
- [53] P. Fayet, Massive Gluinos, Phys. Lett. B **78** (1978) 417.
- [54] G. R. Farrar and P. Fayet, Bounds on R-Hadron Production from Calorimetry Experiments, Phys. Lett. B 79 (1978) 442.
- [55] I. Antoniadis, C. Kounnas and R. Lacaze, Light Gluinos in Deep Inelastic Scattering, Nucl. Phys. B 211 (1983) 216.