

Probing the BFKL dynamics at hadronic colliders

Christophe Royon

IRFU-SPP, CEA Saclay, F91 191 Gif-sur-Yvette cedex, France

We describe different possibilities to probe the BFKL dynamics at hadronic colliders, namely Mueller-Navelet jet, and jet gap jet events. We also discuss briefly the jet veto measurement as performed by the ATLAS collaboration at the LHC.

1 Forward jets and Mueller Navelet jets

In this section, we recall briefly the previous results that we obtained concerning forward jets at HERA and Mueller Navelet jets at the Tevatron or the LHC as a potential test of BFKL dynamics [1, 2]. Forward jets at HERA are an ideal observable to look for BFKL resummation effects. The interval in rapidity between the scattered lepton and the jet in the forward region is large, and when the photon virtuality Q^2 is close to the transverse jet momentum k_T , the DGLAP cross section is small because of the k_T ordering of the emitted gluons. The BFKL NLL formalism leads to a very good description of the forward jet cross section measurements performed by the H1 Collaboration at HERA [2], especially the triple differential cross section of forward jet production as a function of Q^2 , jet p_T and ξ .

Mueller Navelet jets are another ideal processes to study BFKL resummation effects [3]. Two jets with a large interval in rapidity and with similar transverse momenta are considered. A typical observable to look for BFKL effects is the measurement of the azimuthal correlations between both jets. The DGLAP prediction is that this distribution should peak towards π - ie jets are back-to-back- whereas multi-gluon emission via the BFKL mechanism leads to a smoother distribution. The azimuthal correlation is an ideal variable to look for BFKL resummation effects since it is less sensitive to experimental uncertainties such as the jet energy scale as an example [3]. The effect of the energy conservation in the BFKL equation [3] is large when R goes away from 1. The effect is to reduce the effective value of $\Delta\eta$ between the jets and thus the decorrelation effect. However, it is worth noticing that this effect is negligible when the ratio of the jet p_T s is close to 1. It is thus important to perform this measurement as a function of the ratio of the jet p_T .

2 Jet veto measurements in ATLAS

The ATLAS collaboration measured the so-called jet veto cross section [4], namely the events with two high p_T jets, well separated in rapidity and with a veto on jet activity with p_T greater than a given threshold Q_0 between the two jets. The ATLAS collaboration measured the jet veto fraction with respect to the standard dijet cross section, and it was advocated that it might be sensitive to BFKL dynamics. In Ref. [5], we computed the gluon emission at large

angles (which are not considered in usual MC) using the Banfi Marchesini Smye equation, and we showed that the measurement can be effectively described by the gluon resummation and is thus not related to BFKL dynamics. The sensitivity to the BFKL resummation effects appears when one looks for gaps between jets as described in the following section.

3 Jet gap jets at the Tevatron and the LHC

In this section, we describe a new possible measurement which can probe BFKL resummation effects and we compare our predictions with existing D0 and CDF measurements [6].

3.1 BFKL NLL formalism

The production cross section of two jets with a gap in rapidity between them reads

$$\frac{d\sigma^{pp \rightarrow XJJY}}{dx_1 dx_2 dE_T^2} = \mathcal{S} f_{eff}(x_1, E_T^2) f_{eff}(x_2, E_T^2) \frac{d\sigma^{gg \rightarrow gg}}{dE_T^2}, \quad (1)$$

where \sqrt{s} is the total energy of the collision, E_T the transverse momentum of the two jets, x_1 and x_2 their longitudinal fraction of momentum with respect to the incident hadrons, \mathcal{S} the survival probability, and f the effective parton density functions [6]. The rapidity gap between the two jets is $\Delta\eta = \ln(x_1 x_2 s / p_T^2)$.

The cross section is given by

$$\frac{d\sigma^{gg \rightarrow gg}}{dE_T^2} = \frac{1}{16\pi} |A(\Delta\eta, E_T^2)|^2 \quad (2)$$

in terms of the $gg \rightarrow gg$ scattering amplitude $A(\Delta\eta, p_T^2)$.

In the following, we consider the high energy limit in which the rapidity gap $\Delta\eta$ is assumed to be very large. The BFKL framework allows to compute the $gg \rightarrow gg$ amplitude in this regime, and the result is known up to NLL accuracy

$$A(\Delta\eta, E_T^2) = \frac{16N_c\pi\alpha_s^2}{C_F E_T^2} \sum_{p=-\infty}^{\infty} \int \frac{d\gamma}{2i\pi} \frac{[p^2 - (\gamma - 1/2)^2] \exp\{\bar{\alpha}(E_T^2)\chi_{eff}[2p, \gamma, \bar{\alpha}(E_T^2)]\Delta\eta\}}{[(\gamma - 1/2)^2 - (p - 1/2)^2][(\gamma - 1/2)^2 - (p + 1/2)^2]} \quad (3)$$

with the complex integral running along the imaginary axis from $1/2 - i\infty$ to $1/2 + i\infty$, and with only even conformal spins contributing to the sum, and $\bar{\alpha} = \alpha_s N_C / \pi$ the running coupling.

In this study, we performed a parametrised distribution of $d\sigma^{gg \rightarrow gg} / dE_T^2$ so that it can be easily implemented in the Herwig Monte Carlo [7] since performing the integral over γ in particular would be too much time consuming in a Monte Carlo. The implementation of the BFKL cross section in a Monte Carlo is absolutely necessary to make a direct comparison with data. Namely, the measurements are sensitive to the jet size (for instance, experimentally the gap size is different from the rapidity interval between the jets which is not the case by definition in the analytic calculation).




Figure 1: Left: Comparisons between the D0 measurements of the jet-gap-jet event ratio with the NLL- and LL-BFKL calculations. The NLL calculation is in fair agreement with the data. The LL calculation leads to a worse description of the data. Right: Ratio of the jet gap jet to the inclusive jet cross sections at the LHC as a function of jet p_T in double pomeron exchange events where the protons are detected in AFP.

3.2 Comparison with D0 and CDF measurements and predictions for LHC

Let us first notice that the sum over all conformal spins is absolutely necessary. Considering only $p = 0$ in the sum of Equation 3 leads to a wrong normalisation and a wrong jet E_T dependence, and the effect is more pronounced as $\Delta\eta$ diminishes.

The D0 collaboration measured the jet gap jet cross section ratio with respect to the total dijet cross section, requesting for a gap between -1 and 1 in rapidity, as a function of the second leading jet E_T , and $\Delta\eta$ between the two leading jets for two different low and high E_T samples ($15 < E_T < 20$ GeV and $E_T > 30$ GeV). To compare with theory, we compute the following quantity

$$\text{Ratio} = \frac{\text{BFKL NLL HERWIG}}{\text{Dijet Herwig}} \times \frac{\text{LO QCD}}{\text{NLO QCD}} \quad (4)$$

in order to take into account the NLO corrections on the dijet cross sections, where *BFKL NLL HERWIG* and *Dijet Herwig* denote the BFKL NLL and the dijet cross section implemented in HERWIG. The NLO QCD cross section was computed using the NLOJet++ program [8].

The comparison with D0 data [6] is shown in Fig. 1. We find a good agreement between the data and the BFKL calculation. It is worth noticing that the BFKL NLL calculation leads to a better result than the BFKL LL one. The comparison with the CDF data [6] leads to similar conclusions.

Using the same formalism, and assuming a survival probability of 0.03 at the LHC, it is possible to predict the jet gap jet cross section at the LHC. While both LL and NLL BFKL formalisms lead to a weak jet E_T or $\Delta\eta$ dependence, the normalisation is found to be quite

different leading to lower cross section for the BFKL NLL formalism. The ratio of the jet gap jet to the inclusive jet cross sections at the LHC as a function of jet p_T and $\Delta\eta$ is quite flat as shown in Ref. [6].

4 Jet gap jet event in diffractive processes

A new process of detecting jet gap jet events in diffractive double pomeron exchange processes was introduced recently [9]. The idea is to tag the intact protons inside the ATLAS Forward Physics (AFP) detectors [10] located at about 210 m from the ATLAS interaction point on both sides. The advantage of such processes is that they are quite clean since they are not “polluted” by proton remnants and it is possible to go to larger jet separation than for usual jet gap jet events. The normalisation for these processes come from the fit to the D0 discussed in the previous section. The ratio between jet gap jet to inclusive jet events is shown in Fig. 3 requesting protons to be tagged in AFP for both samples. The ratio shows a weak dependence as a function of jet p_T (and also as a function of the difference in rapidity between the two jets). It is worth noticing that the ratio is about 20-30% showing that the jet gap jet events are much more present in the diffractive sample than in the inclusive one as expected.

References

- [1] V.S. Fadin and L.N. Lipatov, Phys. Lett. **B429** (1998) 127; M. Ciafaloni, Phys. Lett. **B429** (1998) 363; M. Ciafaloni and G. Camici, Phys. Lett. **B430** (1998) 349.
- [2] O. Kepka, C. Marquet, R. Peschanski and C. Royon, Phys. Lett. **B655** (2007) 236; Eur. Phys. J. **C55** (2008) 259; C. Marquet and C. Royon, Phys. Rev. D **79** (2009) 034028; A. Sabio Vera and F. Schwennsen, Nucl. Phys. **B776** (2007) 170; Phys. Rev.D **77** (2008) 014001; A. Aktas *et al* [H1 Collaboration], *Eur. Phys. J. C* **46** (2006) 27; H. Navelet, R. Peschanski, C. Royon, S. Wallon, Phys.Lett. **B385** (1996) 357-364; H. Navelet, R. Peschanski, C. Royon, Phys. Lett. **B366** (1996) 329.
- [3] A.H. Mueller and H. Navelet, *Nucl. Phys.* **B282** (1987) 727; C. Marquet, C. Royon, Phys. Rev. D **79** (2009) 034028; B. Ducloué, L. Szymanowski, S. Wallon, JHEP 1305 (2013) 096.
- [4] G. Aad et al, JHEP **1109**, 053 (2011).
- [5] Y. Hatta, C. Marquet, C. Royon, G. Soyez, T. Ueda, D. Werder, Phys. Rev. D **87** (2013) 054016.
- [6] O. Kepka, C. Marquet, C. Royon, Phys. Rev. D **83** (2011) 034036; F. Chevallier, O. Kepka, C. Marquet, C. Royon, Phys. Rev. D **79** (2009) 094019; B. Abbott *et al.*, Phys. Lett. B **440** (1998) 189; F. Abe *et al.*, Phys. Rev. Lett. **80** (1998) 1156.
- [7] G. Marchesini *et al.*, Comp. Phys. Comm. **67** (1992) 465.
- [8] Z. Nagy and Z. Trocsanyi, Phys. Rev. Lett. **87** (2001) 082001.
- [9] C. Marquet, C. Royon, M. Trzebinski, R. Zlebcik, Phys.Rev. D **87** (2013) 3, 034010.
- [10] ATLAS Coll., CERN-LHCC-2011-012.