

A Note on Seiberg Duality and Chiral Symmetry Breaking

Adi Armoni

a.armoni@swan.ac.uk

*Department of Physics, Swansea University
Singleton Park, Swansea, SA2 8PP, UK*

Abstract

Following arXiv:1310.2027 and arXiv:0801.0762, we consider a non-supersymmetric Seiberg duality between electric and magnetic “orientifold field theories”. These theories live on brane configurations of type 0’ string theory. In the electric theory side the scalars acquire a mass and decouple, resulting in an $SU(N_c)$ gauge theory coupled to N_f massless quarks and an additional massless fermion that transforms in the two-index antisymmetric representation. In the magnetic theory side there exists a fundamental meson field that develops a Coleman-Weinberg potential. At the one-loop approximation the potential admits a minimum with chiral symmetry breaking of the form $SU(N_f)_L \times SU(N_f)_R \rightarrow SU(N_f)_V$ and an additional breaking of an axial $U(1)$ symmetry. The resulting theory admits a spectrum whose massless degrees of freedom are N_f^2 Nambu-Goldstone bosons.

1 Introduction

It is widely believed that QCD admits confinement and chiral symmetry breaking due to the formation of the chiral quark condensate $\langle \bar{q}^i q_j \rangle = \delta_j^i \Lambda^3$. Starting with $SU(N_c)$ gauge theory coupled to N_f massless flavors, the IR degrees of freedom of the theory are expected to be the Nambu-Goldstone bosons that correspond to chiral symmetry breaking of the form

$$SU(N_f)_L \times SU(N_f)_R \rightarrow SU(N_f)_V, \quad (1)$$

namely $N_f^2 - 1$ massless mesons. However, even after four decades of intensive studies of QCD, the mechanism behind chiral symmetry breaking is not fully understood.

A promising approach, which led to insights about the IR degrees of freedom of strongly coupled gauge theories, is Seiberg duality [1]. Unfortunately the vast majority of the literature is restricted to super QCD.

In this note we would like to use Seiberg duality between a pair of two non-supersymmetric gauge theories in order to obtain new insights about the IR degrees of freedom of real QCD.

In ref.[2] a Seiberg duality between a pair of two non-supersymmetric gauge theories was proposed. The results of [2] were mainly about the planar limit, where the theory acquires supersymmetry in a well defined sector [3]. The purpose of this note is to focus on the dynamics of the theory at finite N_c , with N_f such that the theory is not conformal in the IR. Our derivations are closely related to those presented recently in [4, 5], see also [6].

The main outcome of the present non-supersymmetric duality is that a QCD-like theory has an IR description in terms of mesons. In fact, contrary to the supersymmetric case, the only massless degrees of freedom are the Nambu-Goldstone mesons that correspond to chiral symmetry breaking of the form (1), plus an additional Nambu-Goldstone boson that corresponds to the breaking of an axial $U(1)$ symmetry.

The paper is organized as follows: in section 2 we present the dual electric and magnetic theories at the classical level. In section 3 we discuss how quantum effects modify the classical picture. Finally in section 4 we summarize our results.

2 Electric and Magnetic theories

We consider a pair of electric and magnetic theories that live on brane configurations of non-critical type 0' string theory [7, 8]. The matter content of the electric-magnetic pair and the duality between them was proposed in [2].

Both the electric and magnetic theories admit an anomaly free $SU(N_f)_L \times SU(N_f)_R \times U(1)_R$ global symmetry. R-symmetry is a name for an anomaly free axial symmetry, borrowed from the supersymmetric jargon. The matter content of both theories is listed in table (1) and table (2) below.

The electric theory is similar to QCD. In addition to the quarks of the theory, there exists a fermion that transforms in the antisymmetric representation of $U(N_c)$ (a “gluino”). In addition there are scalar quarks (“squarks”). As we shall see, these scalar fields will acquire a mass due to quantum corrections and hence will decouple from the low energy theory.

The magnetic theory is a $U(\tilde{N}_c)$ gauge theory, with $\tilde{N}_c = N_f - N_c + 4$. Apart from fields that are charged under the gauge group, it also contains a meson field that transforms in the bifundamental of $SU(N_f)_L \times SU(N_f)_R$. In addition the theory contains “mesino” fields that transform in either the symmetric of $SU(N_f)_L$ or the conjugate symmetric of $SU(N_f)_R$.

The dual pair admits a realization in non-critical type 0' string theory. The duality between the two theories has been derived by using a worldsheet argument [9, 10]. It can also be derived by using the “swapping branes” argument of [11].

We would also like to mention that the duality is supported by 't Hooft matching of global anomalies [2]

$$SU(N_f)^3 \quad N_c d^3(\square), \quad (2)$$

$$SU(N_f)^2 U(1)_R \quad \frac{-N_c^2 + 2N_c}{N_f} d^2(\square), \quad (3)$$

$$U(1)_R^3 \quad N_c(N_c - 1 - 2\frac{(N_c - 2)^3}{N_f^2}), \quad (4)$$

$$U(1)_R \quad -N_c^2 + 3N_c. \quad (5)$$

In the Veneziano large- N limit the electric theory (1) becomes equivalent to the electric $U(N_c)$ SQCD and similarly the magnetic theory (2) becomes equivalent to the magnetic $U(\tilde{N}_c)$ SQCD theory [3, 2]. Hence the large- N_c Seiberg duality between the theories (1) and (2) is straightforward.

Electric Theory				
	$U(N_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_R$
A_μ	adjoint N_c^2	•	•	0
λ	\square $\frac{N_c(N_c-1)}{2}$	•	•	1
$\tilde{\lambda}$	$\bar{\square}$ $\frac{N_c(N_c-1)}{2}$	•	•	1
Φ	$\bar{\square}$ \bar{N}_c	\square N_f	•	$\frac{N_f-N_c+2}{N_f}$
Ψ	\square N_c	\square N_f	•	$\frac{-N_c+2}{N_f}$
$\tilde{\Phi}$	\square N_c	•	$\bar{\square}$ \bar{N}_f	$\frac{N_f-N_c+2}{N_f}$
$\tilde{\Psi}$	$\bar{\square}$ \bar{N}_c	•	$\bar{\square}$ \bar{N}_f	$\frac{-N_c+2}{N_f}$

Table 1: *The matter content of the electric theory.*

Encouraged by the matching of the global anomalies at any N_c and N_f we propose that the duality holds not only at infinite N_c but at finite- N_c as well. As we shall see, quantum effects will lead to the decoupling of certain fields, resulting in a duality between a QCD-like electric theory and a magnetic theory of massless Nambu-Goldstone mesons.

3 Quantum effects

The electric and the magnetic theories are both non-supersymmetric. Therefore scalar field will admit a non-trivial potential, induced by quantum loops.

Magnetic Theory ($\tilde{N}_c = N_f - N_c + 4$)				
	$U(\tilde{N}_c)$	$SU(N_f)_L$	$SU(N_f)_R$	$U(1)_R$
A_μ	adjoint \tilde{N}_c^2	•	•	0
λ	\square $\frac{\tilde{N}_c(\tilde{N}_c-1)}{2}$	•	•	1
$\tilde{\lambda}$	$\overline{\square}$ $\frac{\tilde{N}_c(\tilde{N}_c-1)}{2}$	•	•	1
ϕ	\square \tilde{N}_c	$\overline{\square}$ \tilde{N}_f	•	$\frac{N_c-2}{N_f}$
ψ	$\overline{\square}$ \tilde{N}_c	$\overline{\square}$ \tilde{N}_f	•	$\frac{N_c-N_f-2}{N_f}$
$\tilde{\phi}$	$\overline{\square}$ \tilde{N}_c	•	\square N_f	$\frac{N_c-2}{N_f}$
$\tilde{\psi}$	\square \tilde{N}_c	•	\square N_f	
M	•	\square N_f	$\overline{\square}$ \tilde{N}_f	$\frac{2N_f-2N_c+4}{N_f}$
χ	•	\square $\frac{N_f(N_f+1)}{2}$	•	$\frac{N_f-2N_c+4}{N_f}$
$\tilde{\chi}$	•	•	$\overline{\square}$ $\frac{N_f(N_f+1)}{2}$	$\frac{N_f-2N_c+4}{N_f}$

Table 2: *The matter content of the magnetic theory.*

3.1 Electric theory

The electric theory contains squark fields Φ and $\tilde{\Phi}$. The squark field couples to the gluon and to the antisymmetric gluino. Let us consider the one-

loop correction the the squark mass. There are N_c^2 gluons and $N_c^2 - N_c$ gluinos. Therefore, unlike the supersymmetric case, the imbalance between the bosonic and fermionic degrees of freedom that run in the loop lead to a positive mass square for the squark fields

$$M_{\Phi}^2 = M_{\tilde{\Phi}}^2 = g_e^2 \Lambda^2, \quad (6)$$

where g_e is the electric gauge coupling and Λ is the UV cut-off of the theory. We therefore expect that the squarks will decouple from the low-energy dynamics.

In addition, similarly to $U(N_c)$ SQCD, the coupling of the $U(1)$ gauge boson (the “photon”) flows to a zero value in the IR, due to the coupling of the photon to the quarks.

Thus the low-energy electric theory becomes an $SU(N_c)$ gauge theory coupled to N_f massless quarks and an additional massless Dirac fermion that transforms in the antisymmetric representation. Such a theory is expected to confine and to admit chiral symmetry breaking. According to both Coleman-Witten argument [12] and Vafa-Witten theorem [13], the pattern of the chiral symmetry breaking is given by eq.(1).

3.2 Magnetic Theory

Similarly to the electric theory, the magnetic theory contains scalar quarks (“squarks”). The squarks couple via a gauge interaction (denoted by g_m) to the gluon and to the gluino and via a “Yukawa” interaction (denoted by y) to the meson and to the mesino. The Yukawa interaction is inherited from the supersymmetric theory, due to the superpotential $W = MQ\tilde{Q}$. Moreover, for $SU(N_c)$ SQCD, when $N_c < N_f < \frac{3}{2}N_c$, the couplings g_m and y are related to each other [14]

$$\frac{g_m^2}{y^2} = \frac{(3\frac{N_f}{N_c} - 1)(\frac{N_f}{N_c} - 1)}{(\frac{N_f}{N_c} - 1 - \frac{2}{N_c^2})}. \quad (7)$$

We assume (or postulate) that the relation between g_m and y is the same as in the supersymmetric theory, and in particular $g_m > y$.

We now turn to the calculation of the squark mass. The details of the calculation are very similar to those presented in [5]. The generated mass at the one-loop level is

$$M_{\phi}^2 = M_{\tilde{\phi}}^2 = (g_m^2 - y^2)\Lambda^2, \quad (8)$$

hence $M_\phi^2 = M_{\tilde{\phi}}^2 = cg_m^2 \Lambda^2 > 0$ (with $c \sim 1$). We conclude that the magnetic theory contains N_f massless quarks and N_f massive squarks.

The meson field $M_i^{\tilde{i}}$ couples to the quarks and the squarks. The mass splitting between them generates a Coleman-Weinberg potential for the meson. By using a chiral transformation it is possible to diagonalize the matrix $M_i^{\tilde{i}}$

$$yM_i^{\tilde{i}} = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_{N_f}). \quad (9)$$

The generated one-loop Coleman-Weinberg potential takes the following form

$$\begin{aligned} V(\{\lambda_i\}) &= \quad (10) \\ \tilde{N}_c \left(\text{tr} \int \frac{d^4 p}{(2\pi)^4} \log(p^2 + y^2(MM^\dagger) + cg_m^2 \Lambda^2) - \text{tr} \int \frac{d^4 p}{(2\pi)^4} \log(p^2 + y^2(MM^\dagger)) \right) \\ &= \frac{\tilde{N}_c}{4\pi^2} \sum_{i=1}^{N_f} \{ (\lambda_i^2 + cg_m^2 \Lambda^2)^2 \log(\lambda_i^2 + cg_m^2 \Lambda^2) - (\lambda_i^2)^2 \log(\lambda_i^2) \}. \end{aligned}$$

The potential (10) is plotted in figure (1) below.




Figure 1: The potential for the eigenvalue of the meson field (divided by the cut-off). In this plot we used $cg_m^2 = 0.005$.

The potential admits a unique minimum where all the eigenvalues of $M_i^{\tilde{i}}$ coincide

$$\langle M_i^{\tilde{i}} \rangle \sim \Lambda \delta_i^{\tilde{i}}, \quad (11)$$

namely the vacuum configuration breaks $SU(N_f)_L \times SU(N_f)_R$ to $SU(N_f)_V$ as well as $U(1)_R$ completely.

The breaking

$$SU(N_f)_L \times SU(N_f)_R \times U(1)_R \rightarrow SU(N_f)_V \quad (12)$$

results in N_f^2 Nambu-Goldstone bosons that correspond to flat directions of the potential (10).

By using the SQCD relation between the meson field and the electric variables,

$$M_i^{\tilde{i}} = \tilde{\Phi}_a^{\tilde{i}} \Phi_i^a \quad (13)$$

and the equations of motion for the electric squark fields

$$\Phi_i^a \sim \lambda^{[ab]} \Psi_{bi} \quad (14)$$

$$\tilde{\Phi}_a^{\tilde{i}} \sim \tilde{\lambda}_{[ab]} \tilde{\Psi}^{\tilde{i}b} \quad (15)$$

We conclude that the breaking (12) due to the vacuum expectation value of the meson field (11) can be written in the electric variables as a four-quark condensate of the form

$$\langle \tilde{\lambda}_{[ab]} \tilde{\Psi}^{\tilde{i}b} \lambda^{[ac]} \Psi_{ci} \rangle \sim \Lambda^6 \delta_i^{\tilde{i}}. \quad (16)$$

As a consistency check of our derivation we note that the four-quark condensate carries the same R -charge as the meson field, $R = \frac{2N_f - 2N_c + 4}{N_f}$.

At the minimum of the potential both the quarks and the squarks acquire a (large) mass proportional to $\langle M_i^{\tilde{i}} \rangle$, due to Yukawa terms. The large mass of the matter fields leads to a decoupling of the massless Nambu-Goldstone mesons from the “vector multiplet” (the gluon and the gluino). The $SU(\tilde{N}_c)$ gauge theory will confine and will acquire a mass gap, due to the formation of massive glueballs. In addition there will be a free $U(1)$ “photon”, that can be identified with the corresponding photon of the electric theory.

4 Summary

In this paper we studied a Seiberg duality between a QCD-like $U(N_c)$ electric theory and a $U(\tilde{N}_c)$ magnetic theory. The electric theory flows at low energies to an $SU(N_c)$ gauge theory with N_f massless quarks and an additional massless anti-symmetric fermion (plus an additional “photon”). The

theory admits a $SU(N_f)_L \times SU(N_f)_R \times U(1)_R$ global symmetry. The global symmetry may get broken due a formation of quark condensates.

We analyzed the fate of the global symmetry by using a weakly coupled magnetic theory. We used a one-loop analysis that indicated a breaking pattern of the form (12).

We must stress that the one-loop analysis cannot be fully trusted. At the minimum of the potential the vacuum expectation value of the meson field is large and the magnetic theory becomes strongly coupled. The one-loop Coleman-Weinberg can only be used to argue that the symmetric point $M_i^{\tilde{}} = 0$ is unstable. Nevertheless, our result is consistent with both Coleman-Witten [12] and Vafa-Witten [13].

It will be interesting to explore other aspects of QCD by using the proposed duality.

Acknowledgments. I wish to thank to professor Graham Shore for a careful reading of the manuscript.

References

- [1] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” Nucl. Phys. B **435**, 129 (1995) [hep-th/9411149].
- [2] A. Armoni, D. Israel, G. Moraitis and V. Niarchos, “Non-Supersymmetric Seiberg Duality, Orientifold QCD and Non-Critical Strings,” Phys. Rev. D **77**, 105009 (2008) [arXiv:0801.0762 [hep-th]].
- [3] A. Armoni, M. Shifman and G. Veneziano, “Exact results in nonsupersymmetric large N orientifold field theories,” Nucl. Phys. B **667**, 170 (2003) [hep-th/0302163]; A. Armoni, M. Shifman and G. Veneziano, In *Shifman, M. (ed.) et al.: From fields to strings, vol. 1* 353-444 [hep-th/0403071].
- [4] S. Sugimoto, “Confinement and Dynamical Symmetry Breaking in non-SUSY Gauge Theory from S-duality in String Theory,” Prog. Theor. Phys. **128**, 1175 (2012) [arXiv:1207.2203 [hep-th]].
- [5] A. Armoni, “Non-Supersymmetric Brane Configurations, Seiberg Duality and Dynamical Symmetry Breaking,” arXiv:1310.2027 [hep-th].

- [6] A. Hook and G. Torroba, “S-duality of nonsupersymmetric gauge theories,” arXiv:1309.5948 [hep-th].
- [7] A. Sagnotti, “Some properties of open string theories,” In *Palaiseau 1995, Susy 95* 473-484 [hep-th/9509080].
- [8] A. Sagnotti, “Surprises in open string perturbation theory,” Nucl. Phys. Proc. Suppl. **56B**, 332 (1997) [hep-th/9702093].
- [9] S. Murthy and J. Troost, “D-branes in non-critical superstrings and duality in N=1 gauge theories with flavor,” JHEP **0610**, 019 (2006) [hep-th/0606203].
- [10] S. K. Ashok, S. Murthy and J. Troost, “D-branes in Unoriented Non-critical Strings and Duality in SO(N) and Sp(N) Gauge Theories,” JHEP **0706**, 047 (2007) [hep-th/0703148 [HEP-TH]].
- [11] S. Elitzur, A. Giveon and D. Kutasov, “Branes and N=1 duality in string theory,” Phys. Lett. B **400**, 269 (1997) [hep-th/9702014].
- [12] S. R. Coleman and E. Witten, “Chiral Symmetry Breakdown in Large N Chromodynamics,” Phys. Rev. Lett. **45**, 100 (1980).
- [13] C. Vafa and E. Witten, “Restrictions on Symmetry Breaking in Vector-Like Gauge Theories,” Nucl. Phys. B **234**, 173 (1984).
- [14] R. Oehme, “Reduction of dual theories,” Phys. Rev. D **59**, 105004 (1999) [hep-th/9808054].