

Coarse Graining in Hydrodynamics and Effects of Fluctuations

T. Kodama and T. Koide

Abstract. We address the physical meaning of hydrodynamic approach related with the coarse graining scale in the frame work of variational formulation. We point out that the local thermal equilibrium does not necessarily play a critical role in the description of the collective flow patterns. We further show that the effect of viscosity is also formulated in the form of the variational method including fluctuations.

Instituto de Física, Universidade Federal do Rio de Janeiro, C.P. 68528, 21941-972, Rio de Janeiro, Brazil

E-mail: tkodama@if.ufrj.br, koide@if.ufrj.br

1. Introduction

Hydrodynamic approach on event-by-event (EBE) basis has shown to be very successful to describe the global and collective features of the data from relativistic heavy-ion collisions, particularly of the behavior of the Fourier components of flow pattern $\{v_n\}$ as a function of centrality and transverse momenta data [1]. Such a success of the hydrodynamic approach leads to the expectation that we can determine from a detailed hydrodynamic analysis of experimental data, the initial dynamics of the collisions together with the properties of QCD matter created, such as the equation of state (EoS) and transport coefficients. On the other hand, these successes brought us several new interesting questions and mysteries. The most crucial one is why at all the hydrodynamic approaches work so well for such violent and almost microscopic collisional processes. It is commonly believed that the fundamental hypothesis for the validity of hydrodynamics is the local thermal equilibrium (LTE). If this is true, and the hydrodynamic evolution is the unique scenario for the description of the collision dynamics, then we are led to conclude that the thermalization time and correlation length should be extremely small. These would imply a very important consequence for the further understandings of the QCD dynamics at extreme conditions. For AA collisions this could still be acceptable, but surprisingly, the recent ALICE experiment reports that the behavior of collective flow in the pA data seems to be almost the same as that of AA collisions [2]. This casts a very serious question for the proper physical meaning of hydrodynamic description in pA collisions or, even in AA collisions [3].

2. Relativistic Hydrodynamics and Role of Coarse Graining

Let us denote the conserved four-current density by $n^\mu(x)$, satisfying the continuity equation, $\partial_\mu n^\mu(x) = 0$. We consider the Minkowsky coordinate, $x = \{x^0, x\}$ with metric the $g_{\mu\nu} = \text{diag}\{1, -1, -1, -1\}$. The energy-momentum tensor $T^{\mu\nu}(x)$ of the system also conserves, $\partial_\mu T^{\mu\nu}(x) = 0$. Although these 4 equations are far enough to determine the time evolution of these quantities, but in some special physical situations, the total number of variables reduces

drastically. Suppose that, in the Landau frame (that is, the energy flow rest frame), the spatial part of $T^{\mu\nu}(x)$ becomes isotropic for any x . Furthermore, the rest frames for the matter and energy current coincide. In such a situation, by introducing the Equation of State (EoS) which establishes a functional relation among the local quantities, ε , P and n , as $P = P(\varepsilon, n)$, we obtain the closed set of equations. This is called the ideal hydrodynamics and the explicit forms of the time evolution equations are, $\partial_t \varepsilon + (\vec{v} \cdot \nabla) \varepsilon = -\gamma^{-1} (\varepsilon + P) \theta$, $\partial_t n + (\vec{v} \cdot \nabla) n = -\gamma^{-1} n \theta$, and $d\{(\varepsilon + P)/n\}/dt = -\nabla P/n^*$ where n^μ is the time-like eigenvector of $T^{\mu\nu}(x)$ with $\theta = \partial_\mu u^\mu$, $\gamma = u^0$, $\vec{v} = \vec{u}/\gamma$ and $n^* = \gamma n$.

In the above, the assumption that EoS $P = P(\varepsilon, n)$ is locally satisfied in the strict sense is somewhat a very severe condition. For example, when we consider the hydrodynamic description of the relativistic heavy-ion collisions, the size of the typical fluid element cannot be taken too much smaller than that of the whole system, otherwise the degree of freedom contained in the fluid element becomes too little for any thermodynamical quantities to be defined. Furthermore, the time scale of the collective motion cannot be much larger than the microscopic one because of the very rapid expansion in relativistic heavy-ion collisions. Then, a given fluid element suffers from large fluctuations and inhomogeneity in terms of the microscopic configurations and it will be difficult to ignore the deviation from the thermodynamic limit in EoS. However, we will argue in below that the hydrodynamic responses in, for example, the collective flow do not necessarily require the strictly local EoS in the EBE basis.

For example, let us consider a classical microscopic system which contains a large number of quickly moving point-like particles. Then, the density n_0^* is a sum of the Dirac delta functions. However, we usually do not require a very precise resolution both in space and in time to describe the collective flow behaviors. Thus we introduce an averaged smooth density distribution $\tilde{n}^*(\vec{x}, t)$ from the original distribution $n_0^*(x)$ using a 4 dimensional smoothing kernel with limited support[4] $W(x) = U_\tau(t) \times W_h(\vec{x})$

$$\tilde{n}^*(\vec{x}, t) = \int dt' \int d^3\vec{X} n_0^*(\vec{X}) U_\tau(t' - t) W_h\left(\left|\vec{x} - \vec{r}(t'; \vec{X})\right|\right), \quad (1)$$

Typically U and W are given by the Gaussian distributions with, respectively, width τ and h , which characterize the scales of the time and space resolutions. Similarly, the smoothed spatial current vector $\tilde{\mathbf{j}}(\vec{x}, t)$ can be defined, satisfying the continuity equation, $\partial \tilde{n}^*/\partial t + \nabla_x \cdot \tilde{\mathbf{j}} = 0$.

Using these current and density, the four-current, $\tilde{j}^\mu = (\tilde{n}^*, \tilde{\mathbf{j}})$ and the proper density, $\tilde{n} = \sqrt{\tilde{j}^\nu \tilde{j}_\nu}$ can be composed. The smoothed four-velocity field is then defined as $\tilde{u}^\mu = \tilde{j}^\mu / \tilde{n}$. On the other hand, the smoothed energy-momentum tensor $\tilde{T}^{\mu\nu}$ can be introduced in analogous way as the convolution of the original $T^{\mu\nu}$ using the same smoothing kernel. Such energy-momentum tensor again satisfies the continuity equation, $\partial_\mu \tilde{T}^{\mu\nu} = 0$. From this smoothed energy-momentum tensor, we can calculate the smoothed proper energy density as $\tilde{\varepsilon} \equiv \tilde{u}_\mu \tilde{u}_\nu \tilde{T}^{\mu\nu}$. The smoothed proper energy density defined in this way is an average of the energy density observed in the rest frame of the *matter* flow. The average is taken over all contributions within the range of the coarse-graining scale in space-time. In terms of the hydrodynamic modeling, we take these smoothed quantities as the dynamical variables to represent the coarse-grained system[4].

Let us consider one collision event which is characterized by a microscopic state. Then we can calculate these hydrodynamic variables of this state following the method described above. However, it is obvious that there exist many different microscopic configurations which give the same hydrodynamic response. Let us prepare the set of collision events described by microscopic configurations which gives a given four-current \tilde{j}^μ at the initial time t_0 , and call this set Ω . If we calculate $\tilde{\varepsilon}$ at a space-time point x for each event in Ω , each value of $\tilde{\varepsilon}$ is not same in general. This is true even for the time evolution, $\tilde{j}^\mu(\vec{x}, t)$ with $t > t_0$.

However, if the coarse-graining size is increased, the number of the microscopic configuration in the ensemble Ω at the point x becomes sufficiently large so that $\tilde{\varepsilon}$ and \tilde{n} distribute sharply around their mean-values, $\tilde{\varepsilon}$ and \tilde{n} , respectively, as a consequence of the central limit theorem. Furthermore, since $\tilde{\varepsilon}$ and \tilde{n} are the average energy and matter densities belonging to the same fluid element, we expect that they have a strong correlation, in such a way that $\tilde{\varepsilon}$ can be expressed as a function of \tilde{n} , $\tilde{\varepsilon} = \tilde{\varepsilon}(\tilde{n})$.

Suppose that the fluctuations in $\tilde{\varepsilon}$ and \tilde{j}^μ are not important in the way that the system is characterized basically by the densities \tilde{n}^* and $\tilde{\varepsilon}$. In such a case, we expect that the most promising dynamics will be determined by the optimization of the model action,

$$I = - \int d^4x \tilde{\varepsilon} \left(\frac{1}{\gamma} \tilde{n}^* \right). \quad (2)$$

It is known that this procedure reproduces the model of the ideal hydrodynamics [5]. As a result, the dynamics of the system belonging to a given Ω is described by the hydrodynamic model as the consequence of the coarse graining. There, the realization of LTE is not required for real EBE basis. When the effect of the fluctuations in $\tilde{\varepsilon}$ and \tilde{j}^μ plays an important role, see Sec. 4.

3. Necessity of Real Event by Event Analysis

As shown above, the hydrodynamic description in heavy-ion collisions reduces to a coarse-grained dynamics obtained by the optimization of the model action (2) under the assumption of the existence of an effective EoS, $\tilde{\varepsilon} = \tilde{\varepsilon}(\tilde{n})$. Therefore, the success of ideal hydrodynamic modeling of relativistic heavy ion collisions depends on the consistent choice of the assumed EoS and the model action. These two conditions will be satisfied for a broader range of microscopic configurations than that required by the real "local thermal equilibrium".

On the other hand, the size of Ω depends on the coarse-graining scale. For larger Ω , the two conditions have a better chance to be satisfied. We however lose the better resolution in the space-time recognition for larger coarse graining size. In fact, we cannot observe the inhomogeneities with smaller wavelength than the coarse-graining scale. This affects directly the class of observables that the model can describe. Even though some observables might be insensitive to the inhomogeneities in each event. As an extreme example, we take the situation where the coarse-graining size is larger than the system size and total time evolution. Then the ensemble Ω can be regarded as the statistical ensemble of the whole system itself, and the resultant system is a simple fire-ball model. The thermal model for the particle ratio can be considered in this category.

The coarse-graining size is thus intimately related to the class of observables and the validity of hydrodynamic description. For some observables which do not require a precise space-time resolution, the real EBE hydrodynamics with LTE is not necessary and the effective hydrodynamic description for the statistical ensemble Ω will be sufficient for the understanding of the physics of these observables. As a matter of fact, the experimental observables are usually averaged over collision events classified in terms of the initial configuration rather loosely defined, such as centrality, event plane, etc. In other words, the present collective flow data are still of inclusive nature. In order to claim that the real hydrodynamics with LTE is valid, we need to have observables that reflect the genuine hydrodynamic profile in EBE basis. For example, the remnant of a sharp shock wave propagation, if exists, would be a good evidence and it also tells the possible coarse-graining size of the collective flow, since a shock wave is a genuine local hydrodynamic phenomena. The shock thickness should not be larger than the coarse-graining scale of the collective flow.

The key point is that when we apply the hydrodynamic modeling, we do not know a priori the coarse-graining scale suitable for the flow variables in the real scenario. This puts a certain

limitation in extracting the meaningful information of the initial condition. For this purpose, it is essential to find out the set of observables which carry the information on the inhomogeneities of the initial conditions on the EBE basis. The flow parameters $\{v_n\}$, often called "event-by-event" analysis, in the sense that correlations among different observables measured for each event in coincidence, but there still exists a huge statistical ensemble which gives the same observed correlation. For example, the cumulant method to determine the flow parameters eliminates the information of event plane. In the recent paper, it is pointed out that event plane may differ in low and high p_T domain [4], according to the coarse graining scale. If it can be experimentally measured, it would furnish some information on coarse graining scale in heavy-ion collisions.

4. Fluctuation of Fluid Variables and Stochastic Variational Method

Within the vision that hydrodynamic evolution is an effective dynamics for coarse-grained variables for the energy-momentum tensor, each real collisional event is an element of the statistical ensemble Ω and does not obey a unique time evolution equation due to the difference in the microscopic degrees of freedom to which our macroscopic hydrodynamic variables are blind. When the fluctuation of events in Ω is large, they should be taken into account in the determination of dynamics of coarse-grained hydrodynamic variables. Then the variation procedure in Eq. (2) should be modified so as to include the effect of the fluctuation which was ignored in Sec. 3. The stochastic variational method (SVM) is known as an appropriate approach for such situations[6].

In SVM, we have to introduce two stochastic differential equations (SDE), one for the forward direction in time (FSDE), $d\mathbf{r}(t) = \mathbf{u}dt + \sqrt{2\nu}d\mathbf{W}(t)$, ($dt > 0$), and the other, backward in time (BSDE), which describes the time reversed process of FSDE, $d\mathbf{r}(t) = \tilde{\mathbf{u}}dt + \sqrt{2\nu}d\tilde{\mathbf{W}}(t)$, ($dt < 0$), where ν is the strength of the noise and $\mathbf{W}(t)$ and $\tilde{\mathbf{W}}$ are the independent Wiener processes. \mathbf{u} and $\tilde{\mathbf{u}}$ are the velocity fields for the forward and backward SDEs, respectively. These two SDEs are necessary to accommodate the fixed initial and final boundary conditions in the variational procedure. Therefore, trajectories specified in the two SDE should describe the same physical ensemble. To satisfy this condition, the two Fokker-Planck equations which are derived from FSDE and BSDE should be equivalent. This leads to the consistency condition, $\mathbf{u} = \tilde{\mathbf{u}} + 2\nu\nabla \ln \rho$, where ρ is the particle density which is given by the solution of the Fokker-Planck equation. SVM proposes to determine these velocity fields from an action through the variation principle.

We started from the classical action which leads to the ideal Euler equation for a non-relativistic fluid,

$$I(\mathbf{r}) = \int_{t_a}^{t_b} dt \int d^3\mathbf{R} \left[\frac{\rho_0^m(\mathbf{R})}{2} \left(\frac{d\mathbf{r}(\mathbf{R}, t)}{dt} \right)^2 - J\varepsilon \right], \quad (3)$$

where $\mathbf{r}(\mathbf{R}, t)$ is the Lagrangian coordinate associated with the fluid element \mathbf{R} , and ε is an internal energy density, \mathbf{J} is the Jacobian between \mathbf{r} and \mathbf{R} , and the mass density is defined by $\rho_m = m\rho$. The spatial integral is done over the initial position \mathbf{R} of fluid elements, weighted by the mass density $\rho_0^m(\mathbf{R})$. The corresponding stochastic Lagrangian density is given by

$$\mathcal{L} = \frac{\rho_0^m}{2} \left[\left(\frac{1}{2} + \alpha_2 \right) \left\{ \left(\frac{1}{2} + \alpha_1 \right) \mathbf{u}^2 + \left(\frac{1}{2} - \alpha_1 \right) \tilde{\mathbf{u}}^2 \right\} + \left(\frac{1}{2} - \alpha_2 \right) \tilde{\mathbf{u}} \cdot \mathbf{u} \right] - J\varepsilon. \quad (4)$$

Here α_1 and α_2 is arbitrary constants, come from the ambiguity for the stochastic representation of the kinetic term. The corresponding action is an average over the whole SDE solutions, and the variation is taken with respect to the unknown fields, \mathbf{u} and $\tilde{\mathbf{u}}$, with the constraints that they coincide with the so-called conditional average velocity fields, $D\mathbf{r}$ and $D\tilde{\mathbf{r}}$.

After applying SVM, we obtain

$$\rho^m (\partial_t + \mathbf{u}_m \cdot \nabla) u_m^i - \sum_j \partial_j (\mu \rho_m e_{ij}^m) - 2\kappa \rho^m \sum_j \partial_i (\sqrt{\rho^m}^{-1} \partial_j^2 \sqrt{\rho^m}) = -\nabla P,$$

where $\eta = \alpha_1(1 + 2\alpha_2)\nu\rho^m$, $\kappa = 2\alpha_2\nu^2$ and $e_{ij}^m = \partial_j u_m^i + \partial_i u_m^j$. The pressure P is defined by $(\rho_m)^2 d(\varepsilon/\rho_m)/d\rho_m$. One can see that the second term on the left hand side corresponds to the viscosity and this equation is reduced to the Navier-Stokes (NS) equation when we set $\alpha_2 = 0$. That is, the fluctuation effects which were ignored in Sec. 3 induces the effects of viscosity in accordance with the fluctuation-dissipation theorem¹. For other results of different values of $\alpha's$, see Ref. [6].

5. Concluding remarks

In this work, we studied some fundamental questions of hydrodynamic approach to the description of relativistic heavy ion collisions, in particular, the validity and meaning of LTE. We introduced explicitly the coarse-graining procedure for the hydrodynamic modeling together with its variational formulation. In this picture, the collective flow patterns can be reproduced without requiring the LTE in a strict sense. That is, the hydrodynamic behavior observed in relativistic heavy-ion collisions does not necessarily imply the realization of LTE². We further discussed possible signals for coarse graining scale and genuine hydrodynamic behaviors on event by event basis. For example, the remnant of a sharp shock wave propagation would be a good observable which tells the possible coarse-graining size of the collective flow. Another example is to determine the event plane for different transverse momentum domain. Finally we showed that the coarse-graining is intimately related to the origin of viscosity and this effect can be formulated in the variational method extending dynamical variables to stochastic domain. In order to quantify the questions raised here, it will be useful to perform the analysis of coarse-graining described in this work for a certain microscopic model which gives complete dynamical evolution of the energy-momentum tensor, such as PHSD[8].

This work is supported by CNPq, FAPERJ and PRONEX of Brazil, and also a part of the work has been supported by EMMI-Helmholtz Institute, Germany.

References

- [1] S. Paiva et al., Phys. Rev. C 55, 1455 (1997); C.E. Aguiar, et al., Nucl. Phys. A 698, 639 (2002); R. Andrade, et al., Phys. Rev. Lett. 97, 202302 (2006); R. Andrade, et al., Jr., Braz. J. Phys. 37, 717 (2007); J. Takahashi, et al., Phys. Rev. Lett. 103, 242301 (2009); D.J. Wang, et al., arXiv:1205.4934 (2012); Z. Qiu, et al., Phys. Lett. B 707, 151 (2012); B. Alver and G. Roland, Phys. Rev. C 81, 054905 (2010); D. Teaney and Li Yan, Phys. Rev. C 83, 064904 (2011); B. Schenke, et al., Phys. Rev. Lett. 108, 252301 (2012); A. Dumitru and Y. Nara, Phys. Rev. C 85, 034907 (2012).
- [2] ALICE Collaboration, arXiv:1307.3237.
- [3] M. Gyulassy, talk in 9th International Workshop on Relativistic Aspects of Nuclear Physics, Rio de Janeiro, October, 2013.
- [4] Ph. Mota, et al., Eur. Phys. J. A (2012) 48: 165.
- [5] H-T. Elze, et al., J. Phys. G Nucl. Part. Phys. 25, 1935 (1999).
- [6] See for example Koide T and Kodama T 2012 J. Phys. **A45** 255204 and references therein.
- [7] T. Kodama et al., Int. J. Mod. Phys. E16, 763 (2007); L. McLerran, talk in 9th International Workshop on Relativistic Aspects of Nuclear Physics, Rio de Janeiro, October, 2013.
- [8] See for example, E.L. Bratkovskaya, et al., Nucl.Phys. A856 (2011) 162-182 and references therein.

¹ Exactly speaking, the second coefficient of viscosity vanishes in this equation. To obtain a finite value, we have to consider the variation of entropy.

² This has been already noticed before. See for example, Ref. [7]