



April 4, 2024

Lattice QCD review of charmonium and open-charm spectroscopy

SASA PRELOVSEK¹

*Jozef Stefan Institute, Ljubljana, Slovenia
University of Ljubljana, Slovenia*

Lattice QCD results on the spectroscopy of charmonium(like) states and open charm mesons are reviewed. Near-threshold states $X(3872)$ and $D_{s0}^*(2317)$ were treated rigorously for the first time and the searches for $Z_c^+(3900)$, $X(4140)$ and $cc\bar{u}\bar{d}$ were carried out. The first simulations of the resonances with charm quarks have been performed, including the determination of their strong decay widths. Spectroscopy of highly excited charmed, charmonium and hybrid states has been calculated.

PRESENTED AT

The 6th International Workshop on Charm Physics
(CHARM 2013)
Manchester, UK, 31 August – 4 September, 2013

¹ sasa.prelovsek@ijs.si

1 Introduction and motivation

The masses of states well below the open charm threshold, which starts at $m_D + m_\pi$ for D mesons and at $2m_D$ for charmonium, have been reliably and precisely calculated on the lattice for several years. With exception of [1], the states near threshold or above threshold have been treated using the so-called “single-meson” approach until 2012, which essentially means that effect of the threshold and strong decays of these states were ignored. The impressive spectrum of high lying quark-antiquark and hybrid multiplets was calculated during the past two years using this less-complete treatment. In addition, the first exploratory lattice simulations of the near-threshold states and resonances based on the rigorous treatment were done during the last year. These are the main focus of this talk.

Proper treatment of near-threshold states in lattice QCD is particularly important since most exotic experimental candidates lie near open-charm thresholds. Examples include $X(3872)$ within 1 MeV of $D^0 D^{0*}$ threshold. The structure called $Z_c^+(3900)$ was discovered in $J/\psi \pi^+$ invariant mass in 2013 by BESII [2] and confirmed by Belle and CLEOc [3, 4]. It has quark flavor structure $\bar{c}c\bar{d}u$ and also lies close to DD^* threshold. Other charged charmonium-like states $Z^+(4430)$ and more recent $Z_c^+(4020)$ [5], $Z_c^+(4025)$ [6] have been observed only in one experiment. They also have flavor $\bar{c}c\bar{d}u$ and lie near $D^* D_1$ and $D^* D^*$ thresholds. The only lattice study aimed at $Z_c(3900)$ will be reviewed, while $Z^+(4430)$ was addressed only in [7].

Masses are extracted from the energy levels E_n , and these are obtained from the correlation functions $C_{ij}(t) = \langle O_i(t) O_j(0) \rangle = \sum_n \langle O_i | n \rangle \langle O_j | n \rangle^* e^{-E_n t}$ evaluated on the lattice, where interpolating fields O create and destroy the physical state with given quantum number J^{PC} . The simulations of charmonium(like) states discussed below neglect the charm annihilation Wick contractions in view of OZI suppression. For hadrons with charm quarks it is common to compare $m - m_{reference}$ between lattice and experiment, where the leading discretization errors related to m_c cancel.

2 States well below open charm thresholds

The recipe for extracting a mass of a state well below the strong decay threshold is straightforward. The mass $m = \sqrt{E^2 - P^2}$ in a simulation with $P=0$ is equal to the energy level E . The resulting mass is extrapolated to $a \rightarrow 0$ and $L \rightarrow \infty$, while the quark masses are extrapolated or interpolated to the physical value. Particular care needs to be taken about the discretization errors related to the c quarks and it has been verified that independent methods lead to compatible results in the $a \rightarrow 0$ limit.

The precise masses for low-lying states were obtained recently after extrapolations by HPQCD [11, 12], Briceño *et al.* [13], FNAL/MILC [14] and χ QCD [15], while related studies are in [16, 17, 18, 19]. The resulting masses or mass splittings are




Figure 1: The D (above) and D_s (below) spectra and interpretation from HSC at $m_\pi \simeq 400$ MeV [8] (plot from G. Moir).

typically within 10 MeV from the experimental value.

3 “Single-meson” treatment of excited states

The “single-meson” treatment of states near or above threshold refers to (i) using only quark-antiquark interpolating fields $O \sim \bar{q}q$ for mesons, (ii) assuming that all energy levels correspond to “one-particle” states and (iii) that the mass of the state equals the measured energy level $m = E$. These are strong assumptions for the resonances, which are not asymptotic states. The approach also ignores the effect of the threshold on near-threshold states.

The most extensive D , D_s [8] and $\bar{c}c$ [9] spectrum, shown in Figs. 1 and 2, was extracted by the Hadron Spectrum Collaboration (HSC) on $N_f = 2+1$ anisotropic




Figure 2: The $c\bar{c}$ spectrum and interpretation from the HSC at $m_\pi \approx 400$ MeV [9] (plot from talk of C. Thomas at this meeting).

configurations with $m_\pi \simeq 400$ MeV and two different $L \simeq 2.9$ fm, 1.9 fm. The continuum J^{PC} was reliably identified using advanced spin-identification method. An impressive number of excited states was extracted in each channel with a good accuracy. States are identified with members of $\bar{q}q$ multiplets $1S$, $2S$, $3S$, $1P$, $2P$, $1D$ and $1F$ based on overlaps $\langle O_i | n \rangle$, where interpolators are chosen to resemble multiplet members. There are several remaining states, which are not part of $\bar{q}q$ multiplets and




Figure 3: The D and $\bar{c}c$ spectrum extracted in [10] at $m_\pi \simeq 266$ MeV.

these states show particularly strong overlap with interpolators $O \simeq \bar{q}F_{\mu\nu}q$ containing the gluon field strength tensor. These states are identified as hybrids and also turn out to form several full multiplets. Some of the identified charmonium hybrids have manifestly exotic J^{PC} , while the others have conventional values. The open-charm meson hybrids have non-exotic J^P .

The D meson spectrum in Fig. 3 from a simulation with $m_\pi \simeq 266$ MeV [10] shows reasonable agreement with 2010 Babar [20] and 2013 LHCb [21] excited states, which are shown side by side. The blue crosses result from the “single-meson” treatment, while the red diamonds are resonance masses from the rigorous treatment. The splitting between $1S$ and $2S$ multiplets of the order $\simeq 650$ MeV is in agreement experiments. The new $\bar{c}c$ state 2^{--} [22] agrees with the prediction in Fig. 3.

Other recent related results can be found in [23, 24, 25, 19].

4 Rigorous treatment of resonances

$D_0^*(2400)$ and $D_1(2430)$: Let me illustrate the basics of the rigorous treatment for the example of $D_0^*(2400)$ scalar resonance, which appears in $D\pi$ scattering in s-wave. All states with given quantum number $J^P=0^+$ in principle appear as energy levels:

- Important examples are two-particle states $D(p)\pi(-p)$, where periodic boundary conditions require $p=n\frac{2\pi}{L}$ in absence of the interaction. The corresponding energy $E=\sqrt{m_1^2+p^2}+\sqrt{m_2^2+p^2}$ of non-interacting two-particle states will be marked by horizontal lines in the figures. The meson-meson interpolating fields need to be incorporated to make sure these levels are seen in practice.
- The levels that appear in addition to the expected two-particle states are related to the bound states or resonances.

The first and the third levels in Fig. 4 correspond to the interacting $D\pi$ two-particle states, while the second level is related to $D_0^*(2400)$ resonance. Each energy level E renders an elastic phase shift $\delta(E)$ at that energy via the rigorous Lüscher relation [10]. The three energy levels give the three phase shift points around the $D_0^*(2400)$ resonance region. The Breit-Wigner type fit through these three points leads to the resonance mass in Fig. 3 very close to the experiment, and the resonance width reasonably close as well [10]. The mass and width of the broad $D_1(2430)$ resonance in $D^*\pi$ scattering is also close to experiment (see Fig. 3), but the analysis in this channel relies on few assumptions due to two nearby $D_1(2430)$ and $D_1(2420)$ [10].

χ'_{c0} : An indication for a charmonium $J^{PC}=0^{++}$ resonance with $m_{\chi'_{c0}}=3932\pm 25$ MeV and $\Gamma(\chi'_{c0}\rightarrow\bar{D}D)=36\pm 17$ MeV and additional enhancement of $\sigma(D\bar{D})$ near threshold in a simulation [27] prompts experiments to look for such structures in $D\bar{D}$ invariant mass (see [28, 29] for similar suggestions).




Figure 4: Left: the energy levels and $D\pi$ phase shifts in $D_0^*(2400)$ channel with $I=1/2$ and $J^P=0^+$ [10]. Right: the levels in $D_{s0}^*(2317)$ channel with $I=0$ and $J^P=0^+$ [26].

5 Rigorous treatment of near-threshold states

$D_{s0}(2317)$: Fig. 4 shows lowest three energy levels from the first simulation of $D_{s0}^*(2317)$ that takes into account the effect of the DK threshold by explicitly incorporating DK interpolating fields [26]. The position of the DK threshold is almost physical in this $N_f = 2 + 1$ simulation with nearly physical $m_\pi \simeq 156$ MeV. The second and third levels are related to the interacting DK states, while the first level is related to $D_{s0}^*(2317)$ and it is below threshold as in experiment. The lowest two levels lead to the scattering length $a_0 = -1.33 \pm 0.20$ fm and small effective range r_0 for DK scattering in s-wave. The negative a_0 is an indication for the presence of $D_{s0}^*(2317)$ below threshold [32]. The effective range expansion leads to the position of the pole at $L \rightarrow \infty$, rendering $m_{D_{s0}^*} - \frac{1}{4}(m_{D_s} + 3m_{D_s^*}) = 266 \pm 16$ MeV close to the experimental value 241.5 ± 0.8 MeV.

The “single-meson” treatment using just \bar{sc} interpolators gives only one level in Fig. 4 (orange triangles) with a misleading energy and one can not reliably establish whether this level corresponds to $D_{s0}^*(2317)$ or $D(0)K(0)$.

$X(3872)$: A candidate for the charmonium(like) state $X(3872)$ is found 11 ± 7 MeV below the $D\bar{D}^*$ threshold for $J^{PC} = 1^{++}$, $I = 0$, $N_f = 2$ and $m_\pi \simeq 266$ MeV [30]. This is the first lattice simulation that establishes a candidate for $X(3872)$ (violet stars in Fig. 5) in addition to χ_{c1} (squares) and the nearby scattering states $D\bar{D}^*$ (circles and diamonds) and $J/\psi\omega$ (triangles). The established $X(3872)$ has a large overlap with $\bar{c}c$ as well as $D\bar{D}^*$ interpolating fields [30]. The large and negative




Figure 5: Left: the energy levels in $X(3872)$ channel with $I=0$ and $J^{PC} = 1^{++}$ [30]. Right: the levels in $Z_c(3900)$ channel with $I=1$ and $J^{PC} = 1^{+-}$ [31].

$a_0 = -1.7 \pm 0.4$ fm for $D\bar{D}^*$ scattering is an indication for a shallow bound state $X(3872)$ [32].

The “single-meson” treatment using just $\bar{c}c$ interpolators renders only one level in Fig. 5 (orange triangles) near DD^* threshold just like in previous simulations. One can not reliably establish whether this level is related to $X(3872)$ or $D(0)\bar{D}^*(0)$.

In the $I=1$ channel, only the $D\bar{D}^*$ and $J/\psi \rho$ scattering states are found, and no candidate for $X(3872)$ [30]. This is in agreement with a popular interpretation that $X(3872)$ is dominantly $I=0$, while its small $I=1$ component arises solely from the isospin breaking and is therefore absent in the simulation with $m_u = m_d$.

6 Search for exotic states

Z_c(3900): Three experiments recently reported a discovery of a manifestly exotic $Z_c^+(3900)$ in the decay to $J/\psi \pi^+$ [31]. It has $C = -$, while J and P are experimentally unknown. The first search for this interesting state on the lattice focused on a channel with $J^{PC} = 1^{+-}$ and $I=1$. Fig. 5 indicates that only the expected two-particle states $D\bar{D}^*$ and $J/\psi \pi$ are found. No additional level was found that could be related to $Z_c^+(3900)$. The possible reasons for not finding Z_c^+ may be that its J^{PC} are not 1^{+-} or that the employed interpolating fields are not diverse enough. Simulation with additional types of interpolators will be needed to reach a more definite conclusion.

X(4140): A structure called $X(4140)$ was found in $J/\psi \phi$ invariant mass by CDF

[33] and more recently by CMS and D0 [34, 35]. The s-wave and p-wave $J/\psi \phi$ scattering phase shift in Fig. 6 was extracted in [36] ignoring $\bar{s}s$ annihilation contribution, and no resonant structure was found.




Figure 6: The phase shift for s-wave and p-wave scattering of $J/\psi \phi$ from [36].

$c\bar{c}d\bar{u}$ tetraquark: The potential $V(r)$ between $D = \bar{u}c$ and $D^* = \bar{d}c$ at distance r was extracted using the HALQCD time-dependent method [37]. The resulting potential in Fig. 7 is attractive, but the corresponding DD^* scattering phase shift does not start at $\delta(0) = \pi$, which indicates that there is no $c\bar{c}d\bar{u}$ tetraquark bound state at the simulated m_π .




Figure 7: The DD^* potential and phase shift in double-charm channel [38].

7 Related topics

Interactions of charmed mesons with light pseudoscalars: The scattering length was extracted in five non-resonant channels for four different m_π in a simulation [39]. The simultaneous fit to these channels renders the low-energy constants of the SU(3) Unitarized ChPT. These give indirect prediction for the resonant DK scattering with $I = 0$ and $J^+ = 0^+$, where the pole is found at $m_{D_{s0}^*} = 2315_{-28}^{+18}$ MeV close to the experimental $D_{s0}(2317)$ [39].

Charmonium potential at finite temperature: The $V_C(r)$ part of the potential $V(r) = V_C(r) + \vec{s}_1 \cdot \vec{s}_2 V_S(r)$ between \bar{c} and c at distance r was extracted using the HALQCD time-dependent method at non-zero temperatures in $N_f = 2 + 1$ simulation [40]. The resulting potential in Fig. 8 is temperature dependent and becomes flatter at large r as the temperature increases.




Figure 8: The spin-independent part of the $\bar{c}c$ potential in $N_f = 2 + 1$ simulation [40].

8 Conclusions

The states well below open charm decay threshold are reliably and precisely determined from lattice QCD. During the past two years, the impressive spectrum of D , D_s and $\bar{c}c$ quark-antiquark and hybrid multiplets was calculated within the simplified single-meson approach. Several near-threshold states and resonances were treated rigorously for the first time: candidates for $D_{s0}^*(2317)$, $X(3872)$, $D_0(2400)$ and $D_1(2430)$ were established, while $Z_c(3900)$, $X(4140)$ and bound $cc\bar{d}\bar{u}$ tetraquarks were not found (yet). Precision simulations of these channels in the future would be valuable.

ACKNOWLEDGEMENTS

I would like to thank my collaborators C.B. Lang, L. Leskovec, D. Mohler and R. M. Woloshyn. Thanks to everyone who sent me the material for this review. This work is supported by ARRS project number N1-0020 and FWF project number I1313-N27.

References

- [1] G. S. Bali, S. Collins and C. Ehmman, Phys.Rev. **D84**, 094506 (2011).
- [2] BESIII Collaboration, M. Ablikim *et al.*, Phys.Rev.Lett. **110**, 252001 (2013), [arXiv:1303.5949].
- [3] Belle Collaboration, Z. Liu *et al.*, Phys.Rev.Lett. **110**, 252002 (2013), [arXiv:1304.0121].
- [4] T. Xiao, S. Dobbs, A. Tomaradze and K. K. Seth, [arXiv:1304.3036].
- [5] BESIII Collaboration, M. Ablikim *et al.*, [arXiv:1309.1896].
- [6] BESIII Collaboration, M. Ablikim *et al.*, [arXiv:1308.2760].
- [7] CLQCD Collaboration, G.-Z. Meng *et al.*, Phys.Rev. **D80**, 034503 (2009).
- [8] G. Moir, M. Peardon, S. M. Ryan, C. E. Thomas and L. Liu, JHEP **1305**, 021 (2013), [arXiv:1301.7670].
- [9] Hadron Spectrum Collaboration, L. Liu *et al.*, JHEP **1207**, 126 (2012), [arXiv:1204.5425].
- [10] D. Mohler, S. Prelovsek and R. Woloshyn, Phys.Rev. **D87**, 034501 (2013), [arXiv:1208.4059].
- [11] R. Dowdall, C. Davies, T. Hammant and R. Horgan, Phys.Rev. **D86**, 094510 (2012), [arXiv:1207.5149].
- [12] G. Donald *et al.*, Phys.Rev. **D86**, 094501 (2012), [arXiv:1208.2855].
- [13] R. A. Briceno, H.-W. Lin and D. R. Bolton, Phys.Rev. **D86**, 094504 (2012), [arXiv:1207.3536].
- [14] C. DeTar *et al.*, PoS **LATTICE2012**, 257 (2012), [arXiv:1211.2253].
- [15] χ QCD collaboration, private communication with Y. Yibo and K.F. Liu.
- [16] D. Mohler, [arXiv:1209.5790].

- [17] D. Becirevic and F. Sanfilippo, JHEP **1301**, 028 (2013), [arXiv:1206.1445].
- [18] PACS-CS Collaboration, Y. Namekawa *et al.*, Phys.Rev. **D84**, 074505 (2011).
- [19] M. Kalinowski and M. Wagner, Acta Physica Polonica B Proceedings Supplement vol. **6**, page 991 (2013), [arXiv:1304.7974].
- [20] BaBar Collaboration, P. del Amo Sanchez *et al.*, Phys.Rev. **D82**, 111101 (2010).
- [21] LHCb, R. Aaij *et al.*, JHEP **1309**, 145 (2013), [arXiv:1307.4556].
- [22] Belle Collaboration, V. Bhardwaj *et al.*, Phys.Rev.Lett. **111**, 032001 (2013), [arXiv:1304.3975].
- [23] G. Bali, S. Collins and P. Perez-Rubio, J.Phys.Conf.Ser. **426**, 012017 (2013), [arXiv:1212.0565].
- [24] G. Bali *et al.*, PoS **LATTICE2011**, 135 (2011), [arXiv:1108.6147].
- [25] D. Mohler and R. Woloshyn, Phys.Rev. **D84**, 054505 (2011), [arXiv:1103.5506].
- [26] D. Mohler, C. B. Lang, L. Leskovec, S. Prelovsek and R. Woloshyn, [arXiv:1308.3175].
- [27] S. Prelovsek *et al.*, to be published in proceeding for Lattice 2013.
- [28] D.-Y. Chen *et al.*, Eur.Phys.J. **C72**, 2226 (2012), [arXiv:1207.3561].
- [29] F.-K. Guo and U.-G. Meissner, Phys.Rev. **D86**, 091501 (2012), [arXiv:1208.1134].
- [30] S. Prelovsek and L. Leskovec, [arXiv:1307.5172].
- [31] S. Prelovsek and L. Leskovec, [arXiv:1308.2097].
- [32] S. Sasaki and T. Yamazaki, Phys.Rev. **D74**, 114507 (2006).
- [33] CDF Collaboration, T. Aaltonen *et al.*, Phys.Rev.Lett. **102**, 242002 (2009).
- [34] CMS Collaboration, S. Chatrchyan *et al.*, [arXiv:1309.6920].
- [35] D0 Collaboration, V. M. Abazov *et al.*, [arXiv:1309.6580].
- [36] S. Ozaki and S. Sasaki, Phys.Rev. **D87**, 014506 (2013), [arXiv:1211.5512].
- [37] HAL QCD Collaboration, N. Ishii *et al.*, Phys.Lett. **B712**, 437 (2012), [arXiv:1203.3642].

- [38] HALQCD collaboration, private communication with Y. Ikeda.
- [39] L. Liu, K. Orginos, F.-K. Guo, C. Hanhart and U.-G. Meissner, Phys.Rev. **D87**, 014508 (2013), [arXiv:1208.4535].
- [40] P. W. M. Evans, C. Allton, P. Giudice and J.-I. Skullerud, [arXiv:1309.3415].