

SOME RESULTS ON THE PENALISED NEMATIC LIQUID CRYSTALS DRIVEN BY MULTIPLICATIVE NOISE

ZDZISLAW BRZEŹNIAK, ERIKA HAUSENBLAS, AND PAUL RAZAFIMANDIMBY

ABSTRACT. In this paper we prove several results related to the existence and uniqueness of solution to coupled highly nonlinear stochastic partial differential equations (PDEs). These equations are motivated by the dynamics of nematic liquid crystals under the influence of stochastic external forces. Firstly, we prove the existence of global weak solution (in sense of both stochastic analysis and PDEs). We show the pathwise uniqueness of the solution in 2D domain. Secondly, we establish the existence and uniqueness of local maximal solution which is strong in sense of both PDEs and stochastic analysis. In the 2D case, we show that this solution is global. In contrast to several works in the deterministic setting we replace the Ginzburg-Landau function $\mathbb{1}_{|\mathbf{n}|\leq 1}(|\mathbf{n}|^2 - 1)\mathbf{n}$ by a general polynomial $f(\mathbf{n})$ and we give sufficient conditions on the polynomial f for the aforementioned results to hold.

As a by-product of our investigation we present a general method based on fixed point argument to establish the existence and uniqueness of a local maximal solution of an abstract stochastic evolution equations with coefficients satisfying local Lipschitz condition involving the norms of two different Banach spaces. This general method can be used to treat several stochastic hydrodynamical models such as Navier-Stokes, Magnetohydrodynamic (MHD) equations, and the α -models of Navier-Stokes equations and their MHD counterparts.

1. INTRODUCTION

Nematic liquid crystal is a state of matter that has properties between amorphous liquid and crystalline solid. Molecules of nematic liquid crystals are long and thin, and they tend to align along a common axis. This preferred axis indicates the orientations of the crystalline molecules, hence it is useful to characterize its orientation with a vector field \mathbf{n} which is called the **director**. Since its magnitude has no significance, we shall take \mathbf{n} as a unit vector. We refer to [14] and [20] for a comprehensive treatment of the physics of liquid crystals. To model the dynamics of nematic liquid crystals most scientists use the continuum theory developed by Ericksen [23] and Leslie [36]. From this theory F. Lin and C. Liu [37] derived the most basic and simplest form of dynamical system describing the motion of nematic liquid crystals filling a bounded region $\mathcal{O} \subset \mathbb{R}^d, d = 2, 3$. This system is given by

$$\mathbf{v}_t + (\mathbf{v} \cdot \nabla)\mathbf{v} - \mu \Delta \mathbf{v} + \nabla p = -\lambda \nabla \cdot (\nabla \mathbf{n} \odot \nabla \mathbf{n}), \text{ in } (0, T] \times \mathcal{O} \quad (1.1)$$

$$\operatorname{div} \mathbf{v} = 0, \text{ in } (0, T] \times \mathcal{O} \quad (1.2)$$

$$\mathbf{n}_t + (\mathbf{v} \cdot \nabla)\mathbf{n} = \gamma(\Delta \mathbf{n} + |\nabla \mathbf{n}|^2 \mathbf{n}), \text{ in } (0, T] \times \mathcal{O} \quad (1.3)$$

$$\mathbf{n}(0) = \mathbf{n}_0, \text{ and } \mathbf{v}(0) = \mathbf{v}_0 \text{ in } \mathcal{O} \quad (1.4)$$

$$|\mathbf{n}|^2 = 1, \text{ on } (0, T] \times \mathcal{O}. \quad (1.5)$$

Here p represents the pressure of the fluid and \mathbf{v} its velocity. By the symbol $\nabla \mathbf{n} \odot \nabla \mathbf{n}$ we mean a $d \times d$ -matrix with entries defined by

$$[\nabla \mathbf{n} \odot \nabla \mathbf{n}]_{i,j} = \sum_{k=1}^n \frac{\partial \mathbf{n}^k}{\partial x_i} \frac{\partial \mathbf{n}^k}{\partial x_j}, \quad i, j = 1, \dots, d.$$

In the present work we assume that the boundary of \mathcal{O} is smooth and that

$$\mathbf{v} = 0 \text{ and } \frac{\partial \mathbf{n}}{\partial \nu} = 0 \text{ on } \partial \mathcal{O}, \quad (1.6)$$

where the vector field ν is the unit outward unit normal to $\partial\mathcal{O}$, i.e. at each point x of \mathcal{O} , $\nu(x)$ is perpendicular to the tangent space $T_x\partial\mathcal{O}$, of length 1 and facing outside of \mathcal{O} .

Although the system (1.1)-(1.6) is the most basic and simplest form of equations from the Ericksen-Leslie continuum theory, it retains the most physical significance of the nematic liquid crystals. Moreover it offers many interesting mathematical problems. In fact, on one hand, two of the main mathematical difficulties related to the system (1.1)-(1.6) are non-parabolicity of equation (1.3) and high nonlinearity of the term $\operatorname{div} \sigma^E = -\operatorname{div} (\nabla \mathbf{n} \odot \nabla \mathbf{n})$. The non-parabolicity follows from the fact that

$$\Delta \mathbf{n} + |\nabla \mathbf{n}|^2 \mathbf{n} = \mathbf{n} \times (\Delta \mathbf{n} \times \mathbf{n}), \quad (1.7)$$

so that the linear term $\Delta \mathbf{n}$ in (1.3) is only a tangential part of the full Laplacian. The term $\operatorname{div} (\nabla \mathbf{n} \odot \nabla \mathbf{n})$ makes the problem (1.1)-(1.6) a fully nonlinear and constrained system of PDEs coupled via a quadratic gradient nonlinearity. In addition to the above difficulties, we will certainly have to deal with new challenging and difficult problems due to the introduction of stochastic perturbations. On the other hand, a number of challenging questions about the solutions to Navier-Stokes equations (NSEs) and Geometric Heat equation (GHE) are still open.

In 1995, F. Lin and C. Liu [37] proposed an approximation of the system (1.1)-(1.6) to relax the constraint $|\mathbf{n}|^2 = 1$ and the gradient nonlinearity $|\nabla \mathbf{n}|^2 \mathbf{n}$. More precisely, they studied the following system of equations

$$\mathbf{v}_t + (\mathbf{v} \cdot \nabla) \mathbf{v} - \mu \Delta \mathbf{v} + \nabla p = -\lambda \nabla \cdot (\nabla \mathbf{n} \odot \nabla \mathbf{n}), \text{ in } (0, T] \times \mathcal{O} \quad (1.8)$$

$$\operatorname{div} \mathbf{v} = 0, \text{ in } [0, T] \times \mathcal{O} \quad (1.9)$$

$$\mathbf{n}(0) = \mathbf{n}_0 \text{ and } \mathbf{v}(0) = \mathbf{v}_0 \text{ in } \mathcal{O}, \quad (1.10)$$

$$\mathbf{n}_t + (\mathbf{v} \cdot \nabla) \mathbf{n} = \gamma \left(\Delta \mathbf{n} - \frac{1}{\varepsilon^2} (|\mathbf{n}|^2 - 1) \mathbf{n} \right) \text{ in } (0, T] \times \mathcal{O}, \quad (1.11)$$

where $\varepsilon > 0$ is an arbitrary constant.

Problem (1.8)-(1.11) with boundary conditions (1.6) is much simpler than (1.1)-(1.5) with (1.6), but it is still a difficult and interesting problem. Since the pioneering work [37] the systems (1.8)-(1.11) and (1.1)-(1.5) have been the subject of intensive mathematical studies. We refer, among others, to [26, 28, 37, 38, 39, 40, 53, 16] and references therein for the relevant results.

In this paper we are interested in the mathematical analysis of a stochastic version of problem (1.8)-(1.11). Basically, we will investigate a system of stochastic evolution equations which is obtained by introducing appropriate noise term in (1.1)-(1.5). In contrast to the unpublished manuscript [9] we replace the bounded Ginzburg-Landau function $\mathbb{1}_{|\mathbf{n}| \leq 1} (|\mathbf{n}|^2 - 1) \mathbf{n}$ in the coupled system by a general polynomial function $f(\mathbf{n})$. More precisely, we set $\mu = \lambda = \gamma = 1$ and we consider cylindrical Wiener processes W_1 on a separable Hilbert space K_1 and a standard real-valued Brownian motion W_2 . We assume that W_1 and W_2 are independent. We consider the problem

$$d\mathbf{v}(t) + \left[(\mathbf{v}(t) \cdot \nabla) \mathbf{v}(t) - \Delta \mathbf{v}(t) + \nabla p \right] dt = -\nabla \cdot (\nabla \mathbf{n}(t) \odot \nabla \mathbf{n}(t)) dt + S(\mathbf{v}(t)) dW_1(t), \quad (1.12)$$

$$\operatorname{div} \mathbf{v}(t) = 0, \quad (1.13)$$

$$d\mathbf{n}(t) + (\mathbf{v}(t) \cdot \nabla) \mathbf{n}(t) dt = \left[\Delta \mathbf{n}(t) - f(\mathbf{n}) \right] + (\mathbf{n}(t) \times h) \circ dW_2(t), \quad (1.14)$$

where $(\mathbf{n}(t) \times h) \circ dW_2(t)$ is understood in the Stratonovich sense and f is a polynomial function and the above system holds in $\mathcal{O}_T := (0, T] \times \mathcal{O}$. We equip the system with the boundary condition (1.6) and the initial condition

$$\mathbf{v}(0) = \mathbf{v}_0 \text{ and } \mathbf{n}(0) = \mathbf{n}_0,$$

where \mathbf{v}_0 and \mathbf{n}_0 are given mappings defined on \mathcal{O} . We will give more details about the polynomial f later on.

Our work is motivated by the importance of external perturbation on the dynamics of the director field \mathbf{n} . Indeed, an essential property of nematic liquid crystals is that its director field \mathbf{n} can be

easily distorted. However, it can also be aligned to form a specific pattern under some external perturbations. This pattern formation occurs when a threshold value of the external perturbations is attained; this is the so called Fréedericksz transition. Random external perturbations change a little bit the threshold value for the Fréedericksz transition. For example, it has been found that with the fluctuation of the magnetic field the relaxation time of an unstable state diminishes, *i.e.*, the time for a noisy system to leave an unstable state is much shorter than the unperturbed system. For these results we refer, among others, to [29, 49, 50] and references therein. In all of these works the effect of the hydrodynamic flow has been neglected. However, it is pointed out in [20, Chapter 5] that the fluid flow disturbs the alignment and conversely a change in the alignment will induce a flow in the nematic liquid crystal. Hence for a full understanding of the effect of fluctuating magnetic field on the behavior of the liquid crystals one needs to take into account the dynamics of \mathbf{n} and \mathbf{v} . To initiate this kind of investigation we propose a mathematical study of (1.12)-(1.14) which basically describes an approximation of the system governing the nematic liquid crystals under the influence of fluctuating external forces.

In the present paper we prove several results that are basically the stochastic counterparts of the ones obtained by Lin and Liu in [37]. Our results can be described as follows.

- (i) In section 3 we establish the existence of global martingale solutions (weak in the PDEs sense).

To prove this result we first find a suitable finite dimensional Galerkin approximation of system (1.12)-(1.14) which can be solved locally in time. Our choice of the approximation yields the global existence of the approximating solutions $(\mathbf{v}_m, \mathbf{n}_m)$. For this purpose we derive several important global a priori estimates in higher order Sobolev spaces involving the following two energy functionals

$$\mathcal{E}_1(\mathbf{n}, t) := \|\mathbf{n}(t)\|^q + q \int_0^t \|\mathbf{n}(s)\|^{q-2} \|\nabla \mathbf{n}(s)\|^2 ds + q \int_0^t \|\mathbf{n}(s)\|^{q-2} \|\mathbf{n}(s)\|_{\mathbb{H}^{2N+2}}^{2N+2} ds$$

and

$$\begin{aligned} \mathcal{E}_2(\mathbf{v}, \mathbf{n}, t) &:= \|\mathbf{v}(t)\|^2 + \ell \|\mathbf{n}(t)\|^2 + \|\nabla \mathbf{n}(t)\|^2 + \int_{\mathcal{O}} F(\mathbf{n}(t, x)) dx \\ &\quad + \left(\int_0^t \|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 \right) ds. \end{aligned}$$

Here $F(\cdot)$ is the antiderivative of f such that $F(0) = 0$. These global a priori estimates, the proofs of which are non-trivial and require long and tedious calculation, are very crucial for the proof of the tightness of the distributions of the Galerkin solution $(\mathbf{v}_m, \mathbf{n}_m)$ in appropriate topological spaces such as $L^2(0, T; \mathbb{H}) \times L^2(0, T; \mathbb{H}^1(\mathcal{O}))$. This compactness result will enable us to construct a new probability space on which we also find a new sequence $(\bar{\mathbf{v}}_m, \bar{\mathbf{n}}_m, \bar{W}_1^m, \bar{W}_2^m)$ of solutions of the Galerkin equations. This new sequence is proved to converge to a system $(\mathbf{v}, \mathbf{n}, \bar{W}_1, \bar{W}_2)$ which along with the new probability space will form our martingale solution. To close the first part of our results we show that the weak solution is pathwise unique in the 2-D case.

- (ii) We establish several results about the strong solution to the problem (1.12)-(1.14) in Section 4. We mainly prove the existence of a local and maximal strong solution which is understood in the sense of stochastic calculus and PDEs. In the case $d = 2$ we prove the non-explosion of the maximal solution by a method based on a choice of an appropriate energy functional (or Lyapunov function) and an idea introduced by Schmalfußin [51] to prove the pathwise uniqueness of solution of stochastic Navier-Stokes equations. The estimates for the energy functional are of the form

$$\begin{aligned} \mathcal{E}_3 &= \mathbb{E} \left[\Phi(t \wedge \tau_k) (\|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k))) \right] \\ &\quad + \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \left(2\|\Delta \mathbf{v}(s)\|^2 + \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2 \right) ds, \end{aligned}$$

where $\Phi(\cdot)$ is a certain positive function to be defined later and $\{\tau_k; k = 1, 2, \dots\}$ is a sequence of stopping times, see Section 4 for more details. Schmalfuß trick has so far been used to prove pathwise uniqueness of solution of stochastic PDEs, see for instance [51], [13], [8] and references therein. Our novelty is that we extend the use of this idea to prove the global existence of strong solution of the Ginzburg-Landau approximation of nematic liquid crystals. In particular, our proof gives another proof of the global existence of 2D stochastic Navier-Stokes equations with multiplicative for initial data with finite enstrophy. Thus, our paper can be seen also as a generalization of the results for the existence and uniqueness of local, maximal and global solutions of strong solutions of stochastic Navier-Stokes proved in [25], [19] and [42].

- (iii) In Section 5 we use a fixed point argument and prove several general results on an abstract nonlinear stochastic evolution equations (SEE) with coefficients satisfying local Lipschitz condition involving the norms of two different Banach spaces. We mainly establish the existence and uniqueness of a local and a maximal solution to this SEE and derive a lower estimate for the life-span of the solution. These general results are of independent interest and that is why they are the subject of a separate section. The existence and uniqueness result in Section 4 are corollaries of the general results in the last section. In fact, in Section 4 we consider some functional spaces with more regularity than in the case of weak and martingale solution and show that on these spaces the problem (1.12)-(1.14) with (1.6) falls within the general framework in Section 5. We are also strongly convinced that with these general results it is possible, although has not been done in detail, to prove the existence of strong solution of several stochastic hydrodynamical models such as the NSEs, Magnetohydrodynamic (MHD) equations, α -models for Navier-Stokes (Lagrangian Navier-Stokes α -models, Leray- α models) and the MHD equations associated to them.
- (iv) We prove a maximum principle type theorem in Section 6. More precisely, if we consider $f(\mathbf{n}) = \mathbb{1}_{|\mathbf{n}| \leq 1}(|\mathbf{n}|^2 - 1)\mathbf{n}$ instead and if the initial condition \mathbf{n}_0 satisfies $|\mathbf{n}_0| \leq 1$ almost everywhere, then the solution \mathbf{n} also remains in the unit ball almost everywhere. In contrast to the deterministic case, this result does not follow in straightforward way from well-known results. Here the method of proofs are based on the blending of ideas from [15] and [21].

To the best of our knowledge our work is the first mathematical work which studies the existence and uniqueness of weak martingale of the (1.12)-(1.14). Under the assumption that $f(\cdot)$ is a bounded function, the authors proved in the unpublished manuscript [9] that the system (1.12)-(1.14) has a maximal strong solution which is global for the 2D case. Therefore, the present article is a generalization of [9] in the sense that we allow $f(\cdot)$ to be a general polynomial function. Note that some of the arguments from the last part of the present paper have already been used in the paper [4] which studied the strong solution of some stochastic hydrodynamic equations driven by Lévy noise.

The organization of the present article is as follows. In Section 2 we introduce the notations that are frequently used throughout this paper. In the same section we also state and prove some useful lemmata. By using the scheme we outlined in item (i) above we show in Section 3 that (1.12)-(1.14) admits a weak martingale solution which is pathwise unique in the two dimensional case. In Section 4 we prove the existence of a unique maximal strong solution to (1.12)-(1.14) which turns out to be a global solution when the space dimension is two. The existence and uniqueness of a maximal solution to the stochastic system (1.12)-(1.14) is a corollary of a general theorem given in Section 5. This result is obtained by making use of some fixed point argument and truncation method. In Section 6 a maximum principle type theorem is proved when $f(\mathbf{n}) = \mathbb{1}_{|\mathbf{n}| \leq 1}(|\mathbf{n}|^2 - 1)\mathbf{n}$. In Appendix we prove several crucial estimates about the solution of the stochastic equation for nematic liquid crystals.

2. FUNCTIONAL SPACES AND PREPARATORY LEMMA

Let $d \in \{2, 3\}$ and assume that $\mathcal{O} \subset \mathbb{R}^d$ is a bounded domain with boundary $\partial\mathcal{O}$ of class \mathcal{C}^∞ . For any $p \in [1, \infty)$ and $k \in \mathbb{N}$, $\mathbb{L}^p(\mathcal{O})$ and $\mathbb{W}^{k,p}(\mathcal{O})$ are the well-known Lebesgue and Sobolev spaces,

respectively, of \mathbb{R}^d -valued functions. The corresponding spaces of scalar functions we will denote by standard letter, e.g. $W^{k,p}(\mathcal{O})$. For $p = 2$ we denote $W^{k,2}(\mathcal{O}) = \mathbb{H}^k$ and its norm are denoted by $\|\mathbf{u}\|_k$. By \mathbb{H}_0^1 we mean the space of functions in \mathbb{H}^1 that vanish on the boundary on \mathcal{O} ; \mathbb{H}_0^1 is a Hilbert space when endowed with the scalar product induced by that of \mathbb{H}^1 . The usual scalar product on \mathbb{L}^2 is denoted by $\langle u, v \rangle$ for $u, v \in \mathbb{L}^2$ and its associated norm is denoted by $\|u\|$, $u \in \mathbb{L}^2$. We also introduce the following spaces

$$\begin{aligned}\mathcal{V} &= \left\{ \mathbf{u} \in \mathcal{C}_c^\infty(\mathcal{O}, \mathbb{R}^d) \text{ such that } \nabla \cdot \mathbf{u} = 0 \right\} \\ \mathbb{V} &= \text{closure of } \mathcal{V} \text{ in } \mathbb{H}_0^1(\mathcal{O}) \\ \mathbb{H} &= \text{closure of } \mathcal{V} \text{ in } \mathbb{L}^2(\mathcal{O}).\end{aligned}$$

We endow \mathbb{H} with the scalar product and norm of \mathbb{L}^2 . As usual we equip the space \mathbb{V} with the the scalar product $\langle \nabla \mathbf{u}, \nabla \mathbf{v} \rangle$ which, owing to the Poincaré inequality, is equivalent to the $\mathbb{H}^1(\mathcal{O})$ -scalar product.

Let $\Pi : \mathbb{L}^2 \rightarrow \mathbb{H}$ be the Helmholtz-Leray projection from \mathbb{L}^2 onto \mathbb{H} . We denote by $A = -\Pi\Delta$ the Stokes operator with domain $D(A)$. It is well-known (see for e.g. [57, Chapter I, Section 2.6]) that there exists an orthonormal basis $\{\varphi_i; i = 1, 2, 3, \dots\}$ of \mathbb{H} consisting of the eigenfunctions of the Stokes operator A .

From [45, Proposition 1.24] we can define a self-adjoint operator $A_1 : \mathbb{H}^1 \rightarrow (\mathbb{H}^1)^*$ by

$$\langle A_1 \mathbf{u}, \mathbf{w} \rangle = a(\mathbf{u}, \mathbf{w}) = \int_{\mathcal{O}} \nabla \mathbf{u} \cdot \nabla \mathbf{w} \, dx, \quad \mathbf{u}, \mathbf{w} \in \mathbb{H}^1. \quad (2.1)$$

The Neumann Laplacian acting on \mathbb{R}^n -valued function will be denoted by A_1 , that is,

$$\begin{aligned}D(A_1) &:= \left\{ \mathbf{u} \in \mathbb{H}^2 : \frac{\partial \mathbf{u}}{\partial \nu} = 0 \text{ on } \partial \mathcal{O} \right\}, \\ A_1 \mathbf{u} &:= - \sum_{i=1}^n \frac{\partial^2 \mathbf{u}}{\partial x_i^2}, \quad \mathbf{u} \in D(A_1).\end{aligned} \quad (2.2)$$

It can be shown that $D(A_1) = \{\mathbf{u} \in \mathbb{H}^1 : A_1 \mathbf{u} \in \mathbb{L}^2\}$ and $A_1 \mathbf{u} = A_1 \mathbf{u}$ for $\mathbf{u} \in D(A_1)$. It can also be shown, see e.g. [27, Theorem 5.31], that $\hat{A}_1 = I + A_1$ is a definite positive and self-adjoint operator in the Hilbert space $\mathbb{L}^2 = \mathbb{L}^2(\mathcal{O})$ with compact resolvent. In particular, there exists an ONB $(\phi_k)_{k=1}^\infty$ of \mathbb{L}^2 and an increasing sequence $(\lambda_k)_{k=1}^\infty$ with $\lambda_1 = 0$ and $\lambda_k \nearrow \infty$ as $k \nearrow \infty$ (the eigenvalues of the Neumann Laplacian A_1) such that $A_1 \phi_j = \lambda_j \phi_j$ for any $j \in \mathbb{N}$.

For any $\alpha \in [-\frac{1}{2}, \infty)$ we denote by $\mathbb{X}_\alpha = D(\hat{A}_1^{\frac{1}{2}+\alpha})$, the domain of the fractional power operator $\hat{A}_1^{\frac{1}{2}+\alpha}$. We have the following characterization of the spaces \mathbb{X}_α ,

$$\mathbb{X}_\alpha = \left\{ \mathbf{u} = \sum_{k \in \mathbb{N}} u_k \phi_k : \sum_k (1 + \lambda_k)^{1+2\alpha} |u_k|^2 < \infty \right\}. \quad (2.3)$$

It can be shown that $\mathbb{X}_\alpha \subset \mathbb{H}^{1+2\alpha}$, for all $\alpha \geq 0$ and $\mathbb{X} := \mathbb{X}_0 = \mathbb{H}^1$.

Similarly, for $\beta \in [0, \infty)$, we denote by \mathbb{V}_β the Hilbert space $D(A^\beta)$ endowed with the graph inner product. The Hilbert space $\mathbb{V}_\beta = D(A^\beta)$ for $\beta \in (-\infty, 0)$ can be defined by standard extrapolation methods. In particular, the space $D(A^{-\beta})$ is the dual of \mathbb{V}_β for $\beta \geq 0$. Moreover, for every $\beta, \delta \in \mathbb{R}$ the map A^δ is a linear isomorphism between \mathbb{V}_β and $\mathbb{V}_{\beta-\delta}$. It is also well-known that $\mathbb{V}_{\frac{1}{2}} = \mathbb{V}$.

Throughout this paper \mathbf{B}^* denotes the dual space of a Banach space \mathbf{B} . We denote by $\langle \Psi, \mathbf{b} \rangle$ the value of $\Psi \in \mathbf{B}^*$ on $\mathbf{b} \in \mathbf{B}$.

Given two Hilbert spaces K and H , we denote by $\mathcal{L}(K, H)$ and $\mathcal{T}_2(K, H)$ the space of bounded linear operators and the Hilbert space of all Hilbert-Schmidt operators from K to H , respectively. For $K = H$ we just write $\mathcal{L}(K)$ instead of $\mathcal{L}(K, K)$.

Now, we introduce a trilinear form

$$b(\mathbf{u}, \mathbf{v}, \mathbf{w}) = \sum_{i,j=1}^d \int_{\mathcal{O}} \mathbf{u}^{(i)} \frac{\partial \mathbf{v}^{(j)}}{\partial x_i} \mathbf{w}^{(j)} dx, \quad \mathbf{u} \in \mathbb{L}^p, \mathbf{v} \in \mathbb{W}^{1,q}, \text{ and } \mathbf{w} \in \mathbb{L}^r,$$

with numbers $p, q, r \in [1, \infty]$ satisfying

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} \leq 1.$$

The map b is the trilinear form used in the mathematical analysis of the Navier-Stokes equations, see for instance [57, Chapter II, Section 1.2]. It is well known (see for e.g. [57, Chapter II, Section 1.2]) that one can define a bilinear map \tilde{B} defined on $\mathbb{H}^1 \times \mathbb{H}^1$ with values in $(\mathbb{H}^1)^*$ such that

$$\langle \tilde{B}(\mathbf{u}, \mathbf{v}), \mathbf{w} \rangle = b(\mathbf{u}, \mathbf{v}, \mathbf{w}) \quad \text{for any } \mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{H}^1. \quad (2.4)$$

We can also define a bilinear map B from $\mathbb{V} \times \mathbb{V}$ with values in \mathbb{V}^* such that

$$\langle B(\mathbf{u}, \mathbf{v}), \mathbf{w} \rangle = b(\mathbf{u}, \mathbf{v}, \mathbf{w}) \quad \text{for } \mathbf{w} \in \mathbb{V}, \text{ and } \mathbf{u}, \mathbf{v} \in \mathbb{H}^1. \quad (2.5)$$

Well-known properties of B and \tilde{B} will be given in Appendix.

Let \mathfrak{m} be the trilinear form defined by

$$\mathfrak{m}(\mathbf{n}_1, \mathbf{n}_2, \mathbf{u}) = - \sum_{i,j,k=1}^d \int_{\mathcal{O}} \partial_{x_i} \mathbf{n}_1^{(k)} \partial_{x_j} \mathbf{n}_2^{(k)} \partial_{x_j} \mathbf{u}^{(i)} dx \quad (2.6)$$

for any $\mathbf{n}_1 \in W_1^{1,p}$, $\mathbf{n}_2 \in W_1^{1,q}$ and $\mathbf{u} \in W_1^{1,r}$ with $r, p, q \in (1, \infty)$ satisfying

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} \leq 1.$$

Here $\partial_{x_i} = \frac{\partial}{\partial x_i}$ and ϕ^i is the i -th entry of any vector-valued ϕ . Since $n \leq 4$, the integral in (2.6) is well defined for When $\mathbf{n}_1, \mathbf{n}_2 \in \mathbb{H}^2$ and $\mathbf{u} \in \mathbb{V}$. We have the following lemma.

Lemma 2.1. *There exist a constant $C > 0$ such that*

$$|\mathfrak{m}(\mathbf{n}_1, \mathbf{n}_2, \mathbf{u})| \leq C \|\nabla \mathbf{n}_1\|^{1-\frac{d}{4}} \|\Delta \mathbf{n}_1\|^{\frac{d}{4}} \|\nabla \mathbf{n}_2\|^{1-\frac{d}{4}} \|\Delta \mathbf{n}_2\|^{\frac{d}{4}} \|\nabla \mathbf{u}\|, \quad (2.7)$$

for any $\mathbf{n}_1, \mathbf{n}_2 \in \mathbb{H}^2$ and $\mathbf{u} \in \mathbb{V}$.

Proof of Lemma 2.1. From (2.6) and Hölder's inequality we derive that

$$|\mathfrak{m}(\mathbf{n}_1, \mathbf{n}_2, \mathbf{u})| \leq \int_{\mathcal{O}} |\nabla \mathbf{n}_1| |\nabla \mathbf{n}_2| |\nabla \mathbf{u}| dx.$$

The above integral is well-defined since $\nabla \mathbf{n}_i \in \mathbb{L}^{\frac{2d}{d-2}}$, $i = 1, 2$, $\nabla \mathbf{u} \in \mathbb{L}^2$ and $\frac{d-2}{d} + \frac{1}{2} \leq 1$ for $d \leq 4$. When $d = 2$ we replace $2d/(d-2)$ by any $q \in [4, \infty)$. Note that for $d \leq 4$ we have $|\nabla \mathbf{n}_i| \in \mathbb{L}^4$, $i = 1, 2$. Hence

$$|\mathfrak{m}(\mathbf{n}_1, \mathbf{n}_2, \mathbf{u})| \leq C \|\nabla \mathbf{n}_1\|_{\mathbb{L}^4} \|\nabla \mathbf{n}_2\|_{\mathbb{L}^4} \|\nabla \mathbf{u}\|.$$

This last estimate and Gagliardo-Nirenberg's inequality (A.1) lead us to

$$|\mathfrak{m}(\mathbf{n}_1, \mathbf{n}_2, \mathbf{u})| \leq C \|\nabla \mathbf{n}_1\|^{1-\frac{d}{4}} \|\Delta \mathbf{n}_1\|^{\frac{d}{4}} \|\nabla \mathbf{n}_2\|^{1-\frac{d}{4}} \|\Delta \mathbf{n}_2\|^{\frac{d}{4}} \|\nabla \mathbf{u}\|. \quad (2.8)$$

This concludes the proof of our claim. \square

The above result tells us that, for any $\mathbf{u} \in \mathbb{V}$, $\mathfrak{m}(\mathbf{n}_1, \mathbf{n}_2, \mathbf{u})$ is an element of $\mathcal{L}(\mathbb{V}, \mathbb{R})$ whenever $\mathbf{n}_1, \mathbf{n}_2 \in \mathbb{H}^2$. Now, we state and prove the following proposition.

Proposition 2.2. *There exists a bilinear operator M defined on $\mathbb{H}^2 \times \mathbb{H}^2$ taking values on \mathbb{V}^* such that for any $\mathbf{n}_1, \mathbf{n}_2 \in \mathbb{H}^2$*

$$\langle M(\mathbf{n}_1, \mathbf{n}_2), \mathbf{u} \rangle = \mathbf{m}(\mathbf{n}_1, \mathbf{n}_2, \mathbf{u}) \quad \mathbf{u} \in \mathbb{V}. \quad (2.9)$$

Furthermore, there exists a constant $C > 0$ such that

$$\|M(\mathbf{n}_1, \mathbf{n}_2)\|_{\mathbb{V}^*} \leq C \|\nabla \mathbf{n}_1\|^{1-\frac{d}{4}} \|\Delta \mathbf{n}_1\|^{\frac{d}{4}} \|\nabla \mathbf{n}_2\|^{1-\frac{d}{4}} \|\Delta \mathbf{n}_2\|^{\frac{d}{4}}, \quad (2.10)$$

for any $\mathbf{n}_1, \mathbf{n}_2 \in \mathbb{H}^2$. We also have the following identity

$$\langle \tilde{B}(\mathbf{v}, \mathbf{n}), \Delta \mathbf{n} \rangle = \langle M(\mathbf{n}, \mathbf{n}), \mathbf{v} \rangle, \text{ for any } \mathbf{v} \in \mathbb{V}, \mathbf{n} \in \mathbb{H}^2. \quad (2.11)$$

Proof. The first part and (2.10) follow from Lemma 2.1.

To prove (2.11) we first note that $\langle \tilde{B}(\mathbf{v}, \mathbf{n}_2), \Delta \mathbf{n}_1 \rangle = b(\mathbf{v}, \mathbf{n}_2, \Delta \mathbf{n}_1)$ is well-defined for any $\mathbf{v} \in \mathbb{V}, \mathbf{n}_1, \mathbf{n}_2 \in \mathbb{H}^2$. Thus, taking into account that \mathbf{v} is divergence free and vanishes on the boundary we can perform an integration-by-parts and deduce that

$$\begin{aligned} \langle B(\mathbf{v}, \mathbf{n}), \Delta \mathbf{n} \rangle &= \int_{\mathcal{O}} \mathbf{v}^{(i)} \frac{\partial \mathbf{n}^{(k)}}{\partial x_i} \frac{\partial^2 \mathbf{n}^{(k)}}{\partial x_i \partial x_l} dx \\ &= - \int_{\mathcal{O}} \frac{\partial \mathbf{v}^{(i)}}{\partial x_l} \frac{\partial \mathbf{n}^{(k)}}{\partial x_i} \frac{\partial \mathbf{n}^{(k)}}{\partial x_l} dx - \int_{\mathcal{O}} \mathbf{v}^{(i)} \frac{\partial^2 \mathbf{n}^{(k)}}{\partial x_i \partial x_l} \frac{\partial \mathbf{n}^{(k)}}{\partial x_l} dx \\ &= - \int_{\mathcal{O}} \frac{\partial \mathbf{v}^{(i)}}{\partial x_l} \frac{\partial \mathbf{n}^{(k)}}{\partial x_i} \frac{\partial \mathbf{n}^{(k)}}{\partial x_l} dx - \frac{1}{2} \int_{\mathcal{O}} \mathbf{v}^{(i)} \frac{\partial |\nabla \mathbf{n}|^2}{\partial x_i} dx \\ &= - \int_{\mathcal{O}} \frac{\partial \mathbf{v}^{(i)}}{\partial x_l} \frac{\partial \mathbf{n}^{(k)}}{\partial x_i} \frac{\partial \mathbf{n}^{(k)}}{\partial x_l} dx \\ &= \mathbf{m}(\mathbf{n}, \mathbf{n}, \mathbf{v}) = \langle M(\mathbf{n}, \mathbf{n}), \mathbf{v} \rangle. \end{aligned}$$

In the above chain of equalities summation over repeated indexes are enforced. \square

Remark 2.3. (1) We have $\nabla \cdot (\nabla \mathbf{f} \odot \nabla \mathbf{g}) \in \mathbb{L}^2$ for any $\mathbf{f}, \mathbf{g} \in \mathbb{X}_{\frac{1}{2}} \cap \mathbb{X}_1$. Thus, we infer that for any $\mathbf{f}, \mathbf{g} \in \mathbb{X}_{\frac{1}{2}} \cap \mathbb{X}_1$

$$M(\mathbf{f}, \mathbf{g}) = \Pi[\nabla \cdot (\nabla \mathbf{f} \odot \nabla \mathbf{g})].$$

(2) In some places in this manuscript we use the following shorthand notation:

$$B(\mathbf{u}) := B(\mathbf{u}, \mathbf{u}) \text{ and } M(\mathbf{n}) := M(\mathbf{n}, \mathbf{n}),$$

for any \mathbf{u} and \mathbf{n} such that the above quantities are meaningful.

We assume that $h \in \mathbb{L}^\infty$ is fixed. We can define a linear bounded operator G from \mathbb{L}^2 into itself by

$$G : \mathbb{L}^2 \ni \mathbf{n} \mapsto \mathbf{n} \times h \in \mathbb{L}^2.$$

It is straightforward to check that satisfies

$$\|G(\mathbf{n})\| \leq \|h\|_{\mathbb{L}^\infty} \|\mathbf{n}\|.$$

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space equipped with a filtration $\mathbb{F} = \{\mathcal{F}_t : t \geq 0\}$ satisfying the usual condition. Let $W_2 = (W_2(t))_{t \geq 0}$ be a standard \mathbb{R} -valued Wiener process on $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$. Let us also assume that K_1 is a separable Hilbert space and $W_1 = (W_1(t))_{t \geq 0}$ is a K_1 -cylindrical Wiener process on $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$. Throughout we assume that W_2 and W_1 are independent. Thus we can assume that $W = (W_1(t), W_2(t))$ is K -cylindrical Wiener process on $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, where

$$K = K_1 \times \mathbb{R}.$$

Remark 2.4. If K_2 is a Hilbert space such that the embedding $K_1 \subset K_2$ is Hilbert-Schmidt, then W_1 can be viewed as an K_2 -valued Wiener process. Moreover, there exists a trace class symmetric

nonnegative operator $Q \in \mathcal{L}(K_2)$ such that W_1 has covariance Q . This K_2 -valued K_1 -cylindrical Wiener process is characterised by, for all $t \geq 0$,

$$\mathbb{E} e^{i_{K_2^*} \langle x^*, W(t) \rangle_{K_2}} = e^{-\frac{t}{2} \|x^*\|_{K_1}^2}, \quad x^* \in K_2^*,$$

where K_2^* is the dual space to K_2 such that identifying K_1^* with K_1 we have

$$K_2^* \hookrightarrow K_1^* = K_1 \hookrightarrow K_2.$$

We have the following relation between Stratonovich and Itô's integrals

$$G(\mathbf{n}) \circ dW_2 = \frac{1}{2} G^2(\mathbf{n}) dt + G(\mathbf{n}) dW_2,$$

where $G^2 = G \circ G$ and defined by

$$G^2(\mathbf{n}) = G \circ G(\mathbf{n}) = (\mathbf{n} \times h) \times h, \text{ for any } \mathbf{n} \in \mathbb{L}^2.$$

Now, we rewrite the problem (1.12)-(1.14) in the following abstract form

$$d\mathbf{v}(t) + \left(A\mathbf{v}(t) + B(\mathbf{v}(t), \mathbf{v}(t)) + M(\mathbf{n}(t)) \right) dt = S(\mathbf{v}(t)) dW_1, \quad (2.12)$$

$$d\mathbf{n}(t) + \left(A_1\mathbf{n}(t) + \tilde{B}(\mathbf{v}(t), \mathbf{n}(t)) + f(\mathbf{n}(t)) - \frac{1}{2} G^2(\mathbf{n}(t)) \right) dt = G(\mathbf{n}(t)) dW_2. \quad (2.13)$$

Throughout this paper we impose that the function $f(\cdot)$ satisfies the following set of conditions.

Assumption 2.1. Let I_d be the set defined by

$$I_d = \begin{cases} \mathbb{N} := \{1, 2, 3, \dots\} & \text{if } d = 2, \\ \{1\}, & \text{if } d = 3. \end{cases} \quad (2.14)$$

Throughout this paper we fix $N \in I_d$ and a family of numbers a_k , $k = 0, \dots, N$, with $a_N > 0$. We define a function $\tilde{f} : [0, \infty) \rightarrow \mathbb{R}$ by

$$\tilde{f}(r) = \sum_{k=0}^N a_k r^k, \text{ for any } r \in \mathbb{R}_+.$$

We define a map $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by $f(\mathbf{n}) = \tilde{f}(|\mathbf{n}|^2) \mathbf{n}$ where \tilde{f} is as above.

Let $F : \mathbb{R}^d \rightarrow \mathbb{R}$ be a Fréchet differentiable map such that for any $\mathbf{n} \in \mathbb{R}^d$ and $\mathbf{g} \in \mathbb{R}^d$

$$F'(\mathbf{n})[\mathbf{g}] = f(\mathbf{n}) \cdot \mathbf{g}.$$

Let also \tilde{F} be an antiderivative of \tilde{f} such that $\tilde{F}(0) = 0$. We have

$$\tilde{F}(r) = a_{N+1} r^{N+1} + U(r),$$

where U is a polynomial function of at most degree N and $a_{N+1} > 0$.

Remark 2.5. For any $r \in [0, \infty)$ let $\tilde{f}(r) := r - 1$. Since $1 \in I_d$ then the maps f and F defined on \mathbb{R}^d by $f(\mathbf{n}) := \tilde{f}(|\mathbf{n}|^2) \mathbf{n}$ and $F(\mathbf{n}) := \frac{1}{4} [\tilde{F}(|\mathbf{n}|^2)]^2$ for any $\mathbf{n} \in \mathbb{R}^d$ satisfy the above set of assumptions.

Remark 2.6. We have the following facts

$$|\tilde{f}(r)| \leq \ell_1 (1 + r^N), \quad r > 0, \quad (2.15)$$

$$|\tilde{f}'(r)| \leq \ell_2 (1 + r^{N_1}), \quad r > 0. \quad (2.16)$$

Remark 2.7. Let f be defined in Assumption 2.1. Then, there exist two positive constants $c > 0$ and $\tilde{c} > 0$ such that

$$|f(\mathbf{n})| \leq c(1 + |\mathbf{n}|^{2N+1}) \text{ and } |f'(\mathbf{n})| \leq \tilde{c}(1 + |\mathbf{n}|^{2N}) \text{ for any } \mathbf{n} \in \mathbb{R}^n.$$

Also, by performing elementary calculation we can check that for any $\mathbf{n} \in \mathbb{H}^2$

$$\begin{aligned} \|\Delta \mathbf{n}\|^2 &= \|\Delta \mathbf{n} - f(\mathbf{n}) + f(\mathbf{n})\|^2 \leq 2\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + 2\|f(\mathbf{n})\|^2, \\ &\leq 2\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + C\|\mathbf{n}\|_{\mathbb{L}^{\tilde{q}}}^{\tilde{q}} + C, \end{aligned} \quad (2.17)$$

where $\tilde{q} = 4N + 2$.

Hence there exists a constant $C > 0$ such that

$$\|\mathbf{n}\|_2^2 \leq C(\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{L}^{\tilde{q}}}^{\tilde{q}} + 1), \text{ for any } \mathbf{n} \in \mathbb{H}^2(\mathcal{O}). \quad (2.18)$$

3. EXISTENCE AND UNIQUENESS OF WEAK MARTINGALE SOLUTION

In this section we are going to establish the existence of weak martingale solution of the system (2.12)-(2.13). For this purpose we use the Galerkin approximation to reduce the original system to a system of finite dimensional ordinary stochastic differential equations (SDEs for short). We establish several crucial uniform a priori estimates which will be used to prove the tightness of the family of laws of the sequence of solutions of the system of SDEs on appropriate topological spaces. But before we proceed further we define what we mean by weak martingale solution.

Definition 3.1. Let K_1 be as in Remark 2.4. By a weak martingale solution to (2.12)-(2.13) we mean a system consisting of a complete and filtered probability space

$$(\Omega', \mathcal{F}', \mathbb{P}', \mathbb{F}'),$$

with the filtration $\mathbb{F}' = (\mathcal{F}'_t)_{t \in [0, T]}$ satisfying the usual condition, and progressively measurable stochastic processes

$$(\mathbf{v}(t), \mathbf{n}(t), \bar{W}_1(t), \bar{W}_2(t))_{t \in [0, T]}$$

such that:

- (1) $(\bar{W}_1(t))_{t \in [0, T]}$ (resp. $(\bar{W}_2(t))_{t \in [0, T]}$) is \mathbb{F}' -progressively measurable and K_1 -cylindrical (resp. real-valued) Wiener process,
- (2) $(\mathbf{v}, \mathbf{n}) : [0, T] \times \Omega' \rightarrow \mathbb{V} \times \mathbb{H}^2$ and \mathbb{P}' -a.e. $\omega' \in \Omega'$

$$(\mathbf{v}, \mathbf{n})(\cdot, \omega') \in C([0, T]; \mathbb{V}^{-\beta}) \times C([0, T]; \mathbb{X}_{\beta - \frac{1}{2}}), \text{ for any } \beta \in (0, \frac{1}{2}), \quad (3.1)$$

$$\mathbb{E}' \sup_{0 \leq s \leq T} [\|\mathbf{v}(s, \omega')\| + \|\nabla \mathbf{n}(s, \omega')\|] + \mathbb{E}' \int_0^T (\|\nabla \mathbf{v}(s, \omega')\|^2 + \|\Delta \mathbf{n}(s, \omega')\|^2) ds < \infty, \quad (3.2)$$

- (3) for each $(\Phi, \Psi) \in \mathbb{V} \times \mathbb{L}^2$ we have, for all $t \in [0, T]$, \mathbb{P}' -a.s..

$$\begin{aligned} \langle \mathbf{v}(t) - \mathbf{v}_0, \Phi \rangle + \int_0^t \left\langle A\mathbf{v}(s) + B(\mathbf{v}(s), \mathbf{v}(s)) + M(\mathbf{n}(s)), \Phi \right\rangle ds \\ = \int_0^t \langle S(\mathbf{v}(s)), \Phi \rangle dW_1(s), \end{aligned} \quad (3.3)$$

and

$$\begin{aligned} \langle \mathbf{n}(t) - \mathbf{n}_0, \Psi \rangle + \int_0^t \left\langle A_1 \mathbf{n}(s) + \tilde{B}(\mathbf{v}(s), \mathbf{n}(s)) + f(\mathbf{n}(s)) - \frac{1}{2} G^2(\mathbf{n}(s)), \Psi \right\rangle ds \\ = \int_0^t \langle G(\mathbf{n}(s)), \Psi \rangle dW_2(s). \end{aligned} \quad (3.4)$$

The function S is defined in the next set of hypotheses.

Assumption 3.1. We assume that $S : \mathbb{H} \rightarrow \mathcal{T}_2(K_1, \mathbb{H})$ is a globally Lipschitz map. In particular, there exists $\ell_3 \geq 0$ such that

$$\|S(\mathbf{u})\|_{\mathcal{T}_2}^2 := \|S(\mathbf{u})\|_{\mathcal{T}_2(K_1, \mathbb{H})}^2 \leq \ell_3(1 + \|\mathbf{u}\|^2), \text{ for any } \mathbf{u} \in \mathbb{H}. \quad (3.5)$$

Now we can state our first result in the following theorem.

Theorem 3.2. *If Assumption 3.1 and 2.1 are satisfied, $h \in W^{1,3} \cap L^\infty$, $\mathbf{v}_0 \in \mathbb{H}$, $\mathbf{n}_0 \in \mathbb{H}^1$, and $d = 2, 3$. Then, the system (2.12)-(2.13) has a weak martingale solution in the sense of Definition 3.1.*

Proof. The proof relies on Galerkin approximation and compactness method and is the subject of the subsections 3.1-3.3. \square

Before we state the uniqueness of the weak martingale solution we should make the following remark.

Remark 3.3. We should note that the existence of weak martingale solution stated in Theorem 3.2 still holds if we assume that the map $S(\cdot)$ is only continuous and satisfies a linear growth condition of the form (3.5).

To close this subsection we assume that $d = 2$ and we state the following uniqueness result.

Theorem 3.4. *Let $d = 2$ and assume that $(\mathbf{v}_i, \mathbf{n}_i)$, $i = 1, 2$ are two solution of (2.12) and (2.13) defined on the same stochastic system $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P}, W_1, W_2)$ and with the same initial condition $(\mathbf{v}_0, \mathbf{n}_0) \in \mathbb{H} \times \mathbb{H}^1$. Then, for any $t \in (0, T]$, \mathbb{P} -a.s.*

$$(\mathbf{v}_1(t), \mathbf{n}_1(t)) = (\mathbf{v}_2(t), \mathbf{n}_2(t)).$$

Proof. The proof of this result will be carried out in Subsection 3.4. \square

3.1. Galerkin approximation and a priori uniform estimates. As we mentioned earlier the proof of the existence of weak martingale solution relies on the Galerkin and compactness methods. This subsection will be devoted to the construction of the approximating solutions and the proofs of crucial estimates satisfied by these solutions.

Recall that there exists an orthonormal basis $\{\varphi_i; i = 1, 2, 3, \dots\}$ of \mathbb{H} consisting of the eigenvectors of the Stokes operator A . Recall also that there exists an orthonormal basis $\{\phi_i; i = 1, 2, 3, \dots\}$ of \mathbb{L}^2 consisting of the eigenvectors of the Neumann Laplacian A_1 . For any $m \in \mathbb{N}$ let us define the following finite-dimensional spaces

$$\begin{aligned} \mathbb{H}_m &:= \text{linspan}\{\varphi_1, \dots, \varphi_m\}, \\ \mathbb{L}_m &:= \text{linspan}\{\phi_1, \dots, \phi_m\}. \end{aligned}$$

In this subsection we introduce the finite-dimensional approximation of the system (2.12)-(2.13), justify the existence of solution of such approximation and We are mainly interested in derive uniform estimates for the sequence of approximating solutions. For this we purpose, we denote by π_m (resp. $\hat{\pi}_m$) the projection from \mathbb{H} (resp. \mathbb{L}^2) onto \mathbb{H}_m (resp. \mathbb{L}_m). These operators are self-adjoint. We also consider several mappings:

$$\begin{aligned} B_m &: \mathbb{H}_m \ni \mathbf{u} \mapsto \pi_m B(\mathbf{u}, \mathbf{u}) \in \mathbb{H}_m, \\ S_m &: \mathbb{H}_m \ni \mathbf{u} \mapsto \pi_m \circ S(\mathbf{u}) \in \mathcal{T}_2(K_1, \mathbb{H}_m), \\ M_m &: \mathbb{L}_m \ni \mathbf{n} \mapsto \pi_m M(\mathbf{n}) \in \mathbb{H}_m, \\ f_m &: \mathbb{L}_m \ni \mathbf{n} \mapsto \hat{\pi}_m f(\mathbf{n}) \in \mathbb{L}_m, \\ G_m &: \mathbb{L}_m \ni \mathbf{n} \mapsto \hat{\pi}_m G(\mathbf{n}) \in \mathbb{L}_m, \\ G_m^2 &: \mathbb{L}_m \ni \mathbf{n} \mapsto \hat{\pi}_m G^2(\mathbf{n}) \in \mathbb{L}_m, \\ \tilde{B}_m &: \mathbb{H}_m \times \mathbb{L}_m \ni (\mathbf{u}, \mathbf{n}) \mapsto \hat{\pi}_m \tilde{B}(\mathbf{v}, \mathbf{n}) \in \mathbb{L}_m. \end{aligned}$$

The proof of the following lemma is easy and we omit it.

Lemma 3.5. *The above mappings are locally Lipschitz.*

Let $\mathbf{n}_{0m} = \hat{\pi}_m \mathbf{n}_0$ and $\mathbf{v}_{0m} = \pi_m \mathbf{v}_0$. The Galerkin approximation to (2.12)-(2.13) is

$$d\mathbf{v}_m(t) + [A\mathbf{v}_m(t) + B_m(\mathbf{v}_m(t)) + M_m(\mathbf{n}_m(t))]dt = S_m(\mathbf{v}_m(t))dW_1, \quad (3.6)$$

$$d\mathbf{n}_m(t) + [A_1\mathbf{n}_m(t) + \tilde{B}_m(\mathbf{v}_m(t), \mathbf{n}_m(t)) + f_m(\mathbf{n}_m(t))]dt = \frac{1}{2}G_m^2(\mathbf{n}_m(t)) + G_m(\mathbf{n}_m(t))dW_2. \quad (3.7)$$

The equations (3.6)-(3.7) with initial condition $\mathbf{v}_m(0) = \mathbf{v}_{0m}$ and $\mathbf{n}_m(0) = \mathbf{n}_{0m}$ form a system of stochastic ordinary differential equations with locally Lipschitz coefficients hence owing to [47, Theorem 38, p. 303] it has a unique local maximal solution $(\mathbf{v}_m; \mathbf{n}_m; T_m)$ where T_m is a stopping time.

Remark 3.6. In case we assume that $S(\cdot)$ is only continuous and satisfies (3.5), Lemma 3.5 is no longer true. In fact the mappings defined just above Lemma 3.5 are only continuous and locally bounded. However, with this fact we can still justify the existence, possibly non-unique, of a weak local martingale solution to (3.6)-(3.7) by using results in [30, Chapter IV, Section 2, pp 167-177].

Proposition 3.7. *If all the assumptions of Theorem 3.2 are satisfied, then for any $p \geq 2$ there exists a positive constant C_p such that*

$$\begin{aligned} \sup_{m \in \mathbb{N}} \left(\mathbb{E} \sup_{t \in [0, T_m]} \|\mathbf{n}_m(t)\|^p + p \int_0^{t \wedge T_m} \|\mathbf{n}_m(s)\|^{p-2} \|\nabla \mathbf{n}_m(s)\|^2 ds \right. \\ \left. + p \int_0^{t \wedge T_m} \|\mathbf{n}_m(s)\|^{p-2} \|\mathbf{n}_m(s)\|_{\mathbb{L}^{2N+2}}^{2N+2} ds \right) \leq \mathbb{E} \mathfrak{G}_0(T, p). \end{aligned} \quad (3.8)$$

where

$$\mathfrak{G}_0(T, p) := \|\mathbf{n}_0\|^p (C_p + C_p e^{C_p T}). \quad (3.9)$$

Proof. Note that by the self-adjointness of $\tilde{\pi}_m$

$$\begin{aligned} \langle G_m(\mathbf{n}_m), \mathbf{n}_m \rangle &= \langle \hat{\pi}_m(\mathbf{n}_m \times h), \mathbf{n}_m \rangle, \\ &= \langle \mathbf{n}_m \times h, \mathbf{n}_m \rangle, \\ &= 0. \end{aligned}$$

Since $\mathbf{v}_m(t)$ is a divergence free function it follows from (A.10) that

$$\langle \hat{\pi}_m \tilde{B}(\mathbf{v}_m(t), \mathbf{n}_m(t)), \mathbf{n}_m(t) \rangle = \langle \tilde{B}(\mathbf{v}_m(t), \mathbf{n}_m(t)), \mathbf{n}_m(t) \rangle = 0.$$

Now we can prove the proposition by using the same ideas as in the proof of the estimate (C.1). \square

We also have the following estimates.

Proposition 3.8. *If all the assumptions of Theorem 3.2 are satisfied, then there exists $\ell > 0$ such that for any $p \in [1, \infty)$ the pair $(\mathbf{v}_m, \mathbf{n}_m)$ satisfies*

$$\begin{aligned} \mathbb{E} \left[\sup_{0 \leq s \leq T_m} \left(\|\mathbf{v}_m(s)\|^2 + \|\nabla \mathbf{n}_m(s)\|^2 + \ell \|\mathbf{n}_m(s)\|^2 + \int_{\mathcal{O}} F(\mathbf{n}_m(s, x)) dx \right)^p \right. \\ \left. + \left(\int_0^{T_m} (\|\nabla \mathbf{v}_m(s)\|^2 + \|\Delta \mathbf{n}_m(s) - f(\mathbf{n}_m(s))\|^2) ds \right)^p \right] \leq \mathfrak{G}_1(T, p), \end{aligned} \quad (3.10)$$

and

$$\mathbb{E} \left[\int_0^{T_m} \|\Delta \mathbf{n}_m(s)\|^2 ds \right]^p \leq \mathfrak{G}_1(T, p \cdot (2N + 1)), \quad (3.11)$$

where

$$\begin{aligned} \mathfrak{G}_1(T, p) := \left[\left(\|\mathbf{v}_0\|^2 + \|\mathbf{n}_0\|^2 + \|\nabla \mathbf{n}_0\|^2 + \int_{\mathcal{O}} F(\mathbf{n}_0(x)) dx \right)^p + \kappa T + \kappa \mathfrak{G}_0(T, p) \right] \\ \times \left[1 + \kappa T(T + 1) e^{\kappa(T+1)T} \right], \end{aligned} \quad (3.12)$$

\mathfrak{G}_0 is defined in (3.9) and κ is a positive constant depending only on p .

Proof. The estimates (3.10) follows from Proposition C.2 by taking $\tau_R = T_m$. \square

We state the following corollary.

Corollary 3.9. *The estimates (3.10) and (3.11) remains true with T_m replaced by T .*

Proof. Since the constants on the right hand sides of (3.10) and (3.11) does not depend on m , we can use the same argument as in [1] to show that for any $m \in \mathbb{N}$

$$T_m = T \quad \mathbb{P} - \text{a.s.}$$

This shows that the solution of our Galerkin approximation is globally defined, that is $T_m = T$ and the estimates (3.10) and (3.11) remain valid with T_m replaced by T . \square

In the next proposition we prove two uniform estimates for \mathbf{v}_m and \mathbf{n}_m which are very crucial for our purpose.

Proposition 3.10. *In addition to the assumptions of Theorem 3.2, let $\alpha \in (0, \frac{1}{2})$ and $p \in [2, \infty)$ such that $1 - \frac{d}{4} \geq \alpha - \frac{1}{p}$. Then, there exist positive constants $\bar{\kappa}_5$ and $\bar{\kappa}_6$ such that*

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|\mathbf{v}_m\|_{W^{\alpha,p}(0,T;\mathbb{V}^*)}^2 \leq \bar{\kappa}_5, \quad (3.13)$$

and

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|\mathbf{n}_m\|_{W^{\alpha,p}(0,T;\mathbb{L}^r)}^2 \leq \bar{\kappa}_6, \quad (3.14)$$

where $r = \frac{2N+2}{2N+1} \in (1, 2)$.

Proof. We rewrite the equation for \mathbf{v}_m as

$$\begin{aligned} \mathbf{v}_m(t) &= \mathbf{v}_{0m} - \int_0^t \mathbf{A} \mathbf{v}_m(s) ds - \int_0^t B_m(\mathbf{v}_m(s), \mathbf{v}_m(s)) ds - \int_0^t M_m(\mathbf{n}_m(s)) ds \\ &\quad + \int_0^t S_m(\mathbf{v}_m(s)) dW_1, \\ &= \mathbf{v}_{0m} + \sum_{i=1}^4 I_m^i(t). \end{aligned}$$

Since $\mathbf{A} \in \mathcal{L}(\mathbb{V}, \mathbb{V}^*)$ we easily check that there exists a certain constant $C > 0$ such that

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|I_m^1\|_{W^{1,2}(0,T;\mathbb{V}^*)}^2 \leq C. \quad (3.15)$$

We recall that for any $\alpha \in (0, \frac{1}{2})$ and $p \in [2, \infty)$

$$\mathbb{E} \left\| \int_0^\cdot \zeta(s) dW(s) \right\|_{W^{\alpha,p}(0,T;\mathbb{H})}^p \leq c \mathbb{E} \int_0^T \|\zeta(s)\|_{\mathbb{H}}^p ds; \quad (3.16)$$

see, for instance, [24]. From the above fact we infer that

$$\begin{aligned} \mathbb{E} \|I_m^4\|_{W^{\alpha,p}(0,T;\mathbb{H})}^p &\leq \mathbb{E} \int_0^T \|S_m(\mathbf{v}_m(t))\|_{\mathbb{H}}^p dt, \\ &\leq c \mathbb{E} \int_0^T (\ell_1 + \ell_2 \|\mathbf{v}_m(t)\|^p) ds. \end{aligned}$$

Thanks to (C.4) we derive that

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|I_m^4\|_{W^{\alpha,p}(0,T;\mathbb{H})}^p \leq C. \quad (3.17)$$

Now we treat the term $I_m^3(t)$. As a consequence of Lemma 2.1, we have

$$\mathbb{E} \|M_m(\mathbf{n}_m)\|_{\mathbb{L}^{\frac{d}{4}}(0,T;\mathbb{V}^*)}^2 \leq \left[\mathbb{E} \left(\int_0^T \|\Delta \mathbf{n}_m(t)\|^2 dt \right)^d \right]^{\frac{1}{2}},$$

from which altogether with (C.4) we infer that

$$\mathbb{E} \|I_m^3\|_{W^{1,\frac{d}{4}}(0,T;\mathbb{V}^*)}^2 \leq C. \quad (3.18)$$

Using (A.8) and an argument similar to the proof of the estimate for I_m^3 we conclude that

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|I_m^2(t)\|_{W^{1, \frac{d}{4}}(0, T; \mathbb{V}^*)}^2 \leq C. \quad (3.19)$$

By [55, Section 11, Corollary 19] we have the continuous imbedding

$$W^{1, \frac{d}{4}}(0, T; \mathbb{V}^*) \subset W^{\alpha, p}(0, T; \mathbb{V}^*), \quad (3.20)$$

for $\alpha \in (0, \frac{1}{2})$ and $p \in [2, \infty)$ such that $1 - \frac{d}{4} \geq \alpha - \frac{1}{p}$. Owing to Eqs. (3.15), (3.18), (3.17) and (3.19) and this continuous embedding we infer that (3.13) holds.

The second equations for the Galerkin approximation is written as

$$\begin{aligned} \mathbf{n}_m(t) &= \mathbf{n}_{0m} - \int_0^t A_1 \mathbf{n}_m(s) ds - \int_0^t \hat{\pi}_m [\tilde{B}_m(\mathbf{v}_m(s), \mathbf{n}_m(s))] ds - \int_0^t f_m(\mathbf{n}_m(s)) ds \\ &\quad + \frac{1}{2} \int_0^t G_m^2(\mathbf{n}_m(s)) ds + \int_0^t G_m(\mathbf{n}_m(s)) dW_2, \\ &= \mathbf{n}_{0m} + \sum_{j=1}^5 J_m^j(t). \end{aligned}$$

From (3.11) it is clear that

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|J_m^1\|_{W^{1,2}(0, T; \mathbb{L}^2)}^q \leq C. \quad (3.21)$$

For J_m^2 we have

$$\|\hat{\pi}_m [\tilde{B}_m(\mathbf{v}_m(s), \mathbf{n}_m(s))]\| \leq c \|\mathbf{v}_m(t)\|_{\mathbb{L}^4} \|\nabla \mathbf{n}_m(t)\|_{\mathbb{L}^4}, \text{ for } t \in [0, T].$$

Thanks to Gagliardo-Nirenberg's inequality we have

$$\|\hat{\pi}_m [\tilde{B}_m(\mathbf{v}_m(s), \mathbf{n}_m(s))]\| \leq c (\|\mathbf{v}_m(t)\| \|\nabla \mathbf{n}_m(t)\|)^{\frac{4-d}{4}} (\|\nabla \mathbf{v}_m(t)\| \|\Delta \mathbf{n}_m(t)\|)^{\frac{d}{4}}.$$

Thus

$$\begin{aligned} \|\hat{\pi}_m [\tilde{B}_m(\mathbf{v}_m(s), \mathbf{n}_m(s))]\|_{\mathbb{L}^{\frac{d}{4}}(0, T; \mathbb{L}^2)}^2 &\leq c \sup_{0 \leq t \leq T} (\|\mathbf{v}_m(t)\| \|\nabla \mathbf{n}_m(t)\|)^{\frac{4-d}{2}} \left[\int_0^T \|\nabla \mathbf{v}_m(t)\|^2 dt \right]^{\frac{d}{4}} \\ &\quad \times \left[\int_0^T \|\Delta \mathbf{n}_m(t)\|^2 dt \right]^{\frac{d}{4}}. \end{aligned}$$

Taking the mathematical expectation and using Hölder's inequality lead to

$$\begin{aligned} \sup_{m \in \mathbb{N}} \mathbb{E} \|\hat{\pi}_m [\tilde{B}_m(\mathbf{v}_m(s), \mathbf{n}_m(s))]\|_{\mathbb{L}^{\frac{d}{4}}(0, T; \mathbb{L}^2)}^2 &\leq c \left[\mathbb{E} \sup_{0 \leq t \leq T} \|\mathbf{v}_m(t)\|^{2(4-d)} \right]^{\frac{1}{4}} \left[\mathbb{E} \sup_{0 \leq t \leq T} \|\nabla \mathbf{n}_m(t)\|^{2(4-d)} \right]^{\frac{1}{4}} \\ &\quad \times \left[\mathbb{E} \left(\int_0^T \|\nabla \mathbf{v}_m(t)\|^2 dt \right)^d \right]^{\frac{1}{4}} \end{aligned}$$

Combining this with (3.10) and (3.10) yield

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|J_m^2\|_{W^{1, \frac{d}{4}}(0, T; \mathbb{L}^2)}^2 \leq C. \quad (3.22)$$

For any m and $t \in [0, T]$

$$\begin{aligned} \|G_m^2(\mathbf{n}_m(t))\| &\leq \|h\|_{\mathbb{L}^\infty} \|\mathbf{n}_m(t)\|_{\mathbb{L}^\infty} \|\mathbf{n}_m(t)\|, \\ &\leq \|h\|_{\mathbb{L}^\infty} (\|\mathbf{n}_m(t)\|^2 + \|\mathbf{n}_m(t)\| \|\nabla \mathbf{n}_m(t)\| + \|\mathbf{n}_m(t)\| \|\Delta \mathbf{n}_m(t)\|), \end{aligned}$$

which along with (3.10) and (3.11) yields

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|J_m^4\|_{W^{1,2}(0, T; \mathbb{L}^2)}^2 \leq C. \quad (3.23)$$

For any $h \in \mathbb{L}^\infty(\mathcal{O})$ we have

$$\|h \times \mathbf{n}_m(t)\|^p \leq \|h\|_{\mathbb{L}^\infty}^p \|\mathbf{n}_m(t)\|^p, \quad (3.24)$$

from which with (3.16) and (3.24), we derive that for any $\alpha \in (0, \frac{1}{2})$ and $p \in [2, \infty)$

$$\sup_{m \in \mathbb{N}} \mathbb{E} \|J_m^5\|_{W^{\alpha,p}(0,T;\mathbb{L}^2)}^p \leq C. \quad (3.25)$$

For the polynomial nonlinearity f we have

$$\begin{aligned} \sup_{m \in \mathbb{N}} \mathbb{E} \|J_m^3\|_{W^{\alpha,p}(0,T;\mathbb{L}^r)}^p &\leq \mathbb{E} \int_0^T \int_{\mathcal{O}} |f(\mathbf{n}_m(x,s))|^r dx ds \\ &\leq C + C \mathbb{E} \int_0^T \int_{\mathcal{O}} |\mathbf{n}_m(x,s)|^{2N+2} dx ds \\ &\leq C + C_2, \end{aligned} \quad (3.26)$$

where we have used (3.8). Combining all these estimates complete the proof of our proposition. \square

3.2. Tightness and Compactness results. This subsection is devoted to the study of the tightness of the Galerkin solutions and derive several weak convergence results. The estimates from the previous subsection play an important role in this part of the paper.

Let $p \in [2, \infty)$, $\alpha \in (0, \frac{1}{2})$ and r be as in Proposition 3.10. Let us consider the spaces

$$\begin{aligned} \mathfrak{X}_1 &= L^2(0, T; \mathbb{V}) \cap W^{\alpha,p}(0, T; \mathbb{V}^*), \\ \mathfrak{Y}_1 &= L^2(0, T; \mathbb{H}^2) \cap W^{\alpha,p}(0, T; \mathbb{L}^r) \end{aligned}$$

Recall that \mathbb{V}_β , $\beta \in \mathbb{R}$, is the domain of the of the fractional power operator A^β . Similarly, \mathbb{X}_β is the domain of $(I + A_1)^{\frac{1}{2}+\beta}$. The embedding $\mathbb{V}_\gamma \subset \mathbb{V}_\beta$ (resp. $\mathbb{X}_\gamma \subset \mathbb{X}_\beta$) is compact if that $\gamma > \beta$. We set

$$\begin{aligned} \mathfrak{X}_2 &= L^\infty(0, T; \mathbb{H}) \cap W^{\alpha,p}(0, T; \mathbb{V}^*), \\ \mathfrak{Y}_2 &= L^\infty(0, T; \mathbb{H}^1) \cap W^{\alpha,p}(0, T; \mathbb{L}^2), \end{aligned}$$

and

$$\begin{aligned} \mathfrak{S}_1 &= L^2(0, T; \mathbb{H}) \cap C([0, T]; \mathbb{V}_{-\beta}), \\ \mathfrak{S}_2 &= L^2(0, T; \mathbb{H}^1) \cap C([0, T]; \mathbb{X}_{\beta-\frac{1}{2}}). \end{aligned}$$

We shall prove the following important result.

Theorem 3.11. *Let $p \in [2, \infty)$ and $\alpha \in (0, \frac{1}{2})$ be as in Proposition 3.10 and $\beta \in (0, \frac{1}{2})$ such that $p\beta > 1$. The family of laws $\{\mathcal{L}(\mathbf{v}_m, \mathbf{n}_m), m \in \mathbb{N}\}$ is tight on the Polish space $\mathfrak{S}_1 \times \mathfrak{S}_2$.*

Proof. We firstly prove that $\{\mathcal{L}(\mathbf{v}_m), m \in \mathbb{N}\}$ is tight on $L^2(0, T; \mathbb{H})$. For this aim, we fix an arbitrary number $R > 0$ and prove that

$$\begin{aligned} \mathbb{P}(\|\mathbf{v}_m\|_{\mathfrak{X}_1} > R) &\leq \mathbb{P}\left(\|\mathbf{v}_m\|_{L^2(0,T;\mathbb{V})} > \frac{R}{2}\right) + \mathbb{P}\left(\|\mathbf{v}_m\|_{W^{\alpha,p}(0,T;\mathbb{V}^*)} > \frac{R}{2}\right), \\ &\leq \frac{4}{R^2} \mathbb{E}\left(\|\mathbf{v}_m\|_{L^2(0,T;\mathbb{V})}^2 + \|\mathbf{v}_m\|_{W^{\alpha,p}(0,T;\mathbb{V}^*)}^2\right). \end{aligned}$$

Thanks to (C.4), (3.13), and (3.14) we obtain from the last estimate that

$$\sup_{m \in \mathbb{N}} \mathbb{P}(\|\mathbf{v}_m\|_{\mathfrak{X}_1} > R) \leq \frac{4C}{R^2}. \quad (3.27)$$

Since \mathfrak{X}_1 is compactly embedded into $L^2(0, T; \mathbb{H})$, we conclude that the laws of \mathbf{v}_m form a family of probability measures which is tight on $L^2(0, T; \mathbb{H})$. Secondly, the same argument is used to prove that the laws of \mathbf{n}_m are tight on $L^2(0, T; \mathbb{L}^r)$. Next, let $\beta \in (0, \frac{1}{2})$ and $p \in [2, \infty)$ be satisfying $p\beta > 1$.

By [54, Corollary 5 of Section 8] the spaces \mathfrak{X}_2 and \mathfrak{Y}_2 are compactly imbedded in $C([0, T]; \mathbb{V}_{-\beta})$ and $C([0, T]; \mathbb{X}_{\beta-\frac{1}{2}})$, respectively. Hence the same argument as above provides us with the tightness of $\mathcal{L}(\mathbf{v}_m)$ and $\mathcal{L}(\mathbf{n}_m)$ on $C([0, T]; \mathbb{V}_{-\beta})$ and $C([0, T]; \mathbb{X}_{\beta-\frac{1}{2}})$. Now we can easily conclude the proof of the theorem. \square

Throughout the remaining part of this paper we assume that α , p and β are as in Theorem 3.11. We also use notation from Remark 2.4.

Proposition 3.12. *Let $\mathfrak{S} = \mathfrak{S}_1 \times \mathfrak{S}_2 \times C([0, T]; \mathbb{K}_2) \times C([0, T]; \mathbb{R})$. There exist a Borel probability measure μ on \mathfrak{S} and a subsequence of $(\mathbf{v}_m, \mathbf{n}_m, W_1, W_2)$ such that its laws weakly converge to μ .*

Proof. Thanks to the above lemma the laws of $(\mathbf{v}_m, \mathbf{n}_m, W_1, W_2)$ form a tight family on \mathfrak{S} . Since \mathfrak{S} is a Polish space, we get the result from the application of Prohorov's theorem. \square

The following result relates the above convergence in law to almost sure convergence.

Proposition 3.13. *There exist a complete probability space $(\Omega', \mathcal{F}', \mathbb{P}')$ and a sequence of \mathfrak{S} -valued random variables, denoted by $(\bar{\mathbf{v}}_m, \bar{\mathbf{n}}_m, W_1^m, W_2^m)$, defined on $(\Omega', \mathcal{F}', \mathbb{P}')$ such that its law is equal to the law of $(\mathbf{v}_m, \mathbf{n}_m, W_1, W_2)$ on \mathfrak{S} . Also, there exists an \mathfrak{S} -random variable $(\mathbf{v}, \mathbf{n}, \bar{W}_1, \bar{W}_2)$ defined on $(\Omega', \mathcal{F}', \mathbb{P}')$ such that*

$$\mathcal{L}(\mathbf{v}, \mathbf{n}, \bar{W}_1, \bar{W}_2) = \mu \text{ on } \mathfrak{S}, \quad (3.28)$$

$$\bar{\mathbf{v}}_m \rightarrow \mathbf{v} \text{ in } L^2(0, T; \mathbb{H}) \text{ } \mathbb{P}' - a.s., \quad (3.29)$$

$$\bar{\mathbf{v}}_m \rightarrow \mathbf{v} \text{ in } C([0, T]; \mathbb{V}_{-\beta}) \text{ } \mathbb{P}' - a.s., \quad (3.30)$$

$$\bar{\mathbf{n}}_m \rightarrow \mathbf{n} \text{ in } L^2(0, T; \mathbb{H}^1) \text{ } \mathbb{P}' - a.s., \quad (3.31)$$

$$\bar{\mathbf{n}}_m \rightarrow \mathbf{n} \text{ in } C([0, T]; \mathbb{X}_{\beta-\frac{1}{2}}) \text{ } \mathbb{P}' - a.s., \quad (3.32)$$

$$\bar{W}_1^m \rightarrow \bar{W}_1 \text{ in } C([0, T]; \mathbb{K}_2) \text{ } \mathbb{P}' - a.s., \quad (3.33)$$

$$\bar{W}_2^m \rightarrow \bar{W}_2 \text{ in } C([0, T]; \mathbb{R}) \text{ } \mathbb{P}' - a.s. \quad (3.34)$$

Proof. This is just a consequence of Proposition 3.12 and Skorokhod's Theorem. \square

Let $\mathfrak{X}_3 = L^\infty(0, T; \mathbb{H}) \cap L^2(0, T; \mathbb{V})$ and $\mathfrak{Y}_3 = L^\infty(0, T; \mathbb{H}^1) \cap L^2(0, T; \mathbb{H}^2)$

Proposition 3.14. *If all the assumptions of Theorem 3.2 are verified, then for any $p \geq 2$ $(\bar{\mathbf{v}}_m, \bar{\mathbf{n}}_m)$ satisfies the following estimates on the new probability space $(\Omega', \mathcal{F}', \mathbb{P}')$:*

$$\begin{aligned} \sup_{m \in \mathbb{N}} \left(\mathbb{E} \left[\sup_{t \in [0, T]} \|\bar{\mathbf{n}}_m(t)\|^p + p \int_0^T \|\bar{\mathbf{n}}_m(s)\|^{p-2} \|\nabla \bar{\mathbf{n}}_m(s)\|^2 ds \right. \right. \\ \left. \left. + p \int_0^t \|\bar{\mathbf{n}}_m(s)\|^{p-2} \|\bar{\mathbf{n}}_m(s)\|_{\mathbb{L}_{2N+2}^{2N+2}}^2 ds \right] \right) \leq \mathfrak{G}_0(T, p), \end{aligned} \quad (3.35)$$

$$\begin{aligned} \mathbb{E} \left[\sup_{0 \leq s \leq T} \left(\|\bar{\mathbf{v}}_m(s)\|^2 + \ell \|\bar{\mathbf{n}}_m(s)\|^2 + \|\nabla \bar{\mathbf{n}}_m(s)\|^2 + \int_{\mathcal{O}} F(\bar{\mathbf{n}}_m(s, x)) dx \right)^p \right. \\ \left. + \int_0^T (\|\nabla \bar{\mathbf{v}}_m(s)\|^2 + \|\Delta \bar{\mathbf{n}}_m(s) - f(\mathbf{n}_m(s))\|^2) ds \right]^p \leq \mathfrak{G}_1(T, p), \end{aligned} \quad (3.36)$$

$$\mathbb{E} \left[\int_0^T \|\Delta \mathbf{n}_m(s)\|^2 ds \right]^p \leq \mathfrak{G}_1(T, p \cdot (2N + 1)), \quad (3.37)$$

where $\mathfrak{G}_0(T, p)$, ℓ and $\mathfrak{G}_1(T, p)$ are defined as in Proposition 3.7 and Proposition 3.8, respectively. Furthermore, there exists a constant $C > 0$ such that

$$\sup_{m \in \mathbb{N}} \mathbb{E}' \left[\int_0^T \|B_m(\bar{\mathbf{v}}_m(t), \bar{\mathbf{v}}_m(t))\|_{\mathbb{V}^*}^{\frac{4}{3}} dt \right]^{\frac{d}{2}} \leq C, \quad (3.38)$$

$$\sup_{m \in \mathbb{N}} \mathbb{E}' \left[\int_0^T \|M_m(\bar{\mathbf{n}}_m(t))\|_{\mathbb{V}^*}^{\frac{4}{3}} dt \right]^{\frac{d}{2}} \leq C, \quad (3.39)$$

$$\sup_{m \in \mathbb{N}} \mathbb{E}' \left[\int_0^T \|\tilde{B}_m(\bar{\mathbf{v}}_m(t), \bar{\mathbf{n}}_m(t))\|_{\mathbb{L}^2}^{\frac{4}{3}} dt \right]^{\frac{d}{2}} \leq C, \quad (3.40)$$

$$\sup_{m \in \mathbb{N}} \mathbb{E}' \int_0^T \|f_m(\bar{\mathbf{n}}_m(t))\|_{\mathbb{L}^r}^r dt \leq C, \quad (3.41)$$

where r is defined in Proposition 3.10.

Proof. Consider the function $\Phi(\mathbf{u}, \mathbf{e})$ on $\mathfrak{X}_3 \times \mathfrak{Y}_3 \subset \mathfrak{S}_1 \times \mathfrak{S}_2$ defined by

$$\Phi(\mathbf{u}, \mathbf{e}) = \sup_{0 \leq s \leq T} [\|\mathbf{u}(s)\|^{2p} + \|\nabla \mathbf{e}(s)\|^{2p}] + \tilde{\kappa}_0 \left[\int_0^T (\|\nabla \mathbf{u}(s)\|^2 + \|\Delta \mathbf{e}(s)\|^2) ds \right]^p$$

It is a continuous function, thus Borel measurable, on $\mathfrak{S}_1 \times \mathfrak{S}_2$. Thanks to (3.28) the processes $(\mathbf{v}_m, \mathbf{n}_m)$ and $(\bar{\mathbf{v}}_m, \bar{\mathbf{n}}_m)$ are identical in law. Therefore, we derive that

$$\mathbb{E}\Phi(\mathbf{v}_m, \mathbf{n}_m) = \mathbb{E}'\Phi(\bar{\mathbf{v}}_m, \bar{\mathbf{n}}_m).$$

This identity and the estimate (C.4) implies (3.36). The estimates (3.38), (3.39) and (3.40) can be prove using similar idea to the proof of (3.19), (3.18), (3.22) and (3.26), respectively. \square

We prove several convergences which are of the essence for the proof of our existence result.

Proposition 3.15. *We can extract a subsequence $(\bar{\mathbf{v}}_{m_k}, \bar{\mathbf{n}}_{m_k})$ from $(\bar{\mathbf{v}}_m, \bar{\mathbf{n}}_m)$ such that*

$$\bar{\mathbf{v}}_{m_k} \rightarrow \mathbf{v} \text{ strongly in } L^2(\Omega' \times [0, T]; \mathbb{H}), \quad (3.42)$$

$$\bar{\mathbf{v}}_{m_k} \rightarrow \mathbf{v} \text{ strongly in } L^4(\Omega'; C([0, T]; \mathbb{V}_{-\beta})), \quad (3.43)$$

$$\bar{\mathbf{n}}_{m_k} \rightarrow \mathbf{n} \text{ strongly in } L^2(\Omega' \times [0, T]; \mathbb{H}^1), \quad (3.44)$$

$$\bar{\mathbf{n}}_{m_k} \rightarrow \mathbf{n} \text{ strongly in } L^4(\Omega'; C([0, T]; \mathbb{X}_{\beta-\frac{1}{2}})), \quad (3.45)$$

$$\bar{\mathbf{n}}_{m_k} \rightarrow \mathbf{n} \text{ strongly in } \mathfrak{S}_2 \text{ } \mathbb{P}' - a.s., \quad (3.46)$$

$$\bar{\mathbf{n}}_{m_k} \rightarrow \mathbf{n} \text{ for almost everywhere } (x, t) \text{ and } \mathbb{P}' - a.s.. \quad (3.47)$$

Proof. From (3.36) and Banach-Alaoglu's theorem we infer that there exist a subsequence of $\bar{\mathbf{v}}_{m_k}$ satisfying

$$\bar{\mathbf{v}}_{m_k} \rightarrow \mathbf{v} \text{ weakly in } L^{2p}(\Omega'; L^2(0, T; \mathbb{H})), \quad (3.48)$$

for any $p \in [2, \infty)$. Now let us consider the positive nondecreasing function $\varphi(x) = x^{2p}$, $p \in [2, \infty)$, defined on \mathbb{R}_+ . The function φ obviously satisfies

$$\lim_{x \rightarrow \infty} \frac{\varphi(x)}{x} = \infty. \quad (3.49)$$

Thanks to the estimate $\mathbb{E}' \sup_{t \in [0, T]} \|\bar{\mathbf{v}}_{m_k}\|^{2p} \leq C$ (see (3.36)), we have

$$\sup_{m \geq 1} \mathbb{E}'(\varphi(\|\bar{\mathbf{v}}_{m_k}\|_{L^2(0, T; \mathbb{H})})) < \infty, \quad (3.50)$$

which along with the uniform integrability criteria in [32, Chapter 3, Exercice 6] implies that the family $\{\|\bar{\mathbf{v}}_{m_k}\|_{L^2(0, T; \mathbb{H})} : m \in \mathbb{N}\}$ is uniform integrable with respect to the probability measure.

Thus, we can deduce from Vitali's Convergence Theorem (see, for instance, [32, Chapter 3, Proposition 3.2]) and (3.29) that

$$\mathbb{E}' \|\bar{\mathbf{v}}_{m_k}\|_{L^2(0,T;\mathbb{H})}^2 \rightarrow \mathbb{E}' \|\mathbf{v}\|_{L^2(0,T;\mathbb{H})}^2.$$

Form this and (3.48) we derive that

$$\bar{\mathbf{v}}_{m_k} \rightarrow \mathbf{v} \text{ strongly in } L^2(\Omega' \times [0, T]; \mathbb{H}). \quad (3.51)$$

Thanks to (3.31)-(3.34) in Proposition 3.13 and (3.36) we can use the same argument to show the convergence (3.43), (3.44) and (3.45). Hence by the tightness of the laws of $\bar{\mathbf{n}}_m$ on \mathfrak{S}_2 we can extract a subsequence still denoted by $\bar{\mathbf{n}}_{m_k}$ such that (3.46) and (3.47) hold. \square

The stochastic processes \mathbf{v} and \mathbf{n} satisfies the following properties.

Proposition 3.16. *We have*

$$\mathbb{E}' \sup_{t \in [0, T]} \|\mathbf{v}(t)\|^p < \infty, \quad (3.52)$$

$$\mathbb{E}' \sup_{t \in [0, T]} \|\mathbf{d}(t)\|_{\mathbb{H}^1}^p < \infty, \quad (3.53)$$

for any $p \in [2, \infty)$.

Proof. One can argue exactly as in [7, Proof of (4.12), page 20], so we omit the details. \square

Proposition 3.17. *Let $d \in \{2, 3\}$ and $T \geq 0$. There exist four processes $\mathfrak{B}_1, \mathfrak{M} \in L^2(\Omega'; L^{\frac{4}{4-d}}(0, T; \mathbb{V}^*))$, $\mathfrak{B}_2 \in L^2(\Omega'; L^{\frac{4}{4-d}}(0, T; \mathbb{L}^2))$ and $\mathfrak{f} \in L^{\frac{2N+2}{2N+1}}(\Omega' \times [0, T] \times \mathcal{O})$ such that*

$$B_{m_k}(\bar{\mathbf{v}}_{m_k}, \bar{\mathbf{v}}_{m_k}) \rightarrow \mathfrak{B}_1, \text{ weakly in } L^2(\Omega'; L^{\frac{4}{4-d}}(0, T; \mathbb{V}^*)), \quad (3.54)$$

$$M_{m_k}(\bar{\mathbf{n}}_{m_k}) \rightarrow \mathfrak{M}, \text{ weakly in } L^2(\Omega'; L^{\frac{4}{4-d}}(0, T; \mathbb{V}^*)), \quad (3.55)$$

$$\tilde{B}_{m_k}(\bar{\mathbf{v}}_{m_k}, \bar{\mathbf{n}}_{m_k}) \rightarrow \mathfrak{B}_2, \text{ weakly in } L^2(\Omega'; L^{\frac{4}{4-d}}(0, T; \mathbb{L}^2)), \quad (3.56)$$

$$f_{m_k}(\bar{\mathbf{n}}_{m_k}) \rightarrow \mathfrak{f}, \text{ weakly in } L^{\frac{2N+2}{2N+1}}(\Omega' \times [0, T] \times \mathcal{O}). \quad (3.57)$$

Proof. Note that Proposition 3.14 remains valid with $\bar{\mathbf{n}}_m$ replaced by $\bar{\mathbf{n}}_{m_k}$. Thus, Proposition 3.17 follows from Eqs. (3.38)-(3.41) and application of Banach-Alaoglu's theorem. \square

3.3. Passage to the limit and the end of proof of Theorem 3.2. In this subsection we prove several convergences which will enable us to conclude that the limiting objects that we found in Proposition 3.13 are in fact a weak martingale solution to our problem.

Proposition 3.15 will be used to prove the following result.

Proposition 3.18. *For any process $\Psi \in L^2(\Omega'; L^{\frac{4}{4-d}}(0, T; \mathbb{V}))$, the following identity holds*

$$\begin{aligned} \lim_{m \rightarrow \infty} \mathbb{E}' \int_0^T \mathbb{V}^* \langle B_{m_k}(\bar{\mathbf{v}}_{m_k}(t), \bar{\mathbf{v}}_{m_k}(t)), \Psi(t) \rangle_{\mathbb{V}} dt &= \mathbb{E}' \int_0^T \mathbb{V}^* \langle \mathfrak{B}_1(t), \Psi(t) \rangle_{\mathbb{V}} dt, \\ &= \mathbb{E}' \int_0^T \mathbb{V}^* \langle B(\mathbf{v}(t), \mathbf{v}(t)), \Psi(t) \rangle_{\mathbb{V}} dt. \end{aligned} \quad (3.58)$$

Proof. Let

$$\mathbb{D} = \left\{ \Phi = \sum_{i=1}^k \mathbb{1}_{D_i} \mathbb{1}_{J_i} \psi_i : D_i \subset \Omega, J_i \subset [0, T] \text{ is measurable}, \psi_i \in \mathcal{V} \right\}.$$

Owing to [60, Proposition 21.23] and the density of \mathbb{D} in $L^2(\Omega, \mathbb{P}; L^{\frac{4}{4-d}}(0, T; \mathbb{V}))$ (see, for instance, [48, Theorem 3.2.6]), in order to show that the identity (3.58) holds it is enough to check that

$$\lim_{m \rightarrow \infty} \mathbb{E}' \int_0^T \mathbb{1}_J(t) \mathbb{1}_{D \mathbb{V}^*} \langle B_{m_k}(\bar{\mathbf{v}}_{m_k}(t), \bar{\mathbf{v}}_{m_k}(t)) - B(\mathbf{v}(t), \mathbf{v}(t)), \psi \rangle_{\mathbb{V}} dt = 0,$$

for any $\Phi = \mathbb{1}_D \mathbb{1}_J \psi \in \mathbb{D}$. For this purpose we first note that

$$\begin{aligned} \mathbb{V}^* \langle B_{m_k}(\bar{\mathbf{v}}_{m_k}(t), \bar{\mathbf{v}}_{m_k}(t)) - B(\mathbf{v}(t), \mathbf{v}(t)), \psi \rangle_{\mathbb{V}} &= \mathbb{V}^* \langle \tilde{B}_{m_k}(\bar{\mathbf{v}}_{m_k}(t) - \mathbf{v}(t), \bar{\mathbf{v}}_{m_k}(t)), \psi \rangle_{\mathbb{V}} \\ &\quad + \mathbb{V}^* \langle \tilde{B}_{m_k}(\mathbf{v}(t), \bar{\mathbf{v}}_{m_k}(t) - \mathbf{v}(t)), \psi \rangle_{\mathbb{V}}, \\ &= I_1(t) + I_2(t). \end{aligned}$$

The mapping $\langle B_{m_k}(\mathbf{u}, \cdot), \psi \rangle_{\mathbb{V}^*, \mathbb{V}}$ from $L^2(\Omega'; L^2(0, T; \mathbb{V}))$ into $L^2(\Omega'; L^{\frac{4}{d}}(0, T; \mathbb{R}))$ is linear and continuous. Therefore if $\bar{\mathbf{v}}_{m_k}$ converges to \mathbf{v} weakly in $L^2(\Omega'; L^2(0, T; \mathbb{V}))$ then I_2 converges to 0 weakly in $L^2(\Omega'; L^{\frac{4}{d}}(0, T; \mathbb{R}))$. To deal with I_1 we recall that

$$\begin{aligned} \left| \mathbb{E}' \int_0^T \mathbb{1}_J \mathbb{1}_D(\omega', t) \mathbb{V}^* \langle B_{m_k}(\bar{\mathbf{v}}_{m_k}(t) - \mathbf{v}(t), \bar{\mathbf{v}}_{m_k}(t)), \psi \rangle_{\mathbb{V}} dt \right| &\leq \|\psi\|_{\mathbb{L}^\infty} \left[\mathbb{E}' \int_0^T \|\nabla \bar{\mathbf{v}}_{m_k}(t)\|^2 dt \right]^{\frac{1}{2}} \\ &\quad \times \left[\mathbb{E}' \int_0^T \|\bar{\mathbf{v}}_{m_k}(t) - \mathbf{v}(t)\|^2 dt \right]^{\frac{1}{2}}. \end{aligned}$$

Thanks to (3.36) and the convergence (3.42) we see that the right-hand side of last inequality converges to 0 as m_k goes to infinity. Hence I_1 converges to 0 weakly in $L^2(\Omega'; L^{\frac{4}{d}}(0, T; \mathbb{R}))$. This ends the proof of our proposition. \square

In the next proposition we will prove that \mathfrak{M} coincide with $M(\mathbf{n})$.

Proposition 3.19. *Assume that $n < 4$. For any process $\Psi \in L^2(\Omega'; L^{\frac{4}{4-d}}(0, T; \mathbb{V}))$, the following identity holds*

$$\mathbb{E}' \int_0^T \mathbb{V}^* \langle \mathfrak{M}(t), \Psi(t) \rangle_{\mathbb{V}} dt = \mathbb{E}' \int_0^T \mathbb{V}^* \langle M(\mathbf{n}(t)), \Psi(t) \rangle_{\mathbb{V}} dt. \quad (3.59)$$

Proof. Since π_m strongly converges to the identity operator Id in $L^2(\Omega'; L^{\frac{d}{4}}(0, T; \mathbb{V}^*))$, it is enough to show that (3.59) is true for $M(\bar{\mathbf{n}}_{m_k}(t)) := M(\mathbf{n})$ in place of $M_{m_k}(\bar{\mathbf{n}}_{m_k}(t))$. By the relation (2.6) we have

$$\begin{aligned} \langle M(\bar{\mathbf{n}}_{m_k}(t)) - M(\mathbf{n}(t)), \psi \rangle &= \sum_{i,j,k} \int_{\mathcal{O}} \partial_{x_j} \psi^i \partial_{x_i} \bar{\mathbf{n}}_{m_k}^k(t) \left(\partial_{x_j} \bar{\mathbf{n}}_{m_k}^k(t) - \partial_{x_i} \mathbf{n}^k(t) \right) dx \\ &\quad + \sum_{i,j,k} \int_{\mathcal{O}} \partial_{x_j} \psi^i \partial_{x_j} \mathbf{n}^k(t) \left(\partial_{x_i} \bar{\mathbf{n}}_{m_k}^k(t) - \partial_{x_i} \mathbf{n}^k(t) \right) dx, \end{aligned} \quad (3.60)$$

for any $\psi \in \mathcal{V}$. From this inequality we infer that

$$\begin{aligned} \left| \mathbb{E}' \int_0^T \mathbb{1}_J(\omega', t) \mathbb{V}^* \langle M(\bar{\mathbf{n}}_{m_k}(t)) - M(\mathbf{n}(t)), \psi \rangle_{\mathbb{V}} dt \right| &\leq C \|\nabla \psi\| \left[\mathbb{E}' \int_0^T \|\nabla(\bar{\mathbf{n}}_{m_k}(t) - \mathbf{n}(t))\|^2 dt \right]^{\frac{1}{2}} \\ &\quad \times \left(\left[\mathbb{E}' \sup_{0 \leq t \leq T} \|\nabla \bar{\mathbf{n}}_{m_k}(t)\|^2 \right]^{\frac{1}{2}} + \left[\mathbb{E}' \sup_{0 \leq t \leq T} \|\nabla \mathbf{n}(t)\|^2 \right]^{\frac{1}{2}} \right) \end{aligned} \quad (3.61)$$

Owing to the estimate (3.36) and the convergence (3.44) we infer that the left hand side of the last inequality converges to 0 as m_k goes to infinity. Now we easily conclude the proof. \square

Proposition 3.20. *Let $d \in \{2, 3\}$. Then,*

$$\mathfrak{B}_2 = \tilde{B}(\mathbf{v}, \mathbf{n}) \text{ in } L^2(\Omega'; L^{\frac{d}{4}}(0, T; \mathbb{L}^2)).$$

Proof. The statement in the proposition is equivalent to say that $\tilde{B}_{m_k}(\bar{\mathbf{v}}_{m_k}(t), \bar{\mathbf{n}}_{m_k}(t))$ converges to $\tilde{B}(\mathbf{v}(t), \mathbf{n}(t))$ weakly in $L^2(\Omega'; L^{\frac{d}{4}}(0, T; \mathbb{L}^2))$. To prove this we argue as above, but we consider the set

$$\mathbb{D} = \{\Phi = \mathbb{1}_J \mathbb{1}_D \mathbb{1}_K : J \subset \Omega', D \subset [0, T], K \subset \mathcal{O} \text{ is measurable}\}.$$

For any $\Phi \in \mathbb{D}$ we have

$$\begin{aligned} & \left| \mathbb{E}' \int_{[0,T] \times Q} \tilde{B}_{m_k}(\bar{\mathbf{v}}_{m_k}(t), \bar{\mathbf{n}}_{m_k}(t)) - \tilde{B}(\mathbf{v}(t), \mathbf{n}(t)) \Phi(\omega', t, x) dx dt \right| \\ & \leq \left[\mathbb{E}' \int_0^T \|\bar{\mathbf{v}}_{m_k}(t) - \mathbf{v}(t)\|^2 dt \right]^{\frac{1}{2}} \left[\mathbb{E}' \int_0^T \|\nabla \bar{\mathbf{n}}_{m_k}(t)\|^2 dt \right]^{\frac{1}{2}} \\ & \quad + \left[\mathbb{E}' \int_0^T \|\mathbf{v}(t)\|^2 dt \right]^{\frac{1}{2}} \left[\mathbb{E}' \int_0^T \|\nabla (\bar{\mathbf{v}}_{m_k}(t) - \mathbf{v}(t))\|^2 dt \right]^{\frac{1}{2}} \end{aligned} \quad (3.62)$$

Thanks to (3.36) and (3.44) we deduce that the left hand side of the last inequality converges to 0 as m_k goes to infinity. This proves our claim. \square

The following convergences are also important.

Proposition 3.21. *Let r be as in Proposition 3.10, that is $r = \frac{2N+2}{2N+1} \in (1, 2)$. Then,*

$$\mathbf{f} = f(\mathbf{n}) \text{ in } L^r(\Omega' \times \mathcal{O} \times [0, T]). \quad (3.63)$$

Proof. To prove (3.63), first remark that by definition the embedding $\mathbb{X}_{\beta-\frac{1}{2}} \subset \mathbb{L}^r$ is continuous for any $\beta \in (0, \frac{1}{2})$. The convergence (3.47) implies that for any $k = 0, \dots, N$

$$|\bar{\mathbf{n}}_{m_k}|^{2k} \bar{\mathbf{n}}_{m_k} \rightarrow |\mathbf{n}|^{2k} \mathbf{n} \text{ for almost everywhere } (x, t) \text{ and } \mathbb{P}' - a.s.. \quad (3.64)$$

Since $f(\bar{\mathbf{n}}_{m_k})$ is bounded in $L^r(\Omega' \times \mathcal{O} \times [0, T])$ we can infer from [41, Lemma 1.3, pp. 12] and the convergence (3.64) that

$$f(\bar{\mathbf{n}}_{m_k}) \rightarrow f(\mathbf{n}) \text{ weakly in } L^r(\Omega' \times \mathcal{O} \times [0, T]),$$

which with the uniqueness of weak limit implies the sought result. \square

To simplify notation let us define the processes $\mathcal{M}_k^1(t)$ and $\mathcal{M}_k^2(t)$ by

$$\mathcal{M}_{m_k}^1(t) = \bar{\mathbf{v}}_{m_k}(t) - \bar{\mathbf{v}}_{m_k}(0) + \int_0^t \left(A_1 \bar{\mathbf{v}}_{m_k}(s) + \tilde{B}_{m_k}(\bar{\mathbf{v}}_{m_k}(s), \bar{\mathbf{v}}_{m_k}(s)) - M_{m_k}(\bar{\mathbf{n}}_{m_k}(s)) \right) ds,$$

and

$$\begin{aligned} \mathcal{M}_{m_k}^2(t) &= \bar{\mathbf{n}}_{m_k}(t) - \bar{\mathbf{n}}_{m_k}(0) + \int_0^t \left(A \bar{\mathbf{n}}_{m_k}(s) + \tilde{B}_{m_k}(\bar{\mathbf{v}}_{m_k}(s), \bar{\mathbf{n}}_{m_k}(s)) - f_{m_k}(\bar{\mathbf{n}}_{m_k}(s)) \right) ds \\ &\quad - \int_0^t G_{m_k}^2(\bar{\mathbf{n}}_{m_k}(s)) ds. \end{aligned}$$

Proposition 3.22. *Let $\mathcal{M}_1(t)$ and $\mathcal{M}_2(t)$ be defined by*

$$\mathcal{M}^1(t) = \mathbf{v}(t) - \mathbf{v}_0 + \int_0^t \left(A_1 \mathbf{v}(s) + B(\mathbf{v}(s), \mathbf{v}(s)) - M(\mathbf{n}(s)) \right) ds, \quad (3.65)$$

$$\mathcal{M}^2(t) = \mathbf{n}(t) - \mathbf{n}_0 + \int_0^t \left(A \mathbf{n}(s) + \tilde{B}(\mathbf{v}(s), \mathbf{n}(s)) - f(\mathbf{n}(s)) \right) ds - \int_0^t G^2(\mathbf{n}(s)) ds, \quad (3.66)$$

for any $t \in (0, T]$. Then, for any $t \in (0, T]$

$$\mathcal{M}_{m_k}^1(t) \text{ converges weakly in } L^2(\Omega'; \mathbb{V}^*) \text{ to } \mathcal{M}^1(t),$$

$$\mathcal{M}_{m_k}^2(t) \text{ converges weakly in } L^2(\Omega'; \mathbb{L}^2) \text{ to } \mathcal{M}^2(t),$$

as $m_k \rightarrow \infty$.

Proof. Let $t \in (0, T]$, we first prove that $\mathcal{M}_{m_k}^1(t) \rightarrow \mathcal{M}^1(t)$ weakly in $L^2(\Omega'; \mathbb{V}^*)$ as m goes to infinity. To this end we take an arbitrary $\xi \in L^2(\Omega'; \mathbb{V})$. We have

$$\begin{aligned} \mathbb{E}' \left[\langle \mathcal{M}_{m_k}^1(t), \xi \rangle \right] &= \mathbb{E}' \left[\langle \bar{\mathbf{v}}_{m_k}(t) - \bar{\mathbf{v}}_{m_k}(0), \xi \rangle - \int_0^t \langle \nabla \bar{\mathbf{v}}_{m_k}(s), \nabla \xi \rangle ds - \int_0^t \langle M_{m_k}(\bar{\mathbf{n}}_{m_k}(s)), \xi \rangle ds \right] \\ &\quad + \mathbb{E}' \left[\int_0^t \langle B_{m_k}(\bar{\mathbf{v}}_{m_k}(s), \bar{\mathbf{v}}_{m_k}(s)), \xi \rangle ds \right] \end{aligned}$$

Thanks to the pointwise convergence in $C([0, T]; \mathbb{V}^{-\beta})$, thus in $C([0, T]; \mathbb{V}^*)$, and the convergences (3.54), (3.58), (3.55) and (3.59) we obtain

$$\begin{aligned} \lim_{m \rightarrow \infty} \mathbb{E}' \left[\langle \mathcal{M}_{m_k}^1(t), \xi \rangle \right] &= \mathbb{E}' \left[\langle \mathbf{v}(t) - \mathbf{v}_0, \xi \rangle - \int_0^t \langle \nabla \mathbf{v}(s), \nabla \xi \rangle ds - \int_0^t \langle M(\mathbf{n}(s)), \xi \rangle ds \right] \\ &\quad + \mathbb{E}' \left[\int_0^t \langle B(\mathbf{v}(s), \mathbf{v}(s)), \xi \rangle ds \right], \end{aligned}$$

which proves the sought convergence.

Secondly, we prove that for any $t \in (0, T]$ $\mathcal{M}_{m_k}^2(t) \rightarrow \mathcal{M}^2(t)$ weakly in $L^2(\Omega'; \mathbb{L}^2)$ as m tends to infinity. For this purpose, observe that $G_{m_k}^2(\cdot)$ is a linear mapping from $L^2(\Omega'; C([0, T]; \mathbb{L}^2))$ into itself and it satisfies

$$\mathbb{E}' \|G^2(\mathbf{n}(t))\|_{C([0, T]; \mathbb{L}^2)}^p \leq c \|h\|_{\mathbb{L}^\infty}^2 \mathbb{E}' \|\mathbf{n}(t)\|_{C([0, T]; \mathbb{L}^2)}^p, \quad (3.67)$$

for any $p \in [2, \infty)$. So it is not difficult to show that

$$G_{m_k}(\bar{\mathbf{n}}_{m_k}) \rightarrow G(\mathbf{n}) \text{ strongly in } L^2(\Omega'; C([0, T]; \mathbb{L}^2)). \quad (3.68)$$

Thanks to this observation, the convergences (3.45), (3.63), (3.56) and Proposition (3.20) we can use the same argument as above to show that

$$\lim_{m \rightarrow \infty} \langle \mathcal{M}_{m_k}^2(t), \xi \rangle = \langle \mathcal{M}^2(t), \xi \rangle, \quad (3.69)$$

for any $t \in (0, T]$ and $\xi \in L^2(\Omega'; \mathbb{L}^2)$. This completes the proof of Proposition 3.22. \square

Let \mathcal{N} be the set of null sets \mathcal{F} and for any $t \geq 0$ and $k \in \mathbb{N}$, let

$$\begin{aligned} \hat{\mathcal{F}}_t^{m_k} &:= \sigma \left(\sigma \left((\bar{\mathbf{v}}_{m_k}(s), \bar{\mathbf{n}}_{m_k}(s), \bar{W}_1^{m_k}(s), \bar{W}_2^{m_k}(s)); s \leq t \right) \cup \mathcal{N} \right), \\ \mathcal{F}_t' &:= \sigma \left(\sigma \left((\mathbf{v}(s), \mathbf{n}(s), \bar{W}_1(s), \bar{W}_2(s)); s \leq t \right) \cup \mathcal{N} \right). \end{aligned} \quad (3.70)$$

Let us also define the stochastic processes $\mathfrak{M}_{m_k}^1$ and $\mathfrak{M}_{m_k}^2$ by

$$\begin{aligned} \mathfrak{M}_{m_k}^1(t) &= \int_0^t S_{m_k}(\bar{\mathbf{v}}_{m_k}(s)) d\bar{W}_1^{m_k}(s) \\ \mathfrak{M}_{m_k}^2(t) &= \int_0^t G_{m_k}(\bar{\mathbf{n}}_{m_k}(s)) d\bar{W}_2^{m_k}(s), \end{aligned}$$

for any $t \in [0, T]$.

From Proposition 3.22 we see that (\mathbf{v}, \mathbf{n}) is a solution to our problem if we can show that the processes \bar{W}_1 and \bar{W}_2 defined in Proposition 3.13 are Wiener processes and $\mathcal{M}^1(t)$, $\mathcal{M}^2(t)$ are stochastic integrals with respect to \bar{W}_1 and \bar{W}_2 with integrands $S(\mathbf{v}(t))$ and $G(\mathbf{n}(t))$, respectively. These will be the subjects of the following two propositions.

Proposition 3.23. *We have the following facts:*

- (1) *the stochastic process $(\bar{W}_1(t))_{t \in [0, T]}$ (resp. $(\bar{W}_2(t))_{t \in [0, T]}$) is a K_1 -cylindrical K_2 -valued Wiener process (resp. \mathbb{R} -valued standard Brownian motion) on $(\Omega', \mathcal{F}', \mathbb{P}')$.*

- (2) For any s and t such that $0 \leq s < t \leq T$, the increments $\bar{W}_1(t) - \bar{W}_1(s)$ and $\bar{W}_2(t) - \bar{W}_2(s)$ are independent of the σ -algebra generated by $\mathbf{v}(r)$, $\mathbf{n}(r)$, $\bar{W}_1(r)$, $\bar{W}_2(r)$ for $r \in [0, s]$.
- (3) Finally, \bar{W}_1 and \bar{W}_2 are mutually independent.

Proof. We will just establish the proposition for \bar{W}_1 , the same method applies to \bar{W}_2 . To this end we closely follow [7], but see also [44, Lemma 9.9] for an alternative proof.

Proof of item (1). By Proposition 3.13 the laws of $(\mathbf{v}_{m_k}, \mathbf{n}_{m_k}, W_1, W_2)$ are equal to those of the stochastic process $(\bar{\mathbf{v}}_{m_k}, \bar{\mathbf{n}}_{m_k}, \bar{W}_1^{m_k}, \bar{W}_2^{m_k})$ on \mathfrak{S} . Hence it is easy to check that $\bar{W}_1^{m_k}$ (resp. $\bar{W}_2^{m_k}$) form a sequence of K_1 -cylindrical K_2 -valued Wiener process (resp. \mathbb{R} -valued Wiener process). Moreover, for $0 \leq s < t \leq T$ the increments $\bar{W}_1^{m_k}(t) - \bar{W}_1^{m_k}(s)$ (resp. $\bar{W}_2^{m_k}(t) - \bar{W}_2^{m_k}(s)$) are independent of the σ -algebra generated by the stochastic process $(\bar{\mathbf{v}}_{m_k}(r), \bar{\mathbf{n}}_{m_k}(r), \bar{W}_1^{m_k}(r), \bar{W}_2^{m_k}(r))$, for $r \in [0, s]$.

Now, we will check that \bar{W}_1 is a K_1 -cylindrical K_2 -valued Wiener process by showing that the characteristic function of its finite dimensional distributions is equal to the characteristic function of a Gaussian random variable. For this purpose let $k \in \mathbb{N}$ and $s_0 = 0 < s_1 < \dots < s_k \leq T$ be a partition of $[0, T]$. For each $\mathbf{u} \in K_2^*$ we have

$$\mathbb{E}' \left[e^{i \sum_{j=1}^k \langle \mathbf{u}, \bar{W}_1^{m_k}(s_j) - \bar{W}_1^{m_k}(s_{j-1}) \rangle_{K_2}} \right] = e^{-\frac{1}{2} \sum_{j=1}^k (s_j - s_{j-1}) |\mathbf{u}|_{K_1}^2},$$

where $i^2 = -1$. Thanks to (3.33) and the Lebesgue Dominated Convergence Theorem, we have

$$\begin{aligned} \lim_{m \rightarrow \infty} \mathbb{E}' \left[e^{i \sum_{j=1}^k \langle \mathbf{u}, \bar{W}_1^{m_k}(s_j) - \bar{W}_1^{m_k}(s_{j-1}) \rangle_{K_2}} \right] &= \mathbb{E}' \left[e^{i \sum_{j=1}^k \langle \mathbf{u}, \bar{W}_1(s_j) - \bar{W}_1(s_{j-1}) \rangle_{K_2}} \right] \\ &= e^{-\frac{1}{2} \sum_{j=1}^k (s_j - s_{j-1}) |\mathbf{u}|_{K_1}^2} \end{aligned}$$

from which we infer that the finite dimensional distributions of \bar{W}_1 follow a Gaussian distribution. The same idea can be carried out to prove that the finite dimensional distributions of \bar{W}_2 are Gaussian.

Proof of item (2). Next, we prove that the increments $\bar{W}_1(t) - \bar{W}_1(s)$ and $\bar{W}_2(t) - \bar{W}_2(s)$, $0 \leq s < t \leq T$ are independent of the σ -algebra generated by $(\mathbf{v}(r), \mathbf{n}(r), \bar{W}_1(r), \bar{W}_2(r))$ for $r \in [0, s]$. To this end, let us consider $\{\phi_j : j = 1, \dots, k\} \subset C_b(\mathbb{V}^{-\beta} \times \mathbb{H}^1)$ and $\{\psi_j : j = 1, \dots, k\} \subset C_b(K_2 \times \mathbb{R})$, where for any Banach space \mathbf{B} the space $C_b(\mathbf{B})$ is defined

$$C_b(\mathbf{B}) = \{\phi : \mathbf{B} \rightarrow \mathbb{R}, \phi \text{ is continuous and bounded}\}.$$

Also, let $0 \leq r_1 < \dots < r_k \leq s < t \leq T$, $\psi \in C_b(K_2)$, and $\zeta \in C_b(\mathbb{R})$. For each $m \in \mathbb{M}$, there holds

$$\begin{aligned} &\mathbb{E}' \left[\left(\prod_{j=1}^k \phi_j(\bar{\mathbf{v}}_{m_k}(r_j), \bar{\mathbf{n}}_{m_k}(r_j)) \prod_{j=1}^k \psi_j(\bar{W}_1^{m_k}(r_j), \bar{W}_2^{m_k}(r_j)) \right) \right. \\ &\quad \left. \times \psi(\bar{W}_1^{m_k}(t) - \bar{W}_1^{m_k}(s)) \zeta(\bar{W}_2^{m_k}(t) - \bar{W}_2^{m_k}(s)) \right] \\ &= \mathbb{E}' \left[\prod_{j=1}^k \phi_j(\bar{\mathbf{v}}_{m_k}(r_j), \bar{\mathbf{n}}_{m_k}(r_j)) \prod_{j=1}^k \psi_j(\bar{W}_1^{m_k}(r_j), \bar{W}_2^{m_k}(r_j)) \right] \\ &\quad \times \mathbb{E}'(\zeta(\bar{W}_1^{m_k}(t) - \bar{W}_1^{m_k}(s))) \mathbb{E}'(\psi(\bar{W}_2^{m_k}(t) - \bar{W}_2^{m_k}(s))). \end{aligned}$$

Thanks to (3.30), (3.32), (3.33), (3.34) and the Lebesgue Dominated Convergence Theorem, the same identity is true with $(\mathbf{v}, \mathbf{n}, \bar{W}_1, \bar{W}_2)$ in place of $(\bar{\mathbf{v}}_{m_k}, \bar{\mathbf{n}}_{m_k}, \bar{W}_1^{m_k}, \bar{W}_2^{m_k})$. This completes the proof of the second item of the proposition.

Proof of item (3). One can argue as in the proof of the first part of the proposition, thus we omit the details. □

Proposition 3.24. *For each $t \in (0, T]$ we have*

$$\mathcal{M}^1(t) = \int_0^t S(\mathbf{v}(s)) d\bar{W}_1(s) \text{ in } L^2(\Omega', \mathbb{V}^*), \quad (3.71)$$

$$\mathcal{M}^2(t) = \int_0^t (\mathbf{n}(s) \times h) d\bar{W}_2(s) \text{ in } L^2(\Omega, \mathbb{X}_{\beta-\frac{1}{2}}). \quad (3.72)$$

Proof. We will only prove the identity (3.71) since the same argument can be used to establish (3.72), but see also [7] for an alternative proof.

Let us fixed $t \in (0, T]$ and for any $\varepsilon > 0$ let $\eta_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ a standard mollifier with support in $(0, t)$. For $R \in \{S, S_{m_k}\}$, $\mathbf{u} \in \{\bar{\mathbf{v}}_{m_k}, \mathbf{v}\}$ and $s \in (0, t]$ let us set

$$\begin{aligned} R^\varepsilon(\mathbf{u}(s)) &= (\eta_\varepsilon \star R(\mathbf{u}(\cdot)))(s) \\ &= \int_{-\infty}^{\infty} \eta_\varepsilon(s-r) R(\mathbf{u}(r)) dr. \end{aligned}$$

We recall that, since R is Lipschitz, R^ε is Lipschitz. We also have the following two important facts, see, for instance, [2, Section 1.3]:

(a) for any $p \in [1, \infty)$ there exists a constant $C > 0$ such that for any $\varepsilon > 0$ we have

$$\int_0^t \|R^\varepsilon(\mathbf{u}(s))\|_{\mathcal{T}_2(\mathbb{K}_1, \mathbb{H})}^p ds \leq C \int_0^t \|R(\mathbf{u}(s))\|_{\mathcal{T}_2(\mathbb{K}_1, \mathbb{H})}^p ds. \quad (3.73)$$

(b) For any $p \in [1, \infty)$, we have

$$\lim_{\varepsilon \rightarrow 0} \int_0^t \|R^\varepsilon(\mathbf{u}(s)) - R(\mathbf{u}(s))\|_{\mathcal{T}_2(\mathbb{K}_1, \mathbb{H})}^p ds = 0. \quad (3.74)$$

Now, let $\mathcal{M}_{m_k}^\varepsilon$ and \mathcal{M}^ε be respectively defined by

$$\begin{aligned} \mathcal{M}_{m_k}^\varepsilon(t) &= \int_0^t S_{m_k}^\varepsilon(\bar{\mathbf{v}}_{m_k}(s)) d\bar{W}_1^{m_k}(s), \\ \mathcal{M}^\varepsilon(t) &= \int_0^t S^\varepsilon(\mathbf{v}(s)) d\bar{W}_1(s), \end{aligned}$$

From the Itô isometry, (3.73) and some elementary calculations we infer that there exists a constant $C > 0$ such that for any $\varepsilon > 0$ and $m_k \in \mathbb{N}$

$$\begin{aligned} \mathbb{E}' \|\mathcal{M}_{m_k}(t) - \mathcal{M}_{m_k}^\varepsilon(t)\|^2 &= \mathbb{E}' \int_0^t \|S(\bar{\mathbf{v}}_{m_k}(s)) - S_{m_k}^\varepsilon(\bar{\mathbf{v}}_{m_k}(s))\|_{\mathcal{T}_2(\mathbb{K}_1, \mathbb{H})}^2 ds, \\ &\leq C \mathbb{E}' \int_0^t \|S(\bar{\mathbf{v}}_{m_k}(s)) - S(\mathbf{v}(s))\|_{\mathcal{T}_2(\mathbb{K}_1, \mathbb{H})}^2 ds \end{aligned} \quad (3.75)$$

$$+ C \mathbb{E}' \int_0^t \|S(\mathbf{v}(s)) - S^\varepsilon(\mathbf{v}(s))\|_{\mathcal{T}_2(\mathbb{K}_1, \mathbb{H})}^2 ds. \quad (3.76)$$

From Assumption 3.1 and (3.42) we derive that the first term of in the right hand side of the last estimate converges to 0 as $m_k \rightarrow \infty$. Owing to (3.73) and (3.52) the sequence in the second term of (3.76) is uniformly integrable with respect to the probability measure \mathbb{P}' . Thus, from (3.74) and the Vitali Convergence Theorem we infer that

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E}' \int_0^t \|S(\mathbf{v}(s)) - S^\varepsilon(\mathbf{v}(s))\|_{\mathcal{T}_2(\mathbb{K}_1, \mathbb{H})}^2 ds = 0.$$

Hence, for any $t \in (0, T]$

$$\lim_{\varepsilon \rightarrow 0} \lim_{m_k \rightarrow \infty} \mathbb{E}' \|\mathcal{M}_{m_k}(t) - \mathcal{M}_{m_k}^\varepsilon(t)\|^2 = 0. \quad (3.77)$$

In a similar way, we can prove that

$$\lim_{\varepsilon \rightarrow 0} \lim_{m_k \rightarrow \infty} \mathbb{E}' \|\mathcal{M}^2(t) - \mathcal{M}^\varepsilon(t)\|^2 = 0. \quad (3.78)$$

Next, we will prove that

$$\lim_{\varepsilon \rightarrow 0} \lim_{m_k \rightarrow \infty} \mathbb{E}' \|\mathcal{M}_{m_k}^\varepsilon(t) - \mathcal{M}^\varepsilon(t)\|^2 = 0. \quad (3.79)$$

To this end, we first observe that

$$\begin{aligned} \mathcal{M}_{m_k}^\varepsilon(t) - \mathcal{M}^\varepsilon(t) &= \int_0^t S_{m_k}^\varepsilon(\bar{\mathbf{v}}_{m_k}(s)) \bar{W}_1^{m_k}(s) - \int_0^t S_{m_k}^\varepsilon(\mathbf{v}(s)) d\bar{W}_1(s) \\ &\quad + \int_0^t S_{m_k}^\varepsilon(\mathbf{v}(s)) d\bar{W}_1(s) - \int_0^t S^\varepsilon(\mathbf{v}(s)) d\bar{W}_1(s) \\ &= I_{m_k,1}^\varepsilon + I_{m_k,2}^\varepsilon. \end{aligned}$$

Second, by integration by parts we derive that

$$\begin{aligned} I_{m_k,1}^\varepsilon &= \int_0^t [\eta'_\varepsilon \star S_{m_k}(\mathbf{v}(\cdot))](s) \bar{W}_1(s) ds - \int_0^t [\eta'_\varepsilon \star S_{m_k}(\bar{\mathbf{v}}_{m_k}(\cdot))](s) \bar{W}_1^{m_k}(s) ds \\ &= \int_0^t [\eta'_\varepsilon \star S_{m_k}(\bar{\mathbf{v}}_{m_k}(\cdot))](s) [\bar{W}_1^{m_k}(s) - \bar{W}_1(s)] ds + \int_0^t [S_{m_k}^\varepsilon(\bar{\mathbf{v}}_{m_k}(s)) - S_{m_k}^\varepsilon(\mathbf{v}(s))] d\bar{W}_1(s) \\ &= J_{m_k,1}^\varepsilon + J_{m_k,2}^\varepsilon. \end{aligned}$$

On one hand, by Proposition 3.23 the processes $\bar{W}_1^{m_k}$ and \bar{W}_1 are both K_1 -cylindrical K_2 -valued Wiener processes, thus, for any integer $p \geq 4$ there exists a constant $C > 0$ such that

$$\sup_{m_k \in \mathbb{N}} \mathbb{E}' \sup_{s \in [0, T]} (\|\bar{W}_1^{m_k}(s)\|_{K_2}^p + \|\bar{W}_1(s)\|_{K_2}^p) \leq CQT^{\frac{p}{2}}.$$

Hence, the sequence $\int_0^t \|\bar{W}_1^{m_k}(s) - \bar{W}_1(s)\|_{K_2}^2 ds$ is uniformly integrable with respect to the probability measure \mathbb{P}' , and from (3.33) and the Vitali Convergence Theorem we infer that

$$\lim_{m_k \rightarrow \infty} \mathbb{E}' \int_0^t \|\bar{W}_1^{m_k}(s) - \bar{W}_1(s)\|_{K_2}^2 ds = 0.$$

On the other hand, for any $\varepsilon > 0$ there exists a constant $C(\varepsilon)$ such that

$$\mathbb{E}' \int_0^t \|[\eta'_\varepsilon \star S_{m_k}(\bar{\mathbf{v}}_{m_k}(\cdot))](s)\|_{\mathcal{H}_2(K_1, \mathbb{H})}^2 ds \leq C(\varepsilon) T \mathbb{E}' \sup_{t \in [0, T]} \|S_{m_k}(\bar{\mathbf{v}}_{m_k}(t))\|_{\mathcal{H}_2(K_1, \mathbb{H})}^2,$$

from which along with Assumption 3.1 and (3.35) we infer that for any $\varepsilon > 0$ there exists a constant $C > 0$ such that for any $m_k \in \mathbb{N}$ we have

$$\mathbb{E}' \int_0^t \|[\eta'_\varepsilon \star S_{m_k}(\bar{\mathbf{v}}_{m_k}(\cdot))](s)\|_{\mathcal{H}_2(K_1, \mathbb{H})}^2 ds \leq C(\varepsilon) T.$$

Thus, from these two observation we derive that

$$\lim_{\varepsilon \rightarrow 0} \lim_{m_k \rightarrow \infty} \mathbb{E}' \|J_{m_k,1}^\varepsilon\|^2 = 0.$$

In a straightforward way we can also show that

$$\lim_{\varepsilon \rightarrow 0} \lim_{m_k \rightarrow \infty} (\mathbb{E}' \|J_{m_k,2}^\varepsilon\|^2 + \mathbb{E}' \|I_{m_k,2}^\varepsilon\|^2) = 0.$$

Therefore, we have just proved that

$$\lim_{\varepsilon \rightarrow 0} \lim_{m_k \rightarrow \infty} \mathbb{E}' \|\mathcal{M}_{m_k}^\varepsilon(t) - \mathcal{M}^\varepsilon(t)\|^2 = 0. \quad (3.80)$$

The identities (3.77), (3.78) and (3.80) imply that for any $t \in (0, T]$

$$\lim_{m_k \rightarrow \infty} \mathbb{E}' \|\mathcal{M}_{m_k}^1(t) - \mathcal{M}^1(t)\|^2 = 0. \quad (3.81)$$

To conclude the proof of the proposition we need to show that \mathbb{P}' -a.s.

$$\mathcal{M}_{m_k}^1(t) - \int_0^t S_{m_k}(\bar{\mathbf{v}}_{m_k}(s)) d\bar{W}_1^{m_k}(s) = 0, \quad (3.82)$$

for any $t \in (0, T]$. To this end, let \mathcal{M}_m^1 and $\mathcal{M}_m^\varepsilon$ be the analog of $\mathcal{M}_{m_k}^1$ and $\mathcal{M}_{m_k}^\varepsilon$ with m_k and $\bar{\mathbf{v}}_{m_k}$ replaced by m and $\bar{\mathbf{v}}_m$, respectively. For any $\mathbf{u} \in L^2(0, T; \mathbb{V}^*)$ we set

$$\varphi(\mathbf{u}) = \frac{\int_0^T \|\mathbf{u}(s)\|_{\mathbb{V}^*}^2 ds}{1 + \int_0^T \|\mathbf{u}(s)\|_{\mathbb{V}^*}^2 ds}.$$

Since $(\bar{\mathbf{v}}_{m_k}, \bar{\mathbf{n}}_{m_k}, \bar{W}_1^{m_k})$ and $(\bar{\mathbf{v}}_m, \bar{\mathbf{n}}_m, W_1)$ have the same law and $\varphi(\cdot)$ is continuous as a map from $\mathfrak{S}_1 \times \mathfrak{S}_2 \times C([0, T]; K_1)$ into \mathbb{R} , we infer that

$$\mathbb{E}\varphi(\mathcal{M}_m^1 - \mathcal{M}_m^\varepsilon) = \mathbb{E}'\varphi(\mathcal{M}_{m_k}^1 - \mathcal{M}_{m_k}^\varepsilon).$$

Note that arguing as above we can show that as $\varepsilon \rightarrow 0$ we have

$$\mathbb{E}\varphi(\mathcal{M}_m^1 - \mathfrak{M}_m^1) = \mathbb{E}'\varphi(\mathcal{M}_{m_k}^1 - \mathfrak{M}_{m_k}^1),$$

where

$$\mathfrak{M}_m^1(\cdot) = \int_0^\cdot S_m(\bar{\mathbf{v}}_m(s)) dW_1(s).$$

Since $\bar{\mathbf{v}}_m$ and $\bar{\mathbf{n}}_m$ are the solution of the Galerkin approximation, we have \mathbb{P} -a.s. $\varphi(\mathcal{M}_m^1 - \mathfrak{M}_m^1) = 0$, from which we infer that

$$\mathbb{E}'\varphi(\mathcal{M}_{m_k}^1 - \mathcal{M}_{m_k}) = 0.$$

This last identity implies that \mathbb{P}' -a.s. $\mathcal{M}_{m_k}^1(t) - \mathcal{M}_{m_k}(t) = 0$ for almost everywhere $t \in (0, T]$. Since the maps $\mathcal{M}_{m_k}^1(\cdot)$ $\mathcal{M}_{m_k}(\cdot)$ are continuous in \mathbb{V}^* and agree for almost everywhere $t \in (0, T]$, necessarily they agree for all $t \in (0, T]$. Thus, we have proved the identity (3.82) which along with (3.81) implies the desired equality (3.65). \square

Now we give the promised proof of the existence of weak martingale solution.

Proof of Theorem 3.2. Endowing the complete probability space $(\Omega', \mathcal{F}', \mathbb{P}')$ with the filtration $\mathbb{F}' = (\mathcal{F}'_t)_{t \geq 0}$ which satisfies the usual condition, and combining Propositions 3.22, 3.23 and 3.24 we have just constructed a complete filtered probability space and stochastic processes $\mathbf{v}(t), \mathbf{n}(t), \bar{W}_1(t), \bar{W}_2(t)$ which satisfy all the items of Definition 3.1. \square

3.4. Proof of the pathwise uniqueness of the weak solution in the 2-D case. This subsection is devoted to the proof of the uniqueness stated in Theorem 3.4. Before proceeding to the the actual proof of this pathwise uniqueness, we state and prove the following lemma.

Lemma 3.25. *For any $\alpha_8 > 0$ and $\alpha_9 > 0$ there exist $C(\alpha_8) > 0$, $C_1(\alpha_9) > 0$ and $C_2(\alpha_9) > 0$ such that*

$$|\langle f(\mathbf{n}_1) - f(\mathbf{n}_2), \mathbf{n}_1 - \mathbf{n}_2 \rangle| \leq \alpha_8 \|\nabla \mathbf{n}_1 - \nabla \mathbf{n}_2\|^2 + C(\alpha_8) \|\mathbf{n}_1 - \mathbf{n}_2\|^2 \varphi(\mathbf{n}_1, \mathbf{n}_2), \quad (3.83)$$

$$\begin{aligned} |\langle f(\mathbf{n}_1) - f(\mathbf{n}_2), \Delta \mathbf{n}_1 - \Delta \mathbf{n}_2 \rangle| &\leq \alpha_9 \|\Delta \mathbf{n}_1 - \Delta \mathbf{n}_2\|^2 + C_1(\alpha_9) \|\nabla \mathbf{n}_1 - \nabla \mathbf{n}_2\|^2 \varphi(\mathbf{n}_1, \mathbf{n}_2) \\ &\quad + C_2(\alpha_9) \|\mathbf{n}_1 - \mathbf{n}_2\|^2 \varphi(\mathbf{n}_1, \mathbf{n}_2), \end{aligned} \quad (3.84)$$

where

$$\varphi(\mathbf{n}_1, \mathbf{n}_2) := C \left(1 + \|\mathbf{n}_1\|_{\mathbb{L}^{4N+2}}^{2N} + \|\mathbf{n}_2\|_{\mathbb{L}^{4N+2}}^{2N} \right)^2.$$

Proof of Lemma 3.25. It is enough to prove the estimate (3.83) for the particular case $f(\mathbf{n}) := a_N |\mathbf{n}|^{2N} \mathbf{n}$. For this purpose we recall that

$$|\mathbf{n}_1|^{2N} \mathbf{n}_1 - |\mathbf{n}_2|^{2N} \mathbf{n}_2 = |\mathbf{n}_1|^{2N} (\mathbf{n}_1 - \mathbf{n}_2) + \mathbf{n}_2 (|\mathbf{n}_1| - |\mathbf{n}_2|) \left(\sum_{k=0}^{2N-1} |\mathbf{n}_1|^{2N-k-1} |\mathbf{n}_2|^k \right),$$

from which we easily deduce that

$$|\langle f(\mathbf{n}_1) - f(\mathbf{n}_2), \mathbf{n}_1 - \mathbf{n}_2 \rangle| \leq C \int_{\mathcal{O}} (1 + |\mathbf{n}_1|^{2N} + |\mathbf{n}_2|^{2N}) |\mathbf{n}_1 - \mathbf{n}_2|^2 dx,$$

for any $\mathbf{n}_1, \mathbf{n}_2 \in \mathbb{L}^{2N+2}(\mathcal{O})$. Now, invoking the Hölder, Gagliardo-Nirenberg and Young inequalities we infer that

$$\begin{aligned} |\langle f(\mathbf{n}_1) - f(\mathbf{n}_2), \mathbf{n}_1 - \mathbf{n}_2 \rangle| &\leq C \|\mathbf{n}_1 - \mathbf{n}_2\|_{\mathbb{L}^4}^2 (1 + \|\mathbf{n}_1\|_{\mathbb{L}^{4N+2}}^{2N} + \|\mathbf{n}_2\|_{\mathbb{L}^{4N+2}}^{2N}) \\ &\leq C \|\mathbf{n}_1 - \mathbf{n}_2\| \|\nabla(\mathbf{n}_1 - \mathbf{n}_2)\| (1 + \|\mathbf{n}_1\|_{\mathbb{L}^{4N+2}}^{2N} + \|\mathbf{n}_2\|_{\mathbb{L}^{4N+2}}^{2N}) \\ &\leq \alpha_8 \|\nabla(\mathbf{n}_1 - \mathbf{n}_2)\|^2 + C(\alpha_8) \|\mathbf{n}_1 - \mathbf{n}_2\|^2 (1 + \|\mathbf{n}_1\|_{\mathbb{L}^{4N+2}}^{2N} + \|\mathbf{n}_2\|_{\mathbb{L}^{4N+2}}^{2N})^2. \end{aligned}$$

The last line of the above chain of inequalities implies (3.83).

Using the fact that $\mathbb{H}^1 \subset \mathbb{L}^{4N+2}$ for any $N \in \mathbb{N}$ and the same argument as in the proof of (3.83) we derive that

$$\begin{aligned} |\langle f(\mathbf{n}_1) - f(\mathbf{n}_2), \Delta \mathbf{n}_1 - \Delta \mathbf{n}_2 \rangle| &\leq C \int_{\mathcal{O}} (1 + |\mathbf{n}_1|^{2N} + |\mathbf{n}_2|^{2N}) |\mathbf{n}_1 - \mathbf{n}_2| |\Delta(\mathbf{n}_1 - \mathbf{n}_2)| dx \\ &\leq C \|\mathbf{n}_1 - \mathbf{n}_2\|_{\mathbb{L}^{4N+2}} \|\Delta[\mathbf{n}_1 - \mathbf{n}_2]\| (1 + \|\mathbf{n}_1\|_{\mathbb{L}^{4N+2}}^{2N} + \|\mathbf{n}_2\|_{\mathbb{L}^{4N+2}}^{2N}) \\ &\leq C \|\mathbf{n}_1 - \mathbf{n}_2\|_{\mathbb{H}^1} \|\Delta[\mathbf{n}_1 - \mathbf{n}_2]\| (1 + \|\mathbf{n}_1\|_{\mathbb{L}^{4N+2}}^{2N} + \|\mathbf{n}_2\|_{\mathbb{L}^{4N+2}}^{2N}) \\ &\leq \alpha_9 \|\Delta[\mathbf{n}_1 - \mathbf{n}_2]\|^2 + C(\alpha_9) \|\mathbf{n}_1 - \mathbf{n}_2\|_{\mathbb{H}^1}^2 (1 + \|\mathbf{n}_1\|_{\mathbb{L}^{4N+2}}^{2N} + \|\mathbf{n}_2\|_{\mathbb{L}^{4N+2}}^{2N})^2. \end{aligned}$$

From the last line we easily deduce the proof of (3.84). \square

Now, we give the promised proof of the uniqueness of our solution.

Proof of Theorem 3.4. Let $\mathbf{v} = \mathbf{v}_1 - \mathbf{v}_2$ and $\mathbf{n} = \mathbf{n}_1 - \mathbf{n}_2$. These processes satisfy $(\mathbf{v}(0), \mathbf{n}(0)) = (0, 0)$ and the stochastic equations

$$\begin{aligned} d\mathbf{v}(t) + \left(A\mathbf{v}(t) + B(\mathbf{v}(t), \mathbf{v}_1(t)) + B(\mathbf{v}_2(t), \mathbf{v}(t)) \right) dt &= - \left(M(\mathbf{n}(t), \mathbf{n}_1(t)) + M(\mathbf{n}_2, \mathbf{n}) \right) dt \\ &\quad + [S(\mathbf{v}_1(t)) - S_2(\mathbf{v}_2(t))] dW_1(t), \end{aligned}$$

and

$$\begin{aligned} d\mathbf{n}(t) + \left(A_1\mathbf{n}(t) + \tilde{B}(\mathbf{v}(t), \mathbf{n}_1(t)) + \tilde{B}(\mathbf{v}_2(t), \mathbf{n}(t)) \right) dt &= -[f(\mathbf{n}_2(t)) - f(\mathbf{n}_1(t))] dt \\ &\quad + \frac{1}{2} G^2(\mathbf{n}(t)) dt + G(\mathbf{n}(t)) dW_2(t). \end{aligned}$$

Firstly, recall that from Young's inequality, (A.8), (2.10) and (A.9) we can derive that for any $\alpha_1 > 0$, $\alpha_2 > 0$, $\alpha_3 > 0$, $\alpha_4 > 0$, $\alpha_5 > 0$ and $\alpha_7 > 0$ there exist $C(\alpha_1) > 0$, $C(\alpha_2, \alpha_3) > 0$, $C(\alpha_7, \alpha_4) > 0$ and $C(\alpha_5) > 0$ such that

$$\begin{aligned} |\langle B(\mathbf{v}, \mathbf{v}_1), \mathbf{v} \rangle| &\leq \alpha_1 \|\nabla \mathbf{v}\|^2 + C(\alpha_1) \|\mathbf{v}_1\|^2 \|\nabla \mathbf{v}_1\|^2 \|\mathbf{v}\|^2, \\ |\langle M(\mathbf{n}_2, \mathbf{n}), \mathbf{v} \rangle| &\leq \alpha_2 \|\nabla \mathbf{v}\|^2 + \alpha_3 \|\Delta \mathbf{n}\|^2 + C(\alpha_2, \alpha_3) \|\nabla \mathbf{n}_2\|^2 \|\Delta \mathbf{n}_2\|^2 \|\nabla \mathbf{n}\|^2, \\ |\langle M(\mathbf{n}, \mathbf{n}_1), \mathbf{v} \rangle| &\leq \alpha_7 \|\nabla \mathbf{v}\|^2 + \alpha_4 \|\Delta \mathbf{n}\|^2 + C(\alpha_7, \alpha_4) \|\nabla \mathbf{n}_1\|^2 \|\Delta \mathbf{n}_1\|^2 \|\nabla \mathbf{n}\|^2, \\ |\langle \tilde{B}(\mathbf{v}_2, \mathbf{n}), \Delta \mathbf{n} \rangle| &\leq \alpha_5 \|\Delta \mathbf{n}\|^2 + C(\alpha_5) \|\mathbf{v}_2\|^2 \|\nabla \mathbf{v}_2\|^2 \|\nabla \mathbf{n}\|^2. \end{aligned} \tag{3.85}$$

From Hölder's inequality, Gagliardo-Nirenberg's inequality (A.1) and the Sobolev embedding $\mathbb{H}^2 \subset \mathbb{L}^\infty$ we infer that for any $\alpha_6 > 0$ there exists $C(\alpha_6) > 0$ such that

$$\begin{aligned} |\langle \tilde{B}(\mathbf{v}, \mathbf{n}_1), \mathbf{n} \rangle| &\leq \|\mathbf{v}\| \|\nabla \mathbf{n}_1\| \|\mathbf{n}\|_{\mathbb{L}^\infty}, \\ &\leq \alpha_6 \|\Delta \mathbf{n}\|^2 + C(\alpha_6) \|\mathbf{v}\|^2 \|\nabla \mathbf{n}_1\|^2. \end{aligned}$$

From the proof of Proposition C.2 we see that there exists a constant $C(h) > 0$ such that

$$\begin{aligned} \|\nabla G(\mathbf{n})\|^2 &\leq C(h) (\|\nabla \mathbf{n}\|^2 + \|\mathbf{n}\|^2), \\ \|\nabla G^2(\mathbf{n})\|^2 &\leq C(h) (\|\nabla \mathbf{n}\|^2 + \|\mathbf{n}\|^2), \\ |\langle \nabla G^2(\mathbf{n}), \nabla \mathbf{n} \rangle| &\leq C(h) (\|\nabla \mathbf{n}\|^2 + \|\mathbf{n}\|^2). \end{aligned}$$

Owing to the Lipschitz property of S we have

$$\|S(\mathbf{v}_1) - S(\mathbf{v}_2)\|_{\mathcal{H}_2}^2 \leq C\|\mathbf{v}\|^2.$$

Now, let $\varphi(\mathbf{n}_1, \mathbf{n}_2)$ be as in Lemma 3.25 and

$$\Psi(t) = e^{-\int_0^t (\psi_1(s) + \psi_2(s) + \psi_3(s)) ds}, \text{ for any } t > 0,$$

where

$$\begin{aligned} \psi_1(s) &:= C(\alpha_1)\|\mathbf{v}_1(s)\|^2\|\nabla\mathbf{v}_1(s)\|^2 + C(\alpha_6)\|\nabla\mathbf{n}_1(s)\|^2, \\ \psi_3(s) &:= [C(\alpha_8) + C_2(\alpha_9)]\varphi(\mathbf{n}_1(s), \mathbf{n}_2(s)), \end{aligned}$$

and

$$\begin{aligned} \psi_2(s) &:= C(\alpha_2, \alpha_3)\|\nabla\mathbf{n}_2(s)\|^2\|\Delta\mathbf{n}_2(s)\|^2 + C(\alpha_7, \alpha_4)\|\nabla\mathbf{n}_1(s)\|^2\|\Delta\mathbf{n}_1(s)\|^2 \\ &\quad + C(\alpha_5)\|\mathbf{v}_2(s)\|^2\|\nabla\mathbf{v}_2(s)\|^2 + C_1(\alpha_9)\varphi(\mathbf{n}_1(s), \mathbf{n}_2(s)). \end{aligned}$$

Now applying Itô's formula to $\|\mathbf{n}(t)\|^2$ (as in proof of Proposition C.1) and $\Psi(t)\|\mathbf{n}(t)\|^2$ yield

$$\begin{aligned} d[\Psi(t)\|\mathbf{n}(t)\|^2] &= -2\Psi(t)\|\nabla\mathbf{n}(t)\|^2 dt - 2\Psi(t)\langle\tilde{B}(\mathbf{v}(t), \mathbf{n}_1(t)), \mathbf{n}(t)\rangle \\ &\quad - 2\langle f(\mathbf{n}_2(t)) - f(\mathbf{n}_1(t)), \mathbf{n}(t)\rangle dt + \Psi'(t)\|\mathbf{n}(t)\|^2. \end{aligned}$$

Using the same argument we can show that $\Psi(t)\|\nabla\mathbf{n}(t)\|^2$ and $\Psi(t)\|\mathbf{v}(t)\|^2$ satisfy

$$\begin{aligned} d[\Psi(t)\|\nabla\mathbf{n}(t)\|^2] &= \Psi(t)\left(-\|\Delta\mathbf{n}(t)\|^2 + \langle\tilde{B}(\mathbf{v}(t), \mathbf{n}_1(t)) + \tilde{B}(\mathbf{v}_2(t), \mathbf{n}(t)), \Delta\mathbf{n}(t)\rangle\right) dt \\ &\quad + \Psi(t)\left(2\langle f(\mathbf{n}_2(t)) - f(\mathbf{n}_1(t)), \Delta\mathbf{n}(t)\rangle + \langle\nabla G^2(\mathbf{n}(t)), \nabla\mathbf{n}(t)\rangle\right) dt \\ &\quad + \|G(\mathbf{n}(t))\|^2 dt + \Psi'(t)\|\nabla\mathbf{n}(t)\|^2 dt + 2\Psi(t)\langle\nabla G(\mathbf{n}(t)), \nabla\mathbf{n}(t)\rangle dW_2(t), \end{aligned}$$

and

$$\begin{aligned} d[\Psi(t)\|\mathbf{v}(t)\|^2] &= -2\Psi(t)\left(\|\nabla\mathbf{v}(t)\|^2 + \langle B(\mathbf{v}(t), \mathbf{v}_1(t)) + M(\mathbf{n}(t), \mathbf{n}_1(t)), \mathbf{v}(t)\rangle\right) dt \\ &\quad - 2\Psi(t)\langle M(\mathbf{n}_2(t), \mathbf{n}(t)), \mathbf{v}(t)\rangle dt + \Psi(t)\|S(\mathbf{v}_1(t)) - S(\mathbf{v}_2(t))\|_{\mathcal{H}_2}^2 dt + \Psi'(t)\|\mathbf{v}(t)\|^2 dt \\ &\quad + 2\Psi(t)\langle\mathbf{v}(t), [S(\mathbf{v}_1(t)) - S(\mathbf{v}_2(t))]\rangle dW_1(t). \end{aligned}$$

Summing up these last three equalities side by side and using the inequalities (3.85)-(3.84) imply

$$\begin{aligned} d[\Psi(t)(\|\mathbf{v}(t)\|^2 + \|\mathbf{n}(t)\|^2 + \|\nabla\mathbf{n}(t)\|^2)] &+ 2\Psi(t)\left[\|\nabla\mathbf{v}(t)\|^2 + \|\nabla\mathbf{n}(t)\|^2 + \|\Delta\mathbf{n}(t)\|^2\right] dt \\ &\leq 2\Psi(t)\left(C\left[\|\mathbf{v}(t)\|^2 + \|\mathbf{n}(t)\|^2 + \|\nabla\mathbf{n}(t)\|^2\right] dt + \langle\nabla G(\mathbf{n}(t)), \nabla\mathbf{n}(t)\rangle dW_2(t)\right) \\ &\quad + 2\Psi(t)\left(\langle\mathbf{v}(t), [S(\mathbf{v}_1(t)) - S(\mathbf{v}_2(t))]\rangle dW_1(t) + [\alpha_9 + \sum_{j=3}^6 \alpha_j]\|\Delta\mathbf{n}(t)\|^2\right) \\ &\quad + \Psi(t)\left[\psi_2(t)\|\nabla\mathbf{n}(t)\|^2 + \psi_1(t)\|\mathbf{v}(t)\|^2 + \psi_3(t)\|\mathbf{n}(t)\|^2\right] dt \\ &\quad + \Psi'(t)\left(\|\mathbf{v}(t)\|^2 + \|\mathbf{n}(t)\|^2 + \|\nabla\mathbf{n}(t)\|^2\right) dt \\ &\quad + (\alpha_1 + \alpha_2 + \alpha_7)\|\nabla\mathbf{v}(t)\|^2 + \alpha_8\|\nabla\mathbf{n}(t)\|^2 dt. \end{aligned}$$

Notice that by the choice of Ψ we have

$$\Psi(t)\left[\psi_2(t)\|\nabla\mathbf{n}(t)\|^2 + \psi_1(t)\|\mathbf{v}(t)\|^2 + \psi_3(t)\|\mathbf{n}(t)\|^2\right] + \Psi'(t)\left(\|\mathbf{v}(t)\|^2 + \|\mathbf{n}(t)\|^2 + \|\nabla\mathbf{n}(t)\|^2\right) \leq 0.$$

Hence by choosing $\alpha_j = \alpha_9 = \frac{1}{10}$, $j = 3, \dots, 6$, $\alpha_i = \alpha_7 = \frac{1}{6}$, $i = 2, 3$ and $\alpha_8 = \frac{1}{2}$ we see that

$$\begin{aligned} d[\Psi(t) (\|\mathbf{v}(t)\|^2 + \|\mathbf{n}(t)\|^2 + \|\nabla \mathbf{n}(t)\|^2)] + \Psi(t) [\|\nabla \mathbf{v}(t)\|^2 + \|\Delta \mathbf{n}(t)\|^2 + \|\nabla \mathbf{n}(t)\|^2] dt \\ \leq 2\Psi(t) \left(C \left[\|\mathbf{v}(t)\|^2 + \|\mathbf{n}(t)\|^2 + \|\nabla \mathbf{n}(t)\|^2 \right] dt + \langle \nabla G(\mathbf{n}(t)), \nabla \mathbf{n}(t) \rangle dW_2(t) \right. \\ \left. + \langle \mathbf{v}(t), [S(\mathbf{v}_1(t)) - S(\mathbf{v}_2(t))] \rangle dW_1(t) \right) \end{aligned}$$

Next integrating and taking the mathematical expectation yield

$$\begin{aligned} \mathbb{E} \left[\Psi(t) (\|\mathbf{v}(t)\|^2 + \|\mathbf{n}(t)\|^2 + \|\nabla \mathbf{n}(t)\|^2) \right] + \mathbb{E} \int_0^t \Psi(s) [\|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s)\|^2 + 2\|\nabla \mathbf{n}(s)\|^2] ds \\ \leq C \int_0^t \mathbb{E} \left[\Psi(s) (\|\mathbf{v}(s)\|^2 + \|\mathbf{n}(s)\|^2 + \|\nabla \mathbf{n}(s)\|^2) \right] ds. \end{aligned}$$

Now we can easily conclude the proof by making use of Gronwall's lemma. \square

4. EXISTENCE AND UNIQUENESS OF LOCAL AND GLOBAL STRONG SOLUTION

In this section we are interested in the strong solution to the system (2.12)-(2.13). Here strong solution is both in PDEs and Stochastic Calculus sense. One of the main result, the existence of a local maximal solution, of this section is a corollary of a general framework that will be introduced in Section 5. In this section we will check that the system (2.12)-(2.13) fits into this general framework and hence establish the existence of local maximal solution of the stochastic nematic liquid crystal. We also prove that the maximal solution turns out to be a global one in the two dimensional case. For these ends we will introduce some additional notations and prove several key inequalities related to the nonlinear terms of the stochastic system (2.12)-(2.13).

Hereafter, we put

$$H = \mathbb{H} \times \mathbb{X}_0, \quad V = \mathbb{V}_{\frac{1}{2}} \times \mathbb{X}_{\frac{1}{2}} \text{ and } E = \mathbb{V}_1 \times \mathbb{X}_1. \quad (4.1)$$

Next we denote by $\{S_1(t)\}_{t \geq 0}$ the analytic semigroup generated by $-A$ on \mathbb{H} where A is the Stokes operator. The operator $-A_1$ is the generator of an analytic semigroup $\{\mathbb{T}(t)\}_{t \geq 0}$ on \mathbb{L}^2 satisfying

$$\mathbb{T}(t)\mathbf{u} = \sum_{k \in \mathbb{N}} e^{-\lambda_k t} u_k \phi_k, \quad \mathbf{u} = \sum_{k \in \mathbb{N}} u_k \phi_k \in \mathbb{L}^2. \quad (4.2)$$

By using the representation (2.3) we can show without any difficulty that the space \mathbb{X}_0 is invariant with respect to this semigroup. The restriction of $\{\mathbb{T}(t)\}_{t \geq 0}$ to \mathbb{X}_0 is also an analytic semigroup which will be denoted in the sequel by $\{S_2(t)\}_{t \geq 0}$. The minus infinitesimal generator \tilde{A}_1 of $\{S_2(t)\}_{t \geq 0}$ is the part of A_1 on \mathbb{X}_0 , that is,

$$D(\tilde{A}_1) = \{\mathbf{u} \in D(A_1) : A_1 \mathbf{u} \in \mathbb{X}_0\}, \quad \tilde{A}_1 \mathbf{u} = A_1 \mathbf{u} \text{ for any } \mathbf{u} \in D(\tilde{A}_1).$$

Note that $\mathbb{X}_1 \subset D(\tilde{A}_1)$. With all the above notation, the stochastic equations for nematic liquid crystal (1.12)-(1.14) can be rewritten as the following stochastic evolution equation in the space H ,

$$d\mathbf{y}(t) + \mathbf{A}\mathbf{y}(t)dt + \mathbf{F}(\mathbf{y}(t))dt + \mathbf{L}(\mathbf{y}(t))dt = \mathbf{G}(\mathbf{y}(t))dW(t), \quad (4.3)$$

where, for $\mathbf{y} = (\mathbf{v}, \mathbf{n}) \in E$ and $k = (k_1, k_2) \in \mathbb{K}$,

$$\mathbf{A}\mathbf{y} = \begin{pmatrix} A & 0 \\ 0 & A_1 \end{pmatrix} \begin{pmatrix} \mathbf{v} \\ \mathbf{n} \end{pmatrix}, \quad \mathbf{F}(\mathbf{y}) = \begin{pmatrix} B(\mathbf{v}, \mathbf{v}) + M(\mathbf{n}) \\ \tilde{B}(\mathbf{v}, \mathbf{n}) + f(\mathbf{n}) \end{pmatrix}, \quad (4.4)$$

$$\mathbf{L}(\mathbf{y}) = \begin{pmatrix} 0 \\ -\frac{1}{2}G^2(\mathbf{n}) \end{pmatrix}, \quad \mathbf{G}(\mathbf{y})k = \begin{pmatrix} S(\mathbf{u})k_1 \\ G(\mathbf{n})k_2 \end{pmatrix}. \quad (4.5)$$

Below we will also use the C_0 analytic semigroup $\{S(t)\}_{t \geq 0}$ on $H = \mathbb{H} \times \mathbb{X}_0$ defined by

$$S(t) \begin{pmatrix} \mathbf{v} \\ \mathbf{n} \end{pmatrix} = \begin{pmatrix} S_1(t)\mathbf{v} \\ S_2(t)\mathbf{n} \end{pmatrix}, \quad (\mathbf{v}, \mathbf{n}) \in H.$$

Its infinitesimal generator is $-\mathbf{A}$, where \mathbf{A} is defined in (4.4). Some properties of $\{S(t) : t \geq 0\}$ will be given in Lemmata A.3-A.5.

Assumption 4.1. We assume that $S : \mathbb{H} \rightarrow \mathcal{T}_2(K_1, \mathbb{V})$ is a globally Lipschitz map. In particular, there exists $\ell_5 \geq 0$ such that

$$\|S(\mathbf{u})\|_{\mathcal{T}_2(K_1, \mathbb{V})}^2 \leq \ell_5(1 + \|\mathbf{u}\|^2), \quad \text{for any } \mathbf{u} \in \mathbb{H}.$$

Let us recall the following notations/definition which are borrowed from [35] (see also [6]).

Definition 4.1. For a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with given right-continuous filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$, a stopping time τ is called accessible iff there exists an increasing sequence of stopping times τ_n such that a.s. $\tau_n < \tau$ and $\lim_{n \rightarrow \infty} \tau_n = \tau$, see [35].

Notation. For a stopping time τ we set

$$\Omega_t(\tau) = \{\omega \in \Omega : t < \tau(\omega)\},$$

$$[0, \tau) \times \Omega = \{(t, \omega) \in [0, \infty) \times \Omega : 0 \leq t < \tau(\omega)\}.$$

Definition 4.2. A process $\eta : [0, \tau) \times \Omega \rightarrow X$ (we will also write $\eta(t)$, $t < \tau$), where X is a metric space, is admissible iff

- (i) it is adapted, i.e. $\eta|_{\Omega_t(\tau)} : \Omega_t(\tau) \rightarrow X$ is \mathcal{F}_t measurable, for any $t \geq 0$;
- (ii) for almost all $\omega \in \Omega$, the function $[0, \tau(\omega)) \ni t \mapsto \eta(t, \omega) \in X$ is continuous.

A process $\eta : [0, \tau) \times \Omega \rightarrow X$ is progressively measurable iff, for any $t > 0$, the map

$$[0, t \wedge \tau) \times \Omega \ni (s, \omega) \mapsto \eta(s, \omega) \in X$$

is $\mathcal{B}_{t \wedge \tau} \times \mathcal{F}_{t \wedge \tau}$ measurable.

Two processes $\eta_i : [0, \tau_i) \times \Omega \rightarrow X$, $i = 1, 2$ are called equivalent (we will write $(\eta_1, \tau_1) \sim (\eta_2, \tau_2)$) iff $\tau_1 = \tau_2$ a.s. and for any $t > 0$ the following holds

$$\eta_1(\cdot, \omega) = \eta_2(\cdot, \omega) \text{ on } [0, t]$$

for a.a. $\omega \in \Omega_t(\tau_1) \cap \Omega_t(\tau_2)$.

Note that if processes $\eta_i : [0, \tau_i) \times \Omega \rightarrow X$, $i = 1, 2$ are admissible and for any $t > 0$ $\eta_1(t)|_{\Omega_t(\tau_1)} = \eta_2(t)|_{\Omega_t(\tau_2)}$ a.s. then they are also equivalent.

We now define some concepts of solution to (5.24), see [12, Def. 4.2] or [42, Def. 2.1].

Definition 4.3. Assume that a V -valued \mathcal{F}_0 measurable random variable \mathbf{y}_0 with $\mathbb{E}\|\mathbf{y}_0\|^2 < \infty$ is given. A local mild solution to problem (5.9) (with the initial time 0) is a pair (\mathbf{y}, τ) such that

- (1) τ is an accessible \mathbb{F} -stopping time,
- (2) $\mathbf{y} : [0, \tau) \times \Omega \rightarrow V$ is an admissible process,
- (3) there exists an approximating sequence $(\tau_m)_{m \in \mathbb{N}}$ of finite \mathbb{F} -stopping times such that $\tau_m \nearrow \tau$ a.s. and, for every $m \in \mathbb{N}$ and $t \geq 0$, we have

$$\mathbb{E} \left(\sup_{s \in [0, t \wedge \tau_m]} \|\mathbf{y}(s)\|^2 + \int_0^{t \wedge \tau_m} |\mathbf{y}(s)|_E^2 ds \right) < \infty, \quad (4.6)$$

$$\begin{aligned} \mathbf{y}(t \wedge \tau_m) &= S(t \wedge \tau_m) \mathbf{y}_0 - \int_0^{t \wedge \tau_m} S(t \wedge \tau_m - s) [\mathbf{F}(\mathbf{y}(s)) + \mathbf{L}(\mathbf{y}(s))] ds \\ &+ \int_0^\infty \mathbb{1}_{[0, t \wedge \tau_m)} S(t \wedge \tau_m - s) \mathbf{G}(\mathbf{y}(s)) dW(s). \end{aligned} \quad (4.7)$$

Along the lines of the paper [6], we say that a local solution $\mathbf{y}(t)$, $t < \tau$ is global iff $\tau = \infty$ a.s.

We will check that the nonlinear map \mathbf{F} satisfies the assumption of Proposition 5.12 with $H = \mathbb{H} \times \mathbb{X}_0$, $V = \mathbb{V} \times D(A_1)$ and $E = D(A) \times \mathbb{X}_1$. For this purpose we will prove several estimates.

Lemma 4.4. *There exist some positive constants c_1 and c_2 such that for any $(\mathbf{v}_i, \mathbf{n}_i) \in E$, $i = 1, 2$ we have, with $a = \frac{d}{4}$,*

$$\begin{aligned} \|B(\mathbf{v}_1, \mathbf{v}_1) - B(\mathbf{v}_2, \mathbf{v}_2)\| &\leq c_1 \left(\|\nabla(\mathbf{v}_1 - \mathbf{v}_2)\| \|\nabla \mathbf{v}_1\|^{1-a} \|\Delta \mathbf{v}_1\|^a \right. \\ &\quad \left. + \|\nabla(\mathbf{v}_1 - \mathbf{v}_2)\|^{1-a} \|\Delta(\mathbf{v}_1 - \mathbf{v}_2)\|^a \|\nabla \mathbf{v}_2\| \right) \end{aligned} \quad (4.8)$$

and

$$\begin{aligned} \|M(\mathbf{n}_1) - M(\mathbf{n}_2)\| &\leq c_2 \left(\|\mathbf{n}_1 - \mathbf{n}_2\|_2 \|\mathbf{n}_1\|_2^{1-a} \|\mathbf{n}_1\|_3^a \right. \\ &\quad \left. + \|\mathbf{n}_1 - \mathbf{n}_2\|_2^{1-a} \|\mathbf{n}_1 - \mathbf{n}_2\|_3^a \|\mathbf{n}_2\|_2 \right). \end{aligned} \quad (4.9)$$

Note that we used the shorthand notation $M(\mathbf{n}) := M(\mathbf{n}, \mathbf{n})$ and $B(\mathbf{v}) := B(\mathbf{v}, \mathbf{v})$.

Proof. Set $(\mathbf{w}, \bar{\mathbf{n}}) = (\mathbf{v}_1 - \mathbf{v}_2, \mathbf{n}_1 - \mathbf{n}_2)$. We start with the proof of the estimate (4.8). Notice that the left-hand-side of (4.8) is equal to

$$\|B(\mathbf{w}, \mathbf{v}_1) + B(\mathbf{v}_2, \mathbf{w})\|.$$

Now we estimate the last identity as follows

$$\|B(\mathbf{w}, \mathbf{v}_1) + B(\mathbf{v}_2, \mathbf{w})\| \leq C \|\mathbf{w}\|_{\mathbb{L}^4} \|\nabla \mathbf{v}_1\|_{\mathbb{L}^4} + \|\mathbf{v}_2\|_{\mathbb{L}^4} \|\nabla \mathbf{w}\|_{\mathbb{L}^4},$$

from which along with (A.1) and the embedding (A.3) we easily derive the estimate (4.8).

Next we show that (4.9) holds. From elementary calculi we infer the existence of a constant $C > 0$ such that

$$\|M(\mathbf{f}, \mathbf{g})\| \leq C \|D^2 \mathbf{f}\| \|\nabla \mathbf{g}\|_{\mathbb{L}^\infty} + \|\nabla \mathbf{f}\|_{\mathbb{L}^4} \|D^2 \mathbf{g}\|_{\mathbb{L}^4}.$$

Owing to the embedding (A.3) it is not difficult to check that

$$\|M(\mathbf{f}, \mathbf{g})\| \leq C \|\mathbf{f}\|_2 \left(\|\nabla \mathbf{g}\|_{\mathbb{L}^\infty} + \|D^2 \mathbf{g}\|_{\mathbb{L}^4} \right).$$

Owing to (A.1), (A.4) and the embedding (A.3) we obtain that

$$\|M(\mathbf{f}, \mathbf{g})\| \leq C \|\mathbf{f}\|_2 \|\mathbf{g}\|_2^{1-a} \|\mathbf{g}\|_3^a, \quad a = \frac{d}{4}. \quad (4.10)$$

Note that

$$M(\mathbf{n}_1) - M(\mathbf{n}_2) = M(\mathbf{n}_1 - \mathbf{n}_2, \mathbf{n}_1) + M(\mathbf{n}_2, \mathbf{n}_1 - \mathbf{n}_2).$$

From this last identity and (4.10) we easily deduce the inequality (4.9). This ends the proof of Lemma 4.4. \square

Lemma 4.5. *There exist a constant $c_3 > 0$ such that for any $(\mathbf{v}_i, \mathbf{n}_i) \in E$, $i = 1, 2$ we have, with $a = \frac{d}{4}$,*

$$\begin{aligned} \|\tilde{B}(\mathbf{v}_1, \mathbf{n}_1) - \tilde{B}(\mathbf{v}_2, \mathbf{n}_2)\|_1 &\leq c_3 \left(\|\nabla(\mathbf{v}_1 - \mathbf{v}_2)\| \|\mathbf{n}_1\|_2^{1-a} \|\mathbf{n}_1\|_3^a \right. \\ &\quad \left. + \|(\mathbf{n}_1 - \mathbf{n}_2)\|_2^{1-a} \|(\mathbf{n}_1 - \mathbf{n}_2)\|_3^a \|\nabla \mathbf{v}_2\| \right) \end{aligned} \quad (4.11)$$

Proof. As in the proof of Lemma 4.4 we set $(\mathbf{w}, \bar{\mathbf{n}}) = (\mathbf{v}_1 - \mathbf{v}_2, \mathbf{n}_1 - \mathbf{n}_2)$ and notice that the left hand side of (4.11) is equal to

$$\tilde{B}(\mathbf{w}, \mathbf{n}_1) + \tilde{B}(\mathbf{n}_2, \mathbf{w}) := J_1 + J_2.$$

Now we want to estimate $\|J_i\|_1 = \sqrt{\|J_i\|^2 + \|\nabla J_i\|^2}$ for $i = 1, 2$. Since estimating $\|J_i\|$ is easy we will just focus on the term $\|\nabla J_i\|$. There exists a constant $C > 0$ such that

$$\begin{aligned}\|\nabla J_1\| &\leq C \left(\|\nabla \mathbf{w} \nabla \mathbf{n}_1\| + \|\Delta \mathbf{n}_1 \mathbf{w}\| \right), \\ &\leq C \left(\|\nabla \mathbf{w}\| \|\nabla \mathbf{n}_1\|_{\mathbb{L}^\infty} + \|\mathbf{w}\|_{\mathbb{L}^4} \|\Delta \mathbf{n}_1\|_{\mathbb{L}^4} \right).\end{aligned}$$

Invoking (A.1), (A.2) and the embedding (A.3) we infer that with $a = \frac{d}{4}$,

$$\|\nabla J_1\| \leq C \|\nabla \mathbf{w}\| \left(\|\nabla \mathbf{n}_1\|_1^{1-a} \|\Delta \mathbf{n}_1\|_1^a + \|\Delta \mathbf{n}_1\|^{1-a} \|\mathbf{n}_1\|_3^a \right).$$

This last inequality implies that there exists $\tilde{c} > 0$ such that with $a = \frac{d}{4}$,

$$\|\nabla J_1\| \leq \tilde{c} \|\nabla \mathbf{w}\| \|\mathbf{n}_1\|_2^{1-a} \|\mathbf{n}_1\|_3^a.$$

Using similar argument we can also prove that (again with $a = \frac{d}{4}$)

$$\|\nabla J_2\| \leq \tilde{c} \|\nabla \mathbf{v}_2\| \|\mathbf{n}_1 - \mathbf{n}_2\|_2^{1-a} \|\mathbf{n}_1 - \mathbf{n}_2\|_3^a.$$

The inequality (4.11) easily follows from these last two estimates. \square

Lemma 4.6. *If Assumption 2.1 is verified, then there exists $c_4 > 0$ such that for any $\mathbf{n}_i \in \mathbb{X}_{\frac{1}{2}} \cap \mathbb{X}_1$ with $i = 1, 2$.*

$$\|f(\mathbf{n}_1) - f(\mathbf{n}_2)\|_1 \leq c_4 \left[1 + \|\mathbf{n}_1\|_2^{2N} + \|\mathbf{n}_2\|_2^{2N} \right] \|\mathbf{n}_1 - \mathbf{n}_2\|_2. \quad (4.12)$$

Proof. Since \mathbb{H}^2 is an algebra we have $|\mathbf{n}_1|^{2k} \mathbf{n}_2$ for any $\mathbf{n}_1, \mathbf{n}_2 \in \mathbb{H}^2$ and $k \geq 0$ and there exists $C > 0$ such that

$$\| |\mathbf{n}_1|^{2k} \mathbf{n}_2 \|_2 \leq C \|\mathbf{n}_1\|_2^{2k} \|\mathbf{n}_2\|_2. \quad (4.13)$$

By Young's inequality, there exists $C > 0$ such that

$$\|\mathbf{n}_1\|_2^{2k} \|\mathbf{n}_2\|_2 \leq C \|\mathbf{n}_2\|_2 (1 + \|\mathbf{n}_1\|_2^{2N}), \text{ for } k = 0, \dots, N. \quad (4.14)$$

Thus it is sufficient to prove the lemma for the leading term $a_N |\mathbf{n}|^{2N} \mathbf{n}$.

We have

$$|\mathbf{n}_1|^{2N} \mathbf{n}_1 - |\mathbf{n}_2|^{2N} \mathbf{n}_2 = |\mathbf{n}_1|^{2N} (\mathbf{n}_1 - \mathbf{n}_2) + \mathbf{n}_2 (|\mathbf{n}_1| - |\mathbf{n}_2|) \left(\sum_{k=0}^{2N-1} |\mathbf{n}_1|^{2N-1-k} |\mathbf{n}_2|^k \right),$$

from which along with (4.13) and (4.14) we easily infer that the lemma is true from the leading term $a_N |\mathbf{n}|^{2N} \mathbf{n}$. \square

For two Banach spaces $(B_i, \|\cdot\|_{B_i})$ with $i = 1, 2$ we endow the product space $B_1 \times B_2$ with the norm

$$|(b_1, b_2)| = \sqrt{\|b_1\|_{B_1}^2 + \|b_2\|_{B_2}^2}.$$

Proposition 4.7. *If Assumption 2.1 is satisfied, then there exists a constant $C_0 > 0$ such that for any $\mathbf{y}_i = (\mathbf{v}_i, \mathbf{n}_i)$, $i = 1, 2$, with $\alpha = \frac{d}{4}$, we have*

$$\begin{aligned}\|\mathbf{F}(\mathbf{y}_1) - \mathbf{F}(\mathbf{y}_2)\|_H &\leq C_0 \|\mathbf{y}_1 - \mathbf{y}_2\|_V^{1-\alpha} \left[\|\mathbf{y}_1 - \mathbf{y}_2\|_V^\alpha \|\mathbf{y}_1\|_V^{1-\alpha} \|\mathbf{y}_1\|_E^\alpha + \|\mathbf{y}_1 - \mathbf{y}_2\|_E^\alpha \|\mathbf{y}_2\|_V \right] \\ &\quad + c_4 \|\mathbf{y}_1 - \mathbf{y}_2\|_V \left[1 + \|\mathbf{y}_1\|_V^{2N} + \|\mathbf{y}_2\|_V^{2N} \right].\end{aligned}$$

Proof. The proposition is a consequence of Lemma 4.4, Lemma 4.5 and Lemma 4.6. Its proof is easy and we omit it. \square

For any integer $k > 1$ let

$$\tau_k = \inf\{t \geq 0 : \|\nabla \mathbf{v}(t)\|^2 + \|\Delta \mathbf{n}(t)\|^2 > k^2\}. \quad (4.15)$$

Hereafter, for a vector-valued function $\mathbf{u} : [0, t \wedge \tau_k] \rightarrow \mathbf{B}$ we will write $\int_0^{t \wedge \tau_k} \mathbf{u} \, ds := \int_0^{t \wedge \tau_k} \mathbf{u}(s) \, ds$ for any $t > 0$.

Our first main result is contained in the following theorem. It is basically a corollary of two general theorems (see Theorem 5.15 and Theorem 5.16) that we will state and prove in Section 5.

Theorem 4.8. *If Assumptions 4.1 and 2.1 are satisfied and $(\mathbf{v}_0, \mathbf{n}_0) \in \mathbb{L}^2(\Omega; D(A^{\frac{1}{2}}) \times \mathbb{X}_{\frac{1}{2}})$ is \mathcal{F}_0 -measurable, $h \in \mathbb{H}^2$ and $d = 2, 3$, then the stochastic equation (4.3) for the liquid crystals has a local-maximal strong solution $((\mathbf{v}; \mathbf{n}), \tilde{\tau}_\infty)$ satisfying the following properties:*

- (1) *given $R > 0$ and $\varepsilon > 0$ there exists $\tau(\varepsilon, R) > 0$, such that for every \mathcal{F}_0 -measurable $D(A^{\frac{1}{2}}) \times \mathbb{X}_{\frac{1}{2}}$ -valued random variable $(\mathbf{v}_0, \mathbf{n}_0)$ satisfying $\mathbb{E}\|(\mathbf{v}_0, \mathbf{n}_0)\|_{D(A^{\frac{1}{2}}) \times \mathbb{X}_{\frac{1}{2}}}^2 \leq R^2$, one has*

$$\mathbb{P}(\tilde{\tau}_\infty \geq \tau(\varepsilon, R)) \geq 1 - \varepsilon.$$

- (2) *We also have*

$$\mathbb{P}(\{\tilde{\tau}_\infty < \infty\} \cap \{\|\nabla \mathbf{v}(t)\| + \|\Delta \mathbf{n}(t)\| < \infty\}) = 0, \quad (4.16)$$

$$\limsup_{t \nearrow \tilde{\tau}_\infty} \|\nabla \mathbf{v}(t)\| + \|\Delta \mathbf{n}(t)\| = \infty \text{ a.s. on } \{\tilde{\tau}_\infty < \infty\}. \quad (4.17)$$

Proof. Lemma A.3-A.5 show that $\{\mathbb{S}(t)\}_{t \geq 0}$ on $H = \mathbb{H} \times \mathbb{X}_0$ satisfies Assumption 5.3. Thanks to Proposition 4.7 we can infer from Theorem 5.15 and Theorem 5.16 that problem (4.3) has a local and maximal strong solution satisfying items (1) and (2). This concludes the proof of the theorem. \square

The second result is about global resolvability of the stochastic equation for two dimensional nematic liquid crystal. This result relies on the Khashminskii test for non-explosions (see [34, Theorem III.4.1] for the finite-dimensional case) and some arguments from [11].

Theorem 4.9. *If in addition to Assumptions 4.1 and 3.1, we also suppose that $d = 2$, $h \in \mathbb{H}^2$ and $(\mathbf{v}_0, \mathbf{n}_0) \in \mathbb{L}^2(\Omega; D(A^{\frac{1}{2}}) \times \mathbb{X}_{\frac{1}{2}})$ such that*

$$\mathbb{E}(\mathcal{E}_0^{4N+2}) := \mathbb{E}\left(\|\mathbf{v}_0\|^2 + \|\mathbf{n}_0\|^2 + \|\nabla \mathbf{n}_0\|^2 + \int_{\mathcal{O}} F(\mathbf{n}_0(x)) \, dx\right)^{4N+2} < \infty, \quad (4.18)$$

then the stochastic equation (4.3) for nematic liquid crystals has a global strong solution.

To prove this theorem we need several auxiliary results some of whose proofs will be postponed till the appendix. Let us start with the following 3 lemmata.

Lemma 4.10. *For any $h \in \mathbb{H}^2$, there exists two constants $\kappa_1 > 0$ and $\kappa_2 > 0$, which depends only on the \mathbb{H}^2 -norm of h , such that for any $\mathbf{n} \in \mathbb{H}^1$ satisfying $\Delta \mathbf{n} - f(\mathbf{n}) \in \mathbb{L}^2$ we have*

$$|\langle \Delta \mathbf{n} - f(\mathbf{n}), \Delta \mathbf{n} \times h \rangle + \langle \Delta \mathbf{n} - f(\mathbf{n}), \Delta \mathbf{n} \times \Delta h + (\mathbf{n} \times h) \times \Delta h \rangle| \leq \kappa_1(\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{H}^1}^{4N+2} + 1). \quad (4.19)$$

$$\|\Delta(\mathbf{n} \times h) - f'(\mathbf{n})(\mathbf{n} \times h)\|^2 + |\langle \Delta \mathbf{n} - f(\mathbf{n}), f''(\mathbf{n})[G(\mathbf{n}), G(\mathbf{n})] \rangle| \leq \kappa_2(\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{H}^1}^{4N+2} + 1). \quad (4.20)$$

Proof. The proof of the lemma will be given in Appendix C. \square

Lemma 4.11. *For $N \in I_2$, there exists a constant $\kappa_3 > 0$ such that for any $\mathbf{v} \in \mathbb{H}^1$ satisfying $\Delta \mathbf{n} - f(\mathbf{n}) \in \mathbb{L}^2$ we have*

$$|\langle f'(\mathbf{n})(\Delta \mathbf{n} - f(\mathbf{n})), \Delta \mathbf{n} - f(\mathbf{n}) \rangle| \leq \frac{1}{4} + \kappa_3(\ell_1 + \ell_2 \|\mathbf{n}\|_{\mathbb{H}^1}^{2N})^2 \|\Delta \mathbf{n} - f(\mathbf{n})\|^2, \quad (4.21)$$

where the constants ℓ_1 and ℓ_2 are defined in (2.15) and (2.16), respectively.

Proof. We will carry out the proof of this lemma in Appendix C. \square

Lemma 4.12. *There exists a constant κ_4 such that*

$$2|\langle \nabla \mathbf{n}(\Delta \mathbf{n} - f(\mathbf{n})), A\mathbf{v} \rangle - \langle \nabla \mathbf{n}(\Delta \mathbf{v}), \Delta \mathbf{n} - f(\mathbf{n}) \rangle| \leq \frac{1}{4}\|A\mathbf{v}\|^2 + \frac{1}{4}\|\nabla(\Delta \mathbf{n} - f(\mathbf{n}))\|^2 + \kappa_4\|\nabla \mathbf{n}\|^2\|\Delta \mathbf{n}\|^2\|\Delta \mathbf{n} - f(\mathbf{n})\|^2, \quad (4.22)$$

for any $\mathbf{v} \in D(A)$, $\mathbf{n} \in \mathbb{H}^2$.

Proof. Let us call LHS the right-hand side of the inequality in the statement of the lemma. From the Cauchy-Schwarz inequality we infer that there exists a constant $C > 0$ such that

$$LHS \leq 4C\|A\mathbf{v}\|\|\nabla \mathbf{n}\|_{\mathbb{L}^4}\|\Delta \mathbf{n} - f(\mathbf{n})\|_{\mathbb{L}^4},$$

which, along with the Gagliardo-Nirenberg and the Cauchy inequalities, implies

$$LHS \leq \frac{1}{4}\|A\mathbf{v}\|^2 + 16C^2\|\nabla \mathbf{n}\|\|\Delta \mathbf{n}\|\|\Delta \mathbf{n} - f(\mathbf{n})\|\|\nabla(\Delta \mathbf{n} - f(\mathbf{n}))\|.$$

Invoking again the Cauchy inequality yields

$$LHS \leq \frac{1}{4}\|A\mathbf{v}\|^2 + \frac{1}{4}\|\nabla(\Delta \mathbf{n} - f(\mathbf{n}))\|^2 + 16^2C^4\|\nabla \mathbf{n}\|^2\|\Delta \mathbf{n}\|^2\|\Delta \mathbf{n} - f(\mathbf{n})\|^2,$$

which completes the proof of the lemma. \square

We also need the following lemma.

Lemma 4.13. *There exists a constant κ_5 such that for any $\mathbf{v} \in D(A)$ we have*

$$|(B(\mathbf{v}, \mathbf{v}), A\mathbf{v})| \leq \frac{1}{4}\|A\mathbf{v}\|^2 + \kappa_5\|\mathbf{v}\|^2\|\nabla \mathbf{v}\|^4. \quad (4.23)$$

Proof of Lemma 4.13. First, by Cauchy-Schwarz inequality we obtain

$$|\langle B(\mathbf{v}, \mathbf{v}), A\mathbf{v} \rangle| \leq \|B(\mathbf{v}, \mathbf{v})\|\|A\mathbf{v}\|. \quad (4.24)$$

From the Hölder and the Gagliardo-Nirenberg inequalities, we derive that there exists a constant $C_1 > 0$ such that

$$\|B(\mathbf{v}, \mathbf{v})\| \leq C_1\|\mathbf{v}\|^{1-\frac{d}{4}}\|\nabla \mathbf{v}\|^{\frac{d}{4}}\|\nabla \mathbf{v}\|^{1-\frac{d}{4}}\|\Delta \mathbf{v}\|^{\frac{d}{4}}, \text{ for any } \mathbf{v} \in \mathbb{V}, \mathbf{v} \in D(A),$$

from which along with (4.24) we derive that

$$\begin{aligned} |\langle B(\mathbf{v}, \mathbf{v}), A\mathbf{v} \rangle| &\leq c\|\mathbf{v}\|^{1-\frac{d}{4}}\|\nabla \mathbf{v}\|^{\frac{d}{4}}\|\nabla \mathbf{v}\|^{1-\frac{d}{4}}\|\Delta \mathbf{v}\|^{\frac{d}{4}}\|\Delta \mathbf{v}\|, \\ &\leq c\|\mathbf{v}\|^{1-\frac{d}{4}}\|\nabla \mathbf{v}\|\|\Delta \mathbf{v}\|^{1+\frac{d}{4}}. \end{aligned} \quad (4.25)$$

Let us recall Young's inequality $ab \leq C(\alpha, p, q)a^p + \alpha b^q$ for $p^{-1} + q^{-1} = 1$ and arbitrary $\alpha > 0$. Let us choose $p = \frac{8}{d+4}$ and $q = \frac{8}{4-d}$. Applying Young's inequality with the above p and q in (4.25) we obtain

$$|\langle B(\mathbf{v}, \mathbf{v}), A\mathbf{v} \rangle| \leq \alpha\|A\mathbf{v}\|^2 + C(\alpha, p, q)\|\mathbf{v}\|^2\|\nabla \mathbf{v}\|^{2+\frac{2d}{4-d}}.$$

We easily conclude the proof of the lemma by setting $d = 2$ and $\alpha = \frac{1}{4}$ in the last estimate. \square

Now, we proceed to the statement of proposition which is very crucial for the proof of Theorem 4.9. But before continuing further, let us fix some notations. Let ℓ_1 and ℓ_2 be the positive constants defined in (2.15) and (2.16), respectively. Let $((\mathbf{v}; \mathbf{n}), \tilde{\tau}_\infty)$ be the maximal solution of the problem (4.3) obtained in Theorem 4.8, and let us set

$$\phi(s) = \kappa_3(\ell_1 + \ell_2 \sup_{r \in [0, s]} \|\mathbf{n}(r)\|_{\mathbb{H}^1}^{2N})^2 + \kappa_4\|\nabla \mathbf{n}(s)\|^2\|\Delta \mathbf{n}(s)\|^2 + \kappa_5\|\mathbf{v}(s)\|^2\|\nabla \mathbf{v}(s)\|^2, \quad s \geq 0,$$

$$\Phi(s) = e^{-\int_0^s \phi(r)dr}, \quad s \geq 0,$$

where the constants κ_i , $i = 3, \dots, 5$, are given in Lemma 4.11-4.13. Now, we are ready to state the aforementioned proposition.

Proposition 4.14. *Let all the assumptions of Theorem 4.9 be satisfied and (\mathbf{v}, \mathbf{n}) be the local strong solution of (4.3) obtained in Theorem 4.8. Let*

$$\Psi(\mathbf{n}) = \|\Delta \mathbf{n} - f(\mathbf{n})\|^2.$$

Then, there exist 2 increasing functions $\kappa_0, \kappa_1 : [0, T] \rightarrow (0, \infty)$ such that for any $t > 0$

$$\begin{aligned} \mathbb{E} \left[\Phi(t \wedge \tau_k) (\|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k))) \right] + \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \left(2\|\Delta \mathbf{v}(s)\|^2 + \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2 \right) ds \\ \leq \kappa_0(t) \mathbb{E}(\|\nabla \mathbf{v}_0\|^2 + \Psi(\mathbf{n}_0)) + \kappa_1(t) \mathbb{E}(\mathcal{E}_0^{2N+1}), \end{aligned} \quad (4.26)$$

where $\Phi(\cdot)$ is defined as above and \mathcal{E}_0 is as in (4.18).

Proof. The proof will be given in Appendix C. □

After all these preparations we proceed to the promised proof of Theorem 4.9.

Proof of Theorem 4.9. Let $\{\tau_k; k \in \mathbb{N}\}$ be the sequence of stopping times defined in (4.15). By the item (2) of Theorem 4.8 the solution is global if we are able to show that

$$\mathbb{P}(\tilde{\tau}_\infty < \infty) = 0.$$

For this aim, we first derive the following chain of inequalities

$$\begin{aligned} \mathbb{P}(\tau_k < t) &\leq \mathbb{E} \left(1_{\{\tau_k < t\}} 1_{\{e^{\int_0^{t \wedge \tau_k} \phi(r) dr} \leq k\}} \right) + \mathbb{E} \left(1_{\{\tau_k < t\}} 1_{\{e^{\int_0^{t \wedge \tau_k} \phi(r) dr} > k\}} \right), \\ &\leq \frac{1}{k} \mathbb{E} \left(1_{\{\tau_k < t\}} k^2 e^{-\frac{1}{2} \int_0^{t \wedge \tau_k} \phi(s) ds} \right) + \mathbb{E} \left(1_{\{\tau_k < t\}} 1_{\{\int_0^{t \wedge \tau_k} \phi(s) ds > \log k\}} \right), \\ &\leq \text{I} + \text{II}. \end{aligned}$$

Now, we will estimate I and II separately. First, from the definition of τ_k and Φ we have

$$\begin{aligned} \text{I} &\leq \frac{1}{k} \mathbb{E} (1_{\{\tau_k < t\}} \Phi(t \wedge \tau_k) (\|\mathbf{A} \mathbf{v}(t \wedge \tau_k)\|^2 + \|\Delta \mathbf{n}(t \wedge \tau_k)\|^2)) \\ &\leq \frac{1}{k} \mathbb{E} (\Phi(t \wedge \tau_k) (\|\mathbf{A} \mathbf{v}(t \wedge \tau_k)\|^2 + \|\Delta \mathbf{n}(t \wedge \tau_k)\|^2)) \end{aligned}$$

From Proposition C.2, Remark 2.7, (4.26) and (2.18), we infer that

$$\text{I} \leq \frac{1}{k} \left(\kappa_0(t) \mathbb{E}(\|\nabla \mathbf{v}_0\|^2 + \Psi(\mathbf{n}_0)) + \kappa_1(t) \mathbb{E}(\mathcal{E}_0^{2N+1}) \right),$$

for any $k \in \mathbb{N}$. Second, we estimate II as follows

$$\begin{aligned} \text{II} &= \mathbb{E} \left(1_{\{\tau_k < t\}} 1_{\{\int_0^{t \wedge \tau_k} \phi(r) dr > \log k\}} \right) \\ &\leq \int_{\{\tau_k < t\} \cap \{\int_0^{t \wedge \tau_k} \phi(r) dr > \log k\}} \frac{\int_0^{t \wedge \tau_k} \phi(r) dr}{\log k} d\mathbb{P} \\ &\leq \frac{1}{\log k} \int_{\{\tau_k < t\}} \int_0^{t \wedge \tau_k} \phi(r) dr d\mathbb{P} \\ &\leq \frac{1}{\log k} \mathbb{E} \int_0^{t \wedge \tau_k} \phi(r) dr. \end{aligned}$$

Now, from Proposition C.2 we infer that there exists an increasing function $\tilde{\kappa} : [0, \infty) \rightarrow (0, \infty)$ such that

$$\begin{aligned} \mathbb{E} \int_0^{t \wedge \tau_k} \|\mathbf{v}(s)\|^2 \|\nabla \mathbf{v}(s)\|^2 ds &\leq \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_k} \|\mathbf{v}(s)\|^4 + \mathbb{E} \left[\int_0^{t \wedge \tau_k} \|\nabla \mathbf{v}(s)\|^2 ds \right]^2 \leq \tilde{\kappa}(t)(1 + \mathfrak{G}_1(t, 4N + 2)), \\ \mathbb{E} \int_0^{t \wedge \tau_k} (\ell_1 + \ell_2 \|\mathbf{n}(s)\|_{\mathbb{H}^1}^{2N})^2 ds &\leq \tilde{\kappa}(t)(1 + \mathfrak{G}_1(t, 4N + 2)), \\ \mathbb{E} \int_0^{t \wedge \tau_k} \|\nabla \mathbf{n}(s)\|^2 \|\Delta \mathbf{n}(s)\|^2 ds &\leq \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_k} \|\nabla \mathbf{n}(s)\|^4 + \mathbb{E} \left[\int_0^{t \wedge \tau_k} \|\Delta \mathbf{n}(s)\|^2 ds \right]^2 \leq \tilde{\kappa}(t)(1 + \mathfrak{G}_1(t, 4N + 2)), \end{aligned}$$

where $\mathfrak{G}_1(t, 4N + 2)$ is the constant in (3.12) with $T = t$ and $p = 4N + 2$. Thus, there exists a constant $C > 0$ such that for any $k \in \mathbb{N}$

$$\mathbb{E} \int_0^{t \wedge \tau_k} \phi(s) ds \leq C \tilde{\kappa}(t)(1 + \mathfrak{G}_1(t, 2))$$

and

$$\Pi \leq \frac{1}{\log k} C \tilde{\kappa}(t)(1 + \mathfrak{G}_1(t, 2)).$$

Collecting the information about I and II together, we infer that there exists an increasing function $C : [0, \infty) \rightarrow (0, \infty)$ such that for any $k \in \mathbb{N}$

$$\mathbb{P}(\tau_k < t) \leq C(t) \left[\frac{1}{k} + \frac{1}{\log k} \right],$$

which implies that

$$\lim_{k \rightarrow \infty} \mathbb{P}(\tau_k < t) = 0.$$

Now, we easily deduce from the last identity and Theorem 4.8(2) that $\mathbb{P}(\tilde{\tau}_\infty < \infty) = 0$ which implies the global resolvability of our system for the case $d = 2$. \square

5. STRONG SOLUTION FOR AN ABSTRACT STOCHASTIC EQUATION

The goal of this section is to prove a general result about the existence of local and maximal solution to an abstract stochastic partial differential equations with locally Lipschitz continuous coefficients. This is achieved by using some truncation and fixed point methods.

5.1. Notations and Preliminary. Let V , E and H be separable Banach spaces such that $E \subset V$ continuously. We denote the norm in V by $\|\cdot\|$ and we put

$$X_T := C([0, T]; V) \cap L^2(0, T; E) \quad (5.1)$$

with the norm

$$|u|_{X_T}^2 = \sup_{s \in [0, T]} \|u(s)\|^2 + \int_0^T |u(s)|_E^2 ds. \quad (5.2)$$

Let F and G be two nonlinear mappings satisfying the following sets of conditions.

Assumption 5.1. Suppose that $F : E \rightarrow H$ is such that $F(0) = 0$ and there exist $p, q \geq 1$, $\alpha, \gamma \in [0, 1)$ and $C > 0$ such that

$$\begin{aligned} |F(y) - F(x)|_H &\leq C \left[\|y - x\| \|y\|^{p-\alpha} |y|_E^\alpha + |y - x|_E^\alpha \|y - x\|^{1-\alpha} \|x\|^p \right] \\ &\quad + C \left[\|y - x\| \|y\|^{q-\gamma} |y|_E^\gamma + |y - x|_E^\gamma \|y - x\|^{1-\gamma} \|x\|^q \right], \end{aligned} \quad (5.3)$$

for any $x, y \in E$.

Assumption 5.2. Assume that $G : E \rightarrow V$ such that $G(0) = 0$ and there exists $k \geq 1$, $\beta \in [0, 1)$ and $C_G > 0$ such that

$$\|G(y) - G(x)\| \leq C_G \left[\|y - x\| \|y\|^{k-\beta} |y|_E^\beta + |y - x|_E^\beta \|y - x\|^{1-\beta} \|x\|^k \right], \quad (5.4)$$

for any $x, y \in E$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space equipped with a filtration $\mathbb{F} = \{\mathcal{F}_t : t \geq 0\}$ satisfying the usual condition. By $M^2(X_T)$ we denote the space of all progressively measurable E -values processes whose trajectories belong to X_T almost surely endowed with a norm

$$|u|_{M^2(X_T)}^2 = \mathbb{E} \left[\sup_{s \in [0, T]} \|u(s)\|^2 + \int_0^T |u(s)|_E^2 ds \right]. \quad (5.5)$$

Let us also formulate the following assumptions.

Assumption 5.3. Suppose that the embeddings $E \subset V \subset H$ are continuous. Consider (for simplicity) a one-dimensional Wiener process $W(t)$.

Assume that $S(t)$, $t \in [0, \infty)$, is a family of bounded linear operators on the space H such that there exists two positive constants C_1 and C_2 with the following properties .

(i) For every $T > 0$ and every $f \in L^2(0, T; H)$ a function $u = S * f$ defined by

$$u(t) = \int_0^T S(t-r)f(r) dt, \quad t \in [0, T]$$

belongs to X_T and

$$|u|_{X_T} \leq C_1 |f|_{L^2(0, T; H)}. \quad (5.6)$$

(ii) For every $T > 0$ and every process $\xi \in M^2(0, T; V)$ a process $u = S \diamond \xi$ defined by

$$u(t) = \int_0^T S(t-r)\xi(r) dW(r), \quad t \in [0, T]$$

belongs to $M^2(X_T)$ and

$$|u|_{M^2(X_T)} \leq C_2 |\xi|_{M^2(0, T; V)}.$$

(iii) For every $T > 0$ and every $u_0 \in V$, a function $u = Su_0$ defined by

$$u(t) = S(t)u_0, \quad t \in [0, T]$$

belongs to X_T . Moreover, for every $T_0 > 0$ there exist $C_0 > 0$ such that for all $T \in (0, T_0]$,

$$|u|_{X_T} \leq C_0 \|u_0\|. \quad (5.7)$$

Now let us consider a semigroup $S(t)$, $t \in [0, \infty)$ as above and the abstract SPDEs

$$u(t) = S(t)u_0 + \int_0^t S(t-s)F(u(s))ds + \int_0^t S(t-s)G(u(s))dW(s), \quad \text{for any } t > 0 \quad (5.8)$$

which is a mild version of the problem

$$\begin{cases} du(t) = Au(t) dt + F(u(t)) dt + G(u(t))dW(t), & t > 0, \\ u(0) = u_0. \end{cases} \quad (5.9)$$

Definition 5.1. Assume that a V -valued \mathcal{F}_0 measurable random variable u_0 is given. A local mild solution to problem (5.9) (with the initial time 0) is a pair (u, τ) such that

- (1) τ is an accessible \mathbb{F} -stopping time,
- (2) $u : [0, \tau) \times \Omega \rightarrow V$ is an admissible¹ process,

¹This also follows from condition (3) below.

- (3) there exists an approximating sequence $(\tau_m)_{m \in \mathbb{N}}$ of \mathbb{F} finite stopping times such that $\tau_m \nearrow \tau$ a.s. and, for every $m \in \mathbb{N}$ and $t \geq 0$, we have

$$\mathbb{E} \left(\sup_{s \in [0, t \wedge \tau_m]} \|u(s)\|^2 + \int_0^{t \wedge \tau_m} |u(s)|_E^2 ds \right) < \infty, \quad (5.10)$$

$$\begin{aligned} u(t \wedge \tau_m) &= S(t \wedge \tau_m)u_0 + \int_0^{t \wedge \tau_m} S(t \wedge \tau_m - s)F(u(s))ds \\ &+ \int_0^\infty \mathbb{1}_{[0, t \wedge \tau_m)} S(t \wedge \tau_m - s)G(u(s))dW(s). \end{aligned} \quad (5.11)$$

Along the lines of the paper [6], we said that a local solution $u(t)$, $t < \tau$ is called global iff $\tau = \infty$ a.s.

Remark 5.2. The Definition 5.1 of a local solution is independent of the choice of the sequence (τ_n) . A proof of this fact follows from the continuity of trajectories of the process u (what is a consequence of admissibility of u) and is based on the following three principles.

- (i) If τ is an accessible stopping time, then there exist an increasing sequence τ_n of *discrete* stopping times such that a.s. $\tau_n < \tau$ and $\tau_n \nearrow \tau$;
- (ii) if τ is an accessible stopping time and $\sigma \leq \tau$ is a stopping time, then σ is also accessible.
- (ii) if a pair (u, τ) is a local solution to (5.8), then (5.11) holds with t being any discrete stopping time.

It follows that the following is an equivalent definition of a local solution.

A pair (u, τ) , where τ be an accessible stopping time and $u : [0, \tau) \times \Omega \rightarrow V$ is an admissible process, is a local mild solution to equation (5.9) iff for every accessible stopping time σ such that $\sigma < \tau$, for every $t \geq 0$, a.s.

$$\begin{aligned} u(t \wedge \sigma) &= S(t \wedge \sigma)u_0 + \int_0^{t \wedge \sigma} S(t \wedge \sigma - s)F(u(s))ds \\ &+ \int_0^\infty \mathbb{1}_{[0, t \wedge \sigma)} S(t \wedge \sigma - s)G(u(s))dW(s). \end{aligned} \quad (5.12)$$

Let us first formulate the following useful result.

Proposition 5.3. Assume that a pair (u, τ) is a local mild solution to problem (5.9), where u_0 is an V -valued \mathcal{F}_0 measurable random variable. Then for every finite stopping time σ , a pair $(u|_{[0, \tau \wedge \sigma)} \times \Omega, \tau \wedge \sigma)$ is also a local mild solution to problem (5.9).

Secondly, let us recall the following result, see [22, Lemmata III 6A and 6B].

Lemma 5.4. (The Amalgamation Lemma) Let Δ be a family of accessible stopping times with values in $[0, \infty]$. Then a function

$$\tau := \sup\{\alpha : \alpha \in \Delta\}$$

is an accessible stopping time with values in $[0, \infty]$ and there exists an Δ -valued increasing sequence $\{\alpha_n\}_{n=1}^\infty$ such that τ is the pointwise limit of α_n .

Assume also that for each $\alpha \in \Delta$, $I_\alpha : [0, \alpha) \times \Omega \rightarrow V$ is an admissible process such that for all $\alpha, \beta \in \Delta$ and every $t > 0$,

$$I_\alpha(t) = I_\beta(t) \text{ a.s. on } \Omega_t(\alpha \wedge \beta). \quad (5.13)$$

Then, there exists an admissible process $\mathbf{I} : [0, \tau) \times \Omega \rightarrow V$, such that every $t > 0$,

$$\mathbf{I}(t) = I_\alpha(t) \text{ a.s. on } \Omega_t(\alpha). \quad (5.14)$$

Moreover, if $\tilde{I} : [0, \tau) \times \Omega \rightarrow X$ is any process satisfying (5.14) then the process \tilde{I} is a version of the process \mathbf{I} , i.e. for any $t \in [0, \infty)$

$$\mathbb{P}(\{\omega \in \Omega : t < \tau(\omega), I(t, \omega) \neq \tilde{I}(t, \omega)\}) = 0. \quad (5.15)$$

In particular, if in addition \tilde{I} is an admissible process, then

$$\mathbf{I} = \tilde{I}. \quad (5.16)$$

Remark 5.5. Let us note that because both processes $\mathbf{I} : [0, \tau) \times \Omega \rightarrow V$ and $I_\alpha : [0, \alpha) \times \Omega \rightarrow V$ are admissible (and hence with almost sure continuous trajectories), and since $\alpha \leq \tau$, condition (5.14) is equivalent to the following one:

$$\mathbf{I}|_{[0, \alpha) \times \Omega} = I_\alpha. \quad (5.17)$$

Similarly, condition (5.13) is equivalent to the following one

$$I_\alpha|_{[0, \alpha \wedge \beta) \times \Omega} = I_\beta|_{[0, \alpha \wedge \beta) \times \Omega}. \quad (5.18)$$

Definition 5.6. Consider a family \mathcal{LS} of all local mild solution (u, τ) to the problem (5.9). For two elements $(u, \tau), (v, \sigma) \in \mathcal{LS}$ we write that $(u, \tau) \preceq (v, \sigma)$ iff $\tau \leq \sigma$ a.s. and $v|_{[0, \tau) \times \Omega} \sim u$. Note that if $(u, \tau) \preceq (v, \sigma)$ and $(v, \sigma) \preceq (u, \tau)$, then $(u, \tau) \sim (v, \sigma)$. We write $(u, \tau) \prec (v, \sigma)$ iff $(u, \tau) \preceq (v, \sigma)$ and $(u, \tau) \not\sim (v, \sigma)$. Then, the pair (\mathcal{LS}, \preceq) is a partially ordered set in which, according to the Amalgamation Lemma 5.4, every non-empty chain has an upper bound.

Each such a maximal element (u, τ) in the set (\mathcal{LS}, \preceq) is called a maximal local mild solution to the problem (5.9).

If (u, τ) is a maximal local mild solution to equation (5.9), the stopping time τ is called its lifetime.

A priori, there may be many maximal elements in (\mathcal{LS}, \preceq) and hence many maximal local mild solutions to the problem (5.9). However, as we will see later, if the uniqueness of local solutions holds, then the uniqueness of the maximal local mild solution will follow.

Remark 5.7. The following is an equivalent version of Definition 5.6. For a local mild solution (u, τ) the following conditions are equivalent.

- (nm1) The pair (u, τ) is not a maximal local mild solution to problem (5.9).
- (nm2) There exists a local mild solution (v, σ) to problem (5.9) such that $(u, \tau) \prec (v, \sigma)$.
- (nm3) There exists a local mild solution (v, σ) to problem (5.9) such that $\tau \leq \sigma$ a.s., $v|_{[0, \tau) \times \Omega} \sim u$ and $\mathbb{P}(\tau < \sigma) > 0$.
- (nm4) Every local mild solution (v, σ) to problem (5.9) such that $(u, \tau) \not\sim (v, \sigma)$ satisfies $(u, \tau) \not\prec (v, \sigma)$.

Definition 5.8. A local solution (u, τ) to problem (5.9) is unique iff for any other local solution (v, σ) to (5.9) the restricted processes $u|_{[0, \tau \wedge \sigma) \times \Omega}$ and $v|_{[0, \tau \wedge \sigma) \times \Omega}$ are equivalent.

Proposition 5.9. Suppose that u_0 is a V -valued random variable and \mathcal{F}_0 -measurable. Suppose that there exist at least one local solution (u^0, τ^0) to problem (5.9) and that for any two local solutions (u^1, τ^1) and (u^2, τ^2) , and every $t > 0$,

$$u^1(t) = u^2(t) \text{ a.s. on } \{t < \tau^1 \wedge \tau^2\}. \quad (5.19)$$

Then there exists a maximal local mild solutions to the same problem.

Remark 5.10. Let us note that similarly to Remark 5.5, because both the local solutions u^1 and u^2 are admissible (and hence with almost sure continuous trajectories), condition (5.19) is equivalent to the following one:

$$u^1|_{[0, \tau^1 \wedge \tau^2) \times \Omega} = u^2|_{[0, \tau^1 \wedge \tau^2) \times \Omega}. \quad (5.20)$$

Proof of Proposition 5.9. Consider the family \mathcal{LS} introduced in Definition 5.6. Note that in view of Proposition 5.3, if $(u, \tau) \in \mathcal{LS}$, then for any $T > 0$, $(u|_{[0, T \wedge \tau) \times \Omega}, T \wedge \tau)$ also belongs to \mathcal{LS} . Let \mathcal{LC} be any nonempty chain in \mathcal{LS} containing (u^0, τ^0) . Set

$$\hat{\tau} := \sup \{ \tau : (u, \tau) \in \mathcal{LC} \}.$$

Since the chain \mathcal{LC} is non-empty, from the Amalgamation Lemma 5.4 it follows that $\hat{\tau}$ is an accessible stopping time and that there exists an admissible V -valued process $\hat{u}(t)$, $t < \hat{\tau}$ such that for all $(u, \tau) \in \mathcal{LC}$, $(\hat{u}|_{[0, \tau) \times \Omega}, \tau) \sim (u, \tau)$. In view of the Kuratowski-Zorn Lemma it remains to prove that

the pair $(\hat{u}, \hat{\tau})$ belongs to \mathcal{LS} . To prove this let us consider an \mathcal{LC} -valued sequence, which exists in view of Lemma 5.4, (u^n, α_n) of local mild solutions to problem (5.9) such that $\alpha_n \nearrow \tau$ a.s. Moreover, by the above comment on Proposition 5.3 we can assume that α_n is a bounded from above stopping time. Moreover, since each (u^n, α_n) is a local mild solution, we can find a sequence (σ_n) of local times such that $\sigma_n < \alpha_n$, $\sigma_n \nearrow \tau$ a.s. and for each n , the condition (5.10) and equality (5.11) are satisfied by u^n .

Since $(u^n, \alpha_n) \in \mathcal{LC}$, we infer that $(\hat{u}|_{[0, \alpha_n] \times \Omega}, \alpha_n) \sim (u^n, \alpha_n)$ and thus we infer that also for each n , the condition (5.10) and equality (5.11) are satisfied by \hat{u} . This proves that $(\hat{u}, \hat{\tau}) \in \mathcal{LC}$. The proof is complete. \square

Now we shall prove a counterparts of Corollary 2.29 and Lemma 2.31 from [6].

Proposition 5.11. *Assume that the Assumption 5.1-Assumption 5.3 hold. If a pair (u, τ) is a maximal local solution then*

$$\mathbb{P}\left(\{\omega \in \Omega : \tau(\omega) < \infty, \exists \lim_{t \nearrow \tau(\omega)} u(t)(\omega) \in V\}\right) = 0 \quad (5.21)$$

and

$$\lim_{t \nearrow \tau} |u|_{X_t} = \infty \quad \mathbb{P} - \text{a.s. on } \{\tau < \infty\}. \quad (5.22)$$

Proof of Proposition 5.11 (by contradiction). Part I. Assume that there exists $\Omega_1 \subset \Omega$ such that $\mathbb{P}(\Omega_1) > 0$ and such that for any $\omega \in \Omega_1$ we have $\tau(\omega) < \infty$ and $\lim_{t \nearrow \tau(\omega)} u(t)(\omega) = \bar{v}(\omega) \in V$. Let us take an increasing sequence $\{\sigma_n\}_{n=1}^\infty$ of stopping times such that $\sigma_n < \tau$ a.s. and $\sigma_n \nearrow \tau$ a.s. Since the function $f_n : \Omega \ni (\omega) \mapsto u(\sigma_n(\omega))(\omega) \in V$, is \mathcal{F}_{σ_n} -measurable, see e.g. [33, Proposition I.2.18], and hence \mathcal{F}_τ -measurable, the set

$$\Omega_2 := \{\omega \in \Omega : \tau(\omega) < \infty \text{ and } \lim_n f_n(\omega) := \bar{v}(\omega) \text{ exists}\}$$

belongs to the σ -algebra \mathcal{F}_τ and the function \bar{v} is \mathcal{F}_τ -measurable. Note that obviously $\Omega_1 \subset \Omega_2$ so that $\mathbb{P}(\Omega_2) > 0$.

Let $v(t)$, $\tau(\omega) \leq t < \tau_2(\omega)$, be the solution to (5.9) with initial condition $v_{\tau(\omega)}(\omega) = \bar{v}(\omega)1_{\Omega_2}(\omega)$. Then, by [6, Corollary 2.9] the process $\tilde{u}(t)$ defined by

$$\tilde{u}(t)(\omega) = \begin{cases} u(t)(\omega), & \text{if } t < \tau(\omega), \\ v(t)(\omega), & \text{if } \omega \in \Omega_2 \text{ and } t \in [\tau(\omega), \tau_2(\omega)) \end{cases}$$

is a solution to (5.9). Obviously this contradicts the maximality of the solution $u(t)$.

Part II. Let us assume that there exists $\varepsilon > 0$ such that

$$\mathbb{P}(\{\tau < \infty\} \cap \{\limsup_{t \nearrow \tau} |u|_{X_t} < \infty\}) = 4\varepsilon.$$

Since the function $t \mapsto |u|_{X_t}$ is increasing, it means that

$$\mathbb{P}(\{\tau < \infty\} \cap \{\sup_{t \in [0, \tau)} |u|_{X_t} < \infty\}) = 4\varepsilon > 0.$$

Hence there exists $R > 0$ such that

$$\mathbb{P}(\{\tau < \infty\} \cap \{\sup_{t < \tau_\infty} |u(t)|_{X_t} < R\}) = 3\varepsilon > 0.$$

Define (our definition implies that $\sigma_R = \tau$, possibly ∞ , when the set $\{\omega \in \Omega : |u|_{X_t} < R, t \in [0, \tau)\}$ is empty)

$$\sigma_R = \inf\{t \in [0, \tau] : |u|_{X_t} \geq R\}.$$

It is known that σ_R is a predictable stopping time and the set $\tilde{\Omega} = \{\sigma_R = \tau\} \cap \{\tau < \infty\}$ is \mathcal{F}_{σ_R} -measurable. Note also that $\tilde{\Omega} = \{\tau < \infty\} \cap \{\sup_{t < \tau_\infty} |u(t)|_{X_t} < R\}$ and so

$$\mathbb{P}(\tilde{\Omega}) \geq 3\varepsilon.$$

Next, define two processes x and y by

$$x(t) = 1_{\{t < \sigma_R\}} G(u(t)), \quad y(t) = 1_{\{t < \sigma_R\}} F(u(t)) \quad t \geq 0.$$

Note that $x(t) = 0$ or $|u|_{X_t} < R$.

Since $F(0) = 0$ by assumptions, from (5.3) we infer that

$$|F(x)|_H \leq C \|x\|^{p+1-\alpha} |x|_E^\alpha + C \|x\|^{q+1-\gamma} |x|_E^\gamma, \quad x \in E.$$

Similarly, from (5.4)

$$\|G(x)\| \leq C \left[1 + |x|_E^\beta \|x\|^{k+1-\beta} \right], \quad x \in E.$$

Let us now fix $T > 0$. From elementary calculus we obtain

$$\int_0^T |y(t)|_H^2 dt = \mathbb{E} \int_0^T 1_{[0, \sigma_R)}(t) |F(u(t))|_H^2 dt \leq J_{1,p} + J_{2,q},$$

where

$$J_{1,p} := C^2 \int_0^{T \wedge \sigma_R} \|u(t)\|^{2(p+1-\alpha)} |u(t)|_E^{2\alpha} dt,$$

and

$$J_{2,q} := c^2 \int_0^{T \wedge \sigma_R} \|u(t)\|^{2(q+1-\gamma)} |u(t)|_E^{2\gamma} dt.$$

We can easily check that

$$\begin{aligned} J_{1,p} &\leq C^2 \left[\sup_{t \in [0, T \wedge \sigma_R)} \|u(t)\|^{2(p+1-\alpha)} \int_0^{T \wedge \sigma_R} |u(t)|_E^{2\alpha} dt \right] \\ &\leq C^2 \left[\sup_{t \in [0, T \wedge \sigma_R)} \|u(t)\|^{2(p+1-\alpha)} \left(\int_0^{T \wedge \sigma_R} |u(t)|_E^2 dt \right)^\alpha T^{1-\alpha} \right] \\ &\leq C^2 T^{1-\alpha} \left[(1-\alpha) \sup_{t \in [0, T \wedge \sigma_R)} \|u(t)\|^{2(1+\frac{p}{1-\alpha})} + \alpha \int_0^{T \wedge \sigma_R} |u(t)|_E^2 dt \right] \\ &\leq C^2 T^{1-\alpha} \left[|u|_{X_{T \wedge \sigma_R}}^{2(1+\frac{p}{1-\alpha})} \right] \leq C^2 T^{1-\alpha} R^{2(1+\frac{p}{1-\alpha})}. \end{aligned}$$

With similar argument we can prove that

$$J_{2,q} \leq C^2 T^{1-\gamma} R^{2(1+\frac{q}{1-\gamma})}.$$

In particular, we obtain that

$$\mathbb{E} \int_0^T |y(t)|_H^2 dt \leq \mathbb{E} C^2 [T^{1-\alpha} R^{2(1+\frac{p}{1-\alpha})} + T^{1-\gamma} R^{2(1+\frac{q}{1-\gamma})}] < \infty.$$

In a very similar manner we can show that

$$\mathbb{E} \int_0^T \|x(t)\|_V^2 dt < \infty.$$

This allows us to define a process v by

$$v(t) = S(t)u_0 + \int_0^t S(t-s)y(s)ds + \int_0^\infty 1_{[0,t)} S(t-s)x(s) dW(s), \quad t \geq 0. \quad (5.23)$$

The process v is defined globally and it is V -valued continuous. Moreover, since $y = F(u)$ and $x = G(u)$, on $[0, \sigma_R)$, we infer by [6, Remark 2.27] that $u = v$ on $[0, \sigma_R)$. In particular, there exist, a.s.

$$\lim_{t \nearrow \sigma_R} u(t) \text{ in } V.$$

In particular, the above limit exists on the set $\tilde{\Omega}$. Thus, a.s. on $\tilde{\Omega}$,

$$\lim_{t \nearrow \tau} u(t) \text{ in } V.$$

Clearly, this contradicts the first part of the proposition. \square

5.2. An abstract result. In this subsection we want to establish a general theorem of existence of mild solution to the following abstract SPDEs

$$u(t) = S(t)u_0 + \int_0^t S(t-s)F(u(s))ds + \int_0^t S(t-s)G(u(s))dW(s), \quad \text{for any } t > 0, \quad (5.24)$$

where $S(t)$, $t \in [0, \infty)$ is a semigroup, F and G are nonlinear map satisfying Assumption 5.3, Assumption 5.1, and Assumption 5.2, respectively.

Let $\theta : \mathbb{R}_+ \rightarrow [0, 1]$ be a C_c^∞ non increasing function such that

$$\inf_{x \in \mathbb{R}_+} \theta'(x) \geq -1, \quad \theta(x) = 1 \text{ iff } x \in [0, 1] \quad \text{and } \theta(x) = 0 \text{ iff } x \in [2, \infty). \quad (5.25)$$

and for $n \geq 1$ set $\theta_n(\cdot) = \theta(\frac{\cdot}{n})$. Note that if $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a non decreasing function, then for every $x, y \in \mathbb{R}$,

$$\theta_n(x)h(x) \leq h(2n), \quad |\theta_n(x) - \theta_n(y)| \leq \frac{1}{n}|x - y|. \quad (5.26)$$

Proposition 5.12. *Let F be a nonlinear mapping satisfying Assumption 5.1. Let us consider a map*

$$\Phi_T^n = \Phi_T : X_T \ni u \mapsto \theta_n(|u|_{X_T})F(u) \in L^2(0, T; H).$$

Then Φ_T^n is globally Lipschitz and moreover, for any $u_1, u_2 \in X_T$,

$$|\Phi_T^n(u_1) - \Phi_T^n(u_2)|_{L^2(0, T; H)} \leq C[2nC + 1] \left[(2n)^{p+1}T^{(1-\alpha)/2} + (2n)^{q+1}T^{(1-\gamma)/2} \right] |u_1 - u_2|_{X_T}. \quad (5.27)$$

The proof is based on a proof from the paper [12] which in turn was based on a proof from papers by De Bouard and Debussche [17, 18]. For simplicity, we will write Φ_T instead of Φ_T^n .

Proof. Note that $\Phi_T(0) = 0$. Assume that $u_1, u_2 \in X_T$. Denote, for $i = 1, 2$,

$$\tau_i = \inf\{t \in [0, T] : |u_i|_{X_t} \geq 2n\}.$$

Note that by definition, if the set on the RHS above is empty, then $\tau_i = T$.

Without loss of generality we can assume that $\tau_1 \leq \tau_2$. We have the following chain of inequalities/equalities

$$\begin{aligned} |\Phi_T(u_1) - \Phi_T(u_2)|_{L^2(0, T; H)} &= \left[\int_0^T |\theta_n(|u_1|_{X_t})F(u_1(t)) - \theta_n(|u_2|_{X_t})F(u_2(t))|_H^2 dt \right]^{1/2} \\ &\quad \text{because for } i = 1, 2, \quad \theta_n(|u_i|_{X_t}) = 0 \text{ for } t \geq \tau_i \\ &= \left[\int_0^{\tau_2} |\theta_n(|u_1|_{X_t})F(u_1(t)) - \theta_n(|u_2|_{X_t})F(u_2(t))|_H^2 dt \right]^{1/2} \\ &= \left[\int_0^{\tau_2} |\theta_n(|u_1|_{X_t}) - \theta_n(|u_2|_{X_t})|_H^2 F(u_2(t))^2 + \theta_n(|u_1|_{X_t})[F(u_1(t)) - F(u_2(t))]_H^2 dt \right]^{1/2} \\ &\leq \left[\int_0^{\tau_2} |\theta_n(|u_1|_{X_t}) - \theta_n(|u_2|_{X_t})|_H^2 F(u_2(t))^2 dt \right]^{1/2} \\ &\quad + \left[\int_0^{\tau_2} |\theta_n(|u_1|_{X_t})[F(u_1(t)) - F(u_2(t))]_H^2 dt \right]^{1/2} =: A + B \end{aligned}$$

Next, since θ_n is Lipschitz with Lipschitz constant $2n$ we have

$$\begin{aligned}
A^2 &= \int_0^{\tau_2} |\theta_n(|u_1|_{X_t}) - \theta_n(|u_2|_{X_t})|^2 |F(u_2(t))|^2 dt \\
&\leq 4n^2 C^2 \int_0^{\tau_2} [|u_1|_{X_t} - |u_2|_{X_t}]^2 |F(u_2(t))|_H^2 dt \\
&\quad \text{by Minkowski inequality} \\
&\leq 4n^2 C^2 \int_0^{\tau_2} |u_1 - u_2|_{X_t}^2 |F(u_2(t))|_H^2 dt \leq 4n^2 C^2 \int_0^{\tau_2} |u_1 - u_2|_{X_T}^2 |F(u_2(t))|_H^2 dt \\
&\leq 4n^2 C^2 |u_1 - u_2|_{X_T}^2 \int_0^{\tau_2} |F(u_2(t))|_H^2 dt
\end{aligned}$$

Next, by assumptions

$$\begin{aligned}
\int_0^{\tau_2} |F(u_2(t))|_H^2 dt &\leq C^2 \int_0^{\tau_2} \|u(t)\|^{2p+2-2\alpha} |u(t)|_E^{2\alpha} dt + C^2 \int_0^{\tau_2} \|u(t)\|^{2q+2-2\gamma} |u(t)|_E^{2\gamma} dt \\
&:= I_{1,p} + I_{2,q}.
\end{aligned}$$

By some elementary calculations we obtain

$$\begin{aligned}
I_{1,p} &\leq C^2 \sup_{t \in [0, \tau_2]} \|u(t)\|^{2p+2-2\alpha} \left(\int_0^{\tau_2} |u(t)|_E^2 dt \right)^\alpha \tau_2^{1-\alpha} \\
&\leq C^2 \tau_2^{1-\alpha} |u|_{X_{\tau_2}}^{2p+2} \leq C^2 \tau_2^{1-\alpha} (2n)^{2p+2},
\end{aligned}$$

and

$$\begin{aligned}
I_{2,q} &\leq C^2 \sup_{t \in [0, \tau_2]} \|u(t)\|^{2q+2-2\gamma} \left(\int_0^{\tau_2} |u(t)|_E^2 dt \right)^\gamma \tau_2^{1-\gamma} \\
&\leq C^2 \tau_2^{1-\gamma} |u|_{X_{\tau_2}}^{2q+2} \leq C^2 \tau_2^{1-\gamma} (2n)^{2q+2}.
\end{aligned}$$

Therefore,

$$A \leq C^2 [\tau_2^{(1-\alpha)/2} (2n)^{p+2} + \tau_2^{(1-\gamma)/2} (2n)^{q+2}] |u_1 - u_2|_{X_T}.$$

Also, because $\theta_n(|u_1|_{X_t}) = 0$ for $t \geq \tau_1$, and $\tau_1 \leq \tau_2$, we have

$$\begin{aligned}
B &= \left[\int_0^{\tau_2} |\theta_n(|u_1|_{X_t}) [F(u_1(t)) - F(u_2(t))]|_H^2 dt \right]^{1/2} \\
&= \left[\int_0^{\tau_1} |\theta_n(|u_1|_{X_t}) [F(u_1(t)) - F(u_2(t))]|_H^2 dt \right]^{1/2} \\
&\quad \text{because } \theta_n(|u_1|_{X_t}) \leq 1 \text{ for } t \in [0, \tau_1] \\
&\leq \left[\int_0^{\tau_1} |F(u_1(t)) - F(u_2(t))|_H^2 dt \right]^{1/2} \\
&\leq \tilde{B}_{,p} + \tilde{B}_{,q},
\end{aligned}$$

where

$$\begin{aligned}
\tilde{B}_{,p} &:= C \left[\int_0^{\tau_1} |u_1(t) - u_2(t)|_E^{2\alpha} \|u_1(t) - u_2(t)\|^{2-2\alpha} \|u_2(t)\|^{2p} dt \right]^{1/2} \\
&\quad + C \left[\int_0^{\tau_1} \|u_1(t) - u_2(t)\|^2 \|u_1(t)\|^{2p-2\alpha} |u_1(t)|_E^{2\alpha} dt \right]^{1/2},
\end{aligned}$$

and

$$\begin{aligned}
\tilde{B}_{,q} &:= C \left[\int_0^{\tau_1} |u_1(t) - u_2(t)|_E^{2\gamma} \|u_1(t) - u_2(t)\|^{2-2\gamma} \|u_2(t)\|^{2q} dt \right]^{1/2} \\
&\quad + C \left[\int_0^{\tau_1} \|u_1(t) - u_2(t)\|^2 \|u_1(t)\|^{2p-2\gamma} |u_1(t)|_E^{2\gamma} dt \right]^{1/2}.
\end{aligned}$$

Now we have

$$\begin{aligned}
\tilde{B}_{,p} &\leq C \sup_{t \in [0, \tau_1]} \|u_1(t) - u_2(t)\| \|u_1(t)\|^{p-\alpha} \left[\int_0^{\tau_1} |u_1(t)|_E^{2\alpha} dt \right]^{1/2} \\
&+ C \sup_{t \in [0, \tau_1]} \|u_1(t) - u_2(t)\|^{1-\alpha} \|u_2(t)\|^p \left[\int_0^{\tau_1} |u_1(t) - u_2(t)|_E^{2\alpha} dt \right]^{1/2} \\
&\leq C \sup_{t \in [0, T]} \|u_1(t) - u_2(t)\| \sup_{t \in [0, \tau_1]} \|u_1(t)\|^{p-\alpha} \left[\int_0^{\tau_1} |u_1(t)|_E^2 dt \right]^{\alpha/2} \tau_1^{(1-\alpha)/2} \\
&+ C \sup_{t \in [0, T]} \|u_1(t) - u_2(t)\|^{1-\alpha} \sup_{t \in [0, \tau_1]} \|u_2(t)\|^p \left[\int_0^{\tau_1} |u_1(t) - u_2(t)|_E^2 dt \right]^{\alpha/2} \tau_1^{(1-\alpha)/2} \\
&\leq C |u_1 - u_2|_{X_T} |u_1|_{X_{\tau_1}}^p \tau_1^{(1-\alpha)/2} + C |u_1 - u_2|_{X_T} |u_2|_{X_{\tau_1}}^p \tau_1^{(1-\alpha)/2} \\
&\quad \text{because } |u_1|_{X_{\tau_1}} \leq 2n \text{ and } |u_2|_{X_{\tau_1}} \leq |u_2|_{X_{\tau_2}} \leq 2n \\
&\leq C \tau_1^{(1-\alpha)/2} |u_1 - u_2|_{X_T} \left[|u_1|_{X_{\tau_1}}^p + |u_2|_{X_{\tau_1}}^p \right] \leq C (2n)^{p+1} \tau_1^{(1-\alpha)/2} |u_1 - u_2|_{X_T}.
\end{aligned}$$

In exactly the same way,

$$\tilde{B}_{,q} \leq C (2n)^{q+1} \tau_1^{(1-\gamma)/2} |u_1 - u_2|_{X_T}.$$

Summing up, we proved

$$\begin{aligned}
|\Phi_T(u_1) - \Phi_T(u_2)|_{L^2(0, T; H)} &\leq C^2 [\tau_2^{(1-\alpha)/2} (2n)^{p+2} + \tau_2^{(1-\gamma)/2} (2n)^{q+2}] |u_1 - u_2|_{X_T} \\
&+ C [(2n)^{p+1} \tau_1^{(1-\alpha)/2} + (2n)^{q+1} \tau_1^{(1-\gamma)/2}] |u_1 - u_2|_{X_T} \\
&= C [2nC + 1] \left[(2n)^{p+1} \tau_2^{(1-\alpha)/2} + (2n)^{q+1} \tau_2^{(1-\gamma)/2} \right] |u_1 - u_2|_{X_T}
\end{aligned}$$

The proof is complete. \square

In what follows the function Φ_T^n from Proposition 5.12 will be denoted by $\Phi_{F,T}^n$. The following result is a special case of Proposition 5.12 with $H = V$.

Corollary 5.13. *Let G be a nonlinear mapping satisfying Assumption 5.2. Define a map Φ_G^n by*

$$\Phi_G = \Phi_G^n = \Phi_{G,T}^n : X_T \ni u \mapsto \theta_n(|u|_X) G(u) \in L^2(0, T; V). \quad (5.28)$$

Then $\Phi_{G,T}^n$ is globally Lipschitz and moreover, for any $u_1, u_2 \in X_T$,

$$|\Phi_{G,T}^n(u_1) - \Phi_{G,T}^n(u_2)|_{L^2(0, T; V)} \leq C_G (2n)^{k+1} [2nC_G + 1] T^{(1-\beta)/2} |u_1 - u_2|_{X_T}. \quad (5.29)$$

Proposition 5.14. *Let Assumption 5.1-Assumption 5.3 hold. Consider a map*

$$\Psi_T^n = \Psi_T : M^2(X_T) \ni u \mapsto Su_0 + S * \Phi_{F,T}^n(u) + S \diamond \Phi_{G,T}^n(u) \in M^2(X_T) \quad (5.30)$$

Then Ψ_T is globally Lipschitz and moreover, for any $u_1, u_2 \in M^2(X_T)$,

$$|\Psi_T^n(u_1) - \Psi_T^n(u_2)|_{M^2(X_T)} \leq \hat{C}(n) \left[T^{(1-\alpha)/2} \vee T^{(1-\gamma)/2} \vee T^{(1-\beta)/2} \right] |u_1 - u_2|_{M^2(X_T)}, \quad (5.31)$$

where

$$\hat{C}(n) = C_1 C_F [2nC_F + 1] \left[(2n)^{p+1} + (2n)^{q+1} \right] + C_2 C_G (2n)^{k+1} [2nC_G + 1].$$

Proof. For simplicity of notation we will write Ψ_T instead of Ψ_T^n . We will also write Ψ_F (resp. Ψ_G) instead of $\Psi_{F,T}$ (resp. $\Psi_{G,T}$).

Obviously in view of Assumption 5.3 the map Ψ_T is well defined. Let us fix $u_1, u_2 \in M^2(X_T)$. Then

$$\begin{aligned}
|\Psi_T(u_1) - \Psi_T(u_2)|_{M^2(X_T)} &\leq |S * \Phi_F(u_1) - S * \Phi_F(u_2)|_{M^2(X_T)} \\
&\quad + |S \diamond \Phi_G(u_1) - S \diamond \Phi_G(u_2)|_{M^2(X_T)} \\
&\leq C_1 |\Phi_F(u_1) - \Phi_F(u_2)|_{M^2(0,T;H)} + C_2 |\Phi_G(u_1) - \Phi_G(u_2)|_{M^2(X_T)} \\
&\leq \left[C_1 C_F [2n C_F + 1] \left[(2n)^{p+1} T^{(1-\alpha)/2} + (2n)^{q+1} T^{(1-\gamma)/2} \right] \right. \\
&\quad \left. + C_2 C_G (2n)^{k+1} [2n C_G + 1] T^{(1-\beta)/2} \right] \\
&\leq \hat{C}(n) \left[T^{(1-\alpha)/2} \vee T^{(1-\gamma)/2} \vee T^{(1-\beta)/2} \right],
\end{aligned}$$

where

$$\hat{C}(n) = C_1 C_F [2n C_F + 1] \left[(2n)^{p+1} + (2n)^{q+1} \right] + C_2 C_G (2n)^{k+1} [2n C_G + 1].$$

The proof is complete. \square

The first main result of this subsection is given in the following theorem.

Theorem 5.15. *Suppose that Assumption 5.1-Assumption 5.3 hold. Then for every \mathcal{F}_0 -measurable V -valued square integrable random variable u_0 there exists a local process $u = (u(t), t \in [0, T_1])$ which is the unique local mild solution to our problem. Moreover, given $R > 0$ and $\varepsilon > 0$ there exists $\tau(\varepsilon, R) > 0$, such that for every \mathcal{F}_0 -measurable V -valued random variable u_0 satisfying $\mathbb{E}\|u_0\|^2 \leq R^2$, one has*

$$\mathbb{P}(T_1 \geq \tau(\varepsilon, R)) \geq 1 - \varepsilon.$$

Proof. Without loss of generality we can assume that $p \geq q \vee k$. Owing to Proposition 5.14 we can argue as in [12, Proposition 5.1] to prove the first part of the theorem. For any $n \in \mathbb{N}$, $T > 0$ and $u_0 \in L^2(\Omega, \mathbb{P}; V)$ let Ψ_{T, u_0}^n be the mapping from $M^2(X_T)$ defined by

$$\Psi_{T, u_0}^n(u) = S_t u_0 + \int_0^t S_{t-r} [\theta_n(|u|_{X_r}) F(u(r))] dr + \int_0^t S_{t-r} [\theta_n(|u|_{X_r}) G(u(r))] dW(r).$$

It follows from Proposition 5.12, Corollary 5.13 and Assumption 5.3 that Ψ_{T, u_0}^n maps $M^2(X_T)$ into itself. From Proposition 5.14 we deduce that Ψ_{T, u_0}^n is globally Lipschitz, moreover it is a strict contraction for small T . Therefore we can find $\delta > 0$ such that Ψ_{T, u_0}^n is $\frac{1}{2}$ -contraction. For $k \in \mathbb{N}$ let $(t \wedge \tau_k)_{k \in \mathbb{N}}$ be a sequence of times defined by $t \wedge \tau_k = k\delta$. By the $\frac{1}{2}$ -contraction property of Ψ_{T, u_0}^n we can find $u^{[n,1]} \in M^2(X_\delta)$ such that $u^{[n,1]} = \Psi_{\delta, u_0}^n(u^{[n,1]})$. Since $u^{[n,1]} \in M^2(X_\delta)$ it follows that $u^{[n,1]}(t)$ is \mathcal{F}_t -measurable and $u^{[n,1]}(t) \in L^2(\Omega, \mathbb{P}; V)$ for any $t \in [0, \delta]$. Thus replacing u_0 with $u^{[n,1]}(\delta)$ and using the same argument as above we can find $u^{[n,2]} \in M^2(X_\delta)$ such that $u^{[n,2]} = \Psi_{\delta, u^{[n,1]}(\delta)}^n(u^{[n,2]})$. By induction we can construct a sequence $u^{[n,k]} \in M^2(X_\delta)$ such that $u^{[n,k]} = \Psi_{\delta, u^{[n,k-1]}(\delta)}^n(u^{[n,k]})$. Now let u^n be the process defined by $u^n(t) = u^{[n,1]}(t)$, $t \in [0, \delta]$, and for $k = \lceil \frac{T}{\delta} \rceil + 1$ and $0 \leq t < \delta$, let $u^n(t + k\delta) = u^{[n,k]}(t)$. By construction $u^n \in M^2(X_T)$ and $u^n = \Psi_{T, u_0}^n(u^n)$, consequently u^n is a global solution to the truncated equation

$$u(t) = S_t u_0 + \int_0^t S_{t-r} [\theta_n(|u|_{X_r}) F(u(r))] dr + \int_0^t S_{t-r} [\theta_n(|u|_{X_r}) G(u(r))] dW(r).$$

Arguing exactly as in [12, Theorem 4.9] we can show that it is unique. Now let $(\tau_n)_{n \in \mathbb{N}}$ be the sequence of stopping times defined by

$$\tau_n = \inf\{t \in [0, T] : |u^n|_{X_t} \geq n\}. \quad (5.32)$$

By definition $\theta_n(|u^n|_{X_r}) = 1$ for $r \in [0, t \wedge \tau_n]$, hence

$$\theta_n(|u^n|_{X_r}) F(u^n(r)) = F(u^n(r)), r \in [0, t \wedge \tau_n].$$

From [11, Lemma A.1] we infer that

$$\int_0^{t \wedge \tau_n} S_{t \wedge \tau_n - r} [\theta_n(|u|_{X_r}) G(u(r))] dW(r) = \int_0^{t \wedge \tau_n} S_{t \wedge \tau_n - r} G(u^n(r)) dW(r).$$

From these remarks we easily deduce that u^n satisfies

$$u^n(t \wedge \tau^n) = S_{t \wedge \tau_n} u_0 + \int_0^{t \wedge \tau_n} S_{t \wedge \tau_n - r} F(u^n(r)) dr + \int_0^{t \wedge \tau_n} S_{t \wedge \tau_n - r} G(u^n(r)) dW(r).$$

Since τ_n is an accessible stopping time it follows that the process $(u^n(t), t < \tau_n)$ is a local solution to (5.9). This ends the proof of the first part of the theorem.

For the second part we will follow the lines of [12, Theorem 5.3]. For this we fix $\varepsilon > 0$ and choose N such that $N \geq 2\varepsilon^{-1/2}$. Thanks to Proposition 5.12, Corollary 5.13 and Assumption 5.3 we can deduce that there exist some positive constants $\tilde{C}_i, i = 1, \dots, 4$ such that for any $u \in M^2(X_T)$ we have

$$\begin{aligned} |S * \Phi_T^n(u)|_{M^2(X_T)} &\leq \tilde{C}_1 \tilde{C}_2 (2n \tilde{C}_2 + 1) \left[(2n)^{p+2} T^{(1-\alpha)/2} + (2n)^{q+2} T^{(1-\gamma)/2} \right], \\ |S \diamond \Phi_G^n(u)|_{M^2(X_T)} &\leq \tilde{C}_3 \tilde{C}_4 (2n)^{k+2} [2n \tilde{C}_4 + 1] T^{(1-\beta)/2}. \end{aligned}$$

Since $\alpha, \beta, \gamma \in [0, 1)$ we infer from these inequalities that there exists a sequence $(K_n(T))_n$ of numerical functions with $\lim_{T \rightarrow 0} \sup_n K_n(T) = 0$ and

$$|S * \Phi_T^n(u) + S \diamond \Phi_G^n(u)|_{M^2(X_T)} \leq K_n(T),$$

for any $u \in M^2(X_T)$. Let us put $n = NR$ for some “large” N to be chosen later and choose $\delta_1(\varepsilon, R) > 0$ such that $K_n(\delta_1(\varepsilon, R)) \leq R$. Let Ψ_T^n be the mapping defined by (5.30). Since $\mathbb{E}\|u_0\|^2 \leq R^2$, we infer by the Assumption 5.3 (namely (5.6)) that

$$\begin{aligned} |\Psi_T^n(u)|_{M^2(X_T)} &\leq C_0 R + K_n(T), \\ &\leq (C_0 + 1)R, \end{aligned}$$

for any $T \leq \delta_1(\varepsilon, R)$. That is, for $T \leq \delta_1(\varepsilon, R)$ the range of Ψ_T^n is included in the ball centered at 0 and of radius $(C_0 + 1)R$ of $M^2(X_T)$. Furthermore, Propositions 5.14 implies that there exists $C > 0$ such that for any $u_1, u_2 \in M^2(X_T)$

$$|\Psi_T^n(u_1) - \Psi_T^n(u_2)|_{M^2(X_T)} \leq CN^{p+1} R^{p+1} (NRC + 1) \left[T^{(1-\alpha)/2} \vee T^{(1-\gamma)/2} \vee T^{(1-\beta)/2} \right] |u_1 - u_2|_{M^2(X_T)}.$$

Hence we can find $\delta_2(\varepsilon, R) > 0$ such that Ψ_T^n is a strict contraction for any $T \leq \delta_2(\varepsilon, R)$. Thus if one puts $\tau(\varepsilon, R) = \delta_1(\varepsilon, R) \wedge \delta_2(\varepsilon, R)$, the mapping Ψ_T^n has a unique fixed point u^n which satisfies

$$\mathbb{E}|u^n|_{X_{\tau(\varepsilon, R)}}^2 \leq (C_0 + 1)^2 R^2.$$

As in [12, Proposition 5.1] we can show that $(u^n(t), t \leq \tau_n)$ is a local solution to problem (5.24). By the definition of the stopping time τ_n the set $\{\tau_n \leq \tau(\varepsilon, R)\}$ is contained in the set $\{|u^n|_{X_{\tau(\varepsilon, R)}} \geq n\}$. By Chebychev’s inequality we have

$$\begin{aligned} \mathbb{P}(\tau_n \leq \tau(\varepsilon, R)) &\leq \mathbb{P}(|u^n|_{X_{\tau(\varepsilon, R)}}^2 \geq n), \\ &\leq (C_0 + 1)^2 N^{-2}, \\ &\leq \varepsilon, \end{aligned}$$

for $N \geq (C_0 + 1)\varepsilon^{\frac{1}{2}}$. Therefore for $N \geq (C_0 + 1)\varepsilon^{\frac{1}{2}}$ we have $\mathbb{P}(\tau_n \geq \tau(\varepsilon, R)) \geq 1 - \varepsilon$ and the stopping time $T_1 = \tau_n$ satisfies the requirements of the theorem. This concludes the proof. \square

Let $(\tau_n)_{n \in \mathbb{N}}$ be the sequence of stopping times defined by (5.32) and

$$\tau_\infty(\omega) = \lim_{n \rightarrow \infty} \tau_n(\omega), \quad \omega \in \Omega.$$

Let also $(u(t), t < \tau_\infty)$ be the local process defined by

$$u(t, \omega) = u^n(t, \omega) \text{ if } t < \tau_n(\omega), \omega \in \Omega.$$

The next result is about the existence and uniqueness of a maximal solution and the characterization of its lifespan.

Theorem 5.16. *For every $u_0 \in L^2(\Omega, \mathcal{F}_0, V)$, the process $u = (u(t), t < \tau_\infty)$ defined above is the unique local maximal solution to our equation. Moreover, $\mathbb{P}(\{\tau_\infty < \infty\} \cap \{\sup_{t < \tau_\infty} |u(t)|_V < \infty\}) = 0$ and on $\{\tau_\infty < \infty\}$, $\limsup_{t \rightarrow \tau_\infty} |u(t)|_V = +\infty$ a.s.*

Proof. One can use similar argument as in [5, Theorem 4.10] to show that (u, τ_∞) is a local solution to our problem. Since $\tau_n \nearrow \tau_\infty$ we have $\tau_n \leq \tau_\infty < \infty$ on $\{\tau_\infty < \infty\}$ for any n . This implies in particular that for any $n > 0$ there exists a constant δ such that $|u|_{X_t} \geq |u|_{X_{\tau_n}} \geq n$ whenever $t - \tau_\infty < \delta$ on $\{\tau_\infty < \infty\}$. This yields that $\lim_{t \nearrow \tau_\infty} |u|_{X_t} \rightarrow \infty$ on $\{\tau_\infty < \infty\}$. This concludes that $u = (u(t), t < \tau_\infty)$ is a maximal solution.

To prove the second statement we argue by contradiction. Assume that for some $\varepsilon > 0$,

$$\mathbb{P}(\{\tau_\infty < \infty\} \cap \{t \in [0, \tau_\infty) | u(t)|_V < \infty\}) = 4\varepsilon > 0.$$

Let $R > 0$ and assume that

$$\mathbb{P}(\{\tau_\infty < \infty\} \cap \{|u(t)|_V < R \text{ for all } t \in [0, \tau_\infty)\}) \geq 3\varepsilon.$$

Let $\sigma_R = \inf\{t \in [0, \tau_\infty) : |u(t)|_V \geq R\}$ and $\tilde{\Omega} = \{\sigma_R = \tau_\infty < \infty\}$.

Note that

$$\tilde{\Omega} = \{\tau_\infty < \infty\} \cap \{|u(t)|_V < R \text{ for all } t \in [0, \tau_\infty)\}$$

With this R and previous ε we can choose a number $\tau(\varepsilon, R) > 0$ as in Theorem 5.15. Let us now choose α such that $\alpha = \frac{1}{2}\tau(\varepsilon, R)$. By construction $\tau_n \nearrow \tau_\infty$ almost surely and hence for arbitrary $\delta > 0$ there exists $n_0 > 0$ such that $\mathbb{P}(\Omega_0) \geq (1 - \delta)\mathbb{P}(\tilde{\Omega})$, where $\Omega_0 := \{\omega \in \tilde{\Omega} : \tau_\infty - \tau_{n_0} < \alpha\}$. Choosing $\delta = \frac{1}{3}$ we get $\mathbb{P}(\Omega_0) \geq 2\varepsilon$.

Let $T_0 = \tau_{n_0}$ and

$$y_0 = \begin{cases} u(T_0) & \text{on } \Omega_0, \\ 0 & \text{otherwise.} \end{cases}$$

Note that $\mathbb{E}(|y_0|_V^2) \leq R^2$. Then, thanks to Theorem 5.15 the problem (5.24) with initial condition (starting at T_0) y_0 has a unique local solution denoted by $y(t)$, $t \in [T_0, T_0 + T_1)$. Moreover, the lifespan T_1 of the process $y(\cdot)$ satisfies $\mathbb{P}(T_1 \geq \tau(\varepsilon, R)) > 1 - \varepsilon$. From the first part of our theorem y is the unique maximal solution of (5.24) with initial condition y_0 . Note that $\mathbb{P}(\tau_\infty - T_0 < \frac{1}{2}\tau(\varepsilon, R)) \geq 2\varepsilon$. Hence, we infer that

$$\mathbb{P}(\Omega_1) \geq \varepsilon > 0,$$

where

$$\Omega_1 := \Omega_0 \cap \{T_1 \geq \tau(\varepsilon, R)\}.$$

Next we define a local stochastic process v by

$$v(t, \omega) = \begin{cases} u(t, \omega) & \text{if } \omega \in \Omega_1^c, \\ y(t, \omega) & \text{if } \omega \in \Omega_1 \text{ and } t > T_0, \\ u(t, \omega) & \text{if } \omega \in \Omega_1 \text{ and } t \in [0, T_0], \end{cases}$$

The process $v(\cdot)$ is a local solution of (5.24) with initial condition u_0 . Since

$$\mathbb{P}(T_0 + \frac{1}{2}\tau(\varepsilon, R) > \tau_\infty) \geq \varepsilon,$$

the process $v(\cdot)$ satisfies

$$\begin{aligned} \mathbb{E} \left(|v|_{X_{\tau_\infty + \frac{1}{2}\tau(\varepsilon, R)}} \cdot \mathbb{1}_{\Omega_1} \right) &\leq \mathbb{E} \left(|v|_{X_{T_0 + \tau(\varepsilon, R)}} \cdot \mathbb{1}_{\Omega_1} \right), \\ &\leq \mathbb{E} \left[\left(|u|_{X_{T_0}} + |\mathbb{1}_{[T_0, T_0 + \tau(\varepsilon, R)]} y|_{X_{T_0 + \tau(\varepsilon, R)}} \right) \cdot \mathbb{1}_{\Omega_1} \right], \\ &\leq \mathbb{E} \left(|u|_{X_{T_0}} \mathbb{1}_{\Omega_1} \right) + \mathbb{E} \left(\mathbb{1}_{\Omega_1} |\mathbb{1}_{[T_0, T_0 + \tau(\varepsilon, R)]} y|_{X_{T_0 + \tau(\varepsilon, R)}} \right), \\ &\leq \mathbb{E} \left(|u|_{X_{T_0}} \mathbb{1}_{\Omega_0} \right) + \mathbb{E} \left(\mathbb{1}_{\Omega_1} |y|_{X_{[T_0, T_0 + \tau(\varepsilon, R)]}} \right), \end{aligned}$$

where the space $X_{[a, b]}$ and the norm $|\cdot|_{X_{[a, b]}}$ are defined similarly to the space X_T and norm $|\cdot|_{X_T}$. Since $T_0 = \inf\{t \in [0, \tau_\infty) : |u|_{X_t} \geq n_0\} = \tau_{n_0}$, we infer that T_0 is finite ($T_0 < \tau_\infty$ to be precise) on the set Ω_0 and hence $|u|_{X_{T_0}} = n_0$ on the set Ω_0 . Therefore, the first expected value $\mathbb{E} \left(|u|_{X_{T_0}} \mathbb{1}_{\Omega_0} \right)$ is finite. Since the solution $y(\cdot)$ is such that $\mathbb{1}_{[T_0, T_0 + T_1]} y \in M^2(X_{T_0 + T_1})$ and the lifespan T_1 of $y(\cdot)$ satisfies $\mathbb{P}(T_1 + T_0 \geq T_0 + \tau(\varepsilon, R)) \geq \varepsilon$ we infer that

$$\mathbb{E} \left(|\mathbb{1}_{[T_0, T_0 + \tau(\varepsilon, R)]} y|_{X_{T_0 + \tau(\varepsilon, R)}} \right) \leq \mathbb{E} \left(|\mathbb{1}_{[T_0, T_0 + \tau(\varepsilon, R)]} y|_{X_{T_0 + T_1}}^2 \right) < \infty.$$

Therefore

$$\mathbb{E} \left(|v|_{X_{\tau_\infty + \frac{1}{2}\tau(\varepsilon, R)}} \cdot \mathbb{1}_{\Omega_1} \right) < \infty.$$

This implies in particular that $|v(t)|_{X_{\tau_\infty + \frac{1}{2}\tau(\varepsilon, R)}} < \infty$ on $\Omega_1 \subset \{\tau_\infty < \infty\}$ which contradicts Proposition 5.11. \square

Remark 5.17. (i) The last result is very important since a priori we only know that $|u|_{X_t} \rightarrow \infty$ as $t \nearrow \tau_\infty$ on the set $\{\tau_\infty < \infty\}$.
(ii) The proof of the existence of a global solution could then follow the proof in [12, Theorem 8.12].

6. MAXIMUM PRINCIPLE TYPE THEOREM

In this section we replace in the system (2.12)-(2.13) the general polynomial $f(\mathbf{n})$ by the bounded Ginzburg-Landau function $\mathbb{1}_{|\mathbf{n}| \leq 1}(|\mathbf{n}|^2 - 1)\mathbf{n}$. All our previous result remains true and the analysis are even easier. In the case $f(\mathbf{n}) = \mathbb{1}_{|\mathbf{n}| \leq 1}(|\mathbf{n}|^2 - 1)\mathbf{n}$, we will show that if the initial value \mathbf{n}_0 is in the unit ball, then so are the values of the vector director \mathbf{n} . That is, we must show that $|\mathbf{n}(t)|^2 \leq 1$ almost all $(\omega, t, x) \in \Omega \times [0, T] \times \mathcal{O}$. In fact we have the following theorem.

Theorem 6.1. *Assume that $n \leq 3$ and that a process $(\mathbf{v}, \mathbf{n}) = (\mathbf{v}(t), \mathbf{n}(t))$, $t \in [0, T]$, is a solution to problem (2.12)-(2.13) with initial condition $(\mathbf{v}_0, \mathbf{n}_0)$ such that $|\mathbf{n}_0|^2 \leq 1$ for almost all $(\omega, x) \in \Omega \times \mathcal{O}$. Then $|\mathbf{n}(t)|^2 \leq 1$ for almost all $(\omega, t, x) \in \Omega \times [0, T] \times \mathcal{O}$.*

Proof. We follow the idea in [15, Lemma 2.1] and [21, Proof of Theorem 4, Page 513]. Let $\varphi : \mathbb{R} \rightarrow [0, 1]$ be an increasing function of class \mathcal{C}^∞ such that

$$\begin{aligned} \varphi(s) &= 0 \text{ iff } s \in (-\infty, 1], \\ \varphi(s) &= 1 \text{ iff } s \in [2, +\infty). \end{aligned}$$

Let $\{\varphi_m; m \in \mathbb{N}\}$ and $\{\phi_m; m \in \mathbb{N}\}$ be two sequences of smooth function from \mathbb{R}^n defined by

$$\begin{aligned} \varphi_m(\mathbf{n}) &= \varphi(m(|\mathbf{n}|^2 - 1)), \\ \phi_m(\mathbf{n}) &= (|\mathbf{n}|^2 - 1)\varphi_m(\mathbf{n}), \mathbf{n} \in \mathbb{R}^n. \end{aligned}$$

Define a sequence of function $\{\Psi_m, m \in \mathbb{N}\}$ by

$$\begin{aligned}\Psi_m(\mathbf{n}) &= \|\phi_m(\mathbf{n})\|^2, \\ &= \int_{\mathcal{O}} (|\mathbf{n}|^2 - 1)^2 [\varphi_m(\mathbf{n})]^2 dx, \quad \mathbf{n} \in \mathbb{L}^4(\mathcal{O}),\end{aligned}$$

for any $m \in \mathbb{N}$. It is clear that $\Psi_m : \mathbb{H}^2 \rightarrow \mathbb{R}$ is twice (Fréchet) differentiable and its first and second derivatives satisfy

$$\Psi_m(\mathbf{n})(h) = 4 \int_{\mathcal{O}} ((|\mathbf{n}|^2 - 1) \varphi_m(\mathbf{n}) \mathbf{n} \cdot h) dx + 2m \int_{\mathcal{O}} (|\mathbf{n}|^2 - 1)^2 \varphi_m(\mathbf{n}) (\mathbf{n} \cdot h) dx,$$

and

$$\begin{aligned}\Psi_m''(\mathbf{n})(k, h) &= 8 \int_{\mathcal{O}} \left[\varphi_m(\mathbf{n}) (\mathbf{n} \cdot k) (\mathbf{n} \cdot h) \right] dx + 4 \int_{\mathcal{O}} (\varphi_m(\mathbf{n}) (|\mathbf{n}|^2 - 1) (k \cdot h)) dx \\ &\quad + 16m \int_{\mathcal{O}} ((|\mathbf{n}|^2 - 1) \varphi_m(\mathbf{n}) (\mathbf{n} \cdot k) (\mathbf{n} \cdot h)) dx \\ &\quad + 4m^2 \int_{\mathcal{O}} ((|\mathbf{n}|^2 - 1)^2 \varphi_m''(\mathbf{n}) (\mathbf{n} \cdot k) (\mathbf{n} \cdot h)) dx \\ &\quad + 2m \int_{\mathcal{O}} (|\mathbf{n}|^2 - 1)^2 \varphi_m(\mathbf{n}) (k \cdot h) dx,\end{aligned}$$

for any $\mathbf{n} \in \mathbb{H}^2$ and $h, k \in \mathbb{L}^2(\mathcal{O})$. In particular, for any k, h such that $k \perp \mathbf{n}$ and $h \perp \mathbf{n}$

$$\begin{aligned}\Psi_m(\mathbf{n})(h) &= 0, \\ \Psi_m''(\mathbf{n})(k, h) &= 4 \int_{\mathcal{O}} (|\mathbf{n}|^2 - 1) \varphi_m(\mathbf{n}) (k \cdot h) dx + 2m \int_{\mathcal{O}} (|\mathbf{n}|^2 - 1)^2 \varphi_m(\mathbf{n}) (k \cdot h) dx.\end{aligned}$$

It follows from Itô's formula (see [46, Theorem I.3.3.2, Page 147]) that

$$d[\Psi_m(\mathbf{n})] = \Psi_m(\mathbf{n}) \left(-A_1 \mathbf{n} - \tilde{B}(\mathbf{v}, \mathbf{n}) - \frac{1}{\varepsilon^2} f(\mathbf{n}) + \frac{1}{2} G^2(\mathbf{n}) \right) dt + \frac{1}{2} \Psi_m''(\mathbf{n})(G(\mathbf{n}), G(\mathbf{n})) dt.$$

The integral stochastic vanishes because $G(\mathbf{n}) \perp \mathbf{n}$. Owing to the identity

$$-|\mathbf{n} \times h|^2 = \mathbf{n} \cdot ((\mathbf{n} \times h) \times h),$$

we have

$$\frac{1}{2} \Psi_m''(G(\mathbf{n}), G(\mathbf{n})) + \frac{1}{2} \Psi_m'(G^2(\mathbf{n})) = 0.$$

Hence

$$d[\Psi_m(\mathbf{n})] = \Psi_m(\mathbf{n}) \left(-A_1 \mathbf{n} - \tilde{B}(\mathbf{v}, \mathbf{n}) - \frac{1}{\varepsilon^2} f(\mathbf{n}) \right) dt \quad (6.1)$$

Noticing that from the definition of φ_m and the Lebesgue Dominated Convergence Theorem we have for $\mathbf{n} \in \mathbb{H}^2, h \in \mathbb{L}^2$

$$\begin{aligned}\lim_{m \rightarrow \infty} \Psi_m(\mathbf{n}) &= \|(|\mathbf{n}|^2 - 1)_+\|^2, \\ \lim_{m \rightarrow \infty} \Psi_m(\mathbf{n})(k) &= 4 \int_{\mathcal{O}} [(|\mathbf{n}|^2 - 1)_+ \mathbf{n} \cdot h] dx.\end{aligned}$$

Hence, we obtain from letting $m \rightarrow \infty$ in (6.1) that for almost all $(\omega, t) \in \Omega \times [0, T]$

$$y(t) - y(0) + 4 \int_0^t \left(\int_{\mathcal{O}} \left[+A_1 \mathbf{n} + (\mathbf{v} \cdot \nabla) \mathbf{n} + \frac{1}{\varepsilon^2} f(\mathbf{n}) \right] \cdot \left[\mathbf{n} (|\mathbf{n}|^2 - 1)_+ \right] dx \right) ds = 0,$$

where $y(t) = \|(|\mathbf{n}(t)|^2 - 1)_+\|^2$. Let us set $\xi = (|\mathbf{n}|^2 - 1)_+$, it follows from [3, Exercise 7.1.5, p 283] that $\xi \in \mathbb{H}^1$ if $\mathbf{n} \in \mathbb{H}^1$. Thus, since $\frac{\partial \mathbf{n}}{\partial \mathbf{n}} = 0$ we derive from integration-by-parts that

$$-4 \int_0^t \left(\int_{\mathcal{O}} \Delta \mathbf{n} \cdot \mathbf{n} (|\mathbf{n}|^2 - 1)_+ dx \right) ds = \int_0^t \left(\int_{\mathcal{O}} (2 \nabla(|\mathbf{n}|^2) \cdot \nabla \xi + 4 \xi |\nabla \mathbf{n}|^2) dx \right) ds,$$

Since $\xi \geq 0$ and $|\nabla \mathbf{n}|^2 \geq 0$ a.e. $(t, x) \in \mathcal{O} \times [0, T]$ we easily derive from the above identity that

$$-4 \int_0^t \left(\int_{\mathcal{O}} \Delta \mathbf{n} \cdot \mathbf{n} (|\mathbf{n}|^2 - 1)_+ dx \right) ds \geq 2 \int_0^t \left(\int_{\mathcal{O}} \nabla(|\mathbf{n}|^2 - 1) \cdot \nabla \xi dx \right) ds.$$

Thanks to [3, Exercise 7.1.5, p 283] we have

$$\int_0^t \left(\int_{\mathcal{O}} \nabla(|\mathbf{n}|^2 - 1) \cdot \nabla \xi dx \right) ds = \int_0^t \int_{\mathcal{O}} |\nabla(|\mathbf{n}|^2 - 1)|^2 \mathbb{1}_{\{|\mathbf{n}|^2 > 1\}} dx ds,$$

which implies that

$$-4 \int_0^t \left(\int_{\mathcal{O}} \Delta \mathbf{n} \cdot \mathbf{n} (|\mathbf{n}|^2 - 1)_+ dx \right) ds \geq \int_0^t \int_{\mathcal{O}} |\nabla(|\mathbf{n}|^2 - 1)|^2 \mathbb{1}_{\{|\mathbf{n}|^2 > 1\}} dx ds.$$

We also have

$$\begin{aligned} 4 \int_0^t \left(\int_{\mathcal{O}} [(\mathbf{v} \cdot \nabla) \mathbf{n}] \cdot [\mathbf{n} (|\mathbf{n}|^2 - 1)_+] dx \right) ds &= 2 \int_0^t \left(\int_{\mathcal{O}} [(\mathbf{v} \cdot \nabla)(|\mathbf{n}|^2)] [(|\mathbf{n}|^2 - 1)_+] dx \right) ds, \\ &= \int_0^t \left(\int_{\mathcal{O}} (\mathbf{v} \cdot \nabla) \xi \xi dx \right) ds, \\ &= 0. \end{aligned}$$

Since $f(\mathbf{n}) = 0$ for $|\mathbf{n}|^2 > 1$ and $\xi f(\mathbf{n}) = 0$ for $|\mathbf{n}|^2 \leq 1$ we have

$$4 \int_0^t \left(\int_{\mathcal{O}} \xi f(\mathbf{n}) \cdot \mathbf{n} dx \right) ds = 0.$$

Therefore we see that $y(t)$ satisfies the estimate

$$y(t) + 2 \int_0^t \int_{\{|\mathbf{n}|^2 > 1\}} |\nabla(|\mathbf{n}|^2 - 1)_+|^2 ds \leq y(0),$$

for almost all $(\omega, t) \in \Omega \times [0, T]$. Since the second term in the left hand side of the above inequality is positive and $y(0) = \|(|\mathbf{n}_0|^2 - 1)_+\|^2$ and by assumption $|\mathbf{n}_0|^2 \leq 1$ for almost all $(\omega, t, x) \in \Omega \times [0, T] \times \mathcal{O}$ we derive that

$$y(t) = 0,$$

for almost all $(\omega, t) \in \Omega \times [0, T]$, $T \geq 0$. Hence we have $|\mathbf{n}|^2 \leq 1$ a.e. $(\omega, t, x) \in \Omega \times [0, T] \times \mathcal{O}$, $T \geq 0$. \square

APPENDIX A. SOME IMPORTANT ESTIMATES

In this section we recall or establish some crucial estimates needed for the proof of our mains results.

First, let $d \in \{2, 3\}$ and put $a = \frac{d}{4}$. Then the following estimates, valid for all $\mathbf{u} \in \mathbb{W}^{1,4}$, are special cases of Gagliardo-Nirenberg's inequalities:

$$\|\mathbf{u}\|_{\mathbb{L}^4} \leq \|\mathbf{u}\|^{1-a} \|\nabla \mathbf{u}\|^a, \quad (\text{A.1})$$

$$\|\mathbf{u}\|_{\mathbb{L}^\infty} \leq \|\mathbf{u}\|_{\mathbb{L}^4}^{1-a} \|\nabla \mathbf{u}\|_{\mathbb{L}^4}^a. \quad (\text{A.2})$$

The inequality (A.1) can be written in the spirit of the continuous embedding

$$\mathbb{H}^1 \subset \mathbb{L}^4. \quad (\text{A.3})$$

It follows from (A.2) and (A.3) that for $\mathbf{u} \in D(A_1)$

$$\|\mathbf{u}\|_{\mathbb{L}^\infty} \leq \|\mathbf{u}\|_1^{1-a} \|\nabla \mathbf{u}\|_2^a. \quad (\text{A.4})$$

Next we give some properties of the bilinear form B and \tilde{B} defined in Section 2 (see equations (2.5) and (2.4) on page 6, respectively).

Lemma A.1. *The bilinear map $B(\cdot, \cdot)$ maps continuously $\mathbb{V} \times \mathbb{H}^1$ into \mathbb{V}^* and*

$$\langle B(\mathbf{u}, \mathbf{v}), \mathbf{w} \rangle = b(\mathbf{u}, \mathbf{v}, \mathbf{w}), \text{ for any } \mathbf{u} \in \mathbb{V}, \mathbf{v} \in \mathbb{H}^1, \mathbf{w} \in \mathbb{V}, \quad (\text{A.5})$$

$$\langle B(\mathbf{u}, \mathbf{v}), \mathbf{w} \rangle = -b(\mathbf{u}, \mathbf{w}, \mathbf{v}) \text{ for any } \mathbf{u} \in \mathbb{V}, \mathbf{v} \in \mathbb{H}^1, \mathbf{w} \in \mathbb{V}, \quad (\text{A.6})$$

$$\langle B(\mathbf{u}, \mathbf{v}), \mathbf{v} \rangle = 0 \text{ for any } \mathbf{u} \in \mathbb{V}, \mathbf{v} \in \mathbb{V}, \quad (\text{A.7})$$

$$\|B(\mathbf{u}, \mathbf{v})\|_{\mathbb{V}^*} \leq C_0 \|\mathbf{u}\|^{1-\frac{d}{4}} \|\nabla \mathbf{u}\|^{\frac{d}{4}} \|\mathbf{v}\|^{1-\frac{d}{4}} \|\nabla \mathbf{v}\|^{\frac{d}{4}}, \text{ for all } \mathbf{u} \in \mathbb{V}, \mathbf{v} \in \mathbb{H}^1. \quad (\text{A.8})$$

Proof. This lemma is well-known and we refer to [57, Chapter II, Section 1.2] for its proof. \square

With an abuse of notation, we again denote by $\tilde{B}(\cdot, \cdot)$ the restriction of $\tilde{B}(\cdot, \cdot)$ to $\mathbb{V} \times \mathbb{H}^2$.

Lemma A.2. *The bilinear operator \tilde{B} maps continuously $\mathbb{V} \times \mathbb{H}^2$ into \mathbb{L}^2 and there exists $C_1 > 0$ such that*

$$\|\tilde{B}(\mathbf{v}, \mathbf{n})\| \leq C_1 \|\mathbf{v}\|^{1-\frac{d}{4}} \|\nabla \mathbf{v}\|^{\frac{d}{4}} \|\nabla \mathbf{n}\|^{1-\frac{d}{4}} \|\Delta \mathbf{n}\|^{\frac{d}{4}}, \text{ for any } \mathbf{v} \in \mathbb{V}, \mathbf{n} \in \mathbb{H}^2. \quad (\text{A.9})$$

Moreover, we have

$$\langle \tilde{B}(\mathbf{v}, \mathbf{n}), \mathbf{n} \rangle = 0, \text{ for any } \mathbf{v} \in \mathbb{V}, \mathbf{n} \in \mathbb{H}^2. \quad (\text{A.10})$$

Proof. We can argue as in the proof of (A.8) (see also [57, Chapter II, Section 1.2]) to establish the estimate (A.9). The identity (A.10) easily follows by integration-by-parts and by taking into account that $\nabla \cdot \mathbf{v} = 0$ and \mathbf{v} is zero on the boundary. \square

Next let A and A_1 the self-adjoint linear operators defined in Section 2 on page 4. We recall that the space \mathbb{X}_α , $\alpha \geq 0$ are the domain of the fractional power operator $(I + A_1)^{frac{1}{2} + \alpha}$ and $\mathbb{X}_\alpha \subset \mathbb{H}^{1+2\alpha}$ (see the identity (2.3) on page 5). We will give some properties of the semigroups $\{S_1(t) : t \geq 0\}$ and $\{S_2(t) : t \geq 0\}$ generated by $-A$ on \mathbb{H} and $-A_1$ on $\mathbb{X}_{\frac{1}{2}}$, respectively.

Lemma A.3. *Let $T \in (0, \infty)$, $g \in L^2(0, T; \mathbb{X}_0)$ and*

$$\mathbf{u}(t) = \int_0^T S_2(t-s)g(s)ds, = \sum_{k \in \mathbb{N}} \int_0^T e^{-\lambda_k(t-s)} g_k(s) ds, \quad t \in [0, T]. \quad (\text{A.11})$$

Then

$$\mathbf{u} \in C([0, T]; \mathbb{X}_{\frac{1}{2}}) \cap L^2(0, T; \mathbb{X}_1),$$

and

$$\|\mathbf{u}(t)\|_{C([0, T]; \mathbb{X}_{\frac{1}{2}})} + \|\mathbf{u}(t)\|_{L^2(0, T; \mathbb{X}_1)} \leq (1 + \max(T, 1 + \frac{1}{\lambda_2})) \|g(t)\|_{L^2(0, T; \mathbb{X}_0)}.$$

Proof. This result is well-known. We refer the reader to [46]. \square

Similarly we have the following result, see, for instance, [58].

Lemma A.4. *Let $T \in (0, \infty)$, $\tilde{g} \in L^2(0, T; \mathbb{H})$ and*

$$\mathbf{v}(t) = \int_0^T S_1(t-s)\tilde{g}(s)ds.$$

We have

$$\|\mathbf{v}(t)\|_{C([0, T]; D(A^{\frac{1}{2}}))} + \|\mathbf{v}(t)\|_{L^2(0, T; D(A))} \leq (1 + \frac{1}{\mu_1}) \|\tilde{g}\|_{L^2(0, T; \mathbb{H})},$$

where μ_1 is the smallest eigenvalues of the Stokes operator A .

Let ζ_1 be an element of $M^2(0, T; \mathbb{H})$ (resp. $\zeta_2 \in M^2(0, T; \mathbb{X}_{\frac{1}{2}})$) and consider the stochastic convolutions

$$\mathbf{w}_2(t) = \int_0^t \mathbb{S}_2(t-s) \zeta_2(s) dW_2(s),$$

and

$$\mathbf{w}_1(t) = \int_0^t \mathbb{S}_1(t-s) \zeta_1(s) dW_1(s).$$

Lemma A.5. *There exists $C > 0$ such that*

$$\mathbb{E} \|\mathbf{w}_i(t)\|_{C([0,T]; \tilde{V})} + \mathbb{E} \|\mathbf{w}_i(t)\|_{L^2(0,T; \tilde{E})} \leq C \mathbb{E} \|\zeta_i(t)\|_{L^2(0,T; \tilde{V})},$$

where $(\tilde{V}, \tilde{E}) = (D(A^{\frac{1}{2}}), D(A_1))$ if $i = 1$, and $(\tilde{V}, \tilde{E}) = (D(A_1), \mathbb{X}_1)$ if $i = 2$.

Proof. This result is also well-known, see for example [46, 58]. □

APPENDIX B. PROOF OF THE LEMMA 4.10 AND 4.11

In this section we give the proofs of the Lemma Lemmata 4.10-4.11 and we shall start with that of Lemma 4.10.

Proof of Lemma 4.10. First, we will establish (4.19). For this purpose, we set

$$\begin{aligned} \langle \Delta \mathbf{n} - f(\mathbf{n}), \Delta \mathbf{n} \times h \rangle + \langle \Delta \mathbf{n} - f(\mathbf{n}), \mathbf{n} \times \Delta h + (\mathbf{n} \times h) \times \Delta h \rangle - 2 \langle \Delta \mathbf{n} - f(\mathbf{n}), f'(\mathbf{n})(\mathbf{n} \times h) \times h \rangle \\ =: R_1 + R_2 + R_3. \end{aligned}$$

Owing to Cauchy-Schwarz, Cauchy inequalities, the assumption on h and Remark 2.18 we have that

$$\begin{aligned} |R_1| &\leq C \|\Delta \mathbf{n} - f(\mathbf{n})\| \|\Delta \mathbf{n}\| \|h\|_{\mathbb{L}^\infty} \\ &\leq C \|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + C(h) \|\Delta \mathbf{n}\|^2 \\ &\leq C(\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{L}^{\bar{q}}}^{\bar{q}} + 1), \end{aligned}$$

where $\bar{q} = 4N + 2$. In a similar way we can show that

$$\begin{aligned} |R_2| &\leq C \|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{L}^\infty}^2 (\|\Delta h\|^2 [1 + \|h\|_{\mathbb{L}^\infty}^2]) \\ &\leq C(\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{L}^{\bar{q}}}^{\bar{q}} + 1). \end{aligned}$$

Note that we have used the fact that \mathbb{H}^2 is continuously embedded in \mathbb{L}^∞ . Next, thanks to (2.7) we have

$$\begin{aligned} |R_3| &\leq C \|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + C \int_{\mathcal{O}} |f'(\mathbf{n}(x))| [(\mathbf{n} \times h) \times h] \\ &\leq \|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + C \|h\|_{\mathbb{L}^\infty}^2 \int_{\mathcal{O}} (1 + |\mathbf{n}(x)|^{2N})^2 |\mathbf{n}(x)|^2 dx \\ &\leq C(\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{L}^{\bar{q}}}^{\bar{q}} + 1). \end{aligned}$$

Collecting all these inequalities and using the embedding $\mathbb{H}^1 \subset \mathbb{L}^{4N+2}$ we obtain that

$$|R_1 + R_2 + R_3| \leq C(\|\Delta \mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{H}^1}^{\bar{q}} + 1),$$

which completes the proof of (4.19).

Now, we proceed to the proof of (4.20). To this end, set let us take a look at

$$\|\Delta G(\mathbf{n}) - f'(\mathbf{n})G(\mathbf{n})\|^2 - \langle \Delta \mathbf{n} - f(\mathbf{n}), f''(\mathbf{n})[G(\mathbf{n}), G(\mathbf{n})] \rangle =: J_1 + J_2.$$

By using the fact that $|f''(\mathbf{u})| \leq C(1 + |\mathbf{u}|^{N-2})$ for any $\mathbf{u} \in \mathbb{R}^2$, it is easy to show that

$$\begin{aligned} |J_2| &\leq C\|\Delta\mathbf{n} - f(\mathbf{n})\|^2 + C \int_{\mathcal{O}} |f''(\mathbf{n}(x)) [G(\mathbf{n}(x)), G(\mathbf{n}(x))]|^2 dx \\ &\leq C\|\Delta\mathbf{n} - f(\mathbf{n})\|^2 + C\|h\|_{\mathbb{L}^\infty}^2 \int_{\mathcal{O}} |f''(\mathbf{n}(x))|^2 |\mathbf{n}(x)|^4 dx \\ &\leq C(\|\Delta\mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{L}^q}^{\bar{q}} + 1). \end{aligned} \quad (\text{B.1})$$

To deal with J_1 first remark that

$$\begin{aligned} J_1 &= \|\Delta\mathbf{n} \times h + \mathbf{n} \times \Delta h - f'(\mathbf{n})(\mathbf{n} \times h)\|^2 \\ &\leq C(\|h\|^2 \|\Delta\mathbf{n}\|^2 + \|\mathbf{n}\|_{\mathbb{L}^\infty}^2 \|\Delta h\|^2 + \|f''(\mathbf{n})(\mathbf{n} \times h)\|^2). \end{aligned} \quad (\text{B.2})$$

Now, arguing as in the proof of (4.19) we derive that

$$J_1 \leq C(\|\Delta\mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{L}^q}^{\bar{q}} + 1). \quad (\text{B.3})$$

Owing to (B.1), (B.3) and the Sobolev embedding $\mathbb{H}^1 \subset \mathbb{L}^{4N+2}$ we obtain

$$|J_1 + J_2| \leq C(\|\Delta\mathbf{n} - f(\mathbf{n})\|^2 + \|\mathbf{n}\|_{\mathbb{H}^1}^{\bar{q}} + 1),$$

which completes the proof of (4.20) and Lemma 4.10. \square

Now we shall establish Lemma 4.11.

Proof of Lemma 4.11. For any $\mathbf{u} \in \mathbb{H}^1$, we have

$$\begin{aligned} |\langle f'(\mathbf{n})\mathbf{u}, \mathbf{u} \rangle| &= \int_{\mathcal{O}} \tilde{f}(\mathbf{n}(x)) |\mathbf{u}(x)| dx + 2 \int_{\mathcal{O}} \tilde{f}'(\mathbf{n}(x)) |\mathbf{n}(x) \cdot \mathbf{u}(x)|^2 dx \\ &\leq \left[\int_{\mathcal{O}} |\tilde{f}(\mathbf{n}(x))|^2 dx \right]^{\frac{1}{2}} \left[\int_{\mathcal{O}} |\mathbf{u}(x)|^4 dx \right]^{\frac{1}{2}} + 2 \left[\int_{\mathcal{O}} |\tilde{f}'(\mathbf{n}(x))|^2 |\mathbf{n}(x)|^4 dx \right]^{\frac{1}{2}} \left[\int_{\mathcal{O}} |\mathbf{u}(x)|^4 dx \right]^{\frac{1}{2}} \end{aligned}$$

By Gagliardo-Nirenberg, Cauchy inequalities, (2.15) and (2.16) we derive that

$$|\langle f'(\mathbf{n})\mathbf{u}, \mathbf{u} \rangle| \leq \frac{1}{4} \|\nabla \mathbf{u}\|^2 + (\ell_1 + \ell_2 \|\mathbf{n}\|_{\mathbb{L}^{4N}}^{2N})^2 \|\mathbf{u}\|^2.$$

Putting $\mathbf{u} = \Delta\mathbf{n} - f(\mathbf{n})$ in this last inequality and using the Sobolev embedding $\mathbb{H}^1 \subset \mathbb{L}^{4N}$ we infer that

$$|\langle f'(\mathbf{n})[\Delta\mathbf{n} - f(\mathbf{n})], \Delta\mathbf{n} - f(\mathbf{n}) \rangle| \leq \delta_7 \|\nabla[\Delta\mathbf{n} - f(\mathbf{n})]\|^2 + C(\delta_7)(\ell_1 + \ell_2 \|\mathbf{n}\|_{\mathbb{H}^1}^{2N})^2 \|\Delta\mathbf{n} - f(\mathbf{n})\|^2,$$

from which we easily conclude the proof of (4.21) and of Lemma 4.11. \square

APPENDIX C. BASIC ESTIMATES FOR THE SOLUTION (\mathbf{v}, \mathbf{n})

Throughout this section we will assume that the initial data is random and have finite moments in appropriate functional spaces. These assumptions will be made precise in each of the following four crucial propositions. Obviously, the estimates we derive in this section are still valid for non-random initial data. Throughout, (\mathbf{v}, \mathbf{n}) will denote a weak or a strong solution to (1.14).

C.1. Estimates for $\|\mathbf{n}(t)\|$. Before we proceed further we should note that the forthcoming results are proved for $p \geq 4N + 2$, but they will also hold for $p \in [1, 4N + 2)$.

Proposition C.1. Assume that for some $p \geq 4N + 2$ and $t \geq 0$

$$\mathfrak{G}_0(t, p) < \infty \text{ almost surely,}$$

where $\mathfrak{G}_0(t, p)$ is defined in (3.9). Then, the following estimate holds almost surely

$$\|\mathbf{n}(t)\|^p + p \int_0^t \|\mathbf{n}(s)\|^{p-2} \|\nabla \mathbf{n}(s)\|^2 ds + p \int_0^t \|\mathbf{n}(s)\|^{p-2} \|\mathbf{n}(s)\|_{\mathbb{L}^{2N+2}}^{2N+2} ds \leq \mathfrak{G}_0(t, p). \quad (\text{C.1})$$

Proof. Let $\Psi(\cdot)$ be the mapping defined by $\Psi(\mathbf{n}) = \frac{1}{2}\|\mathbf{n}\|^p$ for any $\mathbf{n} \in \mathbb{L}^2$. This mapping is twice Fréchet differentiable with first and second derivatives defined by

$$\begin{aligned}\Psi'(\mathbf{n})[\mathbf{h}] &= p\|\mathbf{n}\|^{p-2}\langle \mathbf{n}, \mathbf{h} \rangle, \\ \Psi''[\mathbf{h}, \mathbf{k}] &= p(p-2)\|\mathbf{n}\|^{p-4}\langle \mathbf{n}, \mathbf{k} \rangle \langle \mathbf{n}, \mathbf{h} \rangle + p\|\mathbf{n}\|^{p-2}\langle \mathbf{h}, \mathbf{k} \rangle.\end{aligned}$$

by straightforward calculation one can check that if $\mathbf{h} \in \mathbb{L}^2$ and $\mathbf{h} \perp \mathbf{n}$ then $\Psi'(\mathbf{n})[\mathbf{h}] = 0$ and $\Psi''(\mathbf{n})[\mathbf{h}, \mathbf{h}] = p\|\mathbf{n}\|^{p-2}\|\mathbf{h}\|^2$. Moreover, if $\mathbf{n} \in \mathbb{H}^1$ then it is possible to extend $\Psi'(\mathbf{n})[\cdot]$ to the dual space $(\mathbb{H}^1)^*$ of \mathbb{H}^1 by setting

$$\Psi(\mathbf{n})[\mathbf{h}] = p\|\mathbf{n}\|^{p-2}_{(\mathbb{H}^1)^*} \langle \mathbf{h}, \mathbf{n} \rangle_{\mathbb{H}^1} \text{ for any } \mathbf{h} \in (\mathbb{H}^1)^*.$$

Note also that if $\mathbf{n} \in \mathbb{H}^1$ then it follows from the Sobolev continuous embedding $\mathbb{H}^1 \subset \mathbb{L}^{4n+2}$ and Remark 2.7 that $\Psi(\mathbf{n})[f(\mathbf{n})]$ is well defined for any $\mathbf{n} \in \mathbb{H}^1$. From these observation we can apply Itô's formula and derive that

$$\begin{aligned}\Psi(\mathbf{n}(t)) &= \Psi(\mathbf{n}_0) - \int_0^t \Psi'(\mathbf{n}(s))[A\mathbf{n}(s) + \tilde{B}(\mathbf{v}(s), \mathbf{n}(s)) + f(\mathbf{n}(s))] \\ &\quad + \frac{1}{2} \int_0^t \left(\Psi'(\mathbf{n}(s))[G^2(\mathbf{n}(s))] + \Psi''(\mathbf{n}(s))[G(\mathbf{n}(s)), G(\mathbf{n}(s))] \right) ds \\ &\quad + \int_0^t \Psi'(\mathbf{n}(s))[G(\mathbf{n}(s))] dW_2(s)\end{aligned}$$

The stochastic integral vanishes because $\mathbf{n} \times h \perp \mathbf{n}$ and

$$\begin{aligned}\langle G(\mathbf{n}), \mathbf{n} \rangle &= \langle (\mathbf{n} \times h), \mathbf{n} \rangle, \\ &= \langle \mathbf{n} \times h, \mathbf{n} \rangle, \\ &= 0.\end{aligned}$$

Since \mathbf{v} is a divergence free function it follows from (A.10) that

$$\langle \tilde{B}(\mathbf{v}(t), \mathbf{n}(t)), \mathbf{n}(t) \rangle = 0.$$

From the identity

$$\langle (b \times a) \times a, b \rangle = -\|a \times b\|^2,$$

we deduce that

$$\begin{aligned}\Psi'(\mathbf{n}(s))[G^2(\mathbf{n}(s))] + \Psi''(\mathbf{n}(s))[G(\mathbf{n}(s)), G(\mathbf{n}(s))] \\ = 2p\|\mathbf{n}(s)\|^{2(p-1)} \left[\langle G^2(\mathbf{n}(t)), \mathbf{n}(t) \rangle + \|G(\mathbf{n}(t))\|^2 \right] \\ = 0.\end{aligned}$$

Consequently,

$$\begin{aligned}\|\mathbf{n}(t)\|^p &= \|\mathbf{n}_0\|^p - p \int_0^t \|\mathbf{n}(s)\|^{p-2} \|\nabla \mathbf{n}(s)\|^2 ds \\ &\quad - p \int_0^t \|\mathbf{n}(s)\|^{p-2} \langle f(\mathbf{n}(s)), \mathbf{n}(s) \rangle ds,\end{aligned} \tag{C.2}$$

Now, by Assumption 2.1 that there exists a polynomial $\tilde{F}(r) = \sum_{l=1}^{N+1} b_l r^l$ with $\tilde{F}(0) = 0$ and $b_{N+1} > 0$ such that

$$\langle f(\mathbf{n}), \mathbf{n} \rangle = \int_{\mathcal{O}} \tilde{F}(|\mathbf{n}(x)|^2) dx.$$

In fact, it follows from Assumption 2.1 that

$$\begin{aligned}\langle \tilde{f}(|\mathbf{n}|^2)\mathbf{n}, \mathbf{n} \rangle &= \int_{\mathcal{O}} \tilde{f}(|\mathbf{n}(x)|^2) |\mathbf{n}(x)|^2 dx \\ &= \int_{\mathcal{O}} \sum_{k=0}^N a_k (|\mathbf{n}(x)|^2)^{k+1} dx \\ &= \int_{\mathcal{O}} \sum_{l=1}^{N+1} a_{l-1} (|\mathbf{n}(x)|^2)^l dx.\end{aligned}$$

Thanks to this observation we can use [12, Lemma 8.7] to infer that there exists $c > 0$ such that

$$\frac{a_{N+1}}{2} \int_{\mathcal{O}} |\mathbf{n}(x)|^{2N+2} dx - c \int_{\mathcal{O}} |\mathbf{n}(x)|^2 dx \leq \langle f(\mathbf{n}), \mathbf{n} \rangle.$$

From this estimate and (C.2) we deduce that

$$\|\mathbf{n}(t)\|^p + p \int_0^t \|\mathbf{n}(s)\|^{p-2} \|\nabla \mathbf{n}(s)\|^2 ds + p \int_0^t \|\mathbf{n}(s)\|^{p-2} \|\mathbf{n}(s)\|_{\mathbb{L}^{2N+2}}^{2N+2} ds \leq C \int_0^t \|\mathbf{n}(s)\|^p ds + \|\mathbf{n}_0\|^p, \quad (\text{C.3})$$

from which along with an application of the Gronwall lemma we complete the proof of our proposition. \square

C.2. Estimates for $\|\mathbf{v}\|$ and $\|\nabla \mathbf{n}\|$.

Proposition C.2. *Suppose that Assumption 3.1 is verified and that for $p = 1$ and some $p \geq 4N + 2$ and for any $t \geq 0$*

$$\mathbb{E} \mathfrak{G}_1(t, p) < \infty,$$

where $\mathfrak{G}_1(t, p)$ is defined in (3.12). Let $\{\tau_R : R \geq 0\}$ be an arbitrary family of stopping times such that the stochastic process $M_{\cdot \wedge \tau_R}$ defined by

$$M_{t \wedge \tau_R} := \int_0^{t \wedge \tau_R} \langle G(\mathbf{n}(s)), f(\mathbf{n}(s)) - \Delta \mathbf{n}(s) \rangle dW_2(t)$$

is a martingale. Then, there exists $\ell > 0$ such that for any $t \geq 0$

$$\begin{aligned}\mathbb{E} \left[\sup_{0 \leq s \leq t \wedge \tau_R} \left(\|\mathbf{v}(s)\|^2 + \ell \|\mathbf{n}(s)\|^2 + \|\nabla \mathbf{n}(s)\|^2 + \int_{\mathcal{O}} F(\mathbf{n}(s, x)) dx \right)^p \right. \\ \left. + \mathbb{E} \left(\int_0^{t \wedge \tau_R} (\|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2) ds \right)^p \right] \leq \mathbb{E} \mathfrak{G}_1(t, p).\end{aligned} \quad (\text{C.4})$$

Proof. In what follow we fix a time $t \in [0, T \wedge \tau^R]$. Application of Itô's formula to $\Phi(\mathbf{v}(t)) = \frac{1}{2} \|\mathbf{v}(t)\|^2$ yields that for any $t \in [0, T \wedge \tau^R]$ and \mathbb{P} -a.s.

$$\begin{aligned}d\|\mathbf{v}(t)\|^2 &= - \left\langle A\mathbf{v}(t) + B(\mathbf{v}(t)) + M(\mathbf{n}(t)), \mathbf{v}(t) \right\rangle dt \\ &\quad + \frac{1}{2} \|S(\mathbf{v}(t))\|_{\mathbb{H}_2}^2 dt + \langle S(\mathbf{v}(t)), \mathbf{v}(t) \rangle dW_1(t).\end{aligned} \quad (\text{C.5})$$

Now we introduce the function Ψ defined by

$$\Psi(\mathbf{n}) = \frac{1}{2} \|\nabla \mathbf{n}\|^2 + \frac{1}{2} \int_{\mathcal{O}} F(|\mathbf{n}(x)|^2) dx, \mathbf{n} \in \mathbb{H}^1.$$

Arguing as in [12] we can show that the map $\Psi(\cdot)$ is twice Fréchet differentiables and the first and second derivatives of Ψ are given by

$$\begin{aligned}\Psi'(\mathbf{n})\mathbf{g} &= \langle \nabla \mathbf{n}, \nabla \mathbf{g} \rangle + \langle f(\mathbf{n}), \mathbf{g} \rangle, \\ \Psi''(\mathbf{n})(\mathbf{g}, \mathbf{g}) &= \langle \nabla \mathbf{g}, \nabla \mathbf{g} \rangle + \int_{\mathcal{O}} \tilde{f}(\mathbf{n}) |\mathbf{g}|^2 dx + \int_{\mathcal{O}} [\tilde{f}'(\mathbf{n})][\mathbf{n} \cdot \mathbf{g}]^2 dx,\end{aligned}$$

for all $\mathbf{n}, \mathbf{g} \in \mathbb{H}^1$. If $\mathbf{g} \perp \mathbf{n}$ then

$$\Psi''(\mathbf{n})(\mathbf{g}, \mathbf{g}) = \langle \nabla \mathbf{g}, \nabla \mathbf{g} \rangle + \int_{\mathcal{O}} \tilde{f}(\mathbf{n}) |\mathbf{g}|^2 dx.$$

Note also that

$$\Psi'(\mathbf{n})\mathbf{g} = \langle -\Delta \mathbf{n}, \mathbf{g} \rangle + \langle f(\mathbf{n}), \mathbf{g} \rangle,$$

for all $\mathbf{n} \in \mathbb{H}^2$ and $\mathbf{g} \in \mathbb{H}^1$.

Before proceeding further we should recall that it was proved in [12, Lemma 8.9] that there exists $\ell > 0$ such that

$$\|\nabla \mathbf{n}\|^2 + \|\mathbf{n}\|^2 \leq 2\Psi(\mathbf{n}) + \ell \|\mathbf{n}\|^2, \quad (\text{C.6})$$

for any $\mathbf{n} \in \mathbb{H}^1$.

Now, by Itô's formula we have

$$\begin{aligned} d\Psi(\mathbf{n}(t)) &= \left(-\|\Delta \mathbf{n}(t) - f(\mathbf{n}(t))\|^2 + \frac{1}{2} \int_{\mathcal{O}} \tilde{f}(\mathbf{n}(t)) |G(\mathbf{n}(t))|^2 dx \right) dt \\ &\quad + \left\langle \left(\frac{1}{2} G^2(\mathbf{n}(t)) - \tilde{B}(\mathbf{v}(t), \mathbf{n}(t)) \right), f(\mathbf{n}(t)) - \Delta \mathbf{n}(t) \right\rangle dt \\ &\quad + \frac{1}{2} \|\nabla G(\mathbf{n}(t))\|^2 + \langle G(\mathbf{n}(t)), f(\mathbf{n}(t)) - \Delta \mathbf{n}(t) \rangle dW_2(t), \end{aligned}$$

which is equivalent to

$$\begin{aligned} d\Psi(\mathbf{n}(t)) &= \left(-\|\Delta \mathbf{n}(t) - f(\mathbf{n}(t))\|^2 + \int_{\mathcal{O}} [\tilde{f}(\mathbf{n}(t)) |G(\mathbf{n}(t))|^2 dx] dt \right. \\ &\quad \left. + \left(\left\langle \frac{1}{2} G^2(\mathbf{n}(t)), f(\mathbf{n}(t)) - \Delta \mathbf{n}(t) \right\rangle + \langle \tilde{B}(\mathbf{v}(t), \mathbf{n}(t)), \Delta \mathbf{n}(t) \rangle \right) dt \right. \\ &\quad \left. + \frac{1}{2} \|\nabla G(\mathbf{n}(t))\|^2 + \langle G(\mathbf{n}(t)), f(\mathbf{n}(t)) - \Delta \mathbf{n}(t) \rangle dW_2(t) \right). \end{aligned}$$

Note that we have used the fact that

$$\begin{aligned} \langle \mathbf{v} \cdot \nabla \mathbf{n}, f(\mathbf{n}) \rangle &= \sum_{i,j} \int_{\mathcal{O}} \mathbf{v}_i(x) \frac{\partial \mathbf{n}_j(x)}{\partial x_i} \tilde{f}(|\mathbf{n}(x)|^2) \mathbf{n}_j(x) dx \\ &= \frac{1}{2} \int_{\mathcal{O}} \mathbf{v}_i(x) \frac{\partial \tilde{F}(|\mathbf{n}(x)|^2)}{\partial x_i} dx \\ &= \langle \mathbf{v}, \nabla \tilde{F}(|\mathbf{n}|^2) \rangle \\ &= 0, \text{ because } \operatorname{div} \mathbf{v} = 0. \end{aligned}$$

Notice also that

$$\frac{1}{2} \left| \int_{\mathcal{O}} \tilde{f}(|\mathbf{n}(x)|^2) |G(\mathbf{n}(x))|^2 dx \right| \leq \frac{1}{2} \|h\|_{\mathbb{L}^\infty}^2 \int_{\mathcal{O}} |\tilde{f}(|\mathbf{n}(x)|^2)| |\mathbf{n}(x)|^2 dx.$$

By setting $\tilde{f}(r) = \sum_{k=0}^N b_k r^k$ with $b_k = |a_k|$, $k = 0, \dots, N$, we derive that there exists a polynomial \tilde{Q} of degree N such that

$$\tilde{f}(r)r = a_N r^{N+1} + \tilde{Q}(r).$$

From this last identity and the former estimate we derive that

$$\frac{1}{2} \left| \int_{\mathcal{O}} \tilde{f}(|\mathbf{n}|^2) |G(\mathbf{n})|^2 dx \right| \leq \frac{1}{2} \|h\|_{\mathbb{L}^\infty}^2 \left[a_N \int_{\mathcal{O}} |\mathbf{n}(x)|^{2N+2} dx + \left| \int_{\mathcal{O}} \tilde{Q}(|\mathbf{n}(x)|^2) dx \right| \right],$$

from which along with [12, Lemma 8.7] we deduce that

$$\frac{1}{2} \left| \int_{\mathcal{O}} \tilde{f}(|\mathbf{n}|^2) |G(\mathbf{n})|^2 dx \right| \leq C(h) \left(\int_{\mathcal{O}} F(|\mathbf{n}(x)|^2) dx + \|\mathbf{n}\|^2 \right).$$

Thus,

$$\frac{1}{2} \left| \int_{\mathcal{O}} \tilde{f}(|\mathbf{n}|^2) |G(\mathbf{n})|^2 dx \right| \leq C(h) (\Psi(\mathbf{n}) + \|\mathbf{n}\|^2),$$

and

$$\begin{aligned} d\Psi(\mathbf{n}(t)) &\leq \left(-\|\Delta\mathbf{n}(t) - f(\mathbf{n}(t))\|^2 + C(h) [\Psi(\mathbf{n}(t)) + \|\mathbf{n}(t)\|^2] \right) dt \\ &+ \left(\left\langle \frac{1}{2} G^2(\mathbf{n}(t)), f(\mathbf{n}(t)) - \Delta\mathbf{n}(t) \right\rangle + \langle \tilde{B}(\mathbf{v}(t), \mathbf{n}(t)), \Delta\mathbf{n}(t) \rangle \right) dt \\ &+ \frac{1}{2} \|\nabla G(\mathbf{n}(t))\|^2 + \langle G(\mathbf{n}(t)), f(\mathbf{n}(t)) - \Delta\mathbf{n}(t) \rangle dW_2(t). \end{aligned} \quad (\text{C.7})$$

Thanks to (2.11) we derive that

$$\langle \tilde{B}(\mathbf{v}, \mathbf{n}), \Delta\mathbf{n} \rangle - \langle M(\mathbf{n}), \mathbf{v} \rangle = 0.$$

From Lemma A.1 we also derive that

$$\langle B(\mathbf{v}(t)), \mathbf{v}(t) \rangle = 0.$$

Therefore by summing (C.5) and (C.7) side by side and using the two last identities we see that

$$\begin{aligned} d[\Psi(\mathbf{n}(t)) + \frac{1}{2} \|\mathbf{v}(t)\|^2] &+ \left(\|\nabla \mathbf{v}(t)\|^2 + \frac{1}{2} \|\Delta\mathbf{n}(t) - f(\mathbf{n}(t))\|^2 - C(h) [\Psi(\mathbf{n}(t)) + \|\mathbf{n}(t)\|^2] \right) dt \\ &\leq \frac{1}{2} \left(\|S(\mathbf{v}(t))\|_{\mathbb{H}^2}^2 + \|G^2(\mathbf{n}(t))\|^2 + \|\nabla G(\mathbf{n}(t))\|^2 \right) dt \\ &+ \langle \mathbf{v}(t), S(\mathbf{v}(t)) dW_1(t) \rangle + \langle G(\mathbf{n}(t)), f(\mathbf{n}(t)) - \Delta\mathbf{n}(t) \rangle dW_2(t). \end{aligned} \quad (\text{C.8})$$

Now, recall that $G(\mathbf{n}) = \mathbf{n} \times h$. Hence

$$\begin{aligned} \|\nabla G(\mathbf{n}(t))\|^2 &\leq \|\nabla(\mathbf{n}(t) \times h)\|^2 = \|G(\mathbf{n}(t))\|_{\mathbb{H}^1}^2, \\ &\leq \|\nabla(\mathbf{n}(t) \times h)\|^2 = \|\mathbf{n}(t) \times h\|_{\mathbb{H}^1}^2 \\ &\leq \|\nabla(\mathbf{n}(t) \times h)\|^2 + \|\mathbf{n}(t) \times h\|^2 \\ &\leq 2[\|\nabla \mathbf{n}(t) \times h\|^2 + \|\mathbf{n}(t) \times \nabla h\|^2] + \|\mathbf{n}(t) \times h\|^2 \\ &\leq C\|h\|_{\mathbb{L}^\infty}^2 (\|\nabla \mathbf{n}(t)\|^2 + \|\mathbf{n}(t)\|^2) + \|\mathbf{n}(t) \times \nabla h\|^2. \end{aligned} \quad (\text{C.9})$$

From Hölder's inequality and the Sobolev embedding $\mathbb{H}^1 \subset \mathbb{L}^6$ (true for $n = 2, 3$!) we obtain

$$\begin{aligned} \|\mathbf{n}(t) \times \nabla h\|^2 &\leq \|\mathbf{n}(t)\|_{\mathbb{L}^6}^2 \|\nabla h\|_{\mathbb{L}^3}^2, \\ &\leq c(\|\nabla \mathbf{n}(t)\|^2 + \|\mathbf{n}(t)\|^2) \|\nabla h\|_{\mathbb{L}^3}^2. \end{aligned}$$

By plugging this last inequality into (C.9) we infer the existence of a constant $C(h) > 0$ such that

$$\|\nabla G(\mathbf{n}(t))\|^2 \leq C(h) (\|\nabla \mathbf{n}(t)\|^2 + \|\mathbf{n}(t)\|^2).$$

From the definition of G^2 it is easy to see that there exists $C(h) > 0$ such that

$$\|G^2(\mathbf{n}(t))\|^2 \leq C(h) \|\mathbf{n}(t)\|^2 \leq C(h) (\|\nabla \mathbf{n}(t)\|^2 + \|\mathbf{n}(t)\|^2).$$

From the last two estimates, (C.6) and the linear growth assumption (3.1) we derive that there exist two constants $C > 0$ and $C(h) > 0$ such that for any $t \geq 0$

$$\frac{1}{2} \left(\|S(\mathbf{v}(t))\|_{\mathbb{H}^2}^2 + \|G^2(\mathbf{n}(t))\|^2 + \|\nabla G(\mathbf{n}(t))\|^2 \right) \leq C\|\mathbf{v}(t)\|^2 + 2C(h)\Psi(\mathbf{n}(t)) + \ell C(h)\|\mathbf{n}(t)\|^2. \quad (\text{C.10})$$

From this inequality and (C.8) we derive that there exists $C > 0$ such that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} [\Psi(\mathbf{n}(s)) + \frac{1}{2} \|\mathbf{v}(s)\|^2] + \mathbb{E} \int_0^{t \wedge \tau_R} \left(\|\nabla \mathbf{v}(s)\|^2 + \frac{1}{2} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 \right) ds \\ & \leq C \mathbb{E} \int_0^{t \wedge \tau_R} [\|\mathbf{v}(s)\|^2 + \Psi(\mathbf{n}(s))] ds + \mathbb{E} \int_0^{t \wedge \tau_R} \|\mathbf{n}(s)\|^2 ds + \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle \mathbf{v}(r), S(\mathbf{v}(r)) dW_1(r) \rangle \right| \\ & \quad + \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle G(\mathbf{n}(r)), f(\mathbf{n}(r)) - \Delta \mathbf{n}(r) \rangle dW_2(r) \right| + \mathbb{E} \left(\frac{1}{2} \|\mathbf{v}_0\|^2 + \Psi(\mathbf{n}_0) \right). \end{aligned} \quad (\text{C.11})$$

Thanks to the Burkholder-Davis-Gundy, Cauchy-Schwarz and Cauchy inequalities we infer that

$$\begin{aligned} \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle G(\mathbf{n}(r)), f(\mathbf{n}(r)) - \Delta \mathbf{n}(r) \rangle dW_2(r) \right| & \leq C \mathbb{E} \left(\int_0^{t \wedge \tau_R} [\langle G(\mathbf{n}(s)), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle]^2 ds \right)^{\frac{1}{2}} \\ & \leq C \mathbb{E} \left[\sup_{s \in [0, t \wedge \tau_R]} \|G(\mathbf{n}(s))\| \left(\int_0^{t \wedge \tau_R} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \right)^{\frac{1}{2}} \right] \\ & \leq C \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \|G(\mathbf{n}(s))\|^2 + \frac{1}{4} \mathbb{E} \int_0^{t \wedge \tau_R} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \\ & \leq C \|h\|_{\mathbb{L}^\infty}^2 \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \|\mathbf{n}(s)\|^2 + \frac{1}{4} \mathbb{E} \int_0^{t \wedge \tau_R} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds. \end{aligned} \quad (\text{C.12})$$

By making use of a similar argument and the linear growth assumption (3.1) one can prove that

$$\mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle \mathbf{v}(r), S(\mathbf{v}(r)) dW_1(r) \rangle \right| \leq \frac{1}{4} \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \|\mathbf{v}(s)\|^2 + C \mathbb{E} \int_0^{t \wedge \tau_R} \|\mathbf{v}(s)\|^2 ds. \quad (\text{C.13})$$

Note that from (C.6) we easily derive that $\|\mathbf{v}\|^2 + \|\mathbf{n}\|^2 \leq \|\mathbf{v}\|^2 + 2\Psi(\mathbf{n}) + \ell\|\mathbf{n}\|^2$. Hence, using (C.12) and (C.13) in (C.11) we infer that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} [\Psi(\mathbf{n}(s)) + \frac{1}{2} \|\mathbf{v}(s)\|^2] + \mathbb{E} \int_0^{t \wedge \tau_R} \left(\|\nabla \mathbf{v}(s)\|^2 + \frac{1}{2} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 \right) ds \\ & \leq C \mathbb{E} \int_0^{t \wedge \tau_R} (\|\mathbf{v}(s)\|^2 + \Psi(\mathbf{n}(s))) ds + C \varphi(t \wedge \tau_R), \end{aligned}$$

where $\varphi(\cdot)$ is the non-decreasing function defined by

$$\varphi(t) = \mathbb{E} (\|\mathbf{v}_0\|^2 + \Psi(\mathbf{n}_0)) + \mathbb{E} \sup_{s \in [0, t]} \|\mathbf{n}(s)\|^2 + \mathbb{E} \int_0^t \|\mathbf{n}(s)\|^2 ds.$$

Now it follows from Gronwall's lemma that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} [\Psi(\mathbf{n}(s)) + \frac{1}{2} \|\mathbf{v}(s)\|^2] + \mathbb{E} \int_0^{t \wedge \tau_R} \left(\|\nabla \mathbf{v}(s)\|^2 + \frac{1}{2} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 \right) ds \\ & \leq \varphi(t \wedge \tau_R) \left(1 + t \wedge \tau_R e^{Ct \wedge \tau_R} \right). \end{aligned}$$

This completes the proof of our proposition for the case $p = 1$. For the case $p \geq 4N + 2$ we first observe that from (C.8), (C.10) we easily see that

$$\begin{aligned} & \Psi(\mathbf{n}(t)) + \ell \|\mathbf{n}(t)\|^2 + \|\mathbf{v}(t)\|^2 + \int_0^t \left(2\|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 \right) ds \\ & \leq \Psi(\mathbf{n}_0) + \ell \|\mathbf{n}_0\|^2 + \|\mathbf{v}_0\|^2 + C \int_0^t (2\Psi(\mathbf{n}(s)) + \ell \|\mathbf{n}(s)\|^2 + \|\mathbf{v}(s)\|^2) ds \\ & \quad \left| \int_0^t \langle \mathbf{v}(s), S(\mathbf{v}(s)) dW_1(s) \rangle \right| + \left| \int_0^t \langle G(\mathbf{n}(s)), f(\mathbf{n}(s)) - \Delta \mathbf{n}(s) \rangle dW_2(s) \right|. \end{aligned}$$

Rising both sides of this estimate to the power p and taking the supremum over $s \in [0, t \wedge \tau_R]$ and the mathematical expectation imply that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t]} [\psi(s)]^p - \mathbb{E}[\psi(0)]^p + \mathbb{E} \left[\int_0^{t \wedge \tau_R} \left(2\|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 \right) ds \right]^p \\ & \leq Ct \wedge \tau_R \mathbb{E} \int_0^{t \wedge \tau_R} [\psi(s)]^p ds + \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle \mathbf{v}(r), S(\mathbf{v}(r)) dW_1 \rangle \right|^p \\ & \quad + \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle G(\mathbf{n}(r)), f(\mathbf{n}(r)) - \Delta \mathbf{n}(r) \rangle dW_2 \right|^p, \end{aligned} \quad (\text{C.14})$$

where, for the sake of simplicity, we have put $\psi(t) = \Psi(\mathbf{n}(t)) + \ell \|\mathbf{n}(t)\|^2 + \|\mathbf{v}(t)\|^2$.

Now by making use of the Burkholder-Davis-Gundy, Cauchy-Schwarz, Cauchy inequalities and the linear growth assumption (3.1) we derive that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle \mathbf{v}(r), S(\mathbf{v}(r)) dW_1 \rangle \right|^p \leq C \mathbb{E} \left(\int_0^{t \wedge \tau_R} \|\mathbf{v}(s)\|^2 \|S(\mathbf{v}(s))\|_{\mathcal{H}_2}^2 ds \right)^{\frac{p}{2}} \\ & \leq C \mathbb{E} \left[\sup_{s \in [0, t \wedge \tau_R]} [\psi(s)]^{\frac{p}{2}} \left(\int_0^{t \wedge \tau_R} (1 + \|\mathbf{v}(s)\|^2) ds \right)^{\frac{p}{2}} \right] \\ & \leq \frac{1}{4} \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} [\psi(s)]^p + Ct \wedge \tau_R + C \mathbb{E} \int_0^{t \wedge \tau_R} [\psi(s)]^p ds. \end{aligned} \quad (\text{C.15})$$

By using a similar argument we obtain

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle G(\mathbf{n}(r)), f(\mathbf{n}(r)) - \Delta \mathbf{n}(r) \rangle dW_2 \right|^p \leq C \mathbb{E} \left(\int_0^{t \wedge \tau_R} \|G(\mathbf{n}(s))\|^2 \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \right)^{\frac{p}{2}} \\ & \leq C \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \|\mathbf{n}(s) \times h\|^p \\ & \quad + \frac{1}{2} \mathbb{E} \left(\int_0^{t \wedge \tau_R} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \right)^p. \end{aligned}$$

From the last line we easily derive that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \left| \int_0^{s \wedge \tau_R} \langle G(\mathbf{n}(r)), f(\mathbf{n}(r)) - \Delta \mathbf{n}(r) \rangle dW_2 \right|^p - C(h) \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \|\mathbf{n}(s)\|^{2p} \\ & \leq \frac{1}{2} \mathbb{E} \left(\int_0^{t \wedge \tau_R} (2\|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2) ds \right)^p. \end{aligned} \quad (\text{C.16})$$

Inserting (C.15) and (C.16) in (C.14) yields that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} [\psi(s)]^p + \mathbb{E} \left[\int_0^{t \wedge \tau_R} 2 \|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \right] \\ & \leq 2\mathbb{E}[\psi(0)]^p + Ct \wedge \tau_R + C(h) \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \|\mathbf{n}(s)\|^{2p} + C(t \wedge \tau_R + 1) \mathbb{E} \int_0^{t \wedge \tau_R} [\psi(s)]^p ds, \end{aligned}$$

from which altogether with the Gronwall lemma implies that

$$\begin{aligned} & \mathbb{E} \sup_{s \in [0, t]} [\psi(s)]^p + \mathbb{E} \left[\int_0^{t \wedge \tau_R} 2 \|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \right] \\ & \leq \left[2\mathbb{E}[\psi(0)]^p + Ct \wedge \tau_R + C(h) \mathbb{E} \sup_{s \in [0, t \wedge \tau_R]} \|\mathbf{n}(s)\|^{2p} \right] \left(1 + Ct \wedge \tau_R (t \wedge \tau_R + 1) e^{C(t \wedge \tau_R + 1)t \wedge \tau_R} \right). \end{aligned}$$

This completes the proof of the proposition. \square

C.3. Estimate for $\mathbb{E} \int_0^T \|\Delta \mathbf{n}(s)\|^2 ds$.

Proposition C.3. *Suppose that Assumption 2.1 is satisfied and let $p \geq 4N + 2$. If $\mathbb{E} \mathfrak{G}_1(T, q(2N + 1))$ for some $q \in [2, \frac{p}{4N+2}]$ and $T \geq 0$, then there exists a constant $C > 0$ such that*

$$\mathbb{E} \left[\int_0^T \|\Delta \mathbf{n}(s)\|^2 ds \right]^q \leq \mathbb{E} \mathfrak{G}(T, q(2N + 1)) + CT.$$

Proof. The proof follows from (2.17) of Remark 2.7 and Proposition C.2. In fact for $N \in I_d$ the embedding $\mathbb{H}^1 \subset \mathbb{L}^{4N+2}$ is continuous and thanks to (2.17) we derive that

$$\mathbb{E} \left[\int_0^T \|\Delta \mathbf{n}(s)\|^2 ds \right]^q \leq C \mathbb{E} \left[\int_0^T \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \right]^q + C \mathbb{E} \sup_{s \in [0, T]} \|\nabla \mathbf{n}(s)\|^{2q(2N+1)} + C.$$

Now, we easily conclude the proof of our claim by using Proposition C.2 with $p = q(2N + 1)$. \square

C.4. Proof of Proposition 4.14. In this subsection we give the proof of Proposition 4.14, i.e., we will derive the important estimate for $\mathbb{E} \sup_{s \in [0, t \wedge \tau_k]} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2$

Proof of Proposition 4.14. Since the local strong solution $(\mathbf{v}, \mathbf{n}) \in D(A) \times \mathbb{X}_1$ almost surely we can view $\Psi(\mathbf{n})$ as

$$\Psi(\mathbf{n}) = \|-A_1 \mathbf{n} - f(\mathbf{n})\|^2.$$

The functional $\Psi(\mathbf{n})$ is twice differentiable with first and second Fréchet derivatives defined by

$$\Psi'(\mathbf{n})(h) = 2\langle -A_1 \mathbf{n} - f(\mathbf{n}), -A_1 h - f'(\mathbf{n})h \rangle,$$

and

$$\Psi''(\mathbf{n})[h, k] = 2\langle -A_1 k - f'(\mathbf{n})k, -A_1 h - f'(\mathbf{n})h \rangle - 2\langle -A_1 \mathbf{n} - f(\mathbf{n}), f''(\mathbf{n})h k \rangle,$$

for any $h, k \in \mathbb{X}_{\frac{1}{2}}$.

Let us recall that the process $(\mathbf{v}(t), \mathbf{n}(t))$, $t < \tilde{\tau}_\infty$ solves

$$d\mathbf{v}(t) + \left(\Delta \mathbf{v}(t) + B(\mathbf{v}(t), \mathbf{v}(t)) + M(\mathbf{n}(t)) \right) dt = S(\mathbf{v}(t)) dW_1(t),$$

$$d\mathbf{n}(t) + \left(A_1 \mathbf{n}(t) + \tilde{B}(\mathbf{v}(t), \mathbf{n}(t)) + f(\mathbf{n}(t)) - \frac{1}{2} G^2(\mathbf{n}(t)) \right) dt = G(\mathbf{n}(t)) dW_2(t).$$

Since $(\mathbf{v}, \mathbf{n}) \in X_t$ for any $t \in [0, \tau_\infty)$ we infer that

$$\Delta \mathbf{v}(\cdot) + B(\mathbf{v}(\cdot), \mathbf{v}(\cdot)) + M(\mathbf{n}(\cdot)) \in L^2(0, t; \mathbb{H}),$$

and

$$A_1 \mathbf{n}(\cdot) + \tilde{B}(\mathbf{v}(\cdot), \mathbf{n}(\cdot)) + f(\mathbf{n}(\cdot)) - \frac{1}{2} G^2(\mathbf{n}(\cdot)) \in L^2(0, t; \mathbb{X}_0),$$

for any $t \in [0, \tilde{\tau}_\infty)$. Therefore by considering the Gelfand triples $D(A) \subset D(A^{\frac{1}{2}}) \subset \mathbb{H}$ and $\mathbb{X}_1 \subset \mathbb{X}_{\frac{1}{2}} \subset \mathbb{X}_0$ Itô's formula for the functional $\|\nabla \mathbf{v}(t \wedge \tau_k)\|^2$ and $\Psi(\mathbf{n}(t \wedge \tau_k))$ for any integer $k > 0$ are applicable to our situation (see [46, Theorem I.3.3.2, Page 147]). By Itô's formula we have

$$\begin{aligned} \|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 - \|\nabla \mathbf{v}_0\|^2 &= -2 \int_0^{t \wedge \tau_k} \langle \Delta \mathbf{v}(s) + B(\mathbf{v}(s), \mathbf{v}(s)) + \Pi \nabla \cdot (\nabla \mathbf{n}(s) \odot \nabla \mathbf{n}(s)), \Delta \mathbf{v}(s) \rangle ds \\ &\quad + \int_0^{t \wedge \tau_k} \|S(\mathbf{v}(s))\|_{\mathcal{H}_2(K_1, \mathbb{V})}^2 ds + 2 \int_0^{t \wedge \tau_k} \langle A^{\frac{1}{2}} S(\mathbf{v}(s)), A^{\frac{1}{2}} \mathbf{v}(s) \rangle dW_1(s), \end{aligned}$$

where $t \wedge \tau_k := t \wedge \tau_k$. As in Lin and Liu [37] we use the identity

$$\begin{aligned} \langle \Pi \nabla \cdot (\nabla \mathbf{n} \odot \nabla \mathbf{n}), \Delta \mathbf{v} \rangle &= \langle \nabla \mathbf{n} \Delta \mathbf{n}, \Delta \mathbf{v} \rangle + \langle \nabla \left(\frac{|\nabla \mathbf{n}|^2}{2} \right), \Delta \mathbf{v} \rangle, \\ &= \langle \nabla \mathbf{n} \Delta \mathbf{n}, \Delta \mathbf{v} \rangle. \end{aligned}$$

To get the second line of this identity we have used the fact that $\Delta \mathbf{v}$ is divergence free. Now it is easy to see that

$$\langle \Pi \nabla \cdot (\nabla \mathbf{n} \odot \nabla \mathbf{n}), \Delta \mathbf{v} \rangle = \langle \nabla \mathbf{n} (\Delta \mathbf{n} - f(\mathbf{n})), \Delta \mathbf{v} \rangle + \langle \nabla \mathbf{n}, f(\mathbf{n}) \Delta \mathbf{v} \rangle.$$

As a consequence of the last identity we have that

$$\begin{aligned} \|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 - \|\nabla \mathbf{v}_0\|^2 &= -2 \int_0^{t \wedge \tau_k} \langle \Delta \mathbf{v}(s) + B(\mathbf{v}(s), \mathbf{v}(s)) + \nabla \mathbf{n}(s) f(\mathbf{n}(s)), \Delta \mathbf{v}(s) \rangle ds \\ &\quad - 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s) (\Delta \mathbf{n}(s) - f(\mathbf{n}(s))), \Delta \mathbf{v}(s) \rangle ds + \int_0^{t \wedge \tau_k} \|S(\mathbf{v}(s))\|_{\mathcal{H}_2(K_1, \mathbb{V})}^2 ds \quad (\text{C.17}) \\ &\quad + 2 \int_0^{t \wedge \tau_k} \langle \nabla S(\mathbf{v}(s)), \nabla \mathbf{v}(s) \rangle dW_1(s). \end{aligned}$$

At the same time we have

$$\begin{aligned} \Psi(\mathbf{n}(t \wedge \tau_k)) - \Psi(\mathbf{n}_0) &= \int_0^{t \wedge \tau_k} \Psi'(\mathbf{n}(s)) [-\mathbf{v}(s) \cdot \nabla \mathbf{n}(s) - A_1 \mathbf{n}(s) - f(\mathbf{n}(s)) + \frac{1}{2} [G^2(\mathbf{n}(s))]] ds \\ &\quad + \frac{1}{2} \int_0^{t \wedge \tau_k} \Psi''(\mathbf{n}(s)) [G(\mathbf{n}(s)) G(\mathbf{n}(s))] ds + \int_0^{t \wedge \tau_k} \Psi'(\mathbf{n}(s)) [G(\mathbf{n}(s))] dW_2(s). \end{aligned}$$

Using the definition of $\Psi'(\mathbf{n}(s))$ we see from this last equation that

$$\begin{aligned} \Psi(\mathbf{n}(t \wedge \tau_k)) - \Psi(\mathbf{n}_0) &= 2 \int_0^{t \wedge \tau_k} \langle -A_1 \mathbf{n}(s) - f(\mathbf{n}(s)), -A_1 \left(-\mathbf{v}(s) \cdot \nabla \mathbf{n}(s) - A_1 \mathbf{n}(s) - f(\mathbf{n}(s)) \right) \rangle ds \\ &\quad - 2 \int_0^{t \wedge \tau_k} \langle -A_1 \mathbf{n}(s) - f(\mathbf{n}(s)), f'(\mathbf{n}(s)) \left(-\mathbf{v}(s) \cdot \nabla \mathbf{n}(s) - A_1 \mathbf{n}(s) - f(\mathbf{n}(s)) \right) \rangle ds \\ &\quad + \frac{1}{2} \int_0^{t \wedge \tau_k} \Psi'(\mathbf{n}(s)) [G^2(\mathbf{n}(s))] ds + \frac{1}{2} \int_0^{t \wedge \tau_k} \Psi''(\mathbf{n}(s)) [G(\mathbf{n}(s)), G(\mathbf{n}(s))] ds \\ &\quad + \int_0^{t \wedge \tau_k} \Psi'(\mathbf{n}(s)) [G(\mathbf{n}(s))] dW_2(s) \\ &= 2 \sum_{i=1}^4 \int_0^{t \wedge \tau_k} I_i(s) ds + \int_0^{t \wedge \tau_k} \Psi'(\mathbf{n}(s)) [G(\mathbf{n}(s))] dW_2(s). \end{aligned} \quad (\text{C.18})$$

For $(\mathbf{v}, \mathbf{n}) \in D(A) \times \mathbb{X}_1$ almost surely we easily check that $-\mathbf{v} \cdot \nabla \mathbf{n} \in D(A_1)$. Therefore I_1 can be rewritten in the following form

$$I_1(s) = \langle -\nabla \mathbf{n} \Delta \mathbf{v} - \mathbf{v} \cdot \nabla \Delta \mathbf{n}, \Delta \mathbf{n} - f(\mathbf{n}) \rangle + \langle \Delta \mathbf{n} - f(\mathbf{n}), -A_1 (\Delta \mathbf{n} - f(\mathbf{n})) \rangle,$$

where the first term is a product of two function in \mathbb{L}^2 and the second term is understood as the duality pairing between an element of \mathbb{H}^1 and $(\mathbb{H}^1)^*$. Invoking (2.1) we derive that

$$I_1(s) = -\langle \mathbf{v} \cdot \nabla[f(\mathbf{n})], \Delta \mathbf{n} - f(\mathbf{n}) \rangle - \|\nabla(\Delta \mathbf{n} - f(\mathbf{n}))\|^2 + \langle -\nabla \mathbf{n} \Delta \mathbf{v} - \mathbf{v} \cdot \nabla(\Delta \mathbf{n} - f(\mathbf{n})), \Delta \mathbf{n} - f(\mathbf{n}) \rangle. \quad (\text{C.19})$$

We derive from (C.17), (C.18) and (C.19) that

$$\begin{aligned} & \|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k)) - \|\nabla \mathbf{v}_0\|^2 - \Psi(\mathbf{n}_0) + 2 \int_0^{t \wedge \tau_k} [\|\Delta \mathbf{v}(s)\|^2 + \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2] ds \\ &= 2 \int_0^{t \wedge \tau_k} \langle B(\mathbf{v}(s), \mathbf{v}(s)) + \nabla \mathbf{n}(s) f(\mathbf{n}(s)), A \mathbf{v}(s) \rangle ds + 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s) [\Delta \mathbf{n}(s) - f(\mathbf{n}(s))], A \mathbf{v}(s) \rangle ds \\ & \quad - 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s) \Delta \mathbf{v}(s), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds - 2 \int_0^{t \wedge \tau_k} \langle \mathbf{v}(s) \cdot \nabla[f(\mathbf{n}(s))], \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad - 2 \int_0^{t \wedge \tau_k} \langle f'(\mathbf{n}(s)) \left(-\mathbf{v}(s) \cdot \nabla \mathbf{n}(s) + \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \right), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad - 2 \int_0^{t \wedge \tau_k} \langle \mathbf{v}(s) \cdot \nabla[\Delta \mathbf{n}(s) - f(\mathbf{n}(s))], \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds + OT. \end{aligned} \quad (\text{C.20})$$

where the other term OT is defined by

$$\begin{aligned} OT &= \int_0^{t \wedge \tau_k} \|S(\mathbf{v}(s))\|_{\mathcal{H}_2(\mathbf{K}_1, \mathbb{V})}^2 ds + \frac{1}{2} \int_0^{t \wedge \tau_k} \Psi''(\mathbf{n}(s)) [G(\mathbf{n}(s)), G(\mathbf{n}(s))] ds \\ & \quad + 2 \int_0^{t \wedge \tau_k} \langle \nabla S(\mathbf{v}(s)), \nabla \mathbf{v}(s) \rangle dW_1(s) + \int_0^{t \wedge \tau_k} \Psi'(\mathbf{n}(s)) [G(\mathbf{n}(s))] dW_2(s) \\ & \quad + \frac{1}{2} \int_0^{t \wedge \tau_k} \Psi'(\mathbf{n}(s)) [G^2(\mathbf{n}(s))] ds. \end{aligned}$$

Thanks to (A.7) the identity (C.20) can be simplified as follows

$$\begin{aligned} & \|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k)) - \|\nabla \mathbf{v}_0\|^2 - \Psi(\mathbf{n}_0) + 2 \int_0^{t \wedge \tau_k} [\|\Delta \mathbf{v}(s)\|^2 + \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2] ds \\ &= 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s) [\Delta \mathbf{n}(s) - f(\mathbf{n}(s))], A \mathbf{v}(s) \rangle ds - 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s) \Delta \mathbf{v}(s), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad - 2 \int_0^{t \wedge \tau_k} \langle f'(\mathbf{n}(s)) \left(-\mathbf{v}(s) \cdot \nabla \mathbf{n}(s) + \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \right), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad + 2 \int_0^{t \wedge \tau_k} \langle B(\mathbf{v}(s), \mathbf{v}(s)) + \nabla \mathbf{n}(s) f(\mathbf{n}(s)), A \mathbf{v}(s) \rangle ds \\ & \quad - 2 \int_0^{t \wedge \tau_k} \langle \mathbf{v}(s) \cdot \nabla[f(\mathbf{n}(s))], \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds + OT. \end{aligned} \quad (\text{C.21})$$

Now note that for $j = 1, 2$ we have

$$\begin{aligned} [\mathbf{v} \cdot \nabla[f(\mathbf{n})]]_j &= \mathbf{v}_i \cdot \frac{\partial[f(\mathbf{n})]_j}{\partial x_i} \\ &= \tilde{f}(|\mathbf{n}|^2) \mathbf{v}_i \frac{\partial \mathbf{n}_j}{\partial x_i} + 2 \tilde{f}'(|\mathbf{n}|^2) \mathbf{n}_j \mathbf{v}_i \frac{\partial \mathbf{n}_k}{\partial x_i} \mathbf{n}_k \\ &= [\tilde{f}(|\mathbf{n}|^2) \mathbf{v} \cdot \nabla \mathbf{n}]_j + 2 \tilde{f}'(|\mathbf{n}|^2) ([\mathbf{v} \cdot \nabla \mathbf{n}] \cdot \mathbf{n}) \mathbf{n}_j, \end{aligned}$$

where above the summation over repeated indexes are enforced. On the other hand we have

$$f'(\mathbf{n})[\mathbf{v} \cdot \nabla \mathbf{n}] = \tilde{f}(|\mathbf{n}|^2) \mathbf{v} \cdot \nabla \mathbf{n} + 2 \tilde{f}'(|\mathbf{n}|^2) \mathbf{n} ([\mathbf{v} \cdot \nabla \mathbf{n}] \cdot \mathbf{n}).$$

Thus we just proved that

$$\mathbf{v} \cdot \nabla[f(\mathbf{n})] = f'(\mathbf{n})[\mathbf{v} \cdot \nabla \mathbf{n}].$$

From this last identity and (C.21) we deduce that

$$\begin{aligned} & \|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k)) - \|\nabla \mathbf{v}_0\|^2 - \Psi(\mathbf{n}_0) + 2 \int_0^{t \wedge \tau_k} \left[\|\Delta \mathbf{v}(s)\|^2 + \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2 \right] ds \\ &= 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s)[\Delta \mathbf{n}(s) - f(\mathbf{n}(s))], \mathbf{A} \mathbf{v}(s) \rangle ds - 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s) \Delta \mathbf{v}(s), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad - 2 \int_0^{t \wedge \tau_k} \langle f'(\mathbf{n}(s))(\Delta \mathbf{n}(s) - f(\mathbf{n}(s))), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad + 2 \int_0^{t \wedge \tau_k} \langle B(\mathbf{v}(s), \mathbf{v}(s)) + \nabla \mathbf{n}(s) f(\mathbf{n}(s)), \mathbf{A} \mathbf{v}(s) \rangle ds + OT. \end{aligned}$$

But since $\nabla \mathbf{n} f(\mathbf{n}) = \nabla F(\mathbf{n})$ and $\mathbf{A} \mathbf{v}$ is a divergence free function the last identity becomes

$$\begin{aligned} & \|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k)) - \|\nabla \mathbf{v}_0\|^2 - \Psi(\mathbf{n}_0) + 2 \int_0^{t \wedge \tau_k} \left[\|\Delta \mathbf{v}(s)\|^2 + \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2 \right] ds \\ &= 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s)[\Delta \mathbf{n}(s) - f(\mathbf{n}(s))], \mathbf{A} \mathbf{v}(s) \rangle ds - 2 \int_0^{t \wedge \tau_k} \langle \nabla \mathbf{n}(s) \Delta \mathbf{v}(s), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad - 2 \int_0^{t \wedge \tau_k} \langle f'(\mathbf{n}(s))(\Delta \mathbf{n}(s) - f(\mathbf{n}(s))), \Delta \mathbf{n}(s) - f(\mathbf{n}(s)) \rangle ds \\ & \quad + 2 \int_0^{t \wedge \tau_k} \langle B(\mathbf{v}(s), \mathbf{v}(s)), \mathbf{A} \mathbf{v}(s) \rangle ds + OT. \end{aligned} \tag{C.22}$$

To get rid of some bad terms in the right-hand side of the above identity we will apply the Itô's formula to the function

$$\begin{aligned} \Xi(t \wedge \tau_k, \mathbf{v}, \mathbf{n}) &:= \Phi(t \wedge \tau_k) (\|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k))) \\ &= e^{-\int_0^{t \wedge \tau_k} \phi(s) ds} (\|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k))). \end{aligned} \tag{C.23}$$

Before implementing this idea, we shall make few remarks. Since, by assumption, $h \in \mathbb{H}^2$ it is not difficult to check that

$$G^2(\mathbf{n}) = (\mathbf{n} \times h) \times h \in D(A_2),$$

and

$$G(\mathbf{n}) = \mathbf{n} \times h \in D(A_2).$$

Next, observe that

$$\begin{aligned} \frac{1}{2} \Psi'(\mathbf{n})[G^2(\mathbf{n})] &= \langle \Delta \mathbf{n} - f(\mathbf{n}), \Delta \mathbf{n} \times h \rangle + \langle \Delta \mathbf{n} - f(\mathbf{n}), \mathbf{n} \times \Delta h + (\mathbf{n} \times h) \times \Delta h \rangle \\ & \quad - 2 \langle \Delta \mathbf{n} - f(\mathbf{n}), f'(\mathbf{n})(\mathbf{n} \times h) \times h \rangle, \\ \frac{1}{2} \Psi''(\mathbf{n})[G(\mathbf{n}), G(\mathbf{n})] &= \|\Delta G(\mathbf{n}) - f'(\mathbf{n})G(\mathbf{n})\|^2 - \langle \Delta \mathbf{n} - f(\mathbf{n}), f''(\mathbf{n})[G(\mathbf{n}), G(\mathbf{n})] \rangle. \end{aligned}$$

Thus, an application of the Itô formula to the stochastic processes $\Phi(t) \left(\|\nabla \mathbf{v}(t)\|^2 + \Psi(\mathbf{n}(t)) \right)$ and the estimates (4.19), (4.20), (4.21), (4.22) and (4.23) yields

$$\begin{aligned} & \mathbb{E} \Phi(t \wedge \tau_k) \left(\|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k)) \right) \\ & + \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \left(2\|\Delta \mathbf{v}(s)\|^2 + (1 - \delta_7) \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2 \right) ds \\ & \leq E_1 + E_2 + \|\nabla \mathbf{v}_0\|^2 + \Psi(\mathbf{n}_0) + \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \|S(\mathbf{v}(s))\|_{\mathcal{H}_2(\mathbb{K}_1, \mathbb{V})}^2 ds \\ & + C(h) \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \\ & + C \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) (C + C(h) \|\mathbf{n}(s)\|_{\mathbb{H}^1}^{4N+2}) ds, \end{aligned} \quad (\text{C.24})$$

where the terms E_1 and E_2 are defined in (C.25) and (C.27) below. With the help of Burkholder-Davis-Gundy's inequality we have

$$E_1 := 2 \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_k} \left| \int_0^s \Phi(r) \Psi'(\mathbf{n}(r)) [G(\mathbf{n}(r))] dW_2(r) \right| \leq C \mathbb{E} \left(\int_0^{t \wedge \tau_k} [\Phi(s)]^2 |\Psi'(\mathbf{n}(s)) [G(\mathbf{n}(s))]|^2 ds \right)^{\frac{1}{2}}. \quad (\text{C.25})$$

Now invoking the definition of $\Psi'(\mathbf{n})[G(\mathbf{n})]$, the Cauchy-Schwarz and Young inequalities we have

$$\begin{aligned} & 2 \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_k} \left| \int_0^s \Phi(r) \Psi'(\mathbf{n}(r)) [G(\mathbf{n}(r))] dW_2(r) \right| \\ & \leq \mathbb{E} \left[\left(\int_0^s \Phi(r) \|\Delta((\mathbf{n}(r) \times h) \times h) - f'(\mathbf{n}(r))((\mathbf{n}(r) \times h) \times h)\|^2 dr \right)^{\frac{1}{2}} \right. \\ & \quad \left. \times \sup_{0 \leq s \leq t \wedge \tau_k} \sqrt{\Phi(s)} \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\| \right] \\ & \leq C \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) J_1(s) ds + \frac{1}{2} \mathbb{E} \sup_{0 \leq s \leq t} \Phi(s) \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2, \end{aligned}$$

where the process $J_1(s)$, $s < \tilde{\tau}_\infty$ is defined as in (B.2). Thanks to this last inequality and Eqs. (B.3) it is easy to deduce that

$$\begin{aligned} 2 \mathbb{E} \sup_{0 \leq s \leq t} \left| \int_0^{t \wedge \tau_k} \Phi(s) \Psi'(\mathbf{n}(s)) [G(\mathbf{n}(s))] dW_2(s) \right| & \leq C(h) \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 ds \\ & + C(h) t \wedge \tau_k \mathbb{E} \sup_{s \in [0, t \wedge \tau_k]} \Phi(s) \|\mathbf{n}(s)\|_{\mathbb{H}^1}^{4N+2} \\ & + \frac{1}{2} \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_k} \Phi(s) \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2. \end{aligned} \quad (\text{C.26})$$

With similar idea we can prove that

$$\begin{aligned} E_2 := 2 \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_k} \left| \int_0^s \Phi(r) \langle \nabla S(\mathbf{v}(r)), \nabla \mathbf{v}(r) \rangle dW_1(r) \right| & \leq \frac{1}{2} \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \|S(\mathbf{v}(s))\|_{\mathcal{H}_2(\mathbb{K}_1, \mathbb{V})}^2 ds \\ & + \frac{1}{2} \mathbb{E} \sup_{0 \leq s \leq t \wedge \tau_k} \Phi(s) \|\nabla \mathbf{v}(s)\|^2. \end{aligned} \quad (\text{C.27})$$

Plugging (C.26) and (C.27) in (C.24), using the hypotheses on S in Assumption 4.1 and invoking Proposition C.2 with $p = 2N + 1$ we infer that there exists a constant $\tilde{C} > 0$ such that

$$\begin{aligned} & \mathbb{E} \left[e^{-\int_0^{t \wedge \tau_k} \phi(r) dr} \left(\|\nabla \mathbf{v}(t \wedge \tau_k)\|^2 + \Psi(\mathbf{n}(t \wedge \tau_k)) \right) \right] \\ & + \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \left(2\|\Delta \mathbf{v}(s)\|^2 + \|\nabla(\Delta \mathbf{n}(s) - f(\mathbf{n}(s)))\|^2 \right) ds \\ & \leq \|\nabla \mathbf{v}_0\|^2 + \Psi(\mathbf{n}_0) + \tilde{C} \mathbb{E} \int_0^{t \wedge \tau_k} \Phi(s) \left(\|\nabla \mathbf{v}(s)\|^2 + \|\Delta \mathbf{n}(s) - f(\mathbf{n}(s))\|^2 \right) ds \\ & + \tilde{C}(\mathbb{E} \mathfrak{G}_1(t, 2N + 1) + 1), \end{aligned}$$

which altogether with the Gronwall inequality implies the desired estimate (4.26). \square

Acknowledgments. Razafimandimby's research was partially supported by the FWF-Austrian Science through the Stand-Alone project P28010. The research on this paper was initiated during the visit of Razafimandimby to the University of York in October 2012. Part of this work was also carried out when he visited the University of York during October 2013, February 2014. He would like to thank the Mathematics Department at York for hospitality. Z. Brzeźniak presented a lecture on the subject of this paper at the RIMS Symposium on Mathematical Analysis of Incompressible Flow held at Kyoto in February 2013. He would like to thank Professor Toshiaki Hishida for the kind invitation. P. Razafimandimby is very grateful to the organizers of the conference "Nonlinear PDEs in Micromagnetism: Analysis, Numerics and Applications", which was held in ICMS Edinburgh, UK, for their invitation to present a talk at this meeting. He is also very grateful for the financial support he received from the International Centre for Mathematical Sciences (ICMS) Edinburgh.

REFERENCES

- [1] S. Albeverio, Z. Brzeźniak and J.-L. Wu. *Existence of global solutions and invariant measures for stochastic differential equations driven by Poisson type noise with non-Lipschitz coefficients*. J. Math. Anal. Appl. 371(01):309-322, 2010.
- [2] W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander. VECTOR-VALUED LAPLACE TRANSFORMS AND CAUCHY PROBLEMS. Second edition. Monographs in Mathematics, **96**. Birkhäuser/Springer Basel AG, Basel, 2011.
- [3] K. Atkinson and W. Han, THEORETICAL NUMERICAL ANALYSIS. A FUNCTIONAL ANALYSIS FRAMEWORK. Third edition. Volume 39 of *Texts in Applied Mathematics*, Springer, Dordrecht, 2009.
- [4] H. Bessaih, E. Hausenblas and P. R. Razafimandimby. *Strong solutions to stochastic hydrodynamical systems with multiplicative noise of jump type*. NoDEA Nonlinear Differential Equations Appl. **22**(6): 1661–1697, 2015.
- [5] Z. Brzeźniak. *On stochastic convolution in Banach spaces and applications*. Stochastics Stochastics Rep. **61**(3-4): 245–295, 1997.
- [6] Z. Brzeźniak and K.D. Elworthy, *Stochastic differential equations on Banach manifolds*, Methods Funct. Anal. Topology **6**(1):43–84, 2000.
- [7] Z. Brzeźniak, B. Goldys and T. Jegaraj, *Weak solution of a Stochastic Landau-Lifshitz-Gilbert Equation*, Appl. Math. Res. Express, pp. 33, 2012.
- [8] Z. Brzeźniak, E. Hausenblas, and J. Zhu. *2D stochastic Navier-Stokes equations driven by jump noise*. Nonlinear Anal. **79**:122–139, 2013.
- [9] Z. Brzeźniak, E. Hausenblas, and P. Razafimandimby. *Stochastic Nonparabolic dissipative systems modeling the flow of Liquid Crystals: Strong solution*. RIMS Symposium on Mathematical Analysis of Incompressible Flow, February 2013. RIMS Kôkyûroku **1875**: 41–73, 2014.
- [10] Z. Brzeźniak and Y. Li, *Asymptotic compactness and absorbing sets for 2D stochastic Navier-Stokes equations on some unbounded domains*. Trans. Amer. Math. Soc. **358**(12):5587–5629, 2006.
- [11] Z. Brzeźniak, B. Maslowski and J. Seidler, *Stochastic nonlinear beam equations*, Probab. Theory Related Fields, **132**(1):119-149, 2005.
- [12] Z. Brzeźniak and A. Millet. *On the stochastic Strichartz estimates and the stochastic nonlinear Schrödinger equation on a compact Riemannian manifold*. Potential Anal. **41**(2): 269–315, 2014.
- [13] Z. Brzeźniak and E. Motyl. *Existence of a martingale solution of the stochastic Navier-Stokes equations in unbounded 2D and 3D domains*. J. Differential Equations **254**(4): 1627–1685, 2013.
- [14] S. Chandrasekhar, LIQUID CRYSTALS, Cambridge University Press, 1992.
- [15] B. Climent-Ezquerro, F. Guillén-González and M. A. Rojas-Medar, *Reproductivity for a nematic liquid crystal model*, Z. Angew. Math. Phys. **57**(06):984-998, 2006.

- [16] M. Dai and M. Schonbek, *Asymptotic behavior of solutions to the liquid crystal system in $H^m(R^3)$* . SIAM J. Math. Anal. **46**(5): 3131–3150, 2014.
- [17] A. de Bouard and A. Debussche, *A stochastic nonlinear Schrödinger equation with multiplicative noise*, Comm. Math. Phys. **205**(1):161–181, 1999.
- [18] A. de Bouard and A. Debussche, *The stochastic nonlinear Schrödinger equation in H^1* , Stochastic Anal. Appl. **21**(1):97–126, 2003.
- [19] A. Debussche, N. Glatt-Holtz and R. Temam. *Local martingale and pathwise solutions for an abstract fluids model*. Phys. D **240**(14-15): 1123–1144, 2011.
- [20] P. G. de Gennes and J. Prost, *THE PHYSICS OF LIQUID CRYSTALS*. Clarendon Press, Oxford 1993.
- [21] L. Denis, A. Matoussi and L. Stoica, *Maximum principle and comparison theorem for quasi-linear stochastic PDE's*, Electron. J. Probab. **14**(19):500-530, 2009.
- [22] K.D. Elworthy, *STOCHASTIC DIFFERENTIAL EQUATIONS ON MANIFOLDS*, London Math. Soc. LNS v 70, Cambridge University Press, 1982.
- [23] J. L. Ericksen, *Conservation laws for liquid crystals*, Trans. Soc. Rheology, **5**: 23-34, 1961.
- [24] F. Flandoli and D. Gatarek. *Martingale and stationary solutions for stochastic Navier-Stokes equations*. Probab. Theory Related Fields. **102**(03):367-391, 1995.
- [25] N. Glatt-Holtz and M. Ziane. *Strong pathwise solutions of the stochastic Navier-Stokes system*. Adv. Differential Equations **14**(5-6):567–600, 2009.
- [26] F. Guillén-González and M. Rojas-Medar, *Global solution of nematic liquid crystals models*. C.R. Acad. Sci. Paris. Série I, **335**:1085-1090, 2002.
- [27] D. D. Haroske and H. Triebel, *DISTRIBUTIONS, SOBOLEV SPACES, ELLIPTIC EQUATIONS*. EMS Textbooks in Mathematics. European Mathematical Society, Zürich, 2008.
- [28] M.-C. Hong, *Global existence of solutions of the simplified Ericksen-Leslie system in dimension two*. Calculus of Variations, **40**: 15-36, 2011.
- [29] W. Horsthemke and R. Lefever, *NOISE-INDUCED TRANSITIONS. THEORY AND APPLICATIONS IN PHYSICS, CHEMISTRY, AND BIOLOGY*. Springer Series in Synergetics, 15. Springer-Verlag, Berlin, 1984.
- [30] N. Ikeda and S. Watanabe. *STOCHASTIC DIFFERENTIAL EQUATIONS AND DIFFUSION PROCESSES*. Second edition. Volume 24 of North-Holland Mathematical Library, North-Holland Publishing Co., Amsterdam/Kodansha, Ltd., Tokyo, 1989.
- [31] F. Jiang and Z. Tan, *Global weak solution of the flow of liquid crystals system*. Mathematical Methods in the Applied Sciences, **32**: 2243-2266, 2009.
- [32] O. Kallenberg. *FOUNDATIONS OF MODERN PROBABILITY. Probability and its Applications* (New York). Springer-Verlag, New York, 1997.
- [33] I. Karatzas & S. Shreve, *BROWNIAN MOTION AND STOCHASTIC CALCULUS*, 2nd edition, Graduate Texts in Mathematics **113**, Springer Verlag, Berlin Heidelberg New York 1996.
- [34] R.Z. Khasminskii, *STABILITY OF SYSTEMS OF DIFFERENTIAL EQUATIONS UNDER RANDOM PERTURBATIONS OF THEIR PARAMETERS*. Nauka, Moskva, 1969 (Russian); Stochastic stability of differential equations, Sijthoff & Noordhoff, Alphen aan den Rijn, 1980.
- [35] H. Kunita, *STOCHASTIC FLOWS AND STOCHASTIC DIFFERENTIAL EQUATIONS*, Cambridge University Press, 1990.
- [36] F. M. Leslie, *Some constitutive equations for liquid crystals*, Arch. Rational Mech. Anal. **28**(04):265-283, 1968.
- [37] F.-H. Lin and C. Liu, *Nonparabolic dissipative systems modeling the flow of Liquid Crystals*. Communications on Pure and Applied Mathematics, Vol. **XLVIII**: 501-537, 1995.
- [38] F.-H. Lin and C. Liu, *Existence of solutions for the Ericksen-Leslie System*. Arch. Rational Mech. Anal. **154**:135-156, 2000.
- [39] F. Lin and C. Wang, *On the uniqueness of heat flow of harmonic maps and hydrodynamic flow of nematic liquid crystals*. Chinese Annals of Mathematics, Series B. **31B**(6): 921-938, 2010.
- [40] F. Lin, J. Lin and C. Wang, *Liquid crystals in two dimensions*. Arch. Rational Mech. Anal. **197**:297-336, 2010.
- [41] J.L. Lions. *Quelques méthodes de résolution de problèmes aux limites non-linéaires*. Dunod, Paris, 1969.
- [42] R. Mikulevicius. *On strong H^1_2 -solutions of stochastic Navier-Stokes equation in a bounded domain*. SIAM J. Math. Anal. **41**(03):1206-1230, 2009.
- [43] R. Mikulevicius and B. L. Rozovskii, *Stochastic navier-Stokes equations for turbulent flows*. SIAM J. Math. Anal. **35**(5):1250-1310, 2004.
- [44] M. Ondreját. *Stochastic nonlinear wave equations in local Sobolev spaces*. Electron. J. Probab. **15**(33): 1041–1091, 2010.
- [45] E. M. Ouhabaz, *ANALYSIS OF HEAT EQUATIONS ON DOMAINS*. Volume 31 of London Mathematical Society Monographs Series. Princeton University Press, Princeton, 2005.
- [46] E. Pardoux, *Stochastic partial differential equations and filtering of diffusion processes*, Stochastics **3**(02):127-167, 1979.
- [47] P. E. Protter. *Stochastic integration and differential equations*. Second edition. Volume 21 of *Applications of Mathematics*. Stochastic Modelling and Applied Probability. Springer-Verlag, Berlin, 2004.
- [48] A. F. Ruston, *FREDHOLM THEORY IN BANACH SPACES*. Volume 86 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1986.
- [49] M. San Miguel. *Nematic liquid crystals in a stochastic magnetic field: Spatial correlations*. Phys. Rev. A **32**: 3811-3813, 1985.

- [50] F. Sagués and M. San Miguel. *Dynamics of Fréedericksz transition in a fluctuating magnetic field*. Phys. Rev. A. **32**(3): 1843–1851, 1985.
- [51] B. Schmalfuss. *Qualitative properties for the stochastic Navier-Stokes equation*. Nonlinear Anal. **28**(9):1545–1563, 1997.
- [52] J. T. Schwartz. NONLINEAR FUNCTIONAL ANALYSIS. Gordon and Breach Science Publishers, New York, 1969.
- [53] S. Shkoller. *Well-posedness and global attractors for liquid crystal on Riemannian manifolds*. Communication in Partial Differential Equations. **27**(5&6):1103–1137, 2002.
- [54] J. Simon. *Compact sets in the space $L^p(0, T; B)$* . Ann. Mat. Pura Appl.(4) **146**: 65–96, 1987.
- [55] J. Simon. *Sobolev, Besov and Nikol'skii fractional spaces: imbeddings and comparisons for vector valued spaces on an interval*. Ann. Mat. Pura Appl. (4) **157**:117–148, 1990.
- [56] W. A. Strauss, *On continuity of functions with values in various Banach spaces*. Pacific J. Math. **19**: 543–551, 1966.
- [57] R. Temam. NAVIER-STOKES EQUATIONS. North-Holland, 1979.
- [58] M.I. Vishik and A. V. Fursikov, MATHEMATICAL PROBLEMS OF STATISTICAL HYDROMECHANICS (Russian), “Nauka”, Moscow, 1980.
- [59] H. Wen and S. Ding. *Solutions of incompressible hydrodynamic flow of liquid crystals*. Nonlinear Analysis: Real World Applications, **12**:1510–1531, 2011.
- [60] E. Zeidler. *Nonlinear Functional Analysis and its Applications, II/A: Linear Monotone Operators*. Springer-Verlag, New York, 1990.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF YORK, HESLINGTON, YORK YO10 5DD, UK
 E-mail address: `zdzislaw.brzezniak@york.ac.uk`

DEPARTMENT OF MATHEMATICS AND INFORMATION TECHNOLOGY, MONTANUNIVERSITÄT LEOBEN, FR. JOSEFSTR.
 18, 8700 LEOBEN, AUSTRIA
 E-mail address: `erika.hausenblas@unileoben.ac.at`

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF PRETORIA, CORNER OF LYNNWOOD
 ROAD AND ROPER STREET HATFIELD, PRETORIA 0083, SOUTH AFRICA
 E-mail address: `paul.razafimandimby@up.ac.za`