

Entire solutions for a class of elliptic equations involving p-biharmonic operator and Rellich potentials

Mousomi Bhakta *

Abstract

We study existence, multiplicity and qualitative properties of entire solutions for a noncompact problem related to p-biharmonic type equations with weights. More precisely, we deal with the following family of equations

$$\Delta_p^2 u = \lambda |x|^{-2p} |u|^{p-2} u + |x|^{-\beta} |u|^{q-2} u \quad \text{in } \mathbb{R}^N,$$

where $N > 2p$, $p > 1$, $q > p$, $\beta = N - \frac{q}{p}(N - 2p)$ and $\lambda \in \mathbb{R}$ is smaller than the Rellich constant.

Keywords: Caffarelli-Kohn-Nirenberg type inequalities, weighted p-biharmonic operator, Rellich inequality, dilation invariance, breaking symmetry, extremal function.

2010 Mathematics Subject Classification: 26D10, 47F05.

1 Introduction

In this article we study weak solutions to the problem

$$\begin{cases} \Delta_p^2 u = \lambda |x|^{-2p} |u|^{p-2} u + |x|^{-\beta} |u|^{q-2} u & \text{in } \mathbb{R}^N \\ u \in D^{2,p}(\mathbb{R}^N), \quad u \neq 0, \end{cases} \quad (1.1)$$

where $\Delta_p^2 u = \Delta(|\Delta u|^{p-2} \Delta u)$ and

$$p > 1, N > 2p, \quad q > p, \quad \beta = N - \frac{q}{p}(N - 2p), \quad \lambda < \gamma_{N,p}^p \text{ and } \gamma_{N,p} = \frac{N(p-1)(N-2p)}{p^2}. \quad (1.2)$$

Here $D^{2,p}(\mathbb{R}^N)$ is the closure of $C_0^\infty(\mathbb{R}^N)$ with respect to the norm $(\int_{\mathbb{R}^N} |\Delta u|^p dx)^{\frac{1}{p}}$. It is well known that $\gamma_{N,p}^p$ is the best constant in the Rellich inequality

$$\gamma_{N,p}^p \int_{\mathbb{R}^N} |x|^{-2p} |u|^p dx \leq \int_{\mathbb{R}^N} |\Delta u|^p dx \quad \text{for any } u \in D^{2,p}(\mathbb{R}^N). \quad (1.3)$$

*Department of Science and Technology, University of New England, Armidale, NSW-2350, Australia.
Email: mousomi.bhakta@gmail.com

In literature (1.3) with $p = 2$ is considered as the classical Rellich inequality and was proved by Rellich in 1953 (See [18], [19]). Later Davies and Hinz [8] generalized the classical Rellich inequality and showed that (1.3) holds for any $p \in (1, \frac{N}{2})$.

If q coincides with the critical Sobolev exponent

$$p^{**} := \frac{Np}{N-2p},$$

then (1.1) becomes

$$\begin{cases} \Delta_p^2 u = \lambda |x|^{-2p} |u|^{p-2} u + |u|^{p^{**}-2} u & \text{in } \mathbb{R}^N \\ u \in D^{2,p}(\mathbb{R}^N), \quad u \neq 0. \end{cases} \quad (1.4)$$

When $\lambda = 0$, it's well known from the celebrated paper of P.L. Lions [14] that (1.4) has a positive solution U which is the extremal for the Sobolev inequality

$$S^{**} \left(\int_{\mathbb{R}^N} |u|^{p^{**}} dx \right)^{\frac{p}{p^{**}}} \leq \int_{\mathbb{R}^N} |\Delta u|^p dx \quad (1.5)$$

where S^{**} is the Sobolev Constant. To prove that S^{**} is achieved, P.L. Lions had shown that every bounded minimizing sequence is relatively compact up to dilations and translations. Moreover, by using Schwarz symmetrization he showed that, up to a change of sign, any extremal for S^{**} is radially symmetric, nonnegative and decreasing. Later using this information, Hulshof and Van der Vorst [12] proved the uniqueness of extremals for S^{**} , modulo dilations, translations in \mathbb{R}^N and change of sign.

If $p \leq q \leq p^{**}$ and β are as in (1.2), then by interpolating (1.3) and (1.5) via Hölder inequality, it can be easily shown that there exists a constant $C = C(N, p, q) > 0$ such that

$$C \left(\int_{\mathbb{R}^N} |x|^{-\beta} |u|^q dx \right)^{p/q} \leq \int_{\mathbb{R}^N} |\Delta u|^p dx \quad \text{for any } u \in D^{2,p}(\mathbb{R}^N). \quad (1.6)$$

Notice that (1.6) with $p = 2$ is the fourth-order version of the celebrated Caffarelli-Kohn-Nirenberg inequalities [3]. We cite also [5] for a large class of dilation-invariant inequalities on cones.

In recent years problems related with the inequality (1.6) (in the case $p = 2$) and the equation with biharmonic operator have been investigated in several works, we quote [1] [2], [4], [5], [6], [7], [11], [15], [16], [17] and the references there-in. Recently the generalized version of the inequality (1.6) and the extremal of that inequality has been studied by Roberta Musina. (see [16] and [17]).

Note that the choice of β in (1.2) makes (1.1) invariant with respect to the weighted dilation

$$u(x) \mapsto t^{\frac{N-2p}{p}} u(tx), \quad (t > 0). \quad (1.7)$$

As a consequence, the corresponding variational problems exhibit a lack of compactness.

It is clear that the infimum

$$S_q(\lambda) := \inf_{\substack{u \in D^{2,p}(\mathbb{R}^N) \\ u \neq 0}} \frac{\int_{\mathbb{R}^N} |\Delta u|^p dx - \lambda \int_{\mathbb{R}^N} |x|^{-2p} |u|^p dx}{\left(\int_{\mathbb{R}^N} |x|^{-\beta} |u|^q dx \right)^{p/q}}$$

is positive, provided that $\lambda < \gamma_{N,p}^p$. In addition, extremals for $S_q(\lambda)$ give rise to solutions to (1.1) upto a multiplicative constant.

Define,

$$S_q^{\text{rad}}(\lambda) := \inf_{\substack{u \in D^{2,p}(\mathbb{R}^N) \\ u = u(|x|), u \neq 0}} \frac{\int_{\mathbb{R}^N} |\Delta u|^p dx - \lambda \int_{\mathbb{R}^N} |x|^{-2p} |u|^p dx}{\left(\int_{\mathbb{R}^N} |x|^{-\beta} |u|^q dx \right)^{p/q}}$$

which is positive as $S_q^{\text{rad}}(\lambda) \geq S_q(\lambda) > 0$ when $\lambda < \gamma_{N,p}^p$.

We prove the following results.

Theorem 1.1 *Let $N > 2p$, $q > p$, $\lambda < \gamma_{N,p}^p$, and $\beta = N - \frac{q}{p}(N - 2p)$. Then the problem (1.1) has at least one radially symmetric solution u which achieves $S_q^{\text{rad}}(\lambda)$.*

In fact when $p = 2$ and $-(N - 2)^2 \leq \lambda < (\gamma_{N,2})^2$, problem (1.1) has a unique radial solution. (See [2]).

For $\lambda = 0$, Theorem 1.1 was proved in [17, Theorem 1.3]. Following the same procedure as in [17], Theorem 1.1 can be proved in the case $\lambda \neq 0$ as well.

In case $q \leq p^*$ one can use again variational methods to find solutions to (1.1) that are not necessarily radially symmetric.

In Section 2 we prove the next existence result.

Theorem 1.2 *Let $N > 2p$, $q \in (p, p^*]$, $\lambda < \gamma_{N,p}^p$, and $\beta = N - \frac{q}{p}(N - 2p)$. Then*

- (i) *The infimum $S_q(\lambda)$ is achieved for any $q \in (p, p^*)$.*
- (ii) *The infimum $S_{p^*}(\lambda)$ is achieved if and only if $\lambda \geq 0$.*

In Section 3 we wonder whether the solutions in Theorem 1.2 are radially symmetric or breaking symmetry occurs. First, by using rearrangement techniques we prove that extremal for $S_q(\lambda)$ is always radially symmetric provided that $\lambda \geq 0$. In contrast, we show that if $\lambda \ll 0$ and $p \geq 2$ then $S_q(\lambda) < S_q^{\text{rad}}(\lambda)$ i.e break of symmetry occurs. Therefore, if in addition $q \in (p, p^*)$ then problem (1.1) has at least two distinct nontrivial solutions.

2 Existence and non existence of ground state solutions

In this section we will prove Theorem 1.2. A key tool in our argument is the following ε – compactness lemma. This result is an adaptation of a tool already used in previous works, like [2] or [5]. Therefore we omit the proof.

Lemma 2.1 *Let $u_n \in D^{2,p}(\mathbb{R}^N)$ such that $u_n \rightharpoonup 0$ in $\mathcal{D}^{2,p}(\mathbb{R}^N)$ and*

$$\Delta_p^2 u_n - \lambda |x|^{-2p} |u_n|^{p-2} u_n = |x|^{-\beta} |u_n|^{q-2} u_n + f_n \quad (2.1)$$

$$\int_{B_R} |x|^{-\beta} |u_n|^q dx \leq \varepsilon_0 \quad \text{for some } \varepsilon_0, R > 0, \quad (2.2)$$

where $f_n \rightarrow 0$ in the dual space of $D^{2,p}(\mathbb{R}^N)$. If $\varepsilon_0 < S_q(\lambda)^{\frac{q}{q-p}}$, then

$$|x|^{-\beta} |u_n|^q \rightarrow 0 \quad \text{in } L_{\text{loc}}^1(B_R).$$

2.1 Proof of Theorem 1.2 (i)

Step 1: Using Ekeland's variational principle we can choose a minimizing sequence $\{u_n\}$ for $S_q(\lambda)$ such that

$$\begin{aligned} \int_{\mathbb{R}^N} |\Delta u_n|^p - \lambda \int_{\mathbb{R}^N} |x|^{-2p} |u_n|^p &= \int_{\mathbb{R}^N} |x|^{-\beta} |u_n|^q dx \\ &= S_q(\lambda)^{\frac{q}{q-p}} + o(1), \end{aligned} \quad (2.3)$$

$$\Delta_p^2 u_n - \lambda |x|^{-2p} |u_n|^{p-2} u_n = |x|^{-\beta} |u_n|^{q-2} u_n + f_n, \quad (2.4)$$

where $f_n \rightarrow 0$ in the dual space of $D^{2,p}(\mathbb{R}^N)$. Up to a rescaling, we assume that

$$\int_{B_2} |x|^{-\beta} |u_n|^q dx = \frac{1}{2} S_q(\lambda)^{\frac{q}{q-p}}. \quad (2.5)$$

Therefore it can be easily checked that u_n is a bounded sequence in $D^{2,p}(\mathbb{R}^N)$ by Rellich inequality as we have $\lambda < \gamma_{N,p}^p$. Hence we can assume that there exists $u \in D^{2,p}(\mathbb{R}^N)$ such that $u_n \rightharpoonup u$ weakly in $D^{2,p}(\mathbb{R}^N)$.

Claim 1: $u \neq 0$.

We argue by contradiction. Suppose $u_n \rightharpoonup 0$. Then using Lemma (2.1) we obtain

$$o(1) = \int_{B_1} |x|^{-\beta} |u_n|^q dx = \int_{B_2} |x|^{-\beta} |u_n|^q dx - \int_{1 < |x| < 2} |x|^{-\beta} |u_n|^q dx.$$

Thus from (2.5) we infer

$$\int_{1 < |x| < 2} |x|^{-\beta} |u_n|^q dx = \frac{1}{2} S_q(\lambda)^{\frac{q}{q-p}} + o(1) \quad (2.6)$$

which leads to a contradiction by Rellich's compactness theorem, as $q \in (p, p^{**})$. Thus Claim 1 follows.

Claim 2: u is a weak solution of (1.1).

For $p = 2$, it is straight forward. For $p \neq 2$, we choose $w_n \in D^{2,p}(\mathbb{R}^N)$ such that

$$\Delta(|\Delta w_n|^{p-2} \Delta w_n) = f_n; \quad \text{and} \quad w_n \rightarrow 0 \quad \text{in} \quad D^{2,p}(\mathbb{R}^N),$$

where f_n is as in (2.4). Define, $h_n := -|\Delta u_n|^{p-2} \Delta u_n + |\Delta w_n|^{p-2} \Delta w_n$, that is bounded in $L^{\frac{p}{p-1}}(\mathbb{R}^N)$. Also from (2.4), it follows that,

$$-\Delta h_n = \lambda |x|^{-2p} |u_n|^{p-2} u_n + |x|^{-\beta} |u_n|^{q-2} u_n. \quad (2.7)$$

As $p < q < p^{**}$, it is easy to see that 1st term of RHS of (2.7) is bounded in $L^{\frac{q}{q-1}}(\mathbb{R}^N \setminus \{0\})$ and 2nd term of RHS of (2.7) is bounded in $L^{\frac{q}{q-1}}(\mathbb{R}^N \setminus \{0\})$. Since $p < q$ implies $\frac{q}{q-1} < \frac{q}{p-1}$, we can easily conclude that $-\Delta h_n$ is bounded in $L^{\frac{q}{q-1}}(\mathbb{R}^N \setminus \{0\})$, which in turn implies h_n is bounded in $W_{loc}^{2, \frac{q}{q-1}}(\mathbb{R}^N \setminus \{0\})$. Therefore almost everywhere convergence of h_n follows and so of u_n , as $w_n \rightarrow 0$ a.e. Hence using Vitaly's convergence theorem via Holder inequality, we obtain

$$\int_{\mathbb{R}^N} |\Delta u_n|^{p-2} \Delta u_n \Delta \phi \rightarrow \int_{\mathbb{R}^N} |\Delta u|^{p-2} \Delta u \Delta \phi \quad \forall \quad \phi \in C_0^\infty(\mathbb{R}^N).$$

Therefore it is easy to see that the claim follows.

Step 2: Therefore $S_q(\lambda) \left(\int_{\mathbb{R}^N} \frac{|u|^q}{|x|^\beta} dx \right)^{\frac{p}{q}} \leq \int_{\mathbb{R}^N} \left(|\Delta u|^p - \lambda \frac{|u|^p}{|x|^{2p}} \right) dx = \int_{\mathbb{R}^N} \frac{|u|^q}{|x|^\beta} dx$.

Since $u \neq 0$, we obtain $\int_{\mathbb{R}^N} \frac{|u|^q}{|x|^\beta} dx \geq S_q(\lambda)^{\frac{q}{q-p}}$. On the other hand by the lower semicontinuity of the norm in $L^q(\mathbb{R}^N, |x|^{-\beta} dx)$ and (2.3), we have $\int_{\mathbb{R}^N} \frac{|u|^q}{|x|^\beta} dx \leq S_q(\lambda)^{\frac{q}{q-p}}$.

Therefore $\int_{\mathbb{R}^N} \frac{|u|^q}{|x|^\beta} dx = S_q(\lambda)^{\frac{q}{q-p}}$ which in turn implies that $S_q(\lambda)$ is achieved by u . □

2.2 Proof of Theorem 1.2 in the limiting case $q = p^{**}$

We start by pointing out a sufficient condition for existence.

Lemma 2.2 *If $S_{p^{**}}(\lambda) < S^{**}$ then $S_{p^{**}}(\lambda)$ is achieved.*

Proof. As in the proof of Theorem 1.2 (i), we choose a minimizing sequence $\{u_n\}$ for $S_{p^{**}}(\lambda)$ satisfying

$$\begin{aligned} \int_{\mathbb{R}^N} |\Delta u_n|^p dx - \lambda \int_{\mathbb{R}^N} |x|^{-2p} |u_n|^p dx &= \int_{\mathbb{R}^N} |u_n|^{p^{**}} dx + o(1) \\ &= S_{p^{**}}(\lambda)^{N/2p} + o(1) \end{aligned}$$

$$\Delta_p^2 u_n - \lambda |x|^{-2p} |u_n|^{p-2} u_n = |u_n|^{p^{**}-2} u_n + f_n \quad (2.8)$$

$$\int_{B_2} |u_n|^{p^{**}} dx = \frac{1}{2} S_{p^{**}}(\lambda)^{N/2p},$$

where $f_n \rightarrow 0$ in the dual of $D^{2,p}(\mathbb{R}^N)$. In addition, we can assume that u_n weakly converges to u in $D^{2,p}(\mathbb{R}^N)$.

Claim 1: $u \neq 0$.

We will prove the claim by contradiction, thus we assume $u_n \rightharpoonup 0$ in $D^{2,p}(\mathbb{R}^N)$. Arguing as in the proof of Theorem 1.2 (i) we can conclude that (2.6) holds with $q = p^{**}$, that is,

$$\int_{1 < |x| < 2} |u_n|^{p^{**}} dx = \frac{1}{2} S_{p^{**}}(\lambda)^{N/2p} + o(1). \quad (2.9)$$

Now we choose a cut off function ϕ in $C_c^\infty(\mathbb{R}^N \setminus \{0\})$ such that $\phi \equiv 1$ in $B_2 \setminus B_1$. Taking $\phi^p u_n$ as a test function in (2.8) and using Rellich's compactness theorem and Hölder inequality we obtain

$$\int_{\mathbb{R}^N} [|\Delta(\phi u_n)|^p - \lambda |x|^{-2p} |\phi u_n|^p] dx \leq S_{p^{**}}(\lambda) \left(\int_{\mathbb{R}^N} |\phi u_n|^{p^{**}} dx \right)^{\frac{p}{p^{**}}} + o(1)$$

Since ϕ has a compact support in $\mathbb{R}^N \setminus \{0\}$, using Rellich's compactness theorem and the Sobolev inequality and arguing as in the proof of Theorem 1.2 (i) we obtain

$$\int_{\mathbb{R}^N} [|\Delta(\phi u_n)|^p - \lambda |x|^{-2p} |\phi u_n|^p] = \int_{\mathbb{R}^N} |\Delta(\phi u_n)|^p + o(1) \geq S^{**} \left(\int_{\mathbb{R}^N} |\phi u_n|^{p^{**}} dx \right)^{p/p^{**}}.$$

Therefore we have,

$$S^{**} \left(\int_{\mathbb{R}^N} |\phi u_n|^{p^{**}} dx \right)^{p/p^{**}} \leq S_{p^{**}}(\lambda) \left(\int_{\mathbb{R}^N} |\phi u_n|^{p^{**}} dx \right)^{p/p^{**}}.$$

As $S_{p^{**}}(\lambda) < S^{**}$, the above inequality implies $\int_{\mathbb{R}^N} |\phi u_n|^{p^{**}} = o(1)$. Hence $\int_{B_2 \setminus B_1} |u_n|^{p^{**}} dx = o(1)$, since $\phi \equiv 1$ in $B_2 \setminus B_1$. This is a contradiction to (2.9). Thus $u \neq 0$.

It follows by a standard concentration-compactness technique by P. L. Lions (see the proof of [14, Theorem 2.4]) that u_n is relatively compact and therefore up to a subsequence $u_n \rightarrow u$ in $D^{2,p}(\mathbb{R}^N)$. Hence $S_{p^{**}}(\lambda)$ is achieved. □

Proof of Theorem 1.2 - (ii). First we show that $S_{p^{**}}(\lambda) < S^{**}$ holds when $\lambda \in (0, \gamma_{N,p}^p)$. Let U be an extremal of S^{**} which exists by [14]. Therefore if $\lambda \in (0, \gamma_{N,p}^p)$, then we have

$$S_{p^{**}}(\lambda) \leq \frac{\int_{\mathbb{R}^N} [|\Delta U|^p - \lambda |x|^{-2p} |U|^p] dx}{\left(\int_{\mathbb{R}^N} |U|^{p^{**}} dx \right)^{p/p^{**}}} < \frac{\int_{\mathbb{R}^N} |\Delta U|^p dx}{\left(\int_{\mathbb{R}^N} |U|^{p^{**}} dx \right)^{p/p^{**}}} = S^{*}.$$

hence $S_{p^{**}}(\lambda)$ is achieved.

Now it remains to study the case $\lambda < 0$. In this case, it is easy to see that $S_{p^{**}}(\lambda) \geq S^{**}$. Now we choose an arbitrary function u in $C_c^\infty(\mathbb{R}^N \setminus \{0\})$ and set $u_y(x) = u(x+y)$. Therefore

$$\begin{aligned} S_{p^{**}}(\lambda) &\leq \lim_{|y| \rightarrow \infty} \frac{\int [|\Delta u_y|^p - \lambda |x|^{-2p} |u_y|^p]}{\left(\int_{\mathbb{R}^N} |u_y|^{p^{**}} \right)^{p/p^{**}}} = \lim_{|y| \rightarrow \infty} \frac{\int [|\Delta u|^p - \lambda |x-y|^{-2p} |u|^p]}{\left(\int_{\mathbb{R}^N} |u|^{p^{**}} \right)^{p/p^{**}}} \\ &= \frac{\int |\Delta u|^p}{\left(\int_{\mathbb{R}^N} |u|^{p^{**}} \right)^{p/p^{**}}}, \end{aligned}$$

which implies $S_{p^{**}}(\lambda) \leq S^{**}$. Hence $S_{p^{**}}(\lambda) = S^{**}$. Therefore $S_{p^{**}}(\lambda)$ can not be achieved since S^{**} is achieved. □

3 Nonnegativity, symmetry and breaking symmetry

In this section we will study the symmetry, nonnegativity and breaking symmetry of the extremal of $S_q(\lambda)$ depending on the parameter λ . It's known from [14] that when $\lambda = 0$ and $q = p^{**}$, the Sobolev constant S^{**} is achieved by a radially symmetric nonnegative and decreasing function.

Since truncations $u \mapsto u^\pm$ are not allowed in dealing with fourth order differential operators, the nonnegativity of extremals for $S_q(\lambda)$ does not follow by usual arguments.

Theorem 3.1 *Assume $\lambda \neq 0$ or $q < p^{**}$. If $\lambda \geq 0$ then $S_q(\lambda)$ is achieved by a positive function $u \in D^{2,2}(\mathbb{R}^N)$. Moreover, u is radially symmetric about the origin and radially decreasing.*

The proof is based on rearrangement technique which was already used to prove this result in the case $p = 2$ in [2]. This is an easy adaptation of the proof used for $p = 2$.

Proof. Let u be an extremal of $S_q(\lambda)$ and we denote by $(-\Delta u)^*$ the Schwarz symmetrization of $-\Delta u$. Let $v \in D^{2,p}(\mathbb{R}^N)$ such that

$$-\Delta v = (-\Delta u)^*$$

(existence of such function follows from [14]). It turns out that $u^* \leq v$ on \mathbb{R}^N , see for instance Remark II.13 in [14]. If $u = u^*$ then we are done. So assuming $u \neq u^*$ we would like to derive a contradiction. By the theory of symmetrization (see Lieb and Loss [13], Theorem 3.4), we first obtain

$$\int_{\mathbb{R}^N} |\Delta v|^p dx = \int_{\mathbb{R}^N} |(-\Delta u)^*|^p dx = \int_{\mathbb{R}^N} |\Delta u|^p dx.$$

In addition, since we are assuming that $u^* \neq u$, then

$$\int_{\mathbb{R}^N} |x|^{-2p} |u|^p dx < \int_{\mathbb{R}^N} |x|^{-2p} (|u|^p)^* dx = \int_{\mathbb{R}^N} |x|^{-2p} |u^*|^p dx \leq \int_{\mathbb{R}^N} |x|^{-2p} |v|^p dx.$$

Thus we infer that

$$\lambda \int_{\mathbb{R}^N} |x|^{-2p} |u|^p dx \leq \lambda \int_{\mathbb{R}^N} |x|^{-2p} |v|^p dx,$$

and that the strict inequality holds if $\lambda > 0$. Similarly, we find

$$\int_{\mathbb{R}^N} |x|^{-\beta} |u|^q dx \leq \int_{\mathbb{R}^N} |x|^{-\beta} |v|^q dx,$$

and the strict inequality holds if $\beta > 0$, that is, if $q < p^*$. In conclusion, since we are assuming that λ and β are not contemporarily zero, we have that

$$S_q(\lambda) \leq \frac{\int_{\mathbb{R}^N} [|\Delta v|^p - \lambda |x|^{-2p} |v|^p] dx}{\left(\int_{\mathbb{R}^N} |x|^{-\beta} |v|^q dx \right)^{2/q}} < \frac{\int_{\mathbb{R}^N} [|\Delta u|^p - \lambda |x|^{-2p} |u|^p] dx}{\left(\int_{\mathbb{R}^N} |x|^{-\beta} |u|^q dx \right)^{2/q}} = S_q(\lambda),$$

a contradiction. Therefore $u = u^*$, that is, u is a nonnegative and radially symmetric decreasing function.

□

As soon as $\lambda \rightarrow -\infty$, a braking symmetry phenomenon appears. In the next theorem we study the case $q < p^*$, due to the nonexistence result pointed out in the critical case $q = p^*$, $\lambda < 0$. We cite [9], [10] for remarkable breaking symmetry results for similar second-order equations in the case $p = 2$. Also see [2] and [4] for the similar type of results in the case of biharmonic equations.

Theorem 3.2 *If $\lambda \ll 0$ and $2 \leq p < q < p^*$ then $S_q(\lambda) < S_q^{\text{rad}}(\lambda)$ and hence no extremal for $S_q(\lambda)$ is radially symmetric.*

Proof. We already know that $S_q(\lambda) \leq S_q^{\text{rad}}(\lambda)$. We will give an explicit condition on λ to have $S_q(\lambda) < S_q^{\text{rad}}(\lambda)$. Define

$$n(u) = \int_{\mathbb{R}^N} [|\Delta u|^p - \lambda |x|^{-2p} |u|^p] dx, \quad d(u) = \left(\int_{\mathbb{R}^N} |x|^{-\beta} |u|^q dx \right)^{p/q}$$

and $Q(u) = n(u)/d(u)$. Let u be a radially symmetric minimizer of Q on $\mathcal{D}^{2,p}(\mathbb{R}^N)$. Our goal is to show that $-\lambda$ can not be too large. By homogeneity we can assume that $d(u) = 1$. Thus $Q'(u) \cdot v = 0$ and $Q''(u)[v, v] \geq 0$ for all $v \in D^{2,p}(\mathbb{R}^N)$, that is, and

$$\begin{aligned} n'(u) \cdot v &= Q(u) d'(u) \cdot v \\ n''(u)[v, v] &\geq Q(u) d''(u)[v, v] \end{aligned} \tag{3.1}$$

for all $v \in D^{2,p}(\mathbb{R}^N)$.

Let $\varphi_1 \in H^1(\mathbb{S}^{N-1})$ be an Eigenfunction of Laplace-Beltrami operator on \mathbb{S}^{N-1} corresponding to the smallest positive Eigenvalue. Thus

$$-\Delta_\sigma \varphi_1 = (N-1)\varphi_1, \quad \frac{1}{|\mathbb{S}^{N-1}|} \int_{\mathbb{S}^{N-1}} |\varphi_1|^2 d\sigma = 1, \quad \int_{\mathbb{S}^{N-1}} \varphi_1 d\sigma = 0.$$

Now we set the test function v as $v(x) := u(|x|)\varphi_1(\frac{x}{|x|})$. Therefore it turns out that

$$\begin{aligned} d''(u)[v, v] &= p(p-q) \left(\int_{\mathbb{R}^N} |x|^{-\beta} |u|^{q-2} uv \right)^2 + p(q-1) \int_{\mathbb{R}^N} |x|^{-\beta} |u|^{q-2} v^2 \\ &= p(q-1) \int_{\mathbb{R}^N} |x|^{-\beta} |u|^q = p(q-1). \end{aligned} \quad (3.2)$$

Also we see that

$$|\Delta v|^2 = |\Delta(u\varphi_1)|^2 = |\Delta u - (N-1)|x|^{-2}u|^2 \varphi_1^2$$

and therefore as $p \geq 2$, we have

$$\begin{aligned} n''(u)[v, v] &= p(p-1) \int_{\mathbb{R}^N} (|\Delta u|^{p-2} |\Delta v|^2 - \lambda |x|^{-2p} |u|^{p-2} |v|^2) dx \\ &= p(p-1) \left[\int_{\mathbb{R}^N} |\Delta u|^{p-2} |\Delta u - (N-1) \frac{u}{|x|^2}|^2 \varphi_1^2 - \lambda |x|^{-2p} |u|^p \varphi_1^2 \right] dx \\ &= p(p-1) \int_{\mathbb{R}^N} \left[|\Delta u|^{p-2} |\Delta u - (N-1) \frac{u}{|x|^2}|^2 - \lambda |x|^{-2p} |u|^p \right] dx \\ &= p(p-1) \int_{\mathbb{R}^N} \left[|\Delta u|^p + \frac{(N-1)^2 u^2 |\Delta u|^{p-2}}{|x|^4} - \frac{2(N-1) |\Delta u|^{p-2} u \Delta u}{|x|^2} - \lambda \frac{|u|^p}{|x|^{2p}} \right] dx \end{aligned}$$

Since $p \geq 2$, using Hölder inequality and the fact that $d(u) = 1$ we obtain,

$$\begin{aligned} n''(u)[v, v] &\leq p(p-1) \left[n(u) + 2(N-1) \left(\int_{\mathbb{R}^N} |\Delta u|^p dx \right)^{\frac{p-1}{p}} \left(\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{2p}} dx \right)^{\frac{1}{p}} \right] \\ &\quad + p(p-1)(N-1)^2 \left(\int_{\mathbb{R}^N} |\Delta u|^p dx \right)^{\frac{p-2}{p}} \left(\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{2p}} dx \right)^{\frac{2}{p}}. \end{aligned}$$

Therefore from (3.1), (3.2) and the definition of $Q(u) = n(u)$ we obtain

$$\begin{aligned} (q-p)Q(u) &\leq 2(p-1)(N-1) \left(\int_{\mathbb{R}^N} |\Delta u|^p dx \right)^{\frac{p-1}{p}} \left(\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{2p}} dx \right)^{\frac{1}{p}} \\ &\quad + (p-1)(N-1)^2 \left(\int_{\mathbb{R}^N} |\Delta u|^p dx \right)^{\frac{p-2}{p}} \left(\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{2p}} dx \right)^{\frac{2}{p}}. \end{aligned}$$

Thus we have,

$$\begin{aligned} (q-p) \int_{\mathbb{R}^N} |\Delta|^p dx &\leq \lambda(q-p) \int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{2p}} + 2(p-1)(N-1) \left(\int_{\mathbb{R}^N} |\Delta u|^p dx \right)^{\frac{p-1}{p}} \left(\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{2p}} dx \right)^{\frac{1}{p}} \\ &\quad + (p-1)(N-1)^2 \left(\int_{\mathbb{R}^N} |\Delta u|^p dx \right)^{\frac{p-2}{p}} \left(\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^{2p}} dx \right)^{\frac{2}{p}}. \end{aligned}$$

In particular, the quantity

$$X := \left(\frac{\int_{\mathbb{R}^N} |\Delta u|^p}{\int_{\mathbb{R}^N} |x|^{-2p} |u|^p} \right)^{1/p}$$

satisfies the inequality

$$(q-p)X^p \leq \lambda(q-p) + 2(N-1)(p-1)X^{p-1} + (p-1)(N-1)^2 X^{p-2},$$

which implies

$$\lambda \geq \min_{t \in \mathbb{R}} f(t),$$

where

$$f(t) = \left(t^p - \frac{2(N-1)(p-1)}{q-p} t^{p-1} - \frac{(p-1)(N-1)^2}{q-p} t^{p-2} \right).$$

Define

$$t_0 = \frac{\gamma_1(p-1) + \sqrt{\gamma_1^2(p-1)^2 + 4\gamma_2 p(p-2)}}{2p}$$

where $\gamma_1 = \frac{2(p-1)(N-1)}{q-p}$ and $\gamma_2 = \frac{(p-1)(N-1)^2}{q-p}$. By a straight forward calculation we obtain $\gamma \geq f(t_0)$.

Therefore no extremal for $S_q(\lambda)$ is radially symmetric and break of symmetry occurs if $\lambda < f(t_0)$. \square

Remark 3.3 *If q is close enough to p^* then one can obtain a better estimate on the breaking symmetry parameter λ by arguing as follows. Notice that $X > \gamma_{N,p}$ by the Rellich inequality (1.3). Thus, if $\gamma_{N,p} \geq t_0$ that is, if*

$$p + \frac{p(N-1) [p^2(N-1)(p-2) + 2N(p-1)^2(N-2p)]}{N^2(p-1)(N-2p)^2} \leq q \leq p^*,$$

then the radial solution u does not achieve $S_q(\lambda)$ unless

$$\lambda > \min_{t \geq \gamma_{N,p}} f(t) = \gamma_{N,p}^p - \gamma_1(\gamma_{N,p})^{p-1} - \gamma_2(\gamma_{N,p})^{p-2}$$

Conversely, if

$$\lambda \leq \gamma_{N,p}^p - \gamma_1(\gamma_{N,p})^{p-1} - \gamma_2(\gamma_{N,p})^{p-2}$$

then break of symmetry occurs.

Acknowledgements. The Author wishes to thank Prof. Roberta Musina for having suggested the references [15], [16], [17] and for her useful comments. the author also wishes to thank the anonymous referee for careful reading the manuscript and for his/her many valuable suggestions. This research was supported by the fund of Australian Research Council (ARC).

References

- [1] Alves, C. O. ; do Ó, João Marcos, Positive solutions of a fourth-order semilinear problem involving critical growth, *Adv. Nonlinear Stud.*, **2**, (2002), no 4, 437–458.
- [2] Bhakta, Mousomi; Musina, Roberta, Entire solutions for a class of variational problems involving the biharmonic operator and Rellich potentials, *Nonlinear Anal.* **75** (2012), no. 9, 3836–3848.
- [3] Caffarelli, L.; Kohn, R.; Nirenberg, L., First Order Interpolation Inequalities with Weights, *Compositio Math.* **53** (1984), 259–275.
- [4] Caldiroli, P., Radial and non radial ground states for a class of dilation invariant fourth order semilinear elliptic equations on \mathbb{R}^N , *Communications on Pure and Applied Analysis* , Volume 13, (2014), no 2, 811–821.
- [5] Caldiroli, P.; Musina, R., Caffarelli-Kohn-Nirenberg type inequalities for the weighted biharmonic operator in cones, *Milan J. Math.*, **79** (2011), no 2, 657–687.
- [6] Catrina, F.; Wang, Z.-Q., On the Caffarelli-Kohn-Nirenberg inequalities: sharp constants, existence (and nonexistence), and symmetry of extremal functions, *Comm. Pure Appl. Math.* **54** (2001), 229–258.
- [7] Coffman, Ch. D., On the structure of solutions to $\Delta^2 u = \lambda u$ which satisfy the clamped plate conditions on a right angle, *SIAM J. Math. Anal* **13** (1982), 746–757.
- [8] Davies, E. B; Hinz, A. M. Explicit constants for Rellich inequalities in $L^p(\Omega)$, *Math. Z.* **227**, (1998) 511–523.
- [9] Dolbeault, J.; Esteban, M.; Loss, M.; Tarantello, G. On the symmetry of extremals for the Caffarelli-Kohn-Nirenberg inequalities, *Adv. Nonlinear Stud.* **9**, (2009), 713–726.
- [10] Felli, V.; Schneider, M., Perturbation results of critical elliptic equations of Caffarelli-Kohn-Nirenberg type, *J. Diff. Eq.* **191** (2003), 121–142.
- [11] Gazzola, F.; Grunau, H. C., Radial entire solutions for supercritical biharmonic equations, *Math. Ann.* **334** (2006), 905–936.
- [12] Hulshof, J.; Van der Vorst, R.C.A.M., Asymptotic behaviour of ground states, *Proc. Amer. Math. Soc.* **124**, (1996), no. 8, 2423–2431.
- [13] Lieb, H. E., Loss, M., Analysis, Second Edition, *Graduate Studies in Mathematics* Vol 14, American Mathematical Society, 2001.
- [14] Lions, P. L., The concentration-compactness principle in the calculus of variations. The locally compact case. II. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1** (1984), 223–283.
- [15] Mitidieri, E., A simple approach to Hardy inequalities, *Mat. Zametki* **67**, (2000), no. 4, 563–572; translation in *Math. Notes* 67 (2000), no. 3-4, 479–486.
- [16] Musina, Roberta, Optimal Rellich-Sobolev constants and their extremals, *Differential Integral Equations* **27** (2014), no. 5-6, 579–600.
- [17] Musina, Roberta, Weighted Sobolev spaces of radially symmetric functions, *Ann. Mat. Pura Appl.*, (2013) arXiv:1206.6957.

- [18] Rellich, F., Halbbeschränkte Differentialoperatoren höherer Ordnung. In: J.C.H. Gerretsen, J. de Groot (Eds.): Proceedings of the International Congress of Mathematicians 1956, Volume III (pp. 243–250) Groningen: Noordhoff 1956.
- [19] Rellich, F., Perturbation theory of eigenvalue problems, *New York: Courant Institute of Mathematical Sciences, New York University*, (1954).