arXiv:1311.1103v1 [hep-ph] 4 Nov 2013

Charge exchange reaction at high energies

R. Schicker[1], L. Jenkovszky[2], O. Kuprash[3]

[1] *Phys. Inst., Heidelberg*
[2] *BITP, Kiev*
[3] *DESY, Hamburg*

Abstract

The charge exchange reaction pp $\rightarrow$ n Δ^{++} at high energies is examined. The cross section of this reaction is estimated for RHIC and LHC energies based on data taken at the lower energies of the Intersecting Storage Ring (ISR) at CERN. The interest of such measurements for identifying the associated electromagnetic bremsstrahlung radiation is discussed.

1 Introduction

Diffractive reaction channels contribute about 30% to the total hadronic cross section at the energies of the Large Hadron Collider LHC at CERN. A good understanding of hadronic diffraction is therefore necessary for a comprehensive understanding of proton-proton collisions. In single and double diffractive dissociation, one or both of the protons get diffractively excited, and the proton remnants are very much forward focussed[1]. In central diffraction, a hadronic system is formed at mid-rapidity by the fusion of two Pomerons. The above reactions can, however, also be initiated by the exchange of a photon or a Reggeon. Such non-diffractive exchanges represent a potential background in the analysis of diffraction at high energies. A good understanding of electromagnetic processes, as well as of Reggeon exchanges, is therefore mandatory. The analysis of reaction channels which are purely photon or Reggeon induced are therefore of interest for identifying possible background sources in diffraction.

2 Charge exchange reactions

The study of charge exchange mechanism is of interest for an improved understanding of purely Reggeon induced reactions. Reggeon trajectories are defined by bound states of $q\overline{q}$-pairs. The QCD content of these exchanges was studied in Ref.[2], and remains a challenge for theory. The exchange of a charged $q\overline{q}$-pair represents a sudden acceleration of electric charge, accompanied by electromagnetic bremsstrahlung radiation. The measurement of this radiation is of interest as a test of the theorem of Low for soft photon emission as discussed below.

A variety of final states is available in charge exchanges in proton-proton collisions. Charge exchange reactions can be due to the exchange of the pion-, ρ- or A_2-trajectory.

$$pp \to n + \Delta^{++} \to n + p\pi^+ \tag{1}$$
$$pp \to \Delta^0 + \Delta^{++} \to n\pi^0 + p\pi^+ \tag{2}$$
$$pp \to \Delta^0 + \Delta^{++} \to p\pi^- + p\pi^+ \tag{3}$$

The simplest approach for studying these charge exchange reactions is in binary kinematics.




Figure 1: Charge exchange reaction in binary kinematics.

The relevant scattering amplitude at high energies, with a single Regge exchange (pion, ρ, A_2) is

$$A^{\pm}(s,t) = A_i \xi^{\pm}(t)\beta(t)(s/s_i)^{\alpha(t)}, \tag{4}$$

with $A^{\pm}$ the amplitude for l=even and l=odd exchange, respectively. The variables $s = (p_1 + p_2)^2 = (p_3 + p_4)^2$, $t = (p_1 - p_3)^2 = (p_4 + p_2)^2$ are the usual Mandelstam invariants, $\xi^{\pm}(t)$ is the signature factor ($\xi^{\pm}(t) = 1 \pm e^{-i\pi\alpha(t)}$), and $\beta(t)$ is the residue function taken in exponential form $\beta(t) = e^{b_i t}$, where b_i is fitted to each reaction, $pp \to n\Delta^{++}$ or $pp \to \Delta^0\Delta^{++}$. In this type of reactions, the dominant exchange is the pion trajectory at low energies with the ρ-trajectory dominating at high energies. We take a linear form $\alpha(t) = 0.0 + 0.8t$ and $\alpha(t) = 0.5 + 0.9t$ for the pion and ρ-exchange, respectively[3].

Apart from the overall normalization parameter and the standard Regge scale factor s_i, that we set equal to 1 GeV^2, the model contains only one free parameter, namely b that we find from comparison with measured data.

With the a priory unknown normalization

$$\frac{d\sigma}{dt} = \frac{\pi}{s^2}|A(s,t)|^2 \tag{5}$$

one can calculate the cross section. By a comparison with the existing data and fits, e.g.[4], one finds the differential and integrated cross section.

Measurements of the reaction shown in eq.1. above were made at the Argonne National Zero Gradient Synchrotron[5], and at the Intersecting Storage Ring at CERN[6].

The analysis of the cross section of pp $\rightarrow$ nΔ^{++} at momenta P_L up to 11 GeV/c finds good agreement with the Chew-Low one-pion-exchange mechanism taking into account form-factor models[7, 8, 9].

The analysis of the ISR data identifies the ρ- and A_2-contribution which start to dominate the cross section at energies $\sqrt{s} = 31$ GeV[10].

In order to evaluate the cross section at RHIC and LHC energies, we take the ISR data at $\sqrt{s} = 31$, 45 and 53 GeV and fit the data as outlined above.

Table 1: Cross section pp $\rightarrow$ nΔ^{++}

	$\sqrt{s}$ (GeV)	σ (nb)
ISR	31	580 $\pm$90
	45	210$\pm$40
	53	170$\pm$40
RHIC	100	48.5$\pm$5.5
	200	12.2$\pm$1.3
LHC	7×10^3	(10.0$\pm$1.1) $\times10^{-3}$
	14×10^3	(2.4$\pm$0.3) $\times10^{-3}$

The cross sections extrapolated to the RHIC energies of $\sqrt{s} = 100$ and 200 GeV shown in Table 1 are 48.5 and 12.2 nb, respectively. The corresponding values for the LHC energies of $\sqrt{s} = 7$ and 14 TeV are 10.0 and 2.4 pb, respectively. These values can be used for further evaluating experimental prospects of measuring the charge exchange channel pp $\rightarrow$ nΔ^{++} at these collider energies.

3 Electromagnetic radiation in charge exchange reactions

Accelerated electric charge emits bremsstrahlung radiation[11]. The theorem of Low relates the radiative leading and next-to-leading order in photon energy of the bremsstrahlung amplitude to the corresponding non-radiative amplitude shown in Fig.1[12]. The photons emitted from the external lines result in a pole term in the radiative amplitude, and generate the leading k^{-1}-dependence in the photon energy spectrum. Emission from the internal lines gives rise to the next-to-leading constant term in the energy spectrum. The theorem of Low applies to photons in the soft limit, i.e. to photon energies which are smaller than any other momentum scale in the amplitude. The emission of soft photons in the high energy limit was studied in Ref.[13]. This study finds correction in the next-to-leading radiative amplitude due to the internal structure of the

external states. A generalization of the theorem of Low to the production of non-abelian gauge bosons and gravitons is presented in Ref.[14].

4 Summary and outlook

The cross section for the charge exchange reaction pp $\rightarrow$ nΔ^{++} is extrapolated from data at the ISR to the energies of RHIC and LHC. More refined calculations are under way, and will take into account the three body nature of the final state, including the non-resonant continuum at masses beyond the Δ-resonance. The interest for measuring the associated bremsstrahlung photons is outlined. Quantitative predictions for this associated radiation are under study.

5 Acknowledgments

We gratefully acknowledge fruitful discussions with L. Lipatov during the Crimean Conference on issues related to the subject of this paper. This work is partially supported by WP8 of the hadron physics program of the 8th EU program period.

References

[1] L. Jenkovszky et al., Mod.Phys.Lett. A**26**, (2011) 2029.

[2] R. Kirschner, L.N. Lipatov and L. Szymanowski, Nucl. Phys. B **425** (1994) 579; Phys. Rev. D **51** (1995) 838.

[3] A.B. Kaidalov and B.M. Karnakov, Sov. J. Nucl. Phys. **7**, Number 5, (1968) 685.

[4] K.G. Boreskov et al., Sov. J. Nucl. Phys. **15**, Number 2, (1972) 203.

[5] J.D. Mountz et al., Phys. Rev. D **12**, (1975) 1211.

[6] H. De Kerret et al., Phys. Lett. B **69**, (1977) 372.

[7] G.F. Chew and F.E. Low, Phys. Rev. **113** (1959) 1640.

[8] H.P. Dürr and H. Pilkuhn, Nuovo Cimento **40A** (1965) 899.

[9] J. Benecke and H.P. Dürr, Nuovo Cimento **56A** (1968) 269.

[10] K.J.M. Moriarty and H. Navelet, Phys. Lett. B **71** (1977) 208.

[11] J.D. Jackson, Classical Electrodynamics, John Wiley & Sons.

[12] F.E. Low, Phys. Rev. **110** (1958) 974.

[13] V. Del Duca, Nucl. Phys. B **345** (1990) 369.

[14] L.N. Lipatov, Nucl. Phys. B **307** (1988) 705.