

Jet Quenching and Its Azimuthal Anisotropy in AA and possibly High Multiplicity pA and dA Collisions

Xilin Zhang^{1,*} and Jinfeng Liao^{2,3,†}

¹*Institute of Nuclear and Particle Physics and Department of Physics and Astronomy,
Ohio University, Athens, OH 45701, USA.*

²*Physics Department and Center for Exploration of Energy and Matter,
Indiana University, 2401 N Milo B. Sampson Lane, Bloomington, IN 47408, USA.*

³*RIKEN BNL Research Center, Bldg. 510A,
Brookhaven National Laboratory, Upton, NY 11973, USA.*

We report our calculation of jet quenching and its azimuthal anisotropy in the high energy AA and high multiplicity pA and dA collisions. The purpose of this study is twofold. First, we improve our previous event-by-event studies, by properly implementing p_t dependence in the modeling. We show that, within the jet “monography” scenario featuring a strong near-Tc-enhancement of jet energy loss, the computed high- p_t nuclear modification factor R_{AA} and the harmonic coefficients in its azimuthal anisotropy $V_{2,3,4}$, agree well with the available data from both Relativistic Heavy Ion Collider (RHIC) and Large Hadron Collider (LHC). Second, in light of current discussions on possible final state collective behavior in the high-multiplicity pPb and dAu collisions, we examine the implication of final state jet attenuation in such collisions, by applying the same model used in AA collisions and quantifying the corresponding R_{pA} and R_{dA} and their azimuthal anisotropy. The high- p_t V_n are an set of clean indicators of final state energy loss. In particular we find in the most central pPb (5.02 TeV) and dAu (200 GeV) collisions, V_2 is on the order of 0.01 and 0.1 respectively, measurable with current experimental accuracy. In addition, our high- p_t R_{dA} is around 0.6 which is compatible with preliminary dAu results from RHIC.

PACS numbers: 25.75.-q, 12.38.Mh

I. INTRODUCTION

In the high energy heavy ion collisions, a new state of strongly interacting matter, the quark-gluon plasma (QGP) is created. Measuring the properties of this matter is of fundamental interest. Such measurements are now done at a variety of collisional beam energies at both RHIC and LHC.

The highly energetic partons (which ultimately turn into observed jets) born from the very initial hard scatterings provide an invaluable way of probing the matter properties. Along the way penetrating the medium, a jet parton will interact with the medium and lose energy, which can be manifested through comparing the observables (such as leading hadron yield) in the jet kinematic region with the same observables measured from the reference pp collisions. One commonly used observable in jet quenching study is the nuclear modification factor, R_{AA} , defined as the ratio between the hadron production in AA collision and that in NN collision (further scaled by the expected binary collision number). This observable turns out to be a crucial quantity in the establishment of QGP: a significant suppression of high- p_t (transverse momentum) hadron production was first observed at RHIC and then at LHC [1], with R_{AA} reaching ~ 0.2 in the most central collisions. Measurements

of charged particle, identified hadron, heavy flavor, and photon production have coherently pointed to a medium that is extremely opaque to “colored” hard probe. For reviews see e.g. [2].

Another important aspect of jet-quenching analysis, the so-called geometric tomography [3, 4], is about the azimuthal anisotropy of jet energy loss. The geometry of the hot medium created in AA collision event is generally anisotropic in the transverse plane (almond-like), and so is the high-energy parton energy loss moving along different azimuthal angle (ϕ), which leads to a measurable anisotropy in the $R_{AA}(\phi)$ [5–9]. More recently this idea has been extended to systematically quantify the jet response to the initial state fluctuations giving rise to various other harmonic coefficients in the high- p_t region [10–16].

Detailed measurements of jet quenching and its azimuthal anisotropy via leading hadrons have been done over the past decade, with large amounts of data about how R_{AA} depends on hadron’s p_t and azimuthal angle ϕ , the collision centrality class, and the beam energy. It is essential to describe all these characteristics within a unified jet quenching modeling scheme. The first goal of the present paper is to improve our previous studies [9, 14–16] on this subject, and to perform comprehensive computations of the mentioned dependences and compare the results with available data. The jet energy loss model we use here is based on the jet “monography” scenario featuring a strong near-Tc-enhancement (NTcE) of jet-medium interaction. This scenario [6] was first discovered in the effort to explain

*Electronic address: zhangx4@ohio.edu

†Electronic address: liaojji@indiana.edu

the large jet quenching anisotropy (à la V_2 at high p_t) at RHIC [7, 8, 17, 18], and is based on the “magnetic scenario” for sQGP [19–24]. More recently it was shown in Refs. [15, 16] that the NTcE model naturally induces a reduction ($\sim 30\%$) of jet-medium interaction strength from RHIC to LHC. Such reduction is required to explain both RHIC and LHC R_{AA} consistently according to the analysis in Ref. [25]. In our previous works, the NTcE model has been shown to well describe the centrality trend, the anisotropy, and the beam energy dependence of jet quenching. However we’ve used the fractional energy loss (largely motivated by the “flatness” seen at RHIC) leading to no p_t -dependence for the observables. In this study, we introduce the p_t dependence into our jet quenching modeling in a similar way as used in Refs. [11, 26]. As a result, we can compute the jet quenching observables in different p_t bins, which is needed to analyze the jet quenching data from LHC.

The second goal of this study is to examine possible final state jet-medium interaction in the high multiplicity pA and dA collision events. Such events have generated a great deal of interests recently. A di-hadron correlation pattern that is long-range in rapidity while collimated in azimuthal angle (also called as “ridge”) has been observed in high-multiplicity proton-proton (pp at $\sqrt{s_{NN}} = 7$ TeV) [27], proton-lead (pPb at $\sqrt{s_{NN}} = 5.02$ TeV) [28–32], and deuteron-gold collisions (dAu at $\sqrt{s_{NN}} = 200$ GeV) [33]. The same correlation has been regarded as an important indicator of the collective behavior of the matter created in AA collisions at RHIC [34] and LHC [35, 36]. Previous AA studies have concluded that due to the initial state geometry and its fluctuations, different harmonic flows are developed during matter’s expansion, which upon folding together create the observed “ridge” correlation [37–48]. The question still under debate, is whether the “ridge” in pp, pPb, and dAu collisions implies the formation of a thermal QGP in these small systems. Efforts to understand the origin of the “ridge” in these collisions are still ongoing [49]. The hydrodynamic calculations (assuming enough final state interaction to make a nearly thermal medium) have been carried out for pp [50, 51], pPb, and dPb collisions [52–54]. At the same time, there are alternative explanations based on possible intrinsic azimuthal correlations in the early gluon production [55, 56].

While it seems plausible that collective effect is negligible in the pp case, certain degrees of collectivity could be present in the dense partonic system created in the pA and dA collisions (which may be tentatively called “mini-bang”). To approve or disapprove this assertion, it is crucial to have independent probes in addition to the soft-sector observables (e.g. flows, soft “ridge”). We suggest to use the final state interaction induced jet energy loss as an indicator. In this paper, we will make a first analysis of jet quenching and its azimuthal anisotropy in these “mini-bangs” by applying our energy loss modeling (calibrated in AA collisions) and computing R_{pA} and R_{dA} , and their azimuthal anisotropy’s harmonic co-

efficients $V_{1,2,3}$. It is worth mentioning here that the computed R_{dA} compares favorably with recent dAu results [59–61]. In particular as we will show, the high- p_t anisotropy, not easily contaminated by initial state effects, can serve as a “clean” signal of final state jet-medium interaction. (Note there are recent theoretical works [57, 58] about jet quenching in the high multiplicity pp collisions.)

The rest of the paper is organized as follows. Sec. II is devoted to the details of the NTcE model improved by the implementation of p_t dependence. We use central R_{AA} to fix two model parameters, and then compute R_{AA} in different centrality classes and p_t bins for both RHIC and LHC collisions. In Sec. III, we extract the harmonic coefficients $V_{2,3,4}$ from R_{AA} ’s azimuthal anisotropy. In Secs. IV and V, we explore possible jet quenching and its anisotropy in the pPb and dAu collisions and make predictions for future measurements. Finally we summarize the results in Sec. VI.

II. JET QUENCHING IN AA COLLISIONS FROM RHIC TO LHC

In this work, we use the geometric model scheme [5–9, 11, 14–16, 25, 62], in which the geometric features of jet energy loss (e.g. the path-length dependence) are implemented in a phenomenological way. In this class of models, the differential energy loss (dE/dl) experienced by a jet parton through the medium is expressed as

$$dE = -E^\delta \kappa[s(l)] s(l) l^m dl. \quad (2.1)$$

Here $s(l)$ is the local entropy density along the jet path, and $\kappa(s)$ is the local jet quenching strength which as a property of matter should in principle depend on $s(l)$. There are different choices of m and $\kappa(s)$ [7, 8, 11, 25, 62]. We only focus on the mentioned NTcE model which assumes $m = 1$ based on the LPM effect in the radiative energy loss [63], and introduce a strong jet quenching component in the vicinity of T_c (with density s_c and span of s_w) via

$$\kappa(s) = \kappa[1 + \xi \exp(-(s - s_c)^2/s_w^2)]. \quad (2.2)$$

The parameters are $\xi = 6$, $s_c = 7/fm^3$, and $s_w = 2/fm^3$. As demonstrated previously in Refs. [6, 9, 15, 16], this model reproduces the high p_t V_2 at RHIC, and naturally explains the nontrivial reduction of the jet-medium interaction strength (in average) from RHIC to LHC.

In these previous works [6, 9, 15, 16], the fractional energy loss scenario [i.e., setting $\delta = 1$ in Eq. (2.1)] was used, motivated by the weak p_t -dependence of the RHIC data in the jet region. As a result the computed jet quenching observables are all p_t -independent. Such a model fails to describe the LHC R_{AA} data which show a strong p_t -dependence in a significantly larger jet p_t region. Here we improve the model by treating the δ as

a free parameter that needs to be fixed in order to reproduce R_{AA} 's p_t -dependence at both RHIC [64, 65] and LHC [66, 67].

Let's define the ratio between initial and final jet energy as $\mathcal{R}_{\mathbf{P},\mathbf{Ev}}$ in a given AA collision event \mathbf{Ev} for a particular jet path \mathbf{P} , i.e., $E_i = E_f \mathcal{R}_{\mathbf{P},\mathbf{Ev}}(E_f)$. Using Eq. (2.1), we can get ($\delta \neq 1$):

$$\mathcal{R}_{\mathbf{P},\mathbf{Ev}}(E_f) = \left[1 + \frac{(1-\delta)\mathcal{D}_{\mathbf{P},\mathbf{Ev}}}{E_f^{1-\delta}} \right]^{\frac{1}{1-\delta}},$$

$$\mathcal{D}_{\mathbf{P},\mathbf{Ev}} \equiv \int_{\mathbf{P},\mathbf{Ev}} \kappa[s(l)]s(l)l^m dl. \quad (2.3)$$

We focus on mid-rapidity measurements so that $E \approx p_t$. Suppose the reference pp spectrum at the same collision energy is $g_{pp}(p_t)$, based on Eq. (2.3) we can get the formula for the averaged R_{AA} over many events:

$$R_{AA}(p_t, \phi) = \frac{\langle\langle g_{pp}[p_t \mathcal{R}_{\mathbf{P},\mathbf{Ev}}(p_t)] [\mathcal{R}_{\mathbf{P},\mathbf{Ev}}(p_t)]^{1+\delta} \rangle_{\mathbf{P} > \mathbf{Ev}}}{g_{pp}(p_t)}. \quad (2.4)$$

Here, $\langle\langle \rangle_{\mathbf{P} > \mathbf{Ev}}$ means averaging all the possible jet paths from different jet spots in each event and then averaging all the events. The p_t spectrum of pp collision at 200 GeV is expressed as $g_{pp}(p_t) \propto 1/p_t^n$ in the high- p_t region. In Ref. [62], a p_t -dependent index n is introduced. This is supported by the pQCD calculation and also provides an explanation for R_{AA} 's weak p_t -dependence at RHIC (200 GeV). For the $\sqrt{s} = 2.76$ TeV and higher energy at LHC (e.g., $\sqrt{s} = 5.02$ TeV pPb collision), we use the scaling law [68, 69]: $\sqrt{s}^n g_{pp}(p_t) = p_0 [1 + x_T/p_1]^{p_2}$, while $x_T \equiv 2p_t/\sqrt{s}$. The relevant (energy independent) parameters p_1 and p_2 have been fitted to the measurement at $\sqrt{s} = 2.07$ TeV (see Ref. [66] for details).

To study R_{AA} 's centrality and p_t dependences (not azimuthal anisotropy), we use the optical Glauber model [70] to generate initial state density profile. The relation between the initial entropy density and participant density (and collision density) can be found in Ref. [16]. [At LHC (2.76 TeV), the entropy density is roughly doubled from RHIC (200 GeV) case [71].] The total NN cross sections, σ_{NN}^{inel} , at these energies are 42 and 62 mb [70, 72]. In addition, only the longitudinal Bjorken expansion (along the collision axis) [73] is included, while the transverse dynamics is frozen. To fix the parameters, δ and κ , in Eq. (2.1), we use the most central (0 – 5%) high- p_t R_{AA} data from RHIC and LHC (see the plots for central collision in Fig. 1). A χ^2 -analysis has determined the optimal parameters to be: $\delta \approx 0.19$ and $\kappa \approx 0.0065$. (In Eq. (2.1), energy unit is always GeV.) Then we calculate the R_{AA} at other centrality classes. The results for the two collision energies are collected in Fig. 1. The shown data are from Refs. [64–67]. As one can see, the present model describes very well the R_{AA} in different centralities and p_t bins. The fact that the NTcE model

gives correct medium opaqueness evolution from RHIC to LHC is nontrivial, as already emphasized in our previous analysis [15, 16]. In this model, the RHIC fireball with its bulk matter staying closer and longer in the near- T_c region is naturally more opaque than the LHC fireball, and we've quantified the reduction of the jet-medium interaction strength to be $\sim 30\%$ in this opaqueness evolution [16]. Such a reduction has been consistently reported from other studies [25, 74, 75], where either a reduction of jet-medium coupling strength is put in by hand when extrapolating to LHC or a strong running coupling effect is introduced.

III. HIGH p_t AZIMUTHAL ANISOTROPY IN AA COLLISIONS FROM RHIC TO LHC

To study R_{AA} 's anisotropy, we use the Monte-Carlo (MC) Glauber model [70] to simulate the fluctuating initial conditions, and compute the azimuthal-angle ϕ -dependent $R_{AA}(\phi, p_t)$ on an event-by-event basis. Except the initial state, all the other ingredients in this calculation are the same as those in the previous section. The details of implementing the MC-Glauber and a careful analysis of the initial state geometry (e.g., ϵ_n) can be found in Ref. [16]. The medium is again assumed to experience only longitudinal Bjorken expansion [73]. One though may take caution that the transverse expansion in general will reduce the anisotropy of the matter density distribution and thus reduce the hard probe's responses to such anisotropy, as shown in e.g. [74, 76]. A full modeling needs to combine the jet energy loss calculation with a realistic hydrodynamic modeling, which we are currently working on with the results to be reported in future.

Due to the anisotropic and fluctuating matter geometry, the jet energy loss is different in each transverse direction, and for each event there will be an event-wise, azimuthal-angle ϕ -dependent nuclear modification factor $R_{AA}(p_t, \phi)$. Its anisotropy could be quantified by the harmonic coefficients $V_n(p_t)$ from a Fourier decomposition:

$$R_{AA}(\phi, p_t) \equiv R_{AA}(p_t) \times \left(1 + 2 \sum_n V_n(p_t) \cos[n(\phi - \psi_n^J(p_t))] \right). \quad (3.1)$$

Here we extract V_2 , V_3 and V_4 in each event and the corresponding reference angles ψ_2^J , ψ_3^J , and ψ_4^J in different p_t bins. The averaged $V_{2,3,4}$ can be directly compared with experimental data.

Let us first examine the second harmonics, $V_2(p_t)$, at different centralities in Fig. 2. The shown $V_2(p_t)$ is the average value of all the event's V_2 at a given p_t bin. The data are from Refs. [17, 18, 77]. Strictly speaking what the experiments measure would be most close to the V_2 relative to ψ_2^{EP} (the event-plane of final soft hadrons), i.e. $V_2 \cos[2(\psi_2^J - \psi_2^{\text{EP}})]$. Our study shows that ψ_2^J and




FIG. 1: R_{AA} vs p_t in different centrality classes of AA collisions at both RHIC and LHC. The data are from Refs. [64–67].




FIG. 2: V_2 vs p_t in different centrality classes of AA collisions at RHIC and LHC. The data are from Refs. [17, 18, 77]. In the upper panel, “STAR a” and “STAR b” are data based on different analysis methods [17].

ψ_2^{PP} are well correlated: $\langle \cos[2(\psi_2^J - \psi_2^{\text{PP}})] \rangle$ is around 1 in non-central collisions (see details in Ref. [16]) and has a very weak p_t dependence. In Ref. [78], ψ_2^{EP} and ψ_2^{PP} are also shown to be strongly correlated. Because of the two strong correlations, the correction due to the mismatch among the different planes would be minor. It is thus sensible to compare our V_2 with the experimental data for the non-central collisions. Also the impact of medium’s transverse dynamics in the non-central collisions should be smaller than that in central ones because of the shorter jet path in average. We can see the NTcE

model gives correct magnitude and p_t -dependence for V_2 across different centralities at two collision energies. Similar to R_{AA} , the V_2 at RHIC shows a weak p_t -dependence, while this dependence is rather strong at LHC. One does notice that for the LHC case, our model results are consistently higher than data: a plausible explanation could be the omitted transverse expansion effect which would reduce the anisotropy to certain extent and which is expected to be stronger at LHC.

Let us move to the higher harmonics arising from fluctuations. The results of $V_3(p_t)$ and $V_4(p_t)$ at LHC are




FIG. 3: V_3 vs p_t (upper panel) and V_4 vs p_t (lower panel) in different centrality classes of AA collisions at LHC. Two different results are shown here. “NTcE” gives the harmonic component V_n , while “NTcE, pp” is the V_n projected to the corresponding participant planes. See the discussion in text. The data are from Ref. [79].

plotted in Fig. 3 in comparison with the available data from CMS [79]. Two different results are shown in the plots: the “NTcE” results correspond to the averaged V_n ($n = 3, 4$) from Eq.(3.1), while the “NTcE, pp” results are the averaged $V_n \cos[n(\psi_n^J - \psi_n^{PP})]$ ($n = 3, 4$) which account for the mismatch between the initial participant plane angle and the final jet response plane angle. According to Ref. [78], ψ_3^{EP} and ψ_3^{PP} , similar to ψ_2^{EP} and ψ_2^{PP} , correlate quite well in all the centrality classes, but the correlation between ψ_4^{EP} and ψ_4^{PP} is complicated and even changes the sign. Meanwhile by comparing “NTcE” and “NTcE, pp”, we can see the correlation between ψ_3^J and ψ_3^{PP} is also stronger than that between ψ_4^J and ψ_4^{PP} . Keeping in mind the difference between initial participant plane and the final event plane angles, it is nevertheless interesting to compare our results with data, and one finds that the calculated “NTcE, pp” results are in semi-quantitative agreement with the data (with large error bars). In particular for the V_3 our model results are slightly higher than data, a discrepancy that would again be expected due to the omitted transverse expansion. Our future study with a realistic hydrodynamic medium will draw a more definitive answer.

As first pointed out and discussed in details in Refs. [14, 16], a hard-soft dihadron correlation would naturally arise due to the hard sector response (via energy loss) and the soft sector response (via collective expansion)

to the common initial geometric anisotropy:

$$\frac{\int \frac{d\phi_1}{2\pi} \frac{d\phi_2}{2\pi} 2\pi\delta(\phi_2 - \phi_1 - \Delta\phi) < \frac{dN^{hard}}{dyd\phi_1} \frac{dN^{soft}}{dyd\phi_2} >}{< \int \frac{d\phi_1}{2\pi} \frac{dN^{hard}}{dyd\phi_1} > < \int \frac{d\phi_2}{2\pi} \frac{dN^{soft}}{dyd\phi_2} >}, \\ \sim 1 + 2 \sum_{n=1,2,3,\dots} < V_n^s V_n^h > \cos(n\Delta\phi) \quad (3.2)$$

Properly combining the high- p_t V_n from energy loss computations and the low- p_t V_n computed in hydro calculations (e.g., [78]), a significant near-side peak is born out in azimuthal angle, while on the away side a double-hump structure could emerge or not depending on the details of the hierarchy in the V_n vs n spectrum especially with $n = 1, 2$, and 3 [14, 16].

IV. POSSIBLE JET QUENCHING IN HIGH-MULTIPLICITY pPb COLLISIONS AT LHC

As briefly discussed in Sec. I, the nature of high multiplicity events in the LHC pPb collisions (e.g., LHC at $\sqrt{s_{NN}} = 5.02$ TeV) is currently under intensive discussions [52, 80–83]: in particular the focal point is whether significant final state interactions occur and whether a collective bulk medium (close to that created in AA collisions) is ever formed in such events. In Refs. [52–54], the authors have applied hydrodynamic calculations to study the collective expansion of the created matter, assuming its validity for this small system. Their results show sizable elliptical flow which so far is consistent with the available data [31, 53]. While it is conceivable that due to the high parton density a certain degree of collective explosion would develop, how such a small system

with a very short lifetime could thermalize to the extent of justifying hydrodynamics is quite puzzling, especially provided the thermalization process is extremely complicated even for the larger and longer AA fireball [84]. There are also various alternative explanations of measured data based on mostly initial state effects [81–83].

In the present situation, it is of particular interest to find observables other than the soft sector collective expansion that would clearly distinguish the initial state versus final state effects. We here suggest that the hard probe may provide such an opportunity. An extremely dense and (nearly) thermal partonic medium, if indeed formed in high-multiplicity pPb events, will inevitably induce certain amount of final state attenuation to a high energy jet parton traversing the medium. Therefore any measurable effect uniquely from jet energy loss would be a very useful indicator of jet-medium final state interactions. In this section, with the assumption of such a medium in pPb collisions, we explore how large the effects of jet energy loss could be and what would be the best way to detect them, by applying the same NTcE model used in the corresponding AA study.

In this calculation, MC calculation is needed, because of the large initial state fluctuations [52]. In Ref. [85], the authors have shown that different implementations of multiplicity generation in the initial nucleon-nucleon scattering, including Glauber and IP-Glassma models [86, 87], give different initial state eccentricities and different final state elliptical flow. We choose two different energy deposit scenarios in the framework of the MC-Glauber model to generate the initial state [52, 53]: the first assumes a linear relation between entropy density and initial participant density (labeled as “size a”); the second uses collision density instead (labeled as “size b”). In general the spacial “size a” is bigger than “size b”. The difference between the two jet quenching results can be used to estimate the uncertainty of our calculations due to the initial state implementation. This is complementary to the study in Ref. [85]. In Ref. [52], the ratio between entropy and participant density, denoted as s_0 in the following, is estimated to be around 90/fm for both pPb (4.4 TeV) and dPb (3.11 TeV) collisions. In a crude approximation, we use this value in our pPb (5.02 TeV) calculation.

The NN cross section is 68 mb at $\sqrt{s_{NN}} = 5.02$ TeV [53]. The distribution of the participant number N_p has been checked against the one in Ref. [52], based on which we define the most central (i.e. high-multiplicity) events (0% – 4% centrality) as those with $N_p \geq 18$. All the following results are shown for this centrality. Moreover since the scattering is not symmetrical with respect to the two colliding beams, the natural “mid-rapidity” has been properly shifted [88]. Although there is no ideal rapidity “plateau” in the collision’s multiplicity distribution, the data from Ref. [88] indicates that such dependence is weak. We will assume the boost-invariant longitudinal expansion of the medium: this assumption should be no worse for pPb than for PbPb as in this smaller




FIG. 4: R_{pA} vs p_t in the most central pPb collisions. The “size a” and “size b” calculations use different initial state implementation. See discussion in the text.

and shorter-life-time system the transverse expansion is less significant and also the jet’s path length would be much shorter. Finally we use the scaling law formula for the pp reference spectrum in this calculation (see related discussion in Sec. II).

In Fig. 4, the computed R_{pA} (defined in the same way as R_{AA}), based on “size a” and “size b” scenarios, are plotted against the transverse momentum. We see that the jet quenching effect is not negligible (particularly in the region below ~ 20 GeV), and is quite close to the $R_{AA}(p_t)$ in the peripheral collisions at LHC (see Fig. 1 for the 30 – 50% centrality R_{AA} at LHC). Two important points need to be emphasized here if one were to compare this R_{pA} directly with the measurements: 1) This suppression is the effect solely from the assumed final state attenuation—there are however possible initial state effects [e.g. from cold nuclear matter (CNM) on Pb side] that may compensate the suppression of R_{pA} ; 2) Moreover the pp reference spectrum used here is for the minimum-bias events while it is likely that for the high multiplicity events the actual initial spectrum from hard collisions could differ considerably from the one used there—most likely being softer thus reducing the suppression effect. Note that the recently measured R_{pA} by ALICE [32] does not show obvious suppression in minimum-bias events, but we focus on the high multiplicity ones, the nature of which however is still unsettled.

Let us now turn to jet quenching azimuthal anisotropy. Different from the R_{pA} ’s interpretation being complicated, the azimuthal anisotropy should be considered as the “clean” indicator of the final state interaction effect. Fig. 5 shows first three harmonics based on “size a” and “b” initial states. The results already account for the angular difference between ψ_n^J and ψ_n^{PP} . Although the V_2 calculated with “a” and “b” initial states can differ by as large as $\sim 30\%$ (at $p_t = 20$ GeV), our results do indicate a significant $V_2 \sim 0.01$ in high p_t region—this appears




FIG. 5: $V_{1,2,3}$ vs p_t in the most central pPb collisions. Similar to Fig. 4, there are two results (“a” and “b”) based on different initial state implementations. V_1 and V_3 have been rescaled to fit in the plot. Here V_n s are the harmonic components projected to the corresponding participant planes. See the discussion in the text.

encouraging enough to be measurable with current luminosity and accuracy at LHC. [One though should be cautious about applying the current framework to address jets at very high (initial) energy (e.g., 100 GeV) in pPb collisions, because such jets take a major percentage of the system’s total energy [31].] Meanwhile, V_1 and V_3 are much less prominent. Recently, ATLAS has measured V_2 with p_t up to 4 GeV [31]. In the collisions with highest total transverse energy (i.e. highest multiplicity), $V_2 \sim 0.1$ at $p_t = 4$ GeV (based on the two-particle correlation). It is very important and certainly feasible to extend such measurements to higher p_t e.g. ~ 10 GeV. A measured anisotropy at this magnitude in this high- p_t region would be an unambiguous signal of final state jet-medium interaction and thus also a signal of a dense partonic medium in this “mini-bang”.

In addition, similar to the AA situation, combining nonzero high- and low- p_t V_n may produce a hard-soft correlation in pA collisions [see formula (3.2)]. This correlation can be studied by embedding the jet quenching modeling into the hydrodynamical calculation under current assumption about the medium. On the experimental side, the corresponding soft-soft correlation in pA has been measured and used to extract the harmonic flows for the soft sector, and it would be of great interest to see the measurements of hard-trigger soft-associate azimuthal correlations in the future.

V. POSSIBLE JET QUENCHING IN HIGH-MULTIPLICITY dAu COLLISIONS AT RHIC

Motivated by recent observation of the long range correlation in dAu collision (200 GeV) [33], we investigate possible jet quenching in these collisions also. The MC-Glauber simulation follows the study of dPb collision (3.11 TeV) in Ref. [52]. Again we use “size a” and “b” energy deposit scenarios (see Sec. IV), and the NTcE model to compute jet energy loss. The total NN cross




FIG. 6: R_{dA} vs p_t in dAu collisions. Two different centrality classes, 0 – 5% and 0 – 20%, are shown. “size a” and “size b” calculations use different initial state implementations. The data are from PHENIX: (green) squares are π^0 ’s R_{dA} measurement [59] and (blue) circles are π^0 ’s R_{cp} (0 – 20%/60 – 88%) measurements [60, 61]. Besides the uncertainties shown in the plot, the former has a pp reference normalization uncertainty (9.7%), and the latter has a collision number uncertainty (8%). See discussions in the text.

section, the p_t spectrum in NN collision, and the nucleon density in Au nucleus can be found in the AuAu study in Sec. II. To fix the entropy density to participant density ratio s_0 for the dAu collision, we rely on the mid-rapidity multiplicity per participant measurements at RHIC [89] and LHC [88]: $dN_{ch}/(d\eta N_p) = 1.16$ and 2.14. Based on this, we infer the ratio between s_0 in dAu (200 GeV) and in dPb (5.02 TeV) to be $1.16/2.14 \sim 0.54$. We then approximate s_0 in dPb (5.02 TeV) to be around 90/fm [52], and get $s_0 = 49/\text{fm}$ in the dAu collision.

In Fig. 6, we plot the nuclear modification factor R_{dA} ($p_t \geq 5$ GeV) in two different centrality classes, 0 – 5% with $N_p \geq 19$ and 0 – 20% with $N_p \geq 14$. Recent mea-




FIG. 7: $V_{1,2,3}$ vs p_t in dAu collisions. Similar to Fig. 4, there are two types of initial state implementations, “size a” and “size b”, and two centrality classes, 0–5% and 0–20%. V_1 and V_3 have been rescaled to fit in the plot. Here V_n s are the harmonic components projected to the corresponding participant planes. See discussions in the text.

measurements from PHENIX are also shown there, including π^0 ’s R_{dA} in the 0–20% centrality class [59] and the ratio (R_{cp}) between the 0–20% R_{dA} and the 60–88% one [60, 61]. Note that the CNM effects mentioned in Sec. IV also exist in the dAu jet production. Indeed, the measurement [60, 61] reveals that 60–88% R_{dA} rises to about 1.3 in the high- p_t region, which is consistent with the CNM effects. Here we expect that the CNM effects’ impact on R_{cp} is small and the final state jet attenuation in 60–88% R_{dA} is also negligible. As a result, comparing our result with R_{cp} data is more informative than with R_{dA} (0–20%) data. We can see our result agrees reasonably well with the R_{cp} data, although is somewhat lower than the R_{dA} data.

In addition, it is interesting to note that in the $5 \leq p_t \leq 20$ GeV region and 0–5% centrality class, the R_{dA} is smaller than the R_{pA} shown in Fig. 4. The differences between the two collisions include: 1) with the impact parameters set as zero, we find that the averaged entropy densities in the center of the cluster(s) are around 100 and 300 fm^{-3} for dAu and pPb respectively; 2) The medium’s spacial size in the dAu collision is larger than in the pPb collision; 3) The pp collision’s p_t spectrum decreases steeper at 200 GeV than at 5.02 TeV; 4) The NTcE effect is reduced from the dAu to the pPb collision. The competition between the first factor and the latter three gives rise to a smaller R_{dA} than R_{pA} .

As emphasized in Sec. IV, high- p_t azimuthal anisotropy observables are important to determine the final state jet attenuation effect. In Fig. 7, we plot $V_{1,2,3}$ against p_t ($p_t \geq 5$ GeV) in the two different centralities. We see a large $V_2 \sim 0.1, 0.075$ in both two centralities; they are close to the V_2 in non-central AuAu collisions (see Fig. 2). This should encourage the corresponding measurements. For V_1 and V_3 , “size a” and “size b” results can differ by more than 100%. However, the dominance of V_2 in high- p_t azimuthal anisotropy is clear, which reflects the underlying medium’s dipole geometry. In addition, the hard-soft correlation for the dAu collision follows the one for the pPb case (see Sec. IV); thus it is not repeated here.

VI. SUMMARY

In this paper, we have studied the jet quenching and its azimuthal anisotropy both in AA and high multiplicity pA and dA collisions. We focus on the azimuthal-angle dependent nuclear modification factors in the high- p_t region and investigate how the observable depends on the p_t , centrality, collisional beam energies, as well as the different collision systems (AA versus pA and dA).

For the AA collisions, we have improved our previous NTcE model studies [14–16], by implementing proper p_t dependence for the mentioned observables [see Eq. (2.1)]. The calculated $R_{AA}(p_t)$ at RHIC ($5 \leq p_t \leq 20$ GeV) and LHC ($10 \leq p_t \leq 100$ GeV) agrees very well with the data for different centrality classes, in both the p_t dependence and the overall suppression magnitude from 0.2 to 2.76 TeV collisions. These studies further strengthen the case (as we have illustrated previously) that the NTcE model naturally provides a dynamical reduction of the average opaqueness of the created hot matter from RHIC to LHC.

We have further quantified the harmonic coefficients of the ϕ –dependence of R_{AA} in AA collisions. Our results of the second harmonic, V_2 , also agree well with the available data at both RHIC ($5 \leq p_t \leq 20$ GeV) and LHC ($5 \leq p_t \leq 50$ GeV) region. Let us add that, a simultaneous description of both jet quenching and its anisotropy for both RHIC and LHC energies, has been quite a challenge for many other jet quenching modelings [7, 15, 16, 90]. Our results for higher order harmonics, V_3 and V_4 , at LHC also show a fair agreement with the CMS measurements (taking into account the data’s large error bars). It does not escape our attention that the high- p_t V_2 and V_3 results for LHC AA collisions show the consistent pattern of being slightly larger than data, which may be attributed to the two factors, i.e., the bulk medium transverse expansion not accounted for in the present modeling, and the mismatch between initial participant planes and the final event planes of the bulk geometry. To fully address these issues it is necessary to

combine our energy loss modeling with realistic hydrodynamic medium, which is currently being pursued.

Finally, motivated by currently intensive discussions on possibly significant final state interactions in the high multiplicity pPb events at LHC and dAu events at RHIC, we apply our NTcE model to explore possible jet quenching effect in these collisions. We have studied two different initial state implementations in MC-Glauber model. Our R_{pA} and R_{dA} results provide the first quantification of the ballpark for the magnitude of solely final state suppression effect in such collisions. It is worth pointing out that our R_{dA} at high- p_t is consistent with the recent measurements at RHIC. We also quantify the high- p_t anisotropy via the harmonic coefficients $V_{1,2,3}$, and suggest these to be a clean signal of possible final state jet-medium interaction. In particular our calculated V_2

is on the order of 0.01 for the pPb collisions, and 0.1 for the dAu collisions, which should be readily measurable and which, if indeed measured, would be an unambiguous signal of final state jet attenuation in such collisions.

Acknowledgements

X.Z. would like to acknowledge the support from US Department of Energy under grant DE-FG02-93ER-40756. J.L. thank the RIKEN BNL Research Center for partial support. We also thank A. M. Sickles for pointing out the dAu measurements and helpful discussions, and B.G. Zakharov for mentioning his recent work on jet quenching in pp collisions.

-
- [1] K. Aamodt *et al.* [ALICE Collaboration], Phys. Lett. B **696**, 30 (2011). S. Chatrchyan *et al.* [CMS Collaboration], Eur. Phys. J. C **72**, 1945 (2012).
 - [2] M. Gyulassy, I. Vitev, X. N. Wang and B. W. Zhang, arXiv:nucl-th/0302077. P. Jacobs and X. -N. Wang, Prog. Part. Nucl. Phys. **54**, 443 (2005). R. J. Fries and C. Nonaka, Prog. Part. Nucl. Phys. **66**, 607 (2011). J. Casalderrey-Solana, H. Liu, D. Mateos, K. Rajagopal and U. A. Wiedemann, arXiv:1101.0618 [hep-th].
 - [3] M. Gyulassy, I. Vitev and X. N. Wang, Phys. Rev. Lett. **86**, 2537 (2001).
 - [4] X. N. Wang, Phys. Rev. C **63**, 054902 (2001).
 - [5] E. V. Shuryak, Phys. Rev. C **66**, 027902 (2002). A. Drees, H. Feng, J. Jia, Phys. Rev. C **71**, 034909 (2005).
 - [6] J. Liao, E. Shuryak, Phys. Rev. Lett. **102**, 202302 (2009).
 - [7] J. Jia, R. Wei, Phys. Rev. C **82**, 024902 (2010).
 - [8] J. Jia, W. A. Horowitz and J. Liao, Phys. Rev. C **84**, 034904 (2011) [arXiv:1101.0290 [nucl-th]].
 - [9] J. Liao, arXiv:1109.0271 [nucl-th].
 - [10] J. Jia, arXiv:1203.3265 [nucl-th].
 - [11] B. Betz, M. Gyulassy and G. Torrieri, Phys. Rev. C **84**, 024913 (2011) [arXiv:1102.5416 [nucl-th]]. B. Betz, M. Gyulassy, G. Torrieri, arXiv:1106.4564 [nucl-th].
 - [12] S. A. Bass, C. Gale, A. Majumder, C. Nonaka, G. -Y. Qin, T. Renk and J. Ruppert, Phys. Rev. C **79**, 024901 (2009) [arXiv:0808.0908 [nucl-th]].
 - [13] R. Rodriguez, R. J. Fries and E. Ramirez, Phys. Lett. B **693**, 108 (2010). R. J. Fries and R. Rodriguez, Nucl. Phys. A **855**, 424 (2011).
 - [14] X. -L. Zhang and J. -F. Liao, Phys. Lett. B **713**, 35 (2012) [arXiv:1202.1047 [nucl-th]].
 - [15] X. Zhang and J. Liao, arXiv:1208.6361 [nucl-th].
 - [16] X. Zhang and J. Liao, Phys. Rev. C **87**, 044910 (2013) arXiv:1210.1245 [nucl-th].
 - [17] J. Adams *et al.*, [STAR Collaboration], Phys. Rev. C **72**, 014904 (2005).
 - [18] A. Adare *et al.*, [PHENIX Collaboration], Phys. Rev. Lett. **105**, 142301 (2010).
 - [19] J. Liao, E. Shuryak, Phys. Rev. C **75**, 054907 (2007); Phys. Rev. Lett. **101**, 162302 (2008); arXiv:1206.3989 [hep-ph].
 - [20] J. Liao, E. Shuryak, Phys. Rev. D **82**, 094007 (2010); Phys. Rev. C **77**, 064905 (2008); Nucl. Phys. A **775**, 224-234 (2006).
 - [21] M. N. Chernodub and V. I. Zakharov, Phys. Rev. Lett. **98**, 082002 (2007).
 - [22] C. Ratti and E. Shuryak, Phys. Rev. D **80**, 034004 (2009).
 - [23] A. D'Alessandro, M. D'Elia and E. V. Shuryak, Phys. Rev. D **81**, 094501 (2010). A. D'Alessandro and M. D'Elia, Nucl. Phys. B **799**, 241 (2008).
 - [24] V. G. Bornyakov, A. G. Kononenko and A. G. Kononenko, arXiv:1111.0169 [hep-lat]. V. G. Bornyakov and V. V. Braguta, Phys. Rev. D **85**, 014502 (2012).
 - [25] B. Betz and M. Gyulassy, Phys. Rev. C **86**, 024903 (2012) [arXiv:1201.0281 [nucl-th]].
 - [26] W. A. Horowitz, M. Gyulassy, [arXiv:1104.4958 [hep-ph]].
 - [27] V. Khachatryan *et al.* [CMS Collaboration], JHEP **1009**, 091 (2010) [arXiv:1009.4122 [hep-ex]].
 - [28] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **718**, 795 (2013) [arXiv:1210.5482 [nucl-ex]].
 - [29] B. Abelev *et al.* [ALICE Collaboration], Phys. Lett. B **719**, 29 (2013) [arXiv:1212.2001 [nucl-ex]].
 - [30] G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. Lett. **110**, 182302 (2013) [arXiv:1212.5198 [hep-ex]].
 - [31] G. Aad *et al.* [ATLAS Collaboration], arXiv:1303.2084 [hep-ex].
 - [32] B. Abelev *et al.* [ALICE Collaboration], arXiv:1210.4520 [nucl-ex].
 - [33] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **111**, 212301 (2013) [arXiv:1303.1794 [nucl-ex]].
 - [34] B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **80**, 064912 (2009) [arXiv:0909.0191 [nucl-ex]]; J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **95**, 152301 (2005) [nucl-ex/0501016]; B. Alver *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **104**, 062301 (2010) [arXiv:0903.2811 [nucl-ex]].
 - [35] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1107**, 076 (2011) [arXiv:1105.2438 [nucl-ex]]; S. Chatrchyan *et al.* [CMS Collaboration], Phys. Rev. Lett. **109**, 022301 (2012) [arXiv:1204.1850 [nucl-ex]].
 - [36] G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. C **86**, 014907 (2012) [arXiv:1203.3087 [hep-ex]].

- [37] B. Alver and G. Roland, Phys. Rev. C **81**, 054905 (2010) [Erratum-ibid. C **82**, 039903 (2010)] [arXiv:1003.0194 [nucl-th]].
- [38] B. H. Alver, C. Gombeaud, M. Luzum and J. -Y. Ollitrault, Phys. Rev. C **82**, 034913 (2010) [arXiv:1007.5469 [nucl-th]]. M. Luzum, Phys. Lett. B **696**, 499 (2011) [arXiv:1011.5773 [nucl-th]]. R. S. Bhalerao, M. Luzum and J. -Y. Ollitrault, Phys. Rev. C **84**, 034910 (2011) [arXiv:1104.4740 [nucl-th]].
- [39] M. Luzum, J. Phys. G **38**, 124026 (2011) [arXiv:1107.0592 [nucl-th]].
- [40] P. Sorensen, arXiv:0811.2959 [nucl-ex]. A. Mocsy and P. Sorensen, arXiv:1008.3381 [hep-ph]. P. Sorensen, B. Bolliet, A. Mocsy, Y. Pandit and N. Pruthi, Phys. Lett. B **705**, 71 (2011) [arXiv:1102.1403 [nucl-th]].
- [41] D. Teaney and L. Yan, Phys. Rev. C **83**, 064904 (2011) [arXiv:1010.1876 [nucl-th]].
- [42] Z. Qiu and U. W. Heinz, Phys. Rev. C **84**, 024911 (2011) [arXiv:1104.0650 [nucl-th]]. Z. Qiu, C. Shen and U. W. Heinz, Phys. Lett. B **707**, 151 (2012) [arXiv:1110.3033 [nucl-th]].
- [43] P. Staig and E. Shuryak, Phys. Rev. C **84**, 034908 (2011) [arXiv:1008.3139 [nucl-th]]. P. Staig and E. Shuryak, Phys. Rev. C **84**, 044912 (2011) [arXiv:1105.0676 [nucl-th]].
- [44] J. Takahashi, B. M. Tavares, W. L. Qian, R. Andrade, F. Grassi, Y. Hama, T. Kodama and N. Xu, Phys. Rev. Lett. **103**, 242301 (2009) [arXiv:0902.4870 [nucl-th]].
- [45] B. Schenke, S. Jeon and C. Gale, Phys. Rev. Lett. **106**, 042301 (2011) [arXiv:1009.3244 [hep-ph]].
- [46] J. Xu and C. M. Ko, Phys. Rev. C **83**, 021903 (2011) [arXiv:1011.3750 [nucl-th]].
- [47] G. -Y. Qin, H. Petersen, S. A. Bass and B. Muller, Phys. Rev. C **82**, 064903 (2010) [arXiv:1009.1847 [nucl-th]].
- [48] G. -L. Ma and X. -N. Wang, Phys. Rev. Lett. **106**, 162301 (2011) [arXiv:1011.5249 [nucl-th]].
- [49] W. Li, Mod. Phys. Lett. A **27**, 1230018 (2012) [arXiv:1206.0148 [nucl-ex]].
- [50] W. -T. Deng, Z. Xu and C. Greiner, Phys. Lett. B **711**, 301 (2012) [arXiv:1112.0470 [hep-ph]]; E. Avsar, Y. Hatta, C. Flensburg, J. Y. Ollitrault and T. Ueda, J. Phys. G **38**, 124053 (2011) [arXiv:1106.4356 [hep-ph]]; E. Avsar, C. Flensburg, Y. Hatta, J. -Y. Ollitrault and T. Ueda, Phys. Lett. B **702**, 394 (2011) [arXiv:1009.5643 [hep-ph]]; S. M. Troshin and N. E. Tyurin, Mod. Phys. Lett. A **26**, 1095 (2011) [arXiv:1103.0626 [hep-ph]]; A. Kisiel, Phys. Rev. C **84**, 044913 (2011) [arXiv:1012.1517 [nucl-th]]; P. Bozek, Eur. Phys. J. C **71**, 1530 (2011) [arXiv:1010.0405 [hep-ph]].
- [51] E. Shuryak, arXiv:1009.4635 [hep-ph].
- [52] Piotr Bózek, Phys. Rev. C **85**, 014911 (2012).
- [53] Piotr Bózek and Wojciech Broniowski, Phys. Lett. B **720**, 250 (2013)
- [54] P. Bozek and W. Broniowski, Phys. Lett. B **720**, 250 (2013) [arXiv:1301.3314 [nucl-th]].
- [55] E. Levin and A. H. Rezaeian, Phys. Rev. D **84**, 034031 (2011) [arXiv:1105.3275 [hep-ph]]; A. Kovner and M. Lublinsky, Phys. Rev. D **84**, 094011 (2011) [arXiv:1109.0347 [hep-ph]]; A. Kovner and M. Lublinsky, Phys. Rev. D **83**, 034017 (2011) [arXiv:1012.3398 [hep-ph]]; J. Bartels and M. G. Ryskin, arXiv:1105.1638 [hep-ph].
- [56] K. Dusling and R. Venugopalan, Phys. Rev. Lett. **108**, 262001 (2012) [arXiv:1201.2658 [hep-ph]]; A. Dumitru, K. Dusling, F. Gelis, J. Jalilian-Marian, T. Lappi and R. Venugopalan, Phys. Lett. B **697**, 21 (2011) [arXiv:1009.5295 [hep-ph]].
- [57] B. G. Zakharov, arXiv:1307.3674.
- [58] B. G. Zakharov, arXiv:1311.1159 [hep-ph].
- [59] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, 172302 (2007) [nucl-ex/0610036].
- [60] B. Sahlmueller [PHENIX Collaboration], Nucl. Phys. A904-905 **2013**, 795c (2013) [arXiv:1210.5547 [nucl-ex]].
- [61] D. V. Perepelitsa [PHENIX Collaboration], Nucl. Phys. A **904-905**, 1003c (2013).
- [62] W. A. Horowitz and Miklos Gyulassy, Nucl. Phys. A **872**, 265 (2011).
- [63] R. Baier, Y. L. Dokshitzer, A. H. Mueller, S. Peigne and D. Schiff, Nucl. Phys. B **484**, 265 (1997) [hep-ph/9608322].
- [64] J. Adams *et al.*, [STAR Collaboration], Phys. Rev. Lett. **91**, 172302 (2003).
- [65] A. Adare *et al.*, [PHENIX Collaboration], Phys. Rev. Lett. **101**, 232301 (2008).
- [66] S. Chatrchyan *et al.*, [CMS Collaboration], Eur. Phys. J. C **72**, 1945 (2012).
- [67] B. Abelev *et al.*, [ALICE Collaboration] arXiv:1208.2711.
- [68] Francois Arleo, David d'Enterria, and Andre S. Yoon, JHEP **06**, 035 (2010).
- [69] S. Chatrchyan *et al.*, [CMS Collaboration], JHEP **08**, 086 (2011).
- [70] Michael L. Miller, Klaus Reygers, Stephen J. Sanders, and Peter Steinberg, Annu. Rev. Nucl. Part. Sci. **57**, 205 (2007).
- [71] Chun Shen and Ulrich Heinz, Phys. Rev. C **85**, 054902 (2012), Phys. Rev. C **86**, 049903(E) (2012)
- [72] Ulrich Heinz and J. Scott Moreland, Phys. Rev. C **84**, 054905 (2011).
- [73] J. D. Bjorken, Phys. Rev. D **27**, 140 (1983).
- [74] B. Betz and M. Gyulassy, arXiv:1305.6458 [nucl-th].
- [75] A. Buzzatti and M. Gyulassy, Nucl. Phys. A904-905 **2013**, 779c (2013) [arXiv:1210.6417 [hep-ph]].
- [76] D. Molnar and D. Sun, arXiv:1305.1046 [nucl-th].
- [77] S. Chatrchyan *et al.*, [CMS Collaboration], arXiv:1204.1850.
- [78] Z. Qiu and U. W. Heinz, Phys. Rev. C **84**, 024911 (2011).
- [79] <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResults>
- [80] E. Shuryak and I. Zahed, arXiv:1311.0836 [hep-ph].
- [81] K. Dusling and R. Venugopalan, Phys. Rev. D **87**, no. 5, 054014 (2013) [arXiv:1211.3701 [hep-ph]]. K. Dusling and R. Venugopalan, Phys. Rev. D **87**, 094034 (2013) [arXiv:1302.7018 [hep-ph]].
- [82] J. L. Albacete, N. Armesto, R. Baier, G. G. Barnafoldi, J. Barrette, S. De, W. -T. Deng and A. Dumitru *et al.*, Int. J. Mod. Phys. E **22**, 1330007 (2013) [arXiv:1301.3395 [hep-ph]].
- [83] A. Dumitru, T. Lappi and L. McLerran, arXiv:1310.7136 [hep-ph].
- [84] J. -P. Blaizot, F. Gelis, J. -F. Liao, L. McLerran and R. Venugopalan, Nucl. Phys. A **873**, 68 (2012). J. -P. Blaizot, J. Liao and L. McLerran, arXiv:1305.2119 [hep-ph]. X. -G. Huang and J. Liao, arXiv:1303.7214 [nucl-th].
- [85] A. Bzdak, B. Schenke, P. Tribedy and R. Venugopalan, arXiv:1304.3403 [nucl-th].
- [86] B. Schenke, P. Tribedy and R. Venugopalan, Phys. Rev. Lett. **108**, 252301 (2012) [arXiv:1202.6646 [nucl-th]].

- [87] B. Schenke, P. Tribedy and R. Venugopalan, Phys. Rev. C **86**, 034908 (2012) [arXiv:1206.6805 [hep-ph]].
- [88] B. Abelev *et al.* [ALICE Collaboration], Phys. Rev. Lett. **110**, 032301 (2013) [arXiv:1210.3615 [nucl-ex]].
- [89] B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **93**, 082301 (2004) [nucl-ex/0311009].
- [90] D. Molnar and D. Sun, arXiv:1209.2430 [nucl-th]; D. Molnar and D. Sun, arXiv:1305.1046 [nucl-th].