

$B^0 - \bar{B}^0$ mixing in supersymmetry with gauged baryon and lepton numbers

Fei Sun^{a,b*}, Tai-Fu Feng^{a,b,c,d}, Shu-Min Zhao^{a,c,d},
Hai-Bin Zhang^{a,b}, Tie-Jun Gao^c, Jian-Bin Chen^e

^a*Department of Physics, Hebei University, Baoding, 071002, China*

^b*Department of Physics, Dalian University of Technology, Dalian, 116024, China*

^c*Institute of theoretical Physics, Chinese Academy of Sciences, Beijing, 100190, China*

^d*The Key Laboratory of Mathematics-Mechanization (KLMM), Beijing, 100190, China*

^e*Department of Physics, Taiyuan University of Technology, Taiyuan, 030024, China*

Abstract

We perform an analysis on $B^0 - \bar{B}^0$ mixing in the extension of the minimal supersymmetric standard model where baryon and lepton numbers are local gauge symmetries (BLMSSM) by using the effective Hamiltonian method. And the constraint of a 125 GeV Higgs to the parameter space has also been considered. The numerical results indicate that the contributions of the extra particles can be sizeable in $B^0 - \bar{B}^0$ mixing. For certain parameter sets, the theoretical prediction of mass differences Δm_B agrees with the current experimental result. Furthermore, $B^0 - \bar{B}^0$ mixing in the BLMSSM can preliminarily constrain the parameter space. With the development of more precise theoretical analysis and experimental determinations, the $B^0 - \bar{B}^0$ mixing in the BLMSSM will have a clearer picture and the parameter space in this model will also be further constrained.

PACS numbers: 12.60.Jv, 14.40.Nd

Keywords: Supersymmetry, $B^0 - \bar{B}^0$ mixing, Baryon number, Higgs.

* Corresponding author. email:sunfei@mail.dlut.edu.cn

I. INTRODUCTION

The Minimal Supersymmetric Standard Model (MSSM) [1–5], as one of the most appealing options for the physics beyond the Standard Model (SM), has drawn the physicists’ attention for a long time. As the simplest soft broken supersymmetry (SUSY) theory, the MSSM can solve hierarchy problem, ensure that the gauge couplings unify at high energies and provide a good dark matter candidate. To search for new particles predicted by SUSY, the Large Hadron Collider (LHC) has collected huge amounts of data, the CMS [7] and ATLAS [8] experiments now set strong limits on these parameter space [9–12]. However, the present searches are largely based on the assumption of conserved R-parity [6]. Some studies in the low-energy SUSY have been motivated by the results of the LHC [13–23], and R-parity violating scenarios of general MSSM have been proposed [24–48].

A model based on the gauge symmetry group $SU(3) \otimes SU(2) \otimes U(1)_Y \otimes U(1)_B \otimes U(1)_L$ has been investigated at the TeV scale recently [49–52], where B stands for baryon number and L stands for lepton number. In this theory, the baryon and lepton numbers are local gauge symmetries spontaneously broken at the TeV scale. Breaking baryon number can explain the origin of the matter-antimatter asymmetry in the Universe. And breaking lepton number can explain the smallness of neutrino masses [53–57]. Two extensions of the SM where B and L are spontaneously broken gauge symmetries near the weak scale are constructed [58]: model I is a non-supersymmetric extension [59, 60]; model II (BLMSSM) is a supersymmetric extension and is more favoured by the experiments [61]. The BLMSSM has been studied in great detail and could avoid the current LHC bounds on the SUSY mass spectrum [62, 63, 65]. Some further phenomenology analysis based on the BLMSSM coincide with the current experimental data well, the mass and decays of the lightest CP-even Higgs have been investigated in Refs. [65, 66], and the neutron electric dipole moment in CP violating BLMSSM has also been studied [67].

The flavor changing neutral current (FCNC) processes are highly suppressed in the SM, therefore it is a fertile ground to search for physics beyond SM (BSM). FCNC processes such as $b \rightarrow s\gamma$, $K^0 - \bar{K}^0$ and $B^0 - \bar{B}^0$ mixing have played an important role in particle physics over the last four decades. It is well known that CP violation was first observed

in the decays of K_L^0 meson in 1964 [68], and CP violation of the neutral B meson system was observed in 2001 [69]. The first indication of a large top quark mass was also given by $B^0 - \bar{B}^0$ mixing [70, 71]. B -system decays have an advantage over the K -system to provide a direct test of the CP violating of SM and is free of corrections from strong interactions [72–74]. The experiment results of $B^0 - \bar{B}^0$ mixing have been published by the ALEPH [75], DELPHI [76, 77], L3 [78], OPAL [79, 80] BaBar [81], Belle [82], CDF [83], DØ [84], and LHCb [85] collaborations. Current experimental result of mass difference is $\Delta m_B^{Exp} = 0.507 \pm 0.004 \text{ ps}^{-1} = (3.337 \pm 0.033) \times 10^{-13} \text{ GeV}$ [86]. Calculations for $B^0 - \bar{B}^0$ mixing have been done in the SM, the two-Higgs doublet model (2HDM), the MSSM and other models [87–96]. The SM prediction for mass difference is $\Delta m_B^{SM} = 0.543 \pm 0.091 \text{ ps}^{-1}$ [97], which has a good agreement with the experiment. However, the theoretical error is around 17%, which is considerably larger than the experimental error. The running of LHC will resume in 2015 with higher energy and luminosity. Proposals for next-generation B-factories including SuperKEKB in Japan whose target luminosity is $8 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ will start collecting data in the near future [98]. This may also give some hints on physics beyond the SM. So it is important for experimental and theoretical physicist to search for new physics. As a candidate of new physics, the BLMSSM provides new FCNC at loop level in the $B^0 - \bar{B}^0$ mixing. We will carry out our calculations for $B^0 - \bar{B}^0$ mixing in this model.

Our presentation is organized as follows. In Section II, we briefly summarize the main features of the BLMSSM and introduce the superpotential as well as soft breaking terms, then we obtain the mass matrices and couplings needed for $B^0 - \bar{B}^0$ mixing. In Section III, we give the analytical formulae of the $B^0 - \bar{B}^0$ mixing in BLMSSM. The numerical analysis are shown in Section IV. Section V presents our conclusions. Finally, some related formulae are given in Appendix A–B.

II. BLMSSM

In this section, we briefly review some main features of the BLMSSM. In the BLMSSM with gauged baryon (B) and lepton (L), by adding the new quarks with baryon number $B_4 = \frac{3}{2}$ and the new leptons with lepton number $L_4 = \frac{3}{2}$, one can cancel the baryonic and

leptonic anomalies respectively [58]. Compared with the MSSM, the BLMSSM includes many new fields. Tables I–IV list the superfields including the new quarks, new leptons, new Higgs, the exotic superfields $\hat{X}$ and $\hat{X}'$, respectively. As one can see, the left-handed superfields have the same absolute value of $U(1)_B$ as that of the right-handed superfields but with a contrary sign to cancel baryonic anomalies in the quark sector, similarly for the $U(1)_L$ in the leptonic sector to cancel leptonic anomalies.

TABLE I: Superfields including the new quarks in the BLMSSM.

Superfields	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
$\hat{Q}_4$	3	2	1/6	B_4	0
$\hat{U}_4^c$	$\bar{3}$	1	-2/3	$-B_4$	0
$\hat{D}_4^c$	$\bar{3}$	1	1/3	$-B_4$	0
$\hat{Q}_5^c$	$\bar{3}$	2	-1/6	$-(1 + B_4)$	0
$\hat{U}_5$	3	1	2/3	$1 + B_4$	0
$\hat{D}_5$	3	1	-1/3	$1 + B_4$	0

TABLE II: Superfields including the new leptons in the BLMSSM.

Superfields	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
$\hat{L}_4$	1	2	-1/2	0	L_4
$\hat{E}_4^c$	1	1	1	0	$-L_4$
$\hat{N}_4^c$	1	1	0	0	$-L_4$
$\hat{L}_5^c$	1	2	1/2	0	$-(3 + L_4)$
$\hat{E}_5$	1	1	-1	0	$3 + L_4$
$\hat{N}_5$	1	1	0	0	$3 + L_4$

In order to break baryon number spontaneously, we need to introduce the superfields $\hat{\Phi}_B$ and $\hat{\varphi}_B$ to acquire nonzero vacuum expectation values (VEVs), which also generate large

TABLE III: Superfields including the new Higgs in the BLMSSM.

Superfields	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
$\hat{\Phi}_B$	1	1	0	1	0
$\hat{\varphi}_B$	1	1	0	-1	0
$\hat{\Phi}_L$	1	1	0	0	-2
$\hat{\varphi}_L$	1	1	0	0	2

TABLE IV: Superfields avoiding stability for the exotic quarks in the BLMSSM.

Superfields	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$U(1)_B$	$U(1)_L$
$\hat{X}$	1	1	0	$2/3 + B_4$	0
$\hat{X}'$	1	1	0	$-(2/3 + B_4)$	0

mass for the new quarks. Similarly, we introduce the superfields $\hat{\Phi}_L$ and $\hat{\varphi}_L$ to acquire VEVs spontaneously breaking lepton number. Finally, the exotic quarks should be unstable, so the model also includes the superfields $\hat{X}$ and $\hat{X}'$ to avoid the stability for the exotic quarks. Here $\hat{\Phi}_B$ and $\hat{\varphi}_B$ have $U(1)_B$ charge 1 and -1, respectively, $\hat{\Phi}_L$ and $\hat{\varphi}_L$ have $U(1)_L$ charge -2 and 2, respectively. For superfields $\hat{X}$ and $\hat{X}'$, $U(1)_B$ charge is $2/3 + B_4$ and $-(2/3 + B_4)$, respectively. Here the lightest X could be a dark matter candidate.

The superpotential in BLMSSM is written as

$$\mathcal{W}_{BLMSSM} = \mathcal{W}_{MSSM} + \mathcal{W}_B + \mathcal{W}_L + \mathcal{W}_X, \quad (1)$$

where $\mathcal{W}_{MSSM}$ is the superpotential of MSSM, and

$$\begin{aligned}
 \mathcal{W}_B &= \lambda_Q \hat{Q}_4 \hat{Q}_5^c \hat{\Phi}_B + \lambda_U \hat{U}_4^c \hat{U}_5 \hat{\varphi}_B + \lambda_D \hat{D}_4^c \hat{D}_5 \hat{\varphi}_B + \mu_B \hat{\Phi}_B \hat{\varphi}_B \\
 &\quad + Y_{u_4} \hat{Q}_4 \hat{H}_u \hat{U}_4^c + Y_{d_4} \hat{Q}_4 \hat{H}_d \hat{D}_4^c + Y_{u_5} \hat{Q}_5^c \hat{H}_d \hat{U}_5 + Y_{d_5} \hat{Q}_5^c \hat{H}_u \hat{D}_5, \\
 \mathcal{W}_L &= Y_{e_4} \hat{L}_4 \hat{H}_d \hat{E}_4^c + Y_{\nu_4} \hat{L}_4 \hat{H}_u \hat{N}_4^c + Y_{e_5} \hat{L}_5^c \hat{H}_u \hat{E}_5 + Y_{\nu_5} \hat{L}_5^c \hat{H}_d \hat{N}_5 \\
 &\quad + Y_{\nu} \hat{L} \hat{H}_u \hat{N}^c + \lambda_{N^c} \hat{N}^c \hat{N}^c \hat{\varphi}_L + \mu_L \hat{\Phi}_L \hat{\varphi}_L, \\
 \mathcal{W}_X &= \lambda_1 \hat{Q} \hat{Q}_5^c \hat{X} + \lambda_2 \hat{U}^c \hat{U}_5 \hat{X}' + \lambda_3 \hat{D}^c \hat{D}_5 \hat{X}' + \mu_X \hat{X} \hat{X}'.
 \end{aligned} \quad (2)$$

In the superpotential above, the exotic quarks obtain TeV scale masses after Φ_B , φ_B acquiring nonzero VEVs, and the nonzero VEV of φ_L implements the seesaw mechanism for the tiny neutrino masses. Correspondingly, the soft breaking terms are generally given as

$$\begin{aligned}
\mathcal{L}_{soft} = & \mathcal{L}_{soft}^{MSSM} - (m_{\tilde{N}^c}^2)_{IJ} \tilde{N}_I^{c*} \tilde{N}_J^c - m_{\tilde{Q}_4}^2 \tilde{Q}_4^\dagger \tilde{Q}_4 - m_{\tilde{U}_4}^2 \tilde{U}_4^{c*} \tilde{U}_4^c - m_{\tilde{D}_4}^2 \tilde{D}_4^{c*} \tilde{D}_4^c \\
& - m_{\tilde{Q}_5}^2 \tilde{Q}_5^{c\dagger} \tilde{Q}_5^c - m_{\tilde{U}_5}^2 \tilde{U}_5^{c*} \tilde{U}_5^c - m_{\tilde{D}_5}^2 \tilde{D}_5^{c*} \tilde{D}_5^c - m_{\tilde{L}_4}^2 \tilde{L}_4^\dagger \tilde{L}_4 - m_{\tilde{\nu}_4}^2 \tilde{\nu}_4^{c*} \tilde{\nu}_4^c - m_{\tilde{E}_4}^2 \tilde{e}_4^{c*} \tilde{e}_4^c \\
& - m_{\tilde{L}_5}^2 \tilde{L}_5^{c\dagger} \tilde{L}_5^c - m_{\tilde{\nu}_5}^2 \tilde{\nu}_5^{c*} \tilde{\nu}_5^c - m_{\tilde{E}_5}^2 \tilde{e}_5^{c*} \tilde{e}_5^c - m_{\Phi_B}^2 \Phi_B^* \Phi_B - m_{\varphi_B}^2 \varphi_B^* \varphi_B \\
& - m_{\Phi_L}^2 \Phi_L^* \Phi_L - m_{\varphi_L}^2 \varphi_L^* \varphi_L - (m_B \lambda_B \lambda_B + m_L \lambda_L \lambda_L + h.c.) \\
& + \{ A_{u_4} Y_{u_4} \tilde{Q}_4 H_u \tilde{U}_4^c + A_{d_4} Y_{d_4} \tilde{Q}_4 H_d \tilde{D}_4^c + A_{u_5} Y_{u_5} \tilde{Q}_5 H_d \tilde{U}_5^c + A_{d_5} Y_{d_5} \tilde{Q}_5 H_u \tilde{D}_5^c \\
& + A_{BQ} \lambda_Q \tilde{Q}_4 \tilde{Q}_5^c \Phi_B + A_{BU} \lambda_U \tilde{U}_4^c \tilde{U}_5^c \varphi_B + A_{BD} \lambda_D \tilde{D}_4^c \tilde{D}_5^c \varphi_B + B_B \mu_B \Phi_B \varphi_B + h.c. \} \\
& + \{ A_{e_4} Y_{e_4} \tilde{L}_4 H_d \tilde{E}_4^c + A_{N_4} Y_{N_4} \tilde{L}_4 H_u \tilde{N}_4^c + A_{e_5} Y_{e_5} \tilde{L}_5 H_u \tilde{E}_5^c + A_{N_5} Y_{N_5} \tilde{L}_5 H_d \tilde{N}_5^c \\
& + A_N Y_N \tilde{L}_5 H_u \tilde{N}^c + A_{N^c} \lambda_{N^c} \tilde{N}^c \tilde{N}^c \varphi_L + B_L \mu_L \Phi_L \varphi_L + h.c. \} \\
& + \{ A_1 \lambda_1 \tilde{Q} \tilde{Q}_5^c X + A_2 \lambda_2 \tilde{U}^c \tilde{U}_5^c X' + A_3 \lambda_3 \tilde{D}^c \tilde{D}_5^c X' + B_X \mu_X X X' + h.c. \} , \tag{3}
\end{aligned}$$

where $\mathcal{L}_{soft}^{MSSM}$ is the soft breaking terms of MSSM, λ_B , λ_L are gauginos of $U(1)_B$ and $U(1)_L$, respectively. After the $SU(2)_L$ doublets H_u , H_d and $SU(2)_L$ singlets Φ_B , φ_B , Φ_L , φ_L acquiring the nonzero VEVs v_u , v_d , v_B , $\bar{v}_B$, v_L , $\bar{v}_L$

$$\begin{aligned}
H_u &= \begin{pmatrix} H_u^+ \\ \frac{1}{\sqrt{2}}(v_u + H_u^0 + iP_u^0) \end{pmatrix} , \\
H_d &= \begin{pmatrix} \frac{1}{\sqrt{2}}(v_d + H_d^0 + iP_d^0) \\ H_d^- \end{pmatrix} , \\
\Phi_B &= \frac{1}{\sqrt{2}}(v_B + \Phi_B^0 + iP_B^0) , \\
\varphi_B &= \frac{1}{\sqrt{2}}(\bar{v}_B + \varphi_B^0 + i\bar{P}_B^0) , \\
\Phi_L &= \frac{1}{\sqrt{2}}(v_L + \Phi_L^0 + iP_L^0) , \\
\varphi_L &= \frac{1}{\sqrt{2}}(\bar{v}_L + \varphi_L^0 + i\bar{P}_L^0) , \tag{4}
\end{aligned}$$

the local gauge symmetry $SU(2)_L \otimes U(1)_Y \otimes U(1)_B \otimes U(1)_L$ is broken down to the electromagnetic symmetry $U(1)_e$.

After the symmetry breaking, we can obtain the physical spectrum of this model. The chargino mass matrix is as same as the chargino mass matrix in MSSM. Z_+, Z_- are the matrices to diagonalize the chargino mass mixing matrix $M_{\tilde{\chi}^\pm}$

$$Z_-^T \mathcal{M}_{\tilde{\chi}^\pm} Z_+ = \text{diag}(m_{\tilde{\chi}_1^\pm}, m_{\tilde{\chi}_2^\pm}). \quad (5)$$

The exotic bottom quark mass matrix is given by

$$\mathcal{M}_{b'} = \begin{pmatrix} -\frac{1}{\sqrt{2}}\lambda_Q v_B, & -\frac{1}{\sqrt{2}}Y_{d_5} v_u \\ -\frac{1}{\sqrt{2}}Y_{d_4} v_d, & \frac{1}{\sqrt{2}}\lambda_d \bar{v}_B \end{pmatrix}, \quad (6)$$

and this mass matrix is diagonalized by two rotation matrices $W_{b'}$ and $U_{b'}$

$$W_{b'}^\dagger \cdot M_{b'} \cdot U_{b'} = \text{diag}(m_{b'_1}, m_{b'_2}). \quad (7)$$

The mass matrix of the first three families up-type scalar quark is given as follow

$$\mathcal{M}_{\tilde{U}^I}^2 = \begin{pmatrix} m_{\tilde{U}_L^I}^2 + m_{U^I}^2 + D_L^{\tilde{U}^I} + \frac{2}{3}m_{Z_B}^2 \cos 2\beta_B & m_{U^I}(A_{U^I} - \mu^* \cot \beta) \\ m_{U^I}(A_{U^I}^* - \mu \cot \beta) & m_{\tilde{U}_R^I}^2 + m_{U^I}^2 + D_R^{\tilde{U}^I} - \frac{2}{3}m_{Z_B}^2 \cos 2\beta_B \end{pmatrix}, \quad (8)$$

which has some differences from that of MSSM, here $m_{Z_B}^2 = g_B^2(v_B^2 + \bar{v}_B^2)$ is the mass squared of $U(1)_B$ gauge boson Z_B , and the D -terms are

$$\begin{aligned} D_L^{\tilde{U}^I} &= \left(\frac{1}{2} - \frac{2}{3}\sin^2\theta_W\right)m_Z^2 \cos 2\beta, \\ D_R^{\tilde{U}^I} &= -\frac{2}{3}\sin^2\theta_W m_Z^2 \cos 2\beta. \end{aligned} \quad (9)$$

In the basis $(\tilde{Q}_4^{2*}, \tilde{D}_4^c, \tilde{Q}_5^{1c}, \tilde{D}_5)$, the mass term for the exotic bottom scalar quarks in the Lagrangian reads as

$$-\mathcal{L}_{b'}^{mass} = \left(\tilde{Q}_4^{2*}, \tilde{D}_4^c, \tilde{Q}_5^{1c}, \tilde{D}_5 \right) \cdot \mathcal{M}_{b'}^2 \cdot \left(\tilde{Q}_4^{2*}, \tilde{D}_4^c, \tilde{Q}_5^{1c}, \tilde{D}_5 \right)^\dagger, \quad (10)$$

where $\mathcal{M}_{b'}^2$ is a 4×4 matrix, and the matrix elements are listed as follows

$$\begin{aligned} (\mathcal{M}_{b'}^2)_{11} &= m_{\tilde{Q}_4}^2 + \frac{1}{2}Y_{u_4}^2 v_u^2 + \frac{1}{2}Y_{d_4}^2 v_d^2 + \frac{1}{2}\lambda_Q^2 v_B^2 - \left(\frac{1}{2} - \frac{2}{3}s_w^2\right)m_Z^2 \cos 2\beta + \frac{B_4}{2}m_{Z_B}^2 \cos 2\beta_B, \\ (\mathcal{M}_{b'}^2)_{12} &= (\mathcal{M}_{b'}^2)_{21}^* = -\frac{1}{\sqrt{2}}Y_{d_4} v_d A_{d_4} + \frac{1}{\sqrt{2}}Y_{d_4} \mu v_d, \\ (\mathcal{M}_{b'}^2)_{13} &= (\mathcal{M}_{b'}^2)_{31}^* = -\frac{1}{\sqrt{2}}\lambda_Q v_B A_{BQ} + \sqrt{2}\lambda_Q \mu v_B \bar{v}_B, \end{aligned}$$

$$\begin{aligned}
(\mathcal{M}_{\tilde{b}'}^2)_{14} &= (\mathcal{M}_{\tilde{b}'}^2)_{41}^* = -\frac{1}{\sqrt{2}}Y_{d_4}\lambda_d v_d \bar{v}_B + \frac{1}{\sqrt{2}}Y_{d_5}\lambda_Q v_u v_B, \\
(\mathcal{M}_{\tilde{b}'}^2)_{22} &= m_{\tilde{D}_4}^2 + \frac{1}{2}Y_{d_4}^2 v_d^2 + \frac{1}{2}\lambda_d^2 \bar{v}_B^2 - \frac{1}{3}s_w^2 m_Z^2 \cos 2\beta - \frac{B_4}{2}m_{Z_B}^2 \cos 2\beta_B, \\
(\mathcal{M}_{\tilde{b}'}^2)_{23} &= (\mathcal{M}_{\tilde{b}'}^2)_{32}^* = \frac{1}{2}\lambda_Q Y_{d_4} v_d v_B + \frac{1}{2}\lambda_d Y_{d_5} v_u \bar{v}_B, \\
(\mathcal{M}_{\tilde{b}'}^2)_{24} &= (\mathcal{M}_{\tilde{b}'}^2)_{42}^* = -\frac{1}{\sqrt{2}}\lambda_d A_{BD} \bar{v}_B + \frac{1}{\sqrt{2}}\lambda_d \mu_B v_B, \\
(\mathcal{M}_{\tilde{b}'}^2)_{33} &= m_{\tilde{Q}_5}^2 + \frac{1}{2}Y_{u_5}^2 v_d^2 + \frac{1}{2}Y_{d_5}^2 v_u^2 + \frac{1}{2}\lambda_Q^2 v_B^2 - \left(\frac{1}{2} + \frac{1}{3}s_w^2\right)m_Z^2 \cos 2\beta - \frac{1+B_4}{2}m_{Z_B}^2 \cos 2\beta_B, \\
(\mathcal{M}_{\tilde{b}'}^2)_{44} &= m_{\tilde{D}_5}^2 + \frac{1}{2}Y_{d_5}^2 v_u^2 + \frac{1}{2}\lambda_d^2 \bar{v}_B^2 + \frac{1}{3}s_w^2 m_Z^2 \cos 2\beta + \frac{1+B_4}{2}m_{Z_B}^2 \cos 2\beta_B, \\
(\mathcal{M}_{\tilde{b}'}^2)_{34} &= (\mathcal{M}_{\tilde{b}'}^2)_{43}^* = Y_{d_5} A_{d_5} v_u + \frac{1}{\sqrt{2}}Y_{d_5} \mu v_d.
\end{aligned} \tag{11}$$

The mass-squared matrix $\mathcal{M}_{\tilde{b}'}^2$ is diagonalized by the unitary matrix $Z_{\tilde{b}'}$

$$Z_{\tilde{b}'}^\dagger \cdot \mathcal{M}_{\tilde{b}'}^2 \cdot Z_{\tilde{b}'} = \text{diag}(m_{\tilde{b}'_1}^2, m_{\tilde{b}'_2}^2, m_{\tilde{b}'_3}^2, m_{\tilde{b}'_4}^2), \tag{12}$$

and the physical states are related to the gauge states by

$$\begin{pmatrix} \tilde{b}'_1 \\ \tilde{b}'_2 \\ \tilde{b}'_3 \\ \tilde{b}'_4 \end{pmatrix} = Z_{\tilde{b}'}^\dagger \cdot \begin{pmatrix} \tilde{Q}_4^2 \\ \tilde{D}_4^{c*} \\ \tilde{Q}_5^{1c*} \\ \tilde{D}_5^* \end{pmatrix}. \tag{13}$$

The mass squared matrix in the basis (X^*, X') is

$$\mathcal{M}_X^2 = \begin{pmatrix} \mu_X^2 + \frac{1}{2}(\frac{2}{3} + B_4)m_{Z_B}^2 \cos 2\beta_B & -\mu_X^* B_X^* \\ -\mu_X B_X & \mu_X^2 - \frac{1}{2}(\frac{2}{3} + B_4)m_{Z_B}^2 \cos 2\beta_B \end{pmatrix}. \tag{14}$$

Adopting the unitary transformation, the mass eigenstates are

$$\begin{pmatrix} X_1 \\ X_2 \end{pmatrix} = Z_X^\dagger \begin{pmatrix} X \\ X'^* \end{pmatrix}, \tag{15}$$

and the mass squared matrix M_X^2 is diagonalized by

$$Z_X^\dagger \cdot M_X^2 \cdot Z_X = \text{diag}(m_{X_1}^2, m_{X_2}^2). \tag{16}$$

In four-component Dirac spinors, the mass term for superfields $\tilde{X}$ is given by

$$-\mathcal{L}_{\tilde{X}}^{\text{mass}} = \mu_X \tilde{X} \bar{\tilde{X}}, \tag{17}$$

here, we have defined

$$\tilde{X} = \begin{pmatrix} \psi_X \\ \bar{\psi}_{X'} \end{pmatrix}. \quad (18)$$

So the parameter μ_X is the mass of the particle $\tilde{X}$.

In mass basis, we obtain the couplings of quark-exotic quark and the superfields X

$$\mathcal{L}_{Xb'd} = \sum_{\delta, \epsilon=1}^2 \left(-\lambda_1 (W_{b'}^\dagger)_{\delta 1} (Z_X)_{1\epsilon} X_\epsilon \bar{b}_\delta P_L d^I - \lambda_3^* (U_{b'}^\dagger)_{\delta 2} (Z_X)_{2\epsilon} X_\epsilon \bar{b}_\delta P_R d^I \right) + h.c.. \quad (19)$$

We also obtain the couplings of quark-exotic scalar quark and the field $\tilde{X}$

$$\mathcal{L}_{\tilde{X}\tilde{b}'d^I} = \sum_{\rho=1}^4 \left(-\lambda_1 (Z_{\tilde{b}'}^*)_{3\rho} \tilde{b}'_\rho \tilde{X} P_L d^I - \lambda_3^* (Z_{\tilde{b}'}^*)_{4\rho} \tilde{b}'_\rho \tilde{X} P_R d^I \right) + h.c., \quad (20)$$

where λ_1, λ_3 are the coupling coefficients, and δ, ϵ, ρ are the indices of the flavor.

Considering the radiative corrections, the mass squared matrix for the neutral CP-even Higgs in the basis (H_d^0, H_u^0) is written as [99–110]

$$\mathcal{M}_{even}^2 = \begin{pmatrix} M_{11}^2 + \Delta_{11} & M_{12}^2 + \Delta_{12} \\ M_{12}^2 + \Delta_{12} & M_{22}^2 + \Delta_{22} \end{pmatrix}, \quad (21)$$

where

$$\begin{aligned} M_{11}^2 &= m_Z^2 \cos^2 \beta + m_{A^0}^2 \sin^2 \beta, \\ M_{12}^2 &= -(m_Z^2 + m_{A^0}^2) \sin \beta \cos \beta, \\ M_{22}^2 &= m_Z^2 \sin^2 \beta + m_{A^0}^2 \cos^2 \beta, \\ \Delta_{11} &= \Delta_{11}^{MSSM} + \Delta_{11}^B + \Delta_{11}^L, \\ \Delta_{12} &= \Delta_{12}^{MSSM} + \Delta_{12}^B + \Delta_{12}^L, \\ \Delta_{22} &= \Delta_{22}^{MSSM} + \Delta_{22}^B + \Delta_{22}^L, \end{aligned} \quad (22)$$

and the expressions of $\Delta_{11}^{B,L}, \Delta_{12}^{B,L}, \Delta_{22}^{B,L}$ can be found in Refs. [65, 66]. A Higgs around 125 GeV has been observed at the LHC by ATLAS [111] and CMS [112] with the combined significances of 5.9 and 5.0 standard deviations, respectively. So after diagonalizing the mass

squared matrix, the lightest neutral CP even Higgs m_{h_0} should satisfy this constraint. To obtain the Higgs h_0 with mass of 125 GeV gives a strong limit on the parameter space. Considering this constraint, we can also obtain $m_{A^0}^2$ from the inverse solution of Eq. (21). We have

$$m_{A^0}^2 = \frac{m_{h_0}^2 (m_z^2 - m_{h_0}^2 + \Delta_{11} + \Delta_{22}) - m_z^2 \Delta_A + \Delta_{12}^2 - \Delta_{11} \Delta_{22}}{-m_{h_0}^2 + m_z^2 \cos^2 2\beta + \Delta_B}, \quad (23)$$

where

$$\begin{aligned} \Delta_A &= \sin^2 \beta \Delta_{11} + \cos^2 \beta \Delta_{22} + \sin 2\beta \Delta_{12}, \\ \Delta_B &= \cos^2 \beta \Delta_{11} + \sin^2 \beta \Delta_{22} + \sin 2\beta \Delta_{12}. \end{aligned} \quad (24)$$

For the charged Higgs scalars, $H_{1,2}^\pm$ are related to the initial Higgs by the matrix Z_H , and the charged Higgs mass $m_{H_1^\pm}$ satisfy a relation with the pseudo-scalar Higgs mass m_{A^0} at tree-level:

$$m_{H_1^\pm} = \sqrt{m_{A^0}^2 + m_W^2}, \quad (25)$$

Using the Feynman–t’Hooft gauge, another charged Higgs boson $H_2^\pm$ has the same mass as the gauge boson W .

III. $B^0 - \bar{B}^0$ MIXING

When external masses and momenta are neglected, the general form of the effective Hamiltonian for $B^0 - \bar{B}^0$ mixing at the weak scale can be expressed as[113]

$$H_{eff} = \frac{1}{4} \frac{G_F^2}{\pi^2} m_W^2 \sum_{\alpha=1}^8 C_\alpha \mathcal{O}_\alpha, \quad (26)$$

where G_F denotes the Fermi constant, C_α are the corresponding Wilson coefficients, $\mathcal{O}_\alpha$ are the effective operators, which read as

$$\begin{aligned} \mathcal{O}_1 &= \bar{d}\gamma_\mu P_L b \bar{d}\gamma^\mu P_L b, \\ \mathcal{O}_2 &= \bar{d}\gamma_\mu P_L b \bar{d}\gamma^\mu P_R b, \\ \mathcal{O}_3 &= \bar{d}P_L b \bar{d}P_R b, \end{aligned}$$

$$\begin{aligned}
\mathcal{O}_4 &= \bar{d}P_L b \bar{d}P_L b, \\
\mathcal{O}_5 &= \bar{d}\sigma_{\mu\nu}P_L b \bar{d}\sigma^{\mu\nu}P_L b, \\
\mathcal{O}_6 &= \bar{d}\gamma_\mu P_R b \bar{d}\gamma^\mu P_R b, \\
\mathcal{O}_7 &= \bar{d}P_R b \bar{d}P_R b, \\
\mathcal{O}_8 &= \bar{d}\sigma_{\mu\nu}P_R b \bar{d}\sigma^{\mu\nu}P_R b,
\end{aligned} \tag{27}$$

where $P_{R,L} = (1 \pm \gamma_5)/2$ denote the chiral projectors, $\sigma_{\mu\nu} = [\gamma_\mu, \gamma_\nu]/2$, the $SU(3)$ color indices here have omitted for simplicity.




FIG. 1: The box diagram contributions to $B^0 - \bar{B}^0$ mixing in the SM.




FIG. 2: The box diagrams contributing to $B^0 - \bar{B}^0$ mixing in the BLMSSM.

The box diagram contributions to $B^0 - \bar{B}^0$ mixing from the SM are displayed in Fig. 1, and the box diagrams contributing to $B^0 - \bar{B}^0$ mixing in the BLMSSM are shown in Fig. 2. Note that the diagrams including the particles $\tilde{\chi}$ and $\tilde{X}$ should make a Fierz rearrangement to ensure that the operators are color singlet states as follows

$$\begin{aligned}
\mathcal{O}'_1 &= \mathcal{O}_1, \\
\mathcal{O}'_2 &= \mathcal{O}_2, \\
\mathcal{O}'_3 &= -\frac{1}{2}\mathcal{O}_2, \\
\mathcal{O}'_4 &= -\frac{1}{2}\mathcal{O}_4 - \frac{1}{8}\mathcal{O}_5,
\end{aligned}$$

$$\begin{aligned}
\mathcal{O}'_5 &= -6\mathcal{O}_4 + \frac{1}{2}\mathcal{O}_5, \\
\mathcal{O}'_6 &= \mathcal{O}_6, \\
\mathcal{O}'_7 &= -\frac{1}{2}\mathcal{O}_7 - \frac{1}{8}\mathcal{O}_8, \\
\mathcal{O}'_8 &= -6\mathcal{O}_7 + \frac{1}{2}\mathcal{O}_8.
\end{aligned} \tag{28}$$

The operators with a prime stand for the product of two color non-singlet quark current.

After this, the Wilson coefficients are given as follows

$$\begin{aligned}
C_1 &= V_{ib}V_{id}^*V_{jb}V_{jd}^* (f_{p^2}(x_{u_i}, x_W, x_{u_j}, x_W) - 2\frac{x_{u_i}x_{u_j}}{\sin^2\beta}(Z_H^{2k})^2 f_1(x_{u_i}, x_{H_k^-}, x_{u_j}, x_W) \\
&\quad + \frac{1}{4}\frac{x_{u_i}x_{u_j}}{\sin^4\beta}(Z_H^{2k})^2(Z_H^{2l})^2 f_{p^2}(x_{u_i}, x_{H_k^-}, x_{u_j}, x_{H_l^-}) + Z_{i\alpha}^\lambda Z_{j\beta}^{\lambda*} Z_{j\beta}^\eta Z_{i\alpha}^{\eta*} f_{p^2}(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-})) \\
&\quad + \frac{1}{32G_F^2 m_W^4} |\lambda_1|^4 (|(W_{b'}^\dagger)_{\delta,1}|^2 |(W_{b'}^\dagger)_{\varepsilon,1}|^2 |(Z_X)_{1,l}|^2 |(Z_X)_{1,k}|^2 f_{p^2}(x_{b'_\delta}, x_{X_k}, x_{b'_\varepsilon}, x_{X_l}) \\
&\quad + |(Z_{b'}^\dagger)_{3\rho}|^2 |(Z_{b'}^\dagger)_{3\sigma}|^2 f_{p^2}(x_{\tilde{X}}, x_{b'_\rho}, x_{\tilde{X}}, x_{b'_\sigma})) \\
C_2 &= V_{ib}V_{id}^*V_{jb}V_{jd}^* (\frac{1}{4}\frac{\sqrt{x_b x_d}}{\sin^2\beta \cos^2\beta} (x_{u_i} + x_{u_j}) Z_H^{1k} Z_H^{2k} Z_H^{1l} Z_H^{2l} f_{p^2}(x_{u_i}, x_{H_k^-}, x_{u_j}, x_{H_l^-}) \\
&\quad + (Z_{i\alpha}^{d\lambda} Z_{j\beta}^{b\lambda*} Z_{j\beta}^\eta Z_{i\alpha}^{\eta*} + Z_{i\alpha}^\lambda Z_{j\beta}^{\lambda*} Z_{j\beta}^{d\eta} Z_{i\alpha}^{b\eta*}) f_{p^2}(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-}) \\
&\quad - 2\sqrt{x_{\kappa_\lambda^-} x_{\kappa_\eta^-}} (Z_{i\alpha}^\lambda Z_{j\beta}^{b\lambda*} Z_{j\beta}^{d\eta} Z_{i\alpha}^{\eta*} + Z_{i\alpha}^{d\lambda} Z_{j\beta}^{\lambda*} Z_{j\beta}^\eta Z_{i\alpha}^{b\eta*}) f_1(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-})) \\
&\quad + \frac{1}{64G_F^2 m_W^4} |\lambda_1|^2 |\lambda_3|^2 (|(U_{b'}^\dagger)_{\delta 2}|^2 |(W_{b'}^\dagger)_{\varepsilon 1}|^2 (Z_X)_{1,k} (Z_X)_{2,k}^* (Z_X)_{1,l}^* (Z_X)_{2,l} \\
&\quad + |(W_{b'}^\dagger)_{\delta 1}|^2 |(U_{b'}^\dagger)_{\varepsilon 2}|^2 (Z_X)_{1,k}^* (Z_X)_{2,k} (Z_X)_{1,l} (Z_X)_{2,l}^*) f_{p^2}(x_{b'_\delta}, x_{X_k}, x_{b'_\varepsilon}, x_{X_l}) \\
&\quad + 2((Z_{b'}^\dagger)_{3\rho}^* (Z_{b'}^\dagger)_{3\sigma} (Z_{b'}^\dagger)_{4\rho}^* (Z_{b'}^\dagger)_{4\sigma} + (Z_{b'}^\dagger)_{3\rho} (Z_{b'}^\dagger)_{3\sigma}^* (Z_{b'}^\dagger)_{4\rho} (Z_{b'}^\dagger)_{4\sigma}^*) f_{p^2}(x_{\tilde{X}}, x_{b'_\rho}, x_{\tilde{X}}, x_{b'_\sigma}) \\
&\quad - 2(|(Z_{b'}^\dagger)_{3\rho}|^2 |(Z_{b'}^\dagger)_{4\sigma}|^2 + |(Z_{b'}^\dagger)_{3\sigma}|^2 |(Z_{b'}^\dagger)_{4\rho}|^2) x_{\tilde{X}} f_1(x_{\tilde{X}}, x_{b'_\rho}, x_{\tilde{X}}, x_{b'_\sigma}))
\end{aligned}$$

$$\begin{aligned}
C_3 = & V_{ib} V_{id}^* V_{jb} V_{jd}^* (- 2 \frac{\sqrt{x_b x_d}}{\cos^2 \beta} (Z_H^{1k})^2 f_{p^2}(x_{u_i}, x_{H_k^-}, x_{u_j}, x_W) \\
& + \frac{x_{u_i} x_{u_j} \sqrt{x_b x_d}}{\sin^2 \beta \cos^2 \beta} ((Z_H^{1k})^2 (Z_H^{2l})^2 + (Z_H^{2k})^2 (Z_H^{1l})^2) f_1(x_{u_i}, x_{H_k^-}, x_{u_j}, x_{H_l^-}) \\
& + 4 \sqrt{x_{\kappa_\lambda^-} x_{\kappa_\eta^-}} (Z_{i\alpha}^\lambda Z_{j\beta}^{b\lambda*} Z_{j\beta}^{d\eta} Z_{i\alpha}^{\eta*} + Z_{i\alpha}^{d\lambda} Z_{j\beta}^{\lambda*} Z_{j\beta}^\eta Z_{i\alpha}^{b\eta*}) f_1(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-}) \\
& + \frac{1}{16 G_F^2 m_W^4} |\lambda_1|^2 |\lambda_3|^2 (((W_{b'}^\dagger)_{\delta 1} (W_{b'}^\dagger)_{\varepsilon 1}^* (U_{b'}^\dagger)_{\delta 2}^* (U_{b'}^\dagger)_{\varepsilon 2}) (Z_X)_{2,k} |^2 | (Z_X)_{1,l} |^2 \\
& + (W_{b'}^\dagger)_{\delta 1}^* (W_{b'}^\dagger)_{\varepsilon 1} (U_{b'}^\dagger)_{\delta 2} (U_{b'}^\dagger)_{\varepsilon 2}^* | (Z_X)_{1,k} |^2 | (Z_X)_{2,l} |^2) \sqrt{x_{b'_\delta} x_{b'_\varepsilon}} f_1(x_{b'_\delta}, x_{X_k}, x_{b'_\varepsilon}, x_{X_l}) \\
& + (|(Z_{\tilde{b}'}^*)_{3\rho}|^2 |(Z_{\tilde{b}'}^*)_{4\sigma}|^2 + |(Z_{\tilde{b}'}^*)_{3\rho}|^2 |(Z_{\tilde{b}'}^*)_{4\rho}|^2) x_{\tilde{X}} f_1(x_{\tilde{X}}, x_{\tilde{b}'_\rho}, x_{\tilde{X}}, x_{\tilde{b}'_\sigma}))
\end{aligned}$$

$$\begin{aligned}
C_4 = & V_{ib} V_{id}^* V_{jb} V_{jd}^* (\frac{x_{u_i} x_{u_j} x_d}{\sin^2 \beta \cos^2 \beta} Z_H^{1k} Z_H^{2k} Z_H^{1l} Z_H^{2l} f_1(x_{u_i}, x_{H_k^-}, x_{u_j}, x_{H_l^-}) \\
& + \sqrt{x_{\kappa_\lambda^-} x_{\kappa_\eta^-}} Z_{i\alpha}^{d\lambda} Z_{j\beta}^{\lambda*} Z_{j\beta}^{d\eta} Z_{i\alpha}^{\eta*} f_1(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-}) \\
& + \frac{1}{32 G_F^2 m_W^4} \lambda_1^2 \lambda_3^2 (4 (W_{b'}^\dagger)_{\delta 1} (W_{b'}^\dagger)_{\varepsilon 1} (U_{b'}^\dagger)_{\delta 2}^* (U_{b'}^\dagger)_{\varepsilon 2}^* (Z_X)_{1,k} (Z_X)_{2,k}^* (Z_X)_{1,l} (Z_X)_{2,l}^* \\
& \times \sqrt{x_{b'_\delta} x_{b'_\varepsilon}} f_1(x_{b'_\delta}, x_{X_k}, x_{b'_\varepsilon}, x_{X_l}) + (Z_{\tilde{b}'}^*)_{3\rho} (Z_{\tilde{b}'}^*)_{3\sigma} (Z_{\tilde{b}'}^*)_{4\rho} (Z_{\tilde{b}'}^*)_{4\sigma} x_{\tilde{X}} f_1(x_{\tilde{X}}, x_{\tilde{b}'_\rho}, x_{\tilde{X}}, x_{\tilde{b}'_\sigma}))
\end{aligned}$$

$$\begin{aligned}
C_5 = & -\frac{1}{4} V_{ib} V_{id}^* V_{jb} V_{jd}^* \sqrt{x_{\kappa_\lambda^-} x_{\kappa_\eta^-}} Z_{i\alpha}^{d\lambda} Z_{j\beta}^{\lambda*} Z_{j\beta}^{d\eta} Z_{i\alpha}^{\eta*} f_1(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-}) \\
& - \frac{1}{128 G_F^2 m_W^4} \lambda_1^2 \lambda_3^2 (Z_{\tilde{b}'}^*)_{3\rho} (Z_{\tilde{b}'}^*)_{3\sigma} (Z_{\tilde{b}'}^*)_{4\rho} (Z_{\tilde{b}'}^*)_{4\sigma} f_1(x_{\tilde{X}}, x_{\tilde{b}'_\rho}, x_{\tilde{X}}, x_{\tilde{b}'_\sigma})
\end{aligned}$$

$$\begin{aligned}
C_6 = & V_{ib} V_{id}^* V_{jb} V_{jd}^* \left(\frac{1}{4} \frac{x_b x_d}{\sin^2 \beta \cos^2 \beta} (Z_H^{1k})^2 (Z_H^{1l})^2 f_{p^2}(x_{u_i}, x_{H_k^-}, x_{u_j}, x_{H_l^-}) \right. \\
& + Z_{i\alpha}^{d\lambda} Z_{j\beta}^{b\lambda*} Z_{j\beta}^{d\eta} Z_{i\alpha}^{b\eta*} f_{p^2}(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-}) \\
& + \frac{1}{32 G_F^2 m_W^4} |\lambda_3|^4 (|(U_{b'}^\dagger)_{\delta 2}|^2 |(U_{b'}^\dagger)_{\varepsilon 2}|^2 |(Z_X)_{2,k}|^2 |(Z_X)_{2,l}|^2 f_{p^2}(x_{b'_\delta}, x_{X_k}, x_{b'_\varepsilon}, x_{X_l}) \\
& + |(Z_{\tilde{b}'})_{4\rho}|^2 |(Z_{\tilde{b}'})_{4\sigma}|^2 f_{p^2}(x_{\tilde{X}}, x_{\tilde{b}'_\rho}, x_{\tilde{X}}, x_{\tilde{b}'_\sigma}) \left. \right) \\
C_7 = & V_{ib} V_{id}^* V_{jb} V_{jd}^* \left(\frac{x_{u_i} x_{u_j} x_b}{\sin^2 \beta \cos^2 \beta} Z_H^{1k} Z_H^{2k} Z_H^{1l} Z_H^{2l} f_1(x_{u_i}, x_{H_k^-}, x_{u_j}, x_{H_l^-}) \right. \\
& + \sqrt{x_{\kappa_\lambda^-} x_{\kappa_\eta^-}} Z_{i\alpha}^\lambda Z_{j\beta}^{b\lambda*} Z_{j\beta}^\eta Z_{i\alpha}^{b\eta*} f_1(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-}) \\
& + \frac{1}{32 G_F^2 m_W^4} (\lambda_1^*)^2 (\lambda_3^*)^2 (4(W_{b'}^\dagger)_{\delta 1}^* (W_{b'}^\dagger)_{\varepsilon 1}^* (U_{b'}^\dagger)_{\delta 2} (U_{b'}^\dagger)_{\varepsilon 2} (Z_X)_{1,k}^* (Z_X)_{2,k} (Z_X)_{1,l}^* (Z_X)_{2,l} \\
& \times \sqrt{x_{b'_\delta} x_{b'_\varepsilon}} f_1(x_{b'_\delta}, x_{X_k}, x_{b'_\varepsilon}, x_{X_l}) + (Z_{\tilde{b}'})_{3\rho} (Z_{\tilde{b}'})_{3\sigma} (Z_{\tilde{b}'})_{4\rho} (Z_{\tilde{b}'})_{4\sigma} f_1(x_{\tilde{X}}, x_{\tilde{b}'_\rho}, x_{\tilde{X}}, x_{\tilde{b}'_\sigma}) \left. \right) \\
C_8 = & -\frac{1}{4} V_{ib} V_{id}^* V_{jb} V_{jd}^* \sqrt{x_{\kappa_\lambda^-} x_{\kappa_\eta^-}} Z_{i\alpha}^\lambda Z_{j\beta}^{b\lambda*} Z_{j\beta}^\eta Z_{i\alpha}^{b\eta*} f_1(x_{\tilde{u}_\alpha^i}, x_{\tilde{\chi}_\lambda^-}, x_{\tilde{u}_\beta^j}, x_{\tilde{\chi}_\eta^-}) \\
& - \frac{1}{128 G_F^2 m_W^4} (\lambda_1^*)^2 (\lambda_3^*)^2 (Z_{\tilde{b}'})_{3i} (Z_{\tilde{b}'})_{3j} (Z_{\tilde{b}'})_{4i} (Z_{\tilde{b}'})_{4j} f_1(x_{\tilde{X}}, x_{\tilde{b}'_\rho}, x_{\tilde{X}}, x_{\tilde{b}'_\sigma})
\end{aligned} \tag{29}$$

For convenience, we have defined the ratio of mass square as: $x_i = m_i^2/m_W^2$, and here $Z_{i\alpha}^\lambda$, $Z_{i\alpha}^{d\lambda} \dots$ have been defined as

$$\begin{aligned}
Z_{i\alpha}^\lambda = & -Z_{\tilde{U}_\alpha^i}^{1\alpha} Z_+^{1\lambda*} + \frac{\sqrt{2x_{u_i}}}{2 \sin \beta} Z_{\tilde{U}_\alpha^i}^{2\alpha} Z_+^{2\lambda*} \\
Z_{i\alpha}^{d\lambda} = & \frac{\sqrt{2x_d}}{2 \cos \beta} Z_{\tilde{U}_\alpha^i}^{1\alpha} Z_-^{2\lambda} \\
& \dots
\end{aligned} \tag{30}$$

Here f_1 and f_{p^2} are the functions related to the one-loop integral functions.

$$\mu^{2\epsilon} \int \frac{d^D P}{(2\pi)^D} \frac{1}{p^2 - m_1^2} \frac{1}{p^2 - m_2^2} \frac{1}{p^2 - m_3^2} \frac{1}{p^2 - m_4^2} = \frac{1}{16\pi^2 m_W^4} f_1(x_1, x_2, x_3, x_4) \quad (31)$$

$$\mu^{2\epsilon} \int \frac{d^D P}{(2\pi)^D} \frac{1}{p^2 - m_1^2} \frac{1}{p^2 - m_2^2} \frac{1}{p^2 - m_3^2} \frac{1}{p^2 - m_4^2} p^2 = \frac{1}{16\pi^2 m_W^2} f_{p^2}(x_1, x_2, x_3, x_4) \quad (32)$$

The analytical expressions for the functions $f_{p^2}(x_1, x_2, x_3, x_4)$ and $f_1(x_1, x_2, x_3, x_4)$ are listed in Appendix A. It should be noted that we need perform summation over the repeated indices in the calculations.

The matching scale is chosen as $\mu_0 = \mu_W$ in our calculations. Now we should evolve the coefficients from the scale μ_W down to the B -meson scale μ_b

$$\left[\mu \frac{\partial}{\partial \mu} + \beta(\alpha_S) \frac{\partial}{\partial \alpha_S} - \frac{\hat{\gamma}^T}{2} \right] \vec{C}(\mu, \alpha_S) = 0. \quad (33)$$

By solving the renormalization group equation [114], we have

$$\vec{C}(\mu_b) = W(\mu_b, \mu_W) \vec{C}(\mu_W) \quad (34)$$

with

$$W(\mu_b, \mu_W) = \left[\frac{\alpha_S(m_W)}{\alpha_S(m_b)} \right]^{\frac{\gamma^{(0)}}{2\beta_0}}, \quad (35)$$

where $\gamma^{(0)}$ is the anomalous dimensions matrix (ADM) [114, 115], and $\beta_0 = \frac{11N_c - 2n_f}{3}$ with N_c denoting the number of colors and n_f denoting the number of active quark flavors.

The mass difference of $B^0 - \bar{B}^0$ mixing can be expressed as

$$\Delta m_B = \frac{|\langle \bar{B}^0 | H_{\text{eff}}(\Delta B = 2) | B^0 \rangle|}{m_B}. \quad (36)$$

After substituting Eq. (26) into the above equation, at B -meson scale, the mass difference Δm_B can be written by

$$\Delta m_B = \frac{1}{4} \frac{G_F^2}{\pi^2} m_W^2 \sum_{\alpha=1}^8 \frac{|C_\alpha(\mu_b) \langle \bar{B}^0 | \mathcal{O}_\alpha(\mu_b) | B^0 \rangle|}{m_B}, \quad (37)$$

where, the matrix elements $\langle \bar{B}^0 | \mathcal{O}_\alpha | B^0 \rangle$ require non-perturbative QCD calculations by the lattice Monte Carlo estimates. The matrix element is parameterized as $\langle \bar{B}^0 | \mathcal{O}_1 | B^0 \rangle = \frac{2}{3} B_B(\mu) f_B^2 m_B^2$, and the other hadronic matrix elements parameterized are listed in Appendix B.

IV. THE NUMERICAL ANALYSIS

In our calculations for the CKM matrix, we apply the Wolfenstein parametrization and set $A = 0.81$, $\lambda = 0.22$, $\rho = 0.135$, $\eta = 0.349$. For the hadronic matrix element, the recent average of the lattice results is $f_{B_d} \sqrt{B_{B_d}} = 216 \pm 15$ (MeV) [116], and we adopt the central value of the $f_{B_d} \sqrt{B_{B_d}}$ in our calculations. The other SM parameters are chosen as $m_W = 80.385$ GeV, $m_u = 2.3 \times 10^{-3}$ GeV, $m_c = 1.275$ GeV, $m_t = 173.5$ GeV, $m_b = 4.18$ GeV, $m_d = 4.8 \times 10^{-3}$ GeV, $m_B = 5.279$ GeV, $G_F = 1.166 \times 10^{-5}$ GeV $^{-2}$, $\alpha_S(m_W) = 0.12$, $\alpha_S(m_b) = 0.22$ [86].

Now we investigate the numerically behavior of these parameters to the $B^0 - \bar{B}^0$ mixing in BLMSSM. This model contains many parameters. In our following discussions, the parameters needed to study contain $\lambda_{1,3}$, μ_B , m_{Z_B} , m_{D_5} , μ_X . The other parameters are adopted as Refs. [66, 67] which have been analyzed in the signals of decay channels $h \rightarrow \gamma\gamma$ and $h \rightarrow VV^*$ ($V = Z, W$) with the Higgs mass around 125 GeV.

$$\begin{aligned}
m_{\tilde{Q}_{1,2,3}} &= m_{\tilde{U}_{1,2,3}} = m_{\tilde{D}_{1,2,3}} = 1 \text{ TeV}, \\
A_{d,s,b} &= A_{u,c,t} = -1 \text{ TeV}, \\
M_2 &= 750 \text{ GeV}, \\
B_4 &= L_4 = \frac{3}{2}, \\
\tan \beta &= \tan \beta_B = \tan \beta_L = 2, \\
\mu &= -800 \text{ GeV}, \\
m_{\tilde{U}_4} &= m_{\tilde{D}_4} = m_{\tilde{U}_5} = 1 \text{ TeV}, \\
m_{\tilde{L}_4} &= m_{\tilde{\nu}_4} = m_{\tilde{E}_4} = m_{\tilde{L}_5} = m_{\tilde{\nu}_5} = m_{\tilde{E}_5} = 1 \text{ TeV}, \\
A_{\nu_4} &= A_{e_4} = A_{\nu_5} = A_{\nu_5} = A_{u_4} = A_{u_5} = A_{d_4} = A_{d_5} = 550 \text{ GeV},
\end{aligned}$$

$$\begin{aligned}
v_{B_t} &= \sqrt{v_B^2 + \bar{v}_B^2} = 3 \text{ TeV} , \\
A_{BQ} &= A_{BU} = A_{BD} = 1 \text{ TeV}, \\
Y_{u_4} &= 0.76Y_t, \\
Y_{d_4} &= Y_{u_5} = 0.7Y_b, \\
Y_{d_5} &= 0.13Y_t, \\
\lambda_Q &= \lambda_u = \lambda_d = 0.5, \\
m_{\nu_4} &= m_{\nu_5} = 90 \text{ GeV}, \\
m_{e_4} &= m_{e_5} = B_X = 100 \text{ GeV}, \\
m_{\tilde{Q}_4} &= 790 \text{ GeV}.
\end{aligned} \tag{38}$$

In order to see the dependence of the mass difference Δm_B on the parameters space in the BLMSSM, we fix $m_{\tilde{Q}_5} = 1 \text{ TeV}$, $m_{\tilde{D}_5} = 1 \text{ TeV}$, $\mu_B = 500 \text{ GeV}$, $\mu_X = 2.4 \text{ TeV}$, $m_{Z_B} = 1 \text{ TeV}$. From the Wilson coefficients listed in Section III, one can see that the mass difference Δm_B is the continuous function of the parameters λ_1 and λ_3 , and because of the fourth power of $\lambda_{1,3}$ the Δm_B should remarkably increase with the increasing of $|\lambda_1|$ and $|\lambda_3|$. So λ_1 and λ_3 play an important role to the theoretical prediction on Δm_B . Next, the influence of the parameters $\lambda_{1,3}$ to Δm_B will be discussed in detail. We plot the contours corresponding to the mass difference Δm_B in the parameter space of λ_1 and λ_3 in Fig. 3. We can see that Δm_B increases as $|\lambda_{1,3}|$ increases, and sensitively depends on $|\lambda_{1,3}|$ when $|\lambda_1|$ and $|\lambda_3|$ are both larger than 0.2. As one can see, the values of $|\lambda_1|$ and $|\lambda_3|$ that all is larger than 0.25 are disfavored by experiment results under this given assumption.

Next, we investigate the dependence of Δm_B on the parameter m_{Z_B} . In Fig. 4, we plot Δm_B varying with the mass of neutral $U(1)_B$ gauge boson Z_B , when $\lambda_1 = 0.25$ and $\lambda_3 = 0.2$. The figure shows that Δm_B decreases as the m_{Z_B} increases. However, it should be noted that the value of the m_{Z_B} should not be too large, in order to avoid some tachyons appearing, as well as to coincide with the current experimental result on the mass of squarks. Actually, the corrections of some other parameters to Δm_B are small, such as $m_{\tilde{D}_4}$, $m_{\tilde{Q}_4}$ and B_X , which we would not discuss in this paper.

In the following discussions, we choose $\lambda_1 = 0.2$ for simplicity. Now, we investigate the




FIG. 3: (Color online) Contour plots of Δm_B in the parameter space of λ_1 and λ_3 .




FIG. 4: The mass difference Δm_B versus the new gauge boson mass m_{Z_B} .

dependence of Δm_B on the parameter μ_B . Considering that μ_B is the mass parameter of the "brand new" Higgs superfields Φ_B and ϕ_B , the behavior of the Δm_B versus μ_B when $\lambda_3 = 0.25$ is shown in Fig. 5. The numerical result shows that the contribution of the




FIG. 5: The mass difference Δm_B as a function of μ_B .

parameter μ_B to Δm_B is quite small, when μ_B is lighter than 500 GeV. When μ_B is heavier than 500 GeV, Δm_B decreases sharply with the increasing of μ_B .

We plot Δm_B as a function of the exotic right-handed soft-SUSY-breaking squark mass $m_{\tilde{D}_5}$ for three values of λ_3 in Fig. 6, the dotted line corresponds to the result of $\lambda_3 = 0.2$, the dashed line corresponds to the result of $\lambda_3 = 0.25$, the dot-dashed line corresponds to the result of $\lambda_3 = 0.3$. The light gray area denotes the Δm_B^{SM} at 1σ , and the gray area denotes the Δm_B^{Exp} at 1σ . As one can see, Δm_B decreases along with the increasing of $m_{\tilde{D}_5}$ for a given value of λ_3 . Fig. 6 also exhibits that Δm_B has a strong dependence on $m_{\tilde{D}_5}$ for large values of λ_3 . However, this figure indicates that the Δm_B declines slowly with the increasing of $m_{\tilde{D}_5}$, when the value of λ_3 is small. Generally speaking, the influence of the $m_{\tilde{D}_5}$ to Δm_B can be neglected as λ_3 is enough small. Considering the constraint from the Δm_B^{SM} at 1σ , one can see that small values of $m_{\tilde{D}_5}$ can be excluded for large value of λ_3 as well as large values of $m_{\tilde{D}_5}$ can be excluded for small value of λ_3 under the given assumption.

In Fig. 7, we study the dependence of Δm_B on the particle $\tilde{X}$ mass μ_X . The dotted line corresponds to the result when $\lambda_3 = 0.2$, the solid line corresponds to the result when $\lambda_3 = 0.25$, the dashed line corresponds to the result when $\lambda_3 = 0.3$. The light gray area




FIG. 6: The mass difference Δm_B varies with the parameter $m_{\tilde{D}_5}$ for three values of λ_3 . The light gray area denotes the Δm_B^{SM} at 1σ , and the gray area denotes the Δm_B^{Exp} at 1σ .




FIG. 7: The mass difference Δm_B as a function of $\tilde{X}$ mass μ_X for three values of λ_3 . The light gray area denotes the Δm_B^{SM} at 1σ , and the gray area denotes the Δm_B^{Exp} at 1σ .

denotes the Δm_B^{SM} at 1σ , and the gray area denotes the Δm_B^{Exp} at 1σ . It clearly shows a large influence of the new particle $\tilde{X}$ on the mixing of $B^0 - \bar{B}^0$. The mass difference Δm_B decreases with increasing of the μ_X in a very similar manner as that in Fig. 6. We find the mass of the exotic particle $\tilde{X}$ should not be too light for large values of λ_3 , however, the heavy mass of the exotic particle μ_X is also constrained for small values of λ_3 .

V. CONCLUSIONS

With the constraint of a 125 GeV Higgs, we analyze the correction of the extra fermions and scalars to $B^0 - \bar{B}^0$ mixing in the extension of the MSSM where baryon number and lepton number are local gauge symmetries. In this framework, the new particles' LO correction to $B^0 - \bar{B}^0$ mixing is significant in some parameter space. The numerical evaluations indicate that the parameters $\lambda_{1,3}$, $m_{\tilde{D}_5}$ and μ_X are sensitive to the process of $B^0 - \bar{B}^0$ mixing. It is well known that the space that is left for hiding some new physics effects in the $B^0 - \bar{B}^0$ mixing is mainly given by the theoretical error. With the development of more precise theoretical analysis (especially the lattice calculations) and accurate experimental measurements, the $B^0 - \bar{B}^0$ mixing in the BLMSM will have a clearer picture and the parameters space will also be further constrained.

Many experiments have been performed to search for baryon number violation (BNV). Belle and BaBar have obtained the upper limits on the branching fraction of BNV τ decays $\tau^- \rightarrow \Lambda \pi^-$ and $\tau^- \rightarrow \Lambda k^-$ [117, 118]. Some B meson decays $B^0 \rightarrow \Lambda_c^+ l^-$, $B^- \rightarrow \Lambda l^-$ and $B^- \rightarrow \bar{\Lambda} l^-$ have been investigated by BaBar [119]. Charged lepton flavour violation (CLFV) and BNV decays $\tau^- \rightarrow \bar{p} \mu^+ \mu^-$ and $\tau^- \rightarrow p \mu^- \mu^-$ have been carried out by LHCb [120]. Searching for baryon number violation in top-quark decays has been done by CMS [121]. However, these experimental searches for BNV have yield only upper limits. On the other hand, the branching fractions of CLFV process ($\mu \rightarrow e \gamma$, $\mu \rightarrow e e e$, $\tau \rightarrow l \gamma$ and $\tau \rightarrow l l l$ (with $l = e, \mu$), et al.) are predicted very small in the SM. For instance, the SM prediction for branching fractions in muon decays is smaller than 10^{-50} . In the BLMSM, there are some new contributions to these BNV and CLFV processes. And the contributions of BLMSM may significantly enhance these branching fractions. One can have BNV signals from the

decays of squarks and gauginos without conflict with the current experiments. For instance, if the gluino is the lightest supersymmetric particle one could have signals with multitops and multibottoms such as $pp \rightarrow \tilde{g}\tilde{g} \rightarrow ttbbjj$ (j stands for a light jet), which may be observed at the LHC [63, 64]. The projected sensitivity for future experiments that searching for the CLFV processes will be largely improved [122–128]. And the running of LHC will resume in 2015 with higher energy and luminosity. So, it would be interesting to investigate this model. Any observation of BNV or CLFV whose branching fractions is large than that of SM prediction would be a clear sign for BSM physics. Investigating these BNV and CLFV processes can test the BLMSSM and provide constraints on the parameter space.

Acknowledgements

The work has been supported by the National Natural Science Foundation of China (NNSFC) with Grant No. 11275036, No. 11047002, the open project of State Key Laboratory of Mathematics-Mechanization with Grant No. Y3KF311CJ1, the Natural Science Foundation of Hebei province with Grant No. A2013201277, and Natural Science Fund of Hebei University with Grant No. 2011JQ05, No. 2012-242.

Appendix A: Integral function

The functions related to the one-loop integral functions are given as

$$f_1(x_1, x_2, x_3, x_4) = \left\{ \begin{array}{l} -\frac{\log(x_1)x_1}{(x_1-x_2)(x_1-x_3)(x_1-x_4)} + \frac{\log(x_2)x_2}{(x_1-x_2)(x_2-x_3)(x_2-x_4)} \\ + \frac{\log(x_3)x_3}{(x_1-x_3)(x_3-x_2)(x_3-x_4)} + \frac{\log(x_4)x_4}{(x_1-x_4)(x_4-x_2)(x_4-x_3)}, (x_1 \neq x_3 \text{ and } x_2 \neq x_4) \\ \\ -\frac{\log(x_1)x_1}{(x_1-x_2)^2(x_1-x_3)} + \frac{\log(x_3)x_3}{(x_1-x_3)(x_2-x_3)^2} \\ + \frac{\log(x_2)(x_2^2-x_1x_3)}{(x_1-x_2)^2(x_2-x_3)^2} + \frac{1}{(x_1-x_2)(x_2-x_3)}, (x_1 \neq x_3 \text{ and } x_2 = x_4) \\ \\ -\frac{\log(x_2)x_2}{(x_1-x_2)^2(x_2-x_4)} + \frac{\log(x_4)x_4}{(x_1-x_4)^2(x_2-x_4)} \\ + \frac{\log(x_1)(x_1^2-x_2x_4)}{(x_1-x_2)^2(x_1-x_4)^2} - \frac{1}{(x_1-x_2)(x_1-x_4)}, (x_1 = x_3 \text{ and } x_2 \neq x_4) \\ \\ \frac{2\log(x_1)x_1x_2}{(x_1-x_2)^3} - \frac{2\log(x_2)x_1x_2}{(x_1-x_2)^3} - \frac{x_1+x_2}{(x_1-x_2)^2}, (x_1 = x_3 \text{ and } x_2 = x_4) \end{array} \right. \quad (\text{A1})$$

$$f_{p^2}(x_1, x_2, x_3, x_4) = \left\{ \begin{array}{l} -\frac{\log x_1^2}{(x_1-x_2)(x_1-x_3)(x_1-x_4)} + \frac{\log x_2^2}{(x_1-x_2)(x_2-x_3)(x_2-x_4)} \\ + \frac{\log x_3^2}{(x_1-x_3)(x_3-x_2)(x_3-x_4)} + \frac{\log x_4^2}{(x_1-x_4)(x_4-x_2)(x_4-x_3)}, (x_1 \neq x_3 \text{ and } x_2 \neq x_4) \\ \\ -\frac{\log x_1^2}{(x_1-x_2)^2(x_1-x_3)} + \frac{\log x_3^2}{(x_1-x_3)(x_2-x_3)^2} \\ + \frac{\log x_2(x_2^2-x_1x_3)}{(x_1-x_2)^2(x_2-x_3)^2} + \frac{1}{(x_1-x_2)(x_2-x_3)}, (x_1 \neq x_3 \text{ and } x_2 = x_4) \\ \\ -\frac{\log x_2^2}{(x_1-x_2)^2(x_2-x_4)} + \frac{\log x_4^2}{(x_1-x_4)^2(x_2-x_4)} \\ + \frac{\log x_1(x_1^2-x_2x_4)}{(x_1-x_2)^2(x_1-x_4)^2} - \frac{1}{(x_1-x_2)(x_1-x_4)}, (x_1 = x_3 \text{ and } x_2 \neq x_4) \\ \\ \frac{2\log(x_1)x_1x_2}{(x_1-x_2)^3} - \frac{2\log(x_2)x_1x_2}{(x_1-x_2)^3} - \frac{x_1+x_2}{(x_1-x_2)^2}, (x_1 = x_3 \text{ and } x_2 = x_4) \end{array} \right. \quad (\text{A2})$$

Appendix B: Hadronic matrix elements

The hadronic matrix elements can be written as

$$\langle \bar{B}^0 | \mathcal{O}_1 | B^0 \rangle = \frac{2}{3} B_B(\mu) f_B^2 m_B^2$$

$$\langle \bar{B}^0 | \mathcal{O}_2 | B^0 \rangle = -\frac{1}{6} B_B(\mu) f_B^2 m_B^2$$

$$\langle \bar{B}^0 | \mathcal{O}_3 | B^0 \rangle = -\frac{5}{12} B_B(\mu) f_B^2 m_B^2$$

$$\langle \bar{B}^0 | \mathcal{O}_4 | B^0 \rangle = \frac{5}{12} B_B(\mu) f_B^2 m_B^2$$

$$\langle \bar{B}^0 | \mathcal{O}_5 | B^0 \rangle = \frac{1}{2} B_B(\mu) f_B^2 m_B^2$$

$$\langle \bar{B}^0 | \mathcal{O}_6 | B^0 \rangle = \frac{2}{3} B_B(\mu) f_B^2 m_B^2$$

$$\langle \bar{B}^0 | \mathcal{O}_7 | B^0 \rangle = \frac{5}{12} B_B(\mu) f_B^2 m_B^2$$

$$\langle \bar{B}^0 | \mathcal{O}_8 | B^0 \rangle = \frac{1}{2} B_B(\mu) f_B^2 m_B^2.$$

Here f_B is the B -meson decay constant, B_B is the bag parameter.

-
- [1] H.P. Nilles, Phys. Rep. 110 (1984) 1.
 - [2] H.E. Haber and G.L. Kane, Phys. Rept. 117 (1985) 75.
 - [3] J. Rosiek, Phys. Rev. D 41 (1990) 3464 [hep-ph/9511250].
 - [4] H.E. Haber, hep-ph/9306207.
 - [5] S.P. Martin, hep-ph/9709356.
 - [6] G. R. Farrar and P. Fayet, Phys. Lett. B 76 (1978) 575.
 - [7] Supersymmetry Searches at CMS, <http://cms.web.cern.ch/org/cms-papers-and-results>.
 - [8] Supersymmetry Searches at ATLAS, <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPubl>

- [9] P. Bechtle, B. Sarrazin, K. Desch, H. K. Dreiner, P. Wienemann, M. Kramer, C. Robens and B. OLeary, Phys. Rev. D 84 (2011) 011701 [arXiv:1102.4693].
- [10] B. C. Allanach, T. J. Khoo and K. Sakurai, JHEP 1111 (2011) 132 [arXiv:1110.1119].
- [11] O. Buchmuller, R. Cavanaugh, A. De Roeck, M. J. Dolan, J. R. Ellis, H. Flacher, S. Heinemeyer and G. Isidori et al., Eur. Phys. J. C 72 (2012) 1878 [arXiv:1110.3568].
- [12] P. Bechtle, T. Bringmann, K. Desch, H. Dreiner, M. Hamer, C. Hensel, M. Kramer and N. Nguyen et al., JHEP 1206 (2012) 098 [arXiv:1204.4199 [hep-ph]].
- [13] G. Kane, P. Kumar, R. Lu and B. Zheng, Phys. Rev. D 85, 075026 (2012) [arXiv:1112.1059 [hep-ph]].
- [14] H. Baer, V. Barger and A. Mustafayev, Phys. Rev. D 85 (2012) 075010 [arXiv:1112.3017 [hep-ph]].
- [15] J. L. Feng, K. T. Matchev and D. Sanford, Phys.Rev. D 85 (2012) 075007 [arXiv:1112.3021 [hep-ph]].
- [16] S. Heinemeyer, O. Stal and G. Weiglein, Phys. Lett. B 710 (2012) 201 [arXiv:1112.3026 [hep-ph]].
- [17] A. Arbey, M. Battaglia, A. Djouadi, F. Mahmoudi and J. Quevillon, Phys. Lett. B 708(2012) 162-169 [arXiv:1112.3028 [hep-ph]].
- [18] A. Arbey, M. Battaglia and F. Mahmoudi, Eur. Phys. J. C 72 (2012) 1906 [arXiv:1112.3032 [hep-ph]].
- [19] P. Draper, P. Meade, M. Reece and D. Shih, Phys. Rev. D 85, 095007 (2012) [arXiv:1112.3068 [hep-ph]].
- [20] U. Ellwanger, JHEP 03 (2012) 044 [arXiv:1112.3548 [hep-ph]].
- [21] S. Akula, B. Altunkaynak, D. Feldman, P. Nath and G. Peim, Phys. Rev. D 85, 075001 (2012) [arXiv:1112.3645 [hep-ph]].
- [22] M. Kadastik, K. Kannike, A. Racioppi and M. Raidal, JHEP 1205 (2012) 061 [arXiv:1112.3647 [hep-ph]].
- [23] Junjie Cao, Zhaoxia Heng, Dongwei Li, Jin Min Yang, Phys.Lett. B 710 (2012) 665-670 [arXiv:1112.4391 [hep-ph]].
- [24] C.S. Aulakh, R.N. Mohapatra, Phys. Lett. B 119 (1982) 136.

- [25] L.J. Hall, M. Suzuki, Nucl. Phys. B 231 (1984) 419.
- [26] I.H. Lee, Nucl. Phys. B 246 (1984) 120.
- [27] S. Dawson, Nucl. Phys. B 261 (1985) 297.
- [28] G.G. Ross, J.W.F. Valle, Phys. Lett. B 151 (1985) 375.
- [29] R. Barbieri, A. Masiero, Nucl. Phys. B 267 (1986) 679.
- [30] S. Dimopoulos, L.J. Hall, Phys. Lett. B 207 (1987) 210.
- [31] R. Hempfling, Nucl. Phys. B 478 (1996) 3.
- [32] J. Erler, J.L. Feng, N. Polonsky, Phys. Rev. Lett. 78 (1997) 3063.
- [33] S. Roy, B. Mukhopadhyaya, Phys. Rev. D 55 (1997) 7020.
- [34] H.P. Nilles, N. Polonsky, Nucl. Phys. B 484 (1997) 33 [hep-ph/9606388].
- [35] H.K. Dreiner, in Kane, G.L. (ed.): Perspectives on supersymmetry II 565 [hep-ph/9707435].
- [36] M.A. Díaz, J.C. Romao, J.W.F. Valle, Nucl. Phys. B 524 (1998) 23 [hep-ph/9706315].
- [37] M. Hirsch, J.W.F. Valle, Nucl. Phys. B 557 (1999) 60 [hep-ph/9812463].
- [38] F. de Campos et al., hep-ph/9903245.
- [39] C.-H. Chang, T.-F. Feng, Eur. Phys. J. C 12 (2000) 137 [hep-ph/9901260].
- [40] C. Csaki, Y. Grossman and B. Heidenreich, Phys. Rev. D 85 (2012) 095009 [arXiv:1111.1239].
- [41] B.C. Allanach and B. Gripaios, JHEP 1205 (2012) 062 [arXiv:1202.6616].
- [42] H.K. Dreiner, K. Nickel, F. Staub and A. Vicente, Phys. Rev. D 86 (2012) 015003 [arXiv:1204.5925 [hep-ph]].
- [43] P.W. Graham, D. E. Kaplan, S. Rajendran and P. Saraswat, JHEP 1207 (2012) 149 [arXiv:1204.6038 [hep-ph]].
- [44] M. Hanussek and J. S. Kim, Phys. Rev. D 85 (2012) 115021 [arXiv:1205.0019 [hep-ph]].
- [45] H. Dreiner, W. Porod, F. Staub and A. Vicente, arXiv:1205.0557.
- [46] H.-B. Zhang, T.-F. Feng, S.-M. Zhao, T.-J. Gao, Nucl. Phys. B 873 (2013) 300, [arXiv:1304.6248].
- [47] H.-B. Zhang, T.-F. Feng, G.-F. Luo, Z.-F. Ge and S.-M. Zhao, JHEP 1307 (2013) 069 [Erratum *ibid.* 10 (2013) 173], [arXiv:1305.4352].
- [48] H.-B. Zhang, T.-F. Feng, F. Sun, K.-S. Sun, J.-B. Chen and S.-M. Zhao, arXiv:1307.3607.
- [49] P. Fileviez Pérez and M. B. Wise, Phys. Rev. D 82 (2010) 011901 [Erratum-*ibid.* D 82 (2010)

- 079901] [arXiv:1002.1754 [hep-ph]].
- [50] T.R. Dulaney, P. Fileviez Pérez, M. B. Wise, Phys. Rev. D 83 (2011) 023520. [arXiv:1005.0617 [hep-ph]].
 - [51] P.Fileviez Pérez and M. B. Wise, arXiv:1105.3190 [hep-ph].
 - [52] P. Ko and Y. Omura, arXiv:1012.4679 [hep-ph].
 - [53] P. Minkowski, Phys. Lett. 67 B, 421 (1977).
 - [54] T. Yanagida, in Proceedings of the Workshop on the Unified Theory and the Baryon by Number in the Universe, edited by O. Sawada et al. (KEK, Tsukuba, 1979), p. 95.
 - [55] M. Gell-Mann, P. Ramond, and R. Slansky, in Supergravity, edited P. van Nieuwenhuizen et al. (North-Holland, Amsterdam, 1979), p. 315.
 - [56] S.L. Glashow, in Quarks and Leptons, Cargese, edited by M. Le'vy et al. (Plenum, New York, 1980), p. 707.
 - [57] R.N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. 44, 912 (1980).
 - [58] P.F. Pérez, Phys. Lett. B 711 (2012) 353;
 - [59] R. Foot, G. C. Joshi and H. Lew, Phys. Rev. D 40, 2487 (1989).
 - [60] C.D. Carone and H. Murayama, Phys. Rev. D 52, 484 (1995) [arXiv:hep-ph/9501220].
 - [61] P.F. Pérez, Phys. Lett. B 711 (2012) 353.
 - [62] P.F. Pérez and M. B. Wise, Phys. Rev. D 82 (2010) 011901.
 - [63] P.F. Pérez and M. B. Wise, JHEP 1108 (2011) 068.
 - [64] P.F. Pérez and M. B. Wise, Phys. Rev. D 84 (2011) 055015.
 - [65] J. M. Arnold, P.F. Pérez, B. Fornal, and S. Spinner, Phys. Rev. D 85 (2012) 115024.
 - [66] Tai-Fu Feng, Shu-Min Zhao, Hai-Bin Zhang, Yin-Jie Zhang, Yu-Li Yan, Nucl. Phys. B 871 (2013) 223.
 - [67] Shu-Min Zhao, Tai-Fu Feng, Ben Yan, Hai-Bin Zhang, Yin-Jie Zhang, Biao Chen, Xue-Qian Li, JHEP 10 (2013) 020.
 - [68] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, Phys. Rev. Lett. 13 (1964) 138.
 - [69] K. Abe et al. (Belle Collaboration), Phys. Rev. Lett. 87, 091802 (2001).
 - [70] C. Albajar et al. (UA1 collaboration), Phys. Lett. B 186 (1987) 247.
 - [71] J.R. Ellis, J. S. Hagelin, S. Rudaz and D. D. Wu, Nucl. Phys. B 304 (1988) 205.

- [72] K. Abe et al. (Belle Collaboration), hep-ex/0208025.
- [73] B. Aubert et al. (BaBar Collaboration), hep-ex/0207042.
- [74] T. Hurth et al., J. Phys. G G27, 1277 (2001) [hep-ph/0102159].
- [75] D. Buskulic et al. (ALEPH Collab), Z. Phys. C 75, 397(1997).
- [76] J. Abdallah et al. (DELPHI Collab), Eur. Phys. J. C 28, 155 (2003).
- [77] P. Abreu et al. (DELPHI Collab), Z. Phys. C 76, 579 (1997).
- [78] M. Acciarri et al. (L3 Collab), Eur. Phys. J. C 5, 195 (1998).
- [79] G. Alexander et al. (OPAL Collab), Z. Phys. C 72, 377 (1996); K. Ackerstaff et al. (OPAL Collab), Z. Phys. C 76, 417 (1997); G. Abbiendi et al. (OPAL Collab), Phys. Lett. B 493, 266 (2000).
- [80] K. Ackerstaff et al. (OPAL Collab), Z. Phys. C 76, 401 (1997).
- [81] B. Aubert et al. (BaBar Collab), Phys. Rev. Lett. 88, 221802 (2002) and Phys. Rev. D 66, 032003 (2002); Phys. Rev. Lett. 88, 221803 (2002); Phys. Rev. D 67, 072002 (2003); Phys. Rev. D 73, 012004 (2006).
- [82] N.C. Hastings et al. (Belle Collab), Phys. Rev. D 67, 052004 (2003); Y. Zheng et al. (Belle Collab), Phys. Rev. D 67, 092004 (2003); K. Abe et al. (Belle Collab), Phys. Rev. D 71, 072003 (2005).
- [83] F. Abe et al. (CDF Collab), Phys. Rev. Lett. 80, 2057 (1998) and Phys. Rev. D 59, 032001 (1999); Phys. Rev. D 60, 051101 (1999); Phys. Rev. D 60, 072003 (1999); T. Affolder et al. (CDF Collab), Phys. Rev. D 60, 112004 (1999).
- [84] V.M. Abazov et al. (DØ Collab), Phys. Rev. D 74, 112002 (2006).
- [85] R. Aaij et al. (LHCb Collab), Phys. Lett. B 709, 177 (2012).
- [86] J. Beringer et al., Particle Data Group, Phys. Rev. D 86 (2012) 010001.
- [87] A.J. Buras, M. Jamin and R.H. Weisz, Nucl. Phys. B 347 (1990) 491.
- [88] John S. HAGELIN, Nucl. Phys. B 193 (1981) 123-149.
- [89] J. Urban, F. Krauss, U. Jentschura, G. Soft, Nucl. Phys. B 523 (1998) 40-58.
- [90] Tai-Fu Feng, Xue-Qian Li, Wen-Gan Ma, and Feng Zhang, Phys. Rev. D (2001)63, 015013.
- [91] F. Krauss, G. Soff, Nucl. Phys. B 633 (2002) 237-249.
- [92] Soo-hyeon Nam, Phys. Rev. D 66, 055008 (2002)

- [93] Debrupa Chakraverty, Katri Huitu, Anirban Kundu, Phys. Lett. B 558 (2003) 173.
- [94] Javier Virto, JHEP 0911 (2009) 055 [arXiv:0907.5376 [hep-ph]], JHEP 1201 (2012) 120 [arXiv:1111.0940 [hep-ph]].
- [95] FEI SUN et al, Mod. Phys. Lett. A 28, 1350060 (2013).
- [96] FEI SUN et al, Commun. Theor. Phys. 60 (2013) 721.
- [97] A. Lenz and U. Nierste: in Proceedings of CKM2010, Warwick, 2010, edited by Tim Gershon, eConf C 100906 (2010) [arXiv:1102.4274 [hep-ph]].
- [98] Thomas Kuhr, arXiv:1101.1916 [hep-ex].
- [99] S.P. Li and M. Sher, Phys. Lett. B 140 (1984) 33.
- [100] Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. 85 (1991) 1.
- [101] J.R. Ellis, G. Ri-doll and F. Zwirner, Phys. Lett. B 257 (1991) 83; *ibid.* 262 (1991) 477.
- [102] R. Barbieri and M. Frigeni, Phys. Lett. B 258 (1991) 395.
- [103] M. Drees and M. M. Nojiri, Phys. Rev. D 45 (1992) 2482.
- [104] M.A. Diaz and H. E. Haber, Phys. Rev. D 46 (1992) 3086.
- [105] J.A. Casas, J. R. Espinosa, M. Quiros and A. Riotto, Nucl. Phys. B 436(1995)3 [Erratum-*ibid.* B439(1995)466].
- [106] M.S. Carena, M. Quiros and C. E. M. Wagner, Nucl. Phys. B 461 (1996) 407.
- [107] S. Heinemeyer, W. Hollik and G. Weiglein, Eur. Phys. J. C 9 (1999) 343.
- [108] S. Heinemeyer, W. Hollik and G. Weiglein, Comput. Phys. Commun. 24 (2000) 76.
- [109] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, Eur. Phys. J. C 28 (2003) 133.
- [110] M. Frank, T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak and G. Weiglein, JHEP 0702 (2007) 047.
- [111] ATLAS Collaboration, Phys. Lett. B 716 (2012) 1 [arXiv:1207.7214].
- [112] CMS Collaboration, Phys. Lett. B 716 (2012) 30 [arXiv:1207.7235].
- [113] T. Inami and C. S. Lim, Prog. Theor. Phys. 65, 297 (1981).
- [114] M. Ciuchini, E. Franco, V. Lubicz, G. Martinelli, I. Scimemi, and L. Silvestrini, Nucl. Phys. B523, 501 (1998).
- [115] J.A. Bagger, K.T. Matchev and R.J. Zhang, Phys. Lett. B 412 (1997) 77 and references

- therein.
- [116] J. Laiho, E. Lunghi and R. S. Van de Water, Phys. Rev. D 81 (2010) 034503 [arXiv:0910.2928 [hep-ph]].
 - [117] Belle Collaboration, Phys. Lett. B 632 (2006) 51.
 - [118] G.D. Lafferty on behalf of the BABAR Collaboration, Nucl. Phys. B (Proc. Suppl.) 169 (2007) 186.
 - [119] BABAR Collaboration, Phys. Rev. D 83 (2011) 091101(R).
 - [120] LHCb Collaboration, Phys. Lett. B 724 (2013) 36.
 - [121] CMS Collaboration, Phys. Lett. B 731 (2014) 173.
 - [122] B.A. Golden, Ph.D. thesis, University of California, Irvine, 2012, <http://meg.web.psi.ch/docs/theses/BenThesis.pdf>; J. Adam, Ph.D. thesis, ETH Zurich, 2012, http://meg.web.psi.ch/docs/theses/PhDThesis_Adam_2012.pdf.
 - [123] J.L. Hewett et al., arXiv:1205.2671.
 - [124] N. Berger, J. Phys. Conf. Ser. 408, 012070 (2013).
 - [125] A. Kurup (COMET Collaboration), Nucl. Phys. B, Proc. Suppl. 218, 38 (2011); R. Akhmetshin et al. (COMET Collaboration), http://j-parc.jp/researcher/Hadron/en/pac_1203/pdf/COMET-PhaseI-LoI.pdf.
 - [126] R.J. Abrams et al. (Mu2e Collaboration), arXiv:1211.7019; R. K. Kurschke, arXiv:1112.0242; E.C. Dukes, Nucl. Phys. B, Proc. Suppl. 218, 32 (2011).
 - [127] Y. Kuno, Nucl. Phys. B, Proc. Suppl. 149, 376 (2005); R.J. Barow, Nucl. Phys. Proc. Suppl. 218, 44 (2011).
 - [128] K. Hayasaka, J. Phys. Conf. Ser. 171, 012079 (2009).