Search for 2β decay of ¹¹⁶Cd with the help of enriched ¹¹⁶CdWO₄ crystal scintillators

D.V. Poda^{1,a}, A.S. Barabash², P. Belli³, R. Bernabei^{3,4}, F. Cappella^{5,6}, V. Caracciolo⁷, S. Castellano⁷, D.M. Chernyak¹, R. Cerulli⁷, F.A. Danevich¹, S. d'Angelo^{3,4}, A. Incicchitti^{5,6}, V.V. Kobychev¹, S.I. Konovalov², M. Laubenstein⁷, R.B. Podviyanuk¹, O.G. Polischuk^{1,6}, V.N. Shlegel⁸, V.I. Tretyak¹, V.I. Umatov², and Ya.V. Vasiliev⁸

¹ Institute for Nuclear Research, MSP 03680 Kyiv, Ukraine

²Institute of Theoretical and Experimental Physics, 117259 Moscow, Russia

³Dipartimento di Fisica, Università di Roma "Tor Vergata", I-00133 Rome, Italy

⁵Dipartimento di Fisica, Università di Roma "La Sapienza", I-00185 Rome, Italy

⁶INFN, sezione di Roma "La Sapienza", I-00185 Rome, Italy

⁷Laboratori Nazionali del Gran Sasso, INFN, I-67100 Assergi (AQ), Italy

⁸Nikolaev Institute of Inorganic Chemistry, 630090 Novosibirsk, Russia

Abstract. Cadmium tungstate crystal scintillators enriched in ¹¹⁶Cd to 82% (¹¹⁶CdWO₄, total mass of ≈1.2 kg) are used to search for 2 β decay of ¹¹⁶Cd deep underground at the Gran Sasso National Laboratory of the INFN (Italy). The radioactive contamination of the ¹¹⁶CdWO₄ crystals has been studied carefully to reconstruct the background of the detector. The measured half-life of ¹¹⁶Cd relatively to 2 ν 2 β decay is $T_{1/2}^{2\nu2\beta} = [2.8 \pm 0.05(\text{stat.}) \pm 0.4(\text{syst.})] \times 10^{19}$ yr, in agreement with the results of previous experiments. The obtained limit on the 0 ν 2 β decay of ¹¹⁶Cd (considering the data of the last 8696 h run with an advanced background 0.12(2) counts/yr/kg/keV in the energy interval 2.7 – 2.9 MeV) is $T_{1/2} \ge 1.0 \times 10^{23}$ yr at 90% C.L. The sensitivity of the experiment to the 0 ν 2 β process is lim $T_{1/2} = 3 \times 10^{23}$ yr at 90% C.L. over 5 years of the measurements and it can be advanced (by further reduction of the background by a factor 3–30) to the level of lim $T_{1/2} = (0.5-1.5) \times 10^{24}$ yr for the same period of the data taking.

1 Introduction

The rarest nuclear decay among observed - double beta decay (2β) — is a powerful tool to check physics beyond the Standard Model (SM) of particle physics. Allowed in the SM two neutrino double beta decay $(2\nu 2\beta)$ has been registered for only 11 nuclides with a typical half-lives in the range of $10^{18} - 10^{24}$ yr (see e.g. in [1, 2]). Neutrinoless double beta decay $(0\nu 2\beta)$ — forbidden in the SM process — should exist, if neutrino is a Majorana particle, thanks to recently observed non-zero masses of the neutrino [3]. The observation of the $0\nu 2\beta$ decay will testify the lepton number non-conservation, the Majorana nature of the neutrino, an absolute value and hierarchy of the neutrino masses. The $0\nu 2\beta$ decay is also sensitive to an existence of the right-handed currents, hypothetical particles (majorons) and other effects beyond the SM (see the recent reviews [4–9] and references therein).

One of the most promising candidates for the 2β decay search is ¹¹⁶Cd thanks to high 2β energy release ($Q_{2\beta}$ = 2813.50(13) keV [10]), large natural isotopic abundance (δ = 7.49% [11]), availability of enrichment by relatively cheap centrifugal method [12], and promising theoretical estimations (see e.g. [5]). Moreover, existence of excellent detector, cadmium tungstate (CdWO₄) scintillator, already used for 2β decay searches [13–15] allows to realize "source = detector" experiment with a high detection efficiency. In addition, detector based on cadmium tungstate scintillator has several features (see e.g. [15, 16]) which are very important for such kind of studies: low level of intrinsic radioactivity, good scintillation properties, particle identification ability, relatively low cost of the scintillation material, stability for long term operation.

Recently an enriched ¹¹⁶CdWO₄ crystal scintillator with a large mass (1.87 kg) has been developed from deeply purified raw materials, and a new high sensitive 2β experiment has been started in February 2011 at the Gran Sasso National Laboratories (LNGS) of the INFN (Italy) [17]. The results of the last run performed from October 2012 till October 2013 will be reported here.

2 Low-background experiment

The experiment has been performed deep underground (\approx 3600 m w.e.) at the LNGS (Italy). The description of the detector and low background set-up at the previous stages

⁴INFN, sezione di Roma "Tor Vergata", I-00133 Rome, Italy

^ae-mail: poda@kinr.kiev.ua

of the experiment one can find in Refs. [17, 18]. Here we outline the main features of the last stage.

Two ¹¹⁶CdWO₄ crystal scintillators (586 g and 589 g) were fixed inside polystyrene light-guides (070×194 mm). The cavities inside the light-guides were filled with the Ultima Gold AB (PerkinElmer) liquid scintillator (LS). Two high purity quartz light-guides ($\oslash 70 \times 200 \text{ mm}$) were glued to the polystyrene light-guide on both sides. The detectors were viewed from the opposite sides by two low radioactive 3" photomultipliers (PMTs, Hamamatsu R6233MOD). The detector modules, covered by the 3M foil to improve the light collection, were installed inside a low radioactive air-tight Cu box at the DAMA/R&D setup. Copper bricks 5 cm thick were used as an additional passive shield. The Cu box was continuously flushed with high purity nitrogen gas to remove the residual environmental radon. Outer passive shield consists of high purity Cu (10 cm thick), 15 cm of low radioactive lead, 1.5 mm of cadmium and 4 to 10 cm of polyethylene/paraffin. The whole shield is contained inside a Plexiglas box, also continuously flushed by high purity nitrogen.

An event-by-event data acquisition system based on a 1 GS/s 8 bit transient digitizer (Acqiris DC270) was used to record time of each event and scintillation pulseprofiles. The energy scale and the resolution of the detector was calibrated by using the reference ²²Na, ⁶⁰Co, ¹³⁷Cs, and ²²⁸Th γ sources. The energy resolution of the detector was FWHM 4.3(4)% at Q_{26} of ¹¹⁶Cd.

3 Results and discussion

The data at this stage were collected over 8696 h. The applied data analysis is similar to the one described in [17]. The energy spectrum of $\gamma(\beta)$ events accumulated with the ¹¹⁶CdWO₄ detector and selected off-line by the pulseshape and the front edge analyzes is shown in Fig. 1. The background model was built from possible components like internal ⁴⁰K, ⁹⁰Sr-⁹⁰Y, ^{110m}Ag, radionuclides from U/Th chains, external γ from the details of the set-up, $2\nu 2\beta$ decay of ¹¹⁶Cd simulated by using the EGS4 [19] and Decay0 [20] codes. The radiopurity of the ¹¹⁶CdWO₄ crystals is reported in [17, 21, 22]. The radioactive contamination of the PMTs [23] and the sample of Ultima Gold AB liquid scintillator has been measured at the STELLA HPGe facility of the LNGS (Italy). The radiopurity of Cu has been derived from the fit of the background spectrum and it is comparable with the results of the low background measurements with the Cu samples (see e.g. [24]). Table 1 contains radiopurity level of the main sources of the background.

Fit of the background spectrum and the main components of the background are also presented in Fig.1. The fit gives the number of the $2\nu 2\beta$ events of ¹¹⁶Cd as $S_{2\nu 2\beta} = (34927 \pm 614)$ counts for the total distribution. The signal to background ratio in the energy range of 1.6 - 2.4 MeV is $\approx 2:1$. Taking into account the efficiency of the applied pulse-shape discrimination procedure ($\eta_{PSD} = 0.92$), selection of the ²¹²Bi-Po events ($\eta_{BiPo} = 0.99$), detection efficiency to the process searched for



Figure 1. (Color online) The energy spectrum of $\gamma(\beta)$ events measured with the ¹¹⁶CdWO₄ scintillation detector over 8696 h in the low-background set-up (points with error bars) together with the fit (solid black line). The main components of the background are also shown: $2\nu 2\beta$ decay of ¹¹⁶Cd, the distributions of internal ⁴⁰K, ⁹⁰Sr-⁹⁰Y, ²³²Th, ²³⁸U, and cosmogenic ^{110m}Ag, the contribution from external γ quanta from the details of the detector ("LS", "PMT") and set-up ("Cu") under these experimental conditions. The energies are in keV.

 $(\eta_{2\nu2\beta} = 0.986)$, the number of ¹¹⁶Cd nuclei in the used enriched ¹¹⁶CdWO₄ crystals (1162.25 g, enrichment 82.2%, $N_{116} = 1.584 \times 10^{24}$ nuclei), time of measurements (t = 0.992 yr), one can calculate the half-life of ¹¹⁶Cd relatively to the $2\nu2\beta$ decay by using the following formula: $T_{1/2}^{2\nu2\beta} = \ln 2 \cdot \eta_{PSD} \cdot \eta_{BiPo} \cdot \eta_{2\nu2\beta} \cdot N_{116} \cdot t / S_{2\nu2\beta}$. The derived half-life $T_{1/2}^{2\nu2\beta} = [2.8 \pm 0.05(\text{stat.}) \pm 0.4(\text{syst.})] \times 10^{19}$ yr is in agreement with the results of the previous experiments (see Table 2) and the recent world average value $(2.8 \pm 0.2) \times 10^{19}$ yr [31].

In case of the $0\nu 2\beta$ decay of ¹¹⁶Cd to the ground state of ¹¹⁶Sn we should see in the experimental spectrum a peak at the energy ≈ 2.8 MeV. There is no such peculiarity in the spectrum, therefore only lower half-life limit on the process can be set according to formula: $\lim T_{1/2}^{0\nu 2\beta} =$ $\ln 2 \cdot \eta_{PSD} \cdot \eta_{BiPo} \cdot \eta_{0\nu 2\beta} \cdot N_{116} \cdot t / \lim S$. Here $\eta_{0\nu 2\beta}$ is the detection efficiency for the $0\nu 2\beta$ process (equal to 0.947), $\lim S$ is the number of $0\nu 2\beta$ events which can be excluded at a given confidence level (C.L.), and other parameters are described above. The fit gives the area of the $0\nu 2\beta$ peak as (2.3 ± 4.4) counts. It corresponds to $\lim S = 9.2$ counts at 90% C.L. according to the Feldman-Cousins procedure [32]. Substituting all parameters, one can obtain the limit $T_{1/2}^{0\nu 2\beta} \ge 1.0 \times 10^{23}$ yr at 90% C.L., which is on the level of the Solotvina experiment [15] (see Table 3).

One can derive the similar result for the $0v2\beta$ decay of ¹¹⁶Cd by using also the "1 σ approach". For example, there are N = 28 counts in the region of interest (ROI) 2.7 – 2.9 MeV, where the detection efficiency for the $0v2\beta$ process is 0.849. The value lim *S* in such a case can be taken as

Nuclide	Activity (mBq/kg)					
	¹¹⁶ CdWO ₄ No.1	¹¹⁶ CdWO ₄ No.2	LS	PMT	Copper	
²²⁸ Ra				120(20)		
²²⁸ Th	0.031(3)	0.054(5)		83(17)	0.025(14)	
²³⁸ U	0.5(2)	0.7(2)				
²²⁶ Ra	≤ 0.005	≤ 0.005		430(60)	0.011(3)	
²¹⁰ Po	0.6(2)	0.8(2)				
40 K	≤ 0.9	≤ 0.9	1600(200)	540(160)	0.04(1)	
⁹⁰ Sr- ⁹⁰ Y	≤ 0.1	≤ 0.1				
^{110m}Ag	≤ 0.7	≤ 0.7				

Table 1. Radioactive contamination of the details of the ¹¹⁶CdWO4 detector (¹¹⁶CdWO4 crystals No.1 and No.2, liquid scintillator LS, and PMTs) and the inner part of the low-background DAMA/R&D set-up made from Copper.

Table 2. The half-lives of $2\nu 2\beta$ decay of ¹¹⁶Cd (g.s. \rightarrow g.s.) measured in the 2β experiments. The signal to background ratio (S:Bg), main experimental features, and the total exposure are also presented.

$T_{1/2}^{2\nu 2\beta} (10^{19} \text{ yr})$	S:Bg	Experiment	Exposure (kg×yr)	Year [Ref.]
$=2.6^{+0.9}_{-0.5}$	1:3	ELEGANT V, ¹¹⁶ Cd foils	0.02	1995 [25]
$=2.7 + 0.5 \\ -0.4 (stat.) + 0.9 \\ -0.6 (syst.)$	1:1	Solotvina, ¹¹⁶ CdWO ₄ scintillators	0.04	1995 [13]
$=3.75 \pm 0.35(\text{stat.}) \pm 0.21(\text{syst.})$	4:1	NEMO-2, ¹¹⁶ Cd foils	0.11	1995 [26, 27]
$=2.6 \pm 0.1(\text{stat.}) \stackrel{+0.7}{_{-0.4}}(\text{syst.})$	4:1	Solotvina, ¹¹⁶ CdWO ₄ scintillators	0.18	2000 [14]
$=2.9 \pm 0.06(\text{stat.}) \stackrel{+0.4}{_{-0.3}}(\text{syst.})$	3:1	Solotvina, ¹¹⁶ CdWO ₄ scintillators	0.48	2003 [15]
$=2.88 \pm 0.04(\text{stat.}) \pm 0.16(\text{syst.})$	10:1	NEMO-3, ¹¹⁶ Cd foils	1.95	2010 [28-30]
$=2.5 \pm 0.5$	2:1	Present, ¹¹⁶ CdWO ₄ scintillators	1.04	2012 [18]
$=2.8 \pm 0.05$ (stat.) ± 0.4 (syst.)	2:1	Present, ¹¹⁶ CdWO ₄ scintillators	1.15	2013

Table 3. The half-life limits on the $0\nu 2\beta$ decay of ¹¹⁶Cd (g.s. \rightarrow g.s.) established in the 2β experiments. The background rate in the region of interest (Bg in ROI), main experimental features, enrichment, and the total exposure are also presented.

$T_{1/2}^{0\nu 2\beta}$ (yr)	Bg in ROI	Experiment	Enrichment	Exposure	Year [Ref.]
at 90% C.L.	(counts/yr/kg/keV)		(%)	(kg×yr)	
$\geq 2.9 \times 10^{21}$		ELEGANT V, ¹¹⁶ Cd foils	90.7	0.02	1995 [25]
$\geq 2.9\times 10^{22}$	0.53	Solotvina, ¹¹⁶ CdWO ₄ scintillators	83.0	0.08	1995 [13]
$\geq 5.0\times 10^{21}$	0.02	NEMO-2, ¹¹⁶ Cd foils	93.2	0.11	1995 [26, 27]
$\geq 7.0\times 10^{22}$	0.03	Solotvina, ¹¹⁶ CdWO ₄ scintillators	83.0	0.18	2000 [14]
$\geq 1.7\times 10^{23}$	0.04	Solotvina, ¹¹⁶ CdWO ₄ scintillators	83.0	0.53	2003 [15]
$\geq 9.4 \times 10^{19}$	≈9	COBRA, CdZnTe semiconductors	nat.	0.05	2009 [33]
$\geq 1.3\times 10^{23}$	0.005	NEMO-3, ¹¹⁶ Cd foils	93.3	1.95	2010 [28, 30]
$\geq 9.2\times 10^{20}$	≈1	COBRA, CdZnTe semiconductors	nat.	0.23	2013 [34]
$\geq 1.0\times 10^{23}$	0.12	Present, ¹¹⁶ CdWO ₄ scintillators	82.2	1.15	2013

 $1.64 \cdot \sqrt{N} = 8.7$ counts at 90% C.L. So, the calculated halflife limit $T_{1/2}^{0v2\beta} \ge 0.97 \times 10^{23}$ at 90% C.L. is very close to the one obtained from the fit of the background spectrum.

The background 0.12(2) counts/yr/kg/keV in the ROI allows us to reach a sensitivity to the half-life of $0\nu 2\beta$ decay of ¹¹⁶Cd equal to $\lim T_{1/2} = 3 \times 10^{23}$ yr at 90% C.L. over 5 years of measurements. In a case of further reduction of the background by a factor 3–30, the sensitivity can be advanced to the level of $\lim T_{1/2} = (0.5 - 1.5) \times 10^{24}$ yr (it corresponds to the effective neutrino mass $\langle m_{\nu} \rangle = 0.4 - 1.4$ eV).

The main components of the background in the ROI remain 2615 keV γ quanta of ²⁰⁸Tl from contamination of the set-up by ²³²Th, and ²⁰⁸Tl from internal contamination of the ¹¹⁶CdWO₄ crystals by ²²⁸Th. The background could be suppressed by removing of one PMT from each detector module, by installation of an additional light-guide

(quartz) between the crystal and the PMT, by replacing the PMT by another one with a higher level of radiopurity (e.g. Hamamatsu R11065SEL) and by recrystallization of the ¹¹⁶CdWO₄ crystals to decrease thorium thanks to observed low segregation of Th and Ra by cadmium tungstate [21]. The first two modifications have already been applied in October, 2013. In addition, the Ultima Gold LS polluted by ⁴⁰K has been replaced by the Borexino LS and the new measurements are in progress.

4 Conclusions

A low-background experiment using enriched ¹¹⁶CdWO₄ scintillators (total mass of 1.16 kg, enrichment by ¹¹⁶Cd to 82%) is in progress at the LNGS of the INFN (Italy). The exposure of the experiment at the last stage of the data taking (October, 2012 – October, 2013) is 1.15 kg×yr. The

energy resolution of the ¹¹⁶CdWO₄ detector is FWHM = 4.3% at $Q_{2\beta}$ of ¹¹⁶Cd, the background counting rate was reduced to 0.12(2) counts/yr/kg/keV in the region of interest 2.7 – 2.9 MeV, where the $0\nu 2\beta$ peak is expected. The main components of the background in this region remain γ quanta of ²⁰⁸Tl from the contamination of the set-up and internal contamination of the ¹¹⁶CdWO₄ crystals by ²²⁸Th.

The half-life relatively to the $2\nu 2\beta$ decay of 116 Cd to the ground level of 116 Sn was measured as $T_{1/2}^{2\nu 2\beta} = [2.8 \pm 0.05(\text{stat.}) \pm 0.4(\text{syst.})] \times 10^{19}$ yr, in agreement with the results of the previous experiments. The half-life limit on $0\nu 2\beta$ decay of 116 Cd to the ground state of 116 Sn was established as $T_{1/2}^{0\nu 2\beta} \ge 1.0 \times 10^{23}$ at 90% C.L. The current level of the background restricts the sensitivity of the experiment to $0\nu 2\beta$ decay of 116 Cd to lim $T_{1/2} = 3 \times 10^{23}$ yr at 90% C.L. over 5 years of the measurements. The sensitivity can be advanced to the level of lim $T_{1/2} \approx$ $(0.5 - 1.5) \times 10^{24}$ yr (which corresponds to the effective neutrino mass $m_{\nu} = 0.4 - 1.4$ eV) after improvement of the background by factor 3 - 30. An R&D to improve the background conditions of the experiment is in progress.

5 Acknowledgments

D.V. Poda has been partly supported by a Grant of the President of Ukraine for young scientists in 2012 (project GP-F36-110, reg. no. 0112U008078).

References

- V.I. Tretyak and Yu.G. Zdesenko, At. Data Nucl. Data Tables 61, 43–90 (1995); *ibid.* 80, 83–116 (2002)
- [2] A.S. Barabash, Phys. At. Nucl. 73, 162–178 (2010)
- [3] J. Beringer et al. (Particle Data Group), Phys. Rev. D 86, 010001 (2012)
- [4] O. Cremonesi, M. Pavan, arXiv:1310.4692, 1–70 (2013)
- [5] J.D. Vergados, H. Ejiri, and F. Šimkovic, Rep. Prog. Phys. 75, 106301 (2012)
- [6] S.R. Elliott, Mod. Phys. Lett. A 27, 1230009 (2012)
- [7] A. Giuliani and A. Poves, Adv. High En. Phys. 2012, 857016 (2012)
- [8] J.J. Gomez-Cadenas et al., Riv. Nuovo Cim. 35, 29–98 (2012)
- [9] W. Rodejohann, Int. J. Mod. Phys. E 20, 1833–1930 (2011)

- [10] S. Rahaman et al., Phys. Lett. B 703, 412–416 (2011)
- [11] M. Berglund, M.E. Wieser, Pure Appl. Chem. 83, 397–410 (2011)
- [12] A.A. Artyukhov et al., Nucl. Instrum. Meth. Phys. Res. A 401, 281–288 (1997)
- [13] F.A. Danevich et al., Phys. Lett. B 344, 72–78 (1995)
- [14] F.A. Danevich et al., Phys. Rev. C 62, 045501 (2000)
- [15] F.A. Danevich et al., Phys. Rev. C 68, 035501 (2003)
- [16] L. Bardelli et al., Nucl. Instrum. Meth. Phys. Res. A 569, 743–753 (2006)
- [17] A.S. Barabash et al., JINST 6, P08011 (2011)
- [18] A.S. Barabash et al., Proc. NPAE-2012, Kyiv, 353– 356 (2013)
- [19] W.R. Nelson et al., SLAC-Report-265, Stanford, 1– 398 (1985)
- [20] O.A. Ponkratenko et al., Phys. At. Nucl. 63, 1282– 1287 (2000);
 - V.I. Tretyak, to be published
- [21] D. Poda et al., Radiat. Meas. 56, 66–69 (2013)
- [22] F.A. Danevich et al., AIP Conf. Proc. 1549, 201– 2004 (2013)
- [23] R. Bernabei et al., JINST 7, P03009 (2012)
- [24] G. Heusser et al., In: Radioactivity in the Environment (Elsevier, Oxford (UK), 2006) 495–510
- [25] H. Ejiri et al., J. Phys. Soc. Japan 64, 339–343 (1995)
- [26] R. Arnold et al., JETP Lett. 61, 170–174 (1995)
- [27] R. Arnold et al., Z. Phys. C 72, 239–247 (1996)
- [28] R.B. Pahlka, Search for Neutrinoless Double Beta Decay of ¹¹⁶Cd with the NEMO-3 Experiment, Ph.D. thesis, University of Texas at Austin, USA, 1–269 (2010)
- [29] V.I. Tretyak on behalf of the NEMO-3 collaboration, AIP Conf. Proc. 1417, 125–128 (2011)
- [30] R.B. Pahlka for the NEMO-3 and SuperNEMO Collaborations, Phys. Proceedia 37, 1241–1248 (2012)
- [31] A.S. Barabash, Phys. Rev. C 81, 035501 (2010)
- [32] G.J. Feldman, R.D. Cousins, Phys. Rev. D 57, 3873– 3889 (1998)
- [33] J.V. Dawson et al., Phys. Rev. C 80, 025502 (2009)
- [34] D. Gehre for the COBRA Collaboration, On the Search for the Neutrinoless Double Beta Decay with the COBRA-Experiment, talk at the Meeting of the LNGS Scientific Committee, 29 October 2013, LNGS, Assergi (AQ), Italy