

REANALYSIS OF THE $Z_c(4020)$, $Z_c(4025)$, $Z(4050)$ AND $Z(4250)$ AS TETRAQUARK STATES WITH QCD SUM RULES

Zhi-Gang Wang¹

Department of Physics, North China Electric Power University, Baoding 071003, P. R. China

Abstract

In this article, we calculate the contributions of the vacuum condensates up to dimension-10 in the operator product expansion, and study the $C\gamma_\mu - C\gamma_\nu$ type scalar, axial-vector and tensor tetraquark states in details with the QCD sum rules. In calculations, we use the formula $\mu = \sqrt{M_{X/Y/Z}^2 - (2M_c)^2}$ to determine the energy scales of the QCD spectral densities. The predictions $M_{J=2} = (4.02^{+0.09}_{-0.09})$ GeV, $M_{J=1} = (4.02^{+0.07}_{-0.08})$ GeV favor assigning the $Z_c(4020)$ and $Z_c(4025)$ as the $J^{PC} = 1^{+-}$ or 2^{++} diquark-antidiquark type tetraquark states, while the prediction $M_{J=0} = (3.85^{+0.15}_{-0.09})$ GeV disfavors assigning the $Z(4050)$ and $Z(4250)$ as the $J^{PC} = 0^{++}$ diquark-antidiquark type tetraquark states. Furthermore, we discuss the strong decays of the 0^{++} , 1^{+-} , 2^{++} diquark-antidiquark type tetraquark states in details.

PACS number: 12.39.Mk, 12.38.Lg

Key words: Tetraquark state, QCD sum rules

1 Introduction

In 2008, the Belle collaboration reported the first observation of two resonance-like structures in the $\pi^+\chi_{c1}$ invariant mass distribution near 4.1 GeV in the exclusive $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}$ decays, and determined the masses and widths $M_{Z(4050)} = (4051 \pm 14^{+20}_{-41})$ MeV, $M_{Z(4250)} = (4248^{+44+180}_{-29-35})$ MeV, $\Gamma_{Z(4050)} = (82^{+21+47}_{-17-22})$ MeV and $\Gamma_{Z(4250)} = (177^{+54+316}_{-39-61})$ MeV, respectively [1]. There have been several tentative assignments, such as the tetraquark states [2, 3, 4], molecular states [5], etc. In 2011, the BaBar collaboration searched for the $Z^+(4050)$ and $Z^+(4250)$ states in the decays $\bar{B}^0 \rightarrow \chi_{c1}K^-\pi^+$ and $B^+ \rightarrow \chi_{c1}K_S^0\pi^+$ based on the data collected with the BaBar detector at the SLAC PEP-II asymmetric-energy e^+e^- collider, and observed no evidences [6]. The $Z^+(4050)$ and $Z^+(4250)$ still need confirmation.

In 2013, the BESIII collaboration observed the $Z_c^\pm(4025)$ near the $(D^*\bar{D}^*)^\pm$ threshold in the $\pi^\mp$ recoil mass spectrum in the process $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ at a center-of-mass energy of 4.26 GeV, and determined the mass and width $M_{Z_c(4025)} = (4026.3 \pm 2.6 \pm 3.7)$ MeV and $\Gamma_{Z_c(4025)} = (24.8 \pm 5.6 \pm 7.7)$ MeV, respectively [7]. Furthermore, the BESIII collaboration observed the $Z_c(4020)$ in the $\pi^\pm h_c$ mass spectrum in the process $e^+e^- \rightarrow \pi^+\pi^-h_c$ at center-of-mass energies $(3.90 - 4.42)$ GeV, and determined the mass and width $M_{Z_c(4020)} = (4022.9 \pm 0.8 \pm 2.7)$ MeV and $\Gamma_{Z_c(4020)} = (7.9 \pm 2.7 \pm 2.6)$ MeV, respectively [8]. There have been several tentative assignments of the $Z_c(4025)$ and $Z_c(4020)$, such as the re-scattering effects [9], molecular states [10], tetraquark states [11], etc.

The S-wave $D^*\bar{D}^*$ systems have the quantum numbers $J^{PC} = 0^{++}$, 1^{+-} , 2^{++} , the S-wave $\pi^\pm h_c$ systems have the quantum numbers $J^{PC} = 1^{--}$, the S-wave $\pi^\pm\chi_{c1}$ systems have the quantum numbers $J^{PC} = 1^{-+}$ (or $J^{PC} = 1^{--}$). On the other hand, it is also possible for the P-wave $\pi^\pm h_c$ and $\pi^\pm\chi_{c1}$ systems to have the quantum numbers $J^{PC} = 0^{++}$, 1^{+-} , 2^{++} . In Ref.[12], we observe that the predictions based on the QCD sum rules disfavor assigning the $Z_c(4020)$ and $Z_c(4025)$ as the diquark-antidiquark type vector tetraquark states. The $Z_c(4020)$ and $Z_c(4025)$ are potential candidates of the axial-vector and tensor tetraquark states [12]. The $Z(4050)$ and $Z(4250)$ are potential candidates of the scalar tetraquark states [2, 3]. However, we cannot exclude the possibilities that the $Z_c(4025)$, $Z_c(4020)$ and $Z(4050)$ are the same particle with the $J^{PC} = 1^{+-}$ or 2^{++} . The $Z_c(4025)$, $Z_c(4020)$, $Z(4050)$ and $Z(4250)$ are charged charmonium-like states, their quark constituents must be $c\bar{c}u\bar{d}$ or $c\bar{c}d\bar{u}$ irrespective of the diquark-antidiquark type or meson-meson type substructures. The interested reader can consult Ref.[13] for more articles on the exotic X , Y and Z particles.

¹E-mail: zgwang@aliyun.com.

In Refs.[12, 14], we distinguish the charge conjugations of the interpolating currents, calculate the contributions of the vacuum condensates up to dimension-10 in the operator product expansion, study the diquark-antidiquark type axial-vector and vector tetraquark states in a systematic way with the QCD sum rules, make possible assignments of the $X(3872)$, $Z_c(3900)$, $Z_c(3885)$, $Z_c(4020)$, $Z_c(4025)$, $Y(4360)$, $Y(4630)$ and $Y(4660)$. Furthermore, we explore the energy scale dependence of the hidden charmed tetraquark states in details for the first time, and suggest a formula

$$\mu = \sqrt{M_{X/Y/Z}^2 - (2\mathbb{M}_c)^2}, \quad (1)$$

with the effective c -quark mass $\mathbb{M}_c = 1.8 \text{ GeV}$ to determine the energy scales of the QCD spectral densities in the QCD sum rules. The masses of the $J^{PC} = 1^{-+}, 1^{--}$ tetraquark states with symbolic quark structure $c\bar{c}u\bar{d}$ disfavor assigning the $Z_c(4020)$, $Z_c(4025)$ and $Y(4360)$ as the vector tetraquark states; the masses of the vector tetraquark states with symbolic quark structures $c\bar{c}s\bar{s}$ and $c\bar{c}(u\bar{u} + d\bar{d})/\sqrt{2}$ favor assigning the $Y(4660)$ (or $Y(4630)$) as the 1^{--} tetraquark state [12].

In Refs.[2, 3], we study the $C\gamma_\mu - C\gamma^\mu$ and $C\gamma_5 - C\gamma_5$ type scalar tetraquark states with the QCD sum rules by carrying out the operator product expansion to the vacuum condensates up to dimension-10 and setting the energy scale to be $\mu = 1 \text{ GeV}$. In Ref.[11], the Qiao and Tang study the vector and tensor tetraquark states with the QCD sum rules by carrying out the operator product expansion to the vacuum condensates up to dimension-8. They try to assign the $Z_c(4025)$ as the $J^P = 2^+$ tetraquark state (to be more precise, the $J^{PC} = 2^{+-}$ tetraquark state), but do not show (or do not specify) the energy scales of the QCD spectral densities. In Refs.[2, 3, 11], some higher dimension vacuum condensates involving the gluon condensate, mixed condensate and four-quark condensate are neglected, which maybe impair the predictive ability.

In this article, we distinguish the charge conjugations of the interpolating currents, calculate the contributions of the vacuum condensates up to dimension-10 in a consistent way, study the scalar, axial-vector, tensor hidden charmed tetraquark states with the $C\gamma_\mu - C\gamma_\nu$ type interpolating currents in a systematic way, make tentative assignments of the $Z_c(4020)$, $Z_c(4025)$, $Z(4050)$ and $Z(4250)$ based on the QCD sum rules. Furthermore, we explore validity of the formula on how to determine the energy scales of the QCD spectral densities [12].

The diquarks have five Dirac tensor structures, scalar $C\gamma_5$, pseudoscalar C , vector $C\gamma_\mu\gamma_5$, axial vector $C\gamma_\mu$ and tensor $C\sigma_{\mu\nu}$. The structures $C\gamma_\mu$ and $C\sigma_{\mu\nu}$ are symmetric, the structures $C\gamma_5$, C and $C\gamma_\mu\gamma_5$ are antisymmetric. The attractive interactions of one-gluon exchange favor formation of the diquarks in color antitriplet $\bar{3}_c$, flavor antitriplet $\bar{3}_f$ and spin singlet 1_s (or flavor sextet 6_f and spin triplet 3_s) [15], the favored configurations are the scalar and axial-vector diquark states. We take the diquark states as the basic constituents following Jaffe and Wilczek [16]. If additional partial derivative is not introduced, the scalar $C\gamma_5$ diquark-antidiquark pair can form scalar tetraquark states only, while the axial-vector $C\gamma_\mu$ diquark-antidiquark pair can form scalar, axial-vector, tensor tetraquark states. The scalar and axial-vector heavy-light diquark states have almost degenerate masses from the QCD sum rules [17]. In this article, we choose the $C\gamma_\mu - C\gamma_\nu$ type interpolating currents to study the tetraquark states.

The article is arranged as follows: we derive the QCD sum rules for the masses and pole residues of the scalar, axial-vector, tensor tetraquark states in section 2; in section 3, we present the numerical results and discussions; section 4 is reserved for our conclusion.

2 QCD sum rules for the scalar, axial-vector and tensor tetraquark states

In the following, we write down the two-point correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ and $\Pi(p)$ in the QCD sum rules,

$$\Pi_{\mu\nu\alpha\beta}(p) = i \int d^4x e^{ip \cdot x} \langle 0 | T \{ \eta_{\mu\nu}(x) \eta_{\alpha\beta}^\dagger(0) \} | 0 \rangle, \quad (2)$$

$$\Pi(p) = i \int d^4x e^{ip \cdot x} \langle 0 | T \{ \eta(x) \eta^\dagger(0) \} | 0 \rangle, \quad (3)$$

$$\eta_{\mu\nu}(x) = \frac{\epsilon^{ijk}\epsilon^{imn}}{\sqrt{2}} \{ u^j(x) C \gamma_\mu c^k(x) \bar{d}^m(x) \gamma_\nu C \bar{c}^n(x) + t u^j(x) C \gamma_\nu c^k(x) \bar{d}^m(x) \gamma_\mu C \bar{c}^n(x) \}, \quad (4)$$

$$\eta(x) = \epsilon^{ijk}\epsilon^{imn} u^j(x) C \gamma_\mu c^k(x) \bar{d}^m(x) \gamma^\mu C \bar{c}^n(x), \quad (5)$$

where the i, j, k, m, n are color indexes, the C is the charge conjugation matrix. Under charge conjugation transform $\hat{C}$, the currents $\eta_{\mu\nu}(x)$ and $\eta(x)$ have the following properties,

$$\begin{aligned} \hat{C} \eta_{\mu\nu}(x) \hat{C}^{-1} &= \pm \eta_{\mu\nu}(x) |_{u \leftrightarrow d} \quad \text{for } t = \pm 1, \\ \hat{C} \eta(x) \hat{C}^{-1} &= \eta(x) |_{u \leftrightarrow d}, \end{aligned} \quad (6)$$

which originate from the charge conjugation property of the axial-vector diquark states,

$$\hat{C} [\epsilon^{ijk} q^j C \gamma_\mu c^k] \hat{C}^{-1} = \epsilon^{ijk} \bar{q}^j \gamma_\mu C \bar{c}^k. \quad (7)$$

The charged currents $\eta_{\mu\nu}(x)$ and $\eta(x)$ change their charge signs under charge conjugation. The neutral partners $\tilde{\eta}_{\mu\nu}$ and $\tilde{\eta}$,

$$\begin{aligned} \tilde{\eta}_{\mu\nu} &= \frac{\epsilon^{ijk}\epsilon^{imn}}{2} \{ u^j C \gamma_\mu c^k \bar{u}^m \gamma_\nu C \bar{c}^n + d^j C \gamma_\mu c^k \bar{d}^m \gamma_\nu C \bar{c}^n \\ &\quad + t u^j C \gamma_\nu c^k \bar{u}^m \gamma_\mu C \bar{c}^n + t d^j C \gamma_\nu c^k \bar{d}^m \gamma_\mu C \bar{c}^n \}, \end{aligned} \quad (8)$$

$$\tilde{\eta} = \frac{\epsilon^{ijk}\epsilon^{imn}}{\sqrt{2}} \{ u^j C \gamma_\mu c^k \bar{u}^m \gamma^\mu C \bar{c}^n + d^j C \gamma_\mu c^k \bar{d}^m \gamma^\mu C \bar{c}^n \}, \quad (9)$$

are eigenstates of the charge conjugation. The currents $\tilde{\eta}_{\mu\nu}$ and $\eta_{\mu\nu}$ ($\tilde{\eta}$ and η) have the same charge conjugations. We take the currents $\eta(x)$, $\eta_{\mu\nu}^{t=-}(x)$ and $\eta_{\mu\nu}^{t=+}(x)$ to interpolate the scalar, axial-vector and tensor tetraquark states, respectively. The $Z_c(4020)$ and $Z_c(4025)$ are potential candidates of the axial-vector and tensor tetraquark states, the $Z(4050)$ and $Z(4250)$ are potential candidates of the scalar tetraquark states. Other possibilities are not excluded.

At the hadronic side, we can insert a complete set of intermediate hadronic states with the same quantum numbers as the current operators $\eta_{\mu\nu}(x)$ and $\eta(x)$ into the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ and $\Pi(p)$ to obtain the hadronic representation [18, 19]. After isolating the ground state contributions of the scalar, axial-vector and tensor tetraquark states, we get the following results,

$$\begin{aligned} \Pi_{\mu\nu\alpha\beta}^{J=2}(p) &= \Pi_{J=2}(p) \left(\frac{\tilde{g}_{\mu\alpha}\tilde{g}_{\nu\beta} + \tilde{g}_{\mu\beta}\tilde{g}_{\nu\alpha}}{2} - \frac{\tilde{g}_{\mu\nu}\tilde{g}_{\alpha\beta}}{3} \right) + \Pi_s(p) g_{\mu\nu} g_{\alpha\beta}, \\ &= \frac{\lambda_Z^2}{M_Z^2 - p^2} \left(\frac{\tilde{g}_{\mu\alpha}\tilde{g}_{\nu\beta} + \tilde{g}_{\mu\beta}\tilde{g}_{\nu\alpha}}{2} - \frac{\tilde{g}_{\mu\nu}\tilde{g}_{\alpha\beta}}{3} \right) + \dots, \end{aligned} \quad (10)$$

$$\begin{aligned} \Pi_{\mu\nu\alpha\beta}^{J=1}(p) &= \Pi_{J=1}(p) (-\tilde{g}_{\mu\alpha}p_\nu p_\beta - \tilde{g}_{\nu\beta}p_\mu p_\alpha + \tilde{g}_{\mu\beta}p_\nu p_\alpha + \tilde{g}_{\nu\alpha}p_\mu p_\beta) + \Pi_s(p) (g_{\mu\alpha}g_{\nu\beta} - g_{\mu\beta}g_{\nu\alpha}), \\ &= \frac{\lambda_Z^2}{M_Z^2 - p^2} (-\tilde{g}_{\mu\alpha}p_\nu p_\beta - \tilde{g}_{\nu\beta}p_\mu p_\alpha + \tilde{g}_{\mu\beta}p_\nu p_\alpha + \tilde{g}_{\nu\alpha}p_\mu p_\beta) + \dots, \end{aligned} \quad (11)$$

$$\Pi^{J=0}(p) = \Pi_{J=0}(p) = \frac{\lambda_Z^2}{M_Z^2 - p^2} + \dots, \quad (12)$$

where the notation $\tilde{g}_{\mu\nu} = g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2}$, the components $\Pi_s(p)$ are irrelevant in the present analysis [20], we add the superscripts and subscripts $J = 2, 1, 0$ to denote the total angular momentum. The pole residues λ_Z are defined by

$$\begin{aligned}\langle 0|\eta_{\mu\nu}^{t=+}(0)|Z_{J=2}(p)\rangle &= \lambda_Z \varepsilon_{\mu\nu}, \\ \langle 0|\eta_{\mu\nu}^{t=-}(0)|Z_{J=1}(p)\rangle &= \lambda_Z (\varepsilon_\mu p_\nu - \varepsilon_\nu p_\mu), \\ \langle 0|\eta(0)|Z_{J=0}(p)\rangle &= \lambda_Z,\end{aligned}\tag{13}$$

the $\varepsilon_{\mu\nu}$ and ε_μ are the polarization vectors of the tensor and axial-vector tetraquark states respectively with the following properties,

$$\begin{aligned}\sum_\lambda \varepsilon_{\alpha\beta}^*(\lambda, p) \varepsilon_{\mu\nu}(\lambda, p) &= \frac{\tilde{g}_{\alpha\mu} \tilde{g}_{\beta\nu} + \tilde{g}_{\alpha\nu} \tilde{g}_{\beta\mu}}{2} - \frac{\tilde{g}_{\alpha\beta} \tilde{g}_{\mu\nu}}{3}, \\ \sum_\lambda \varepsilon_\mu^*(\lambda, p) \varepsilon_\nu(\lambda, p) &= -\tilde{g}_{\mu\nu}.\end{aligned}\tag{14}$$

The tensor current $\eta_{\alpha\beta}^{t=-}$ has no coupling with the $J^P = 2^+$ (or tensor) tetraquark states, as the Lorentz indexes α and β are antisymmetric.

In the following, we briefly outline the operator product expansion for the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ and $\Pi(p)$ in perturbative QCD. We contract the u , d and c quark fields in the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ and $\Pi(p)$ with Wick theorem, and obtain the results:

$$\begin{aligned}\Pi_{\mu\nu\alpha\beta}(p) &= \frac{i\epsilon^{ijk}\epsilon^{imn}\epsilon^{i'j'k'}\epsilon^{i'm'n'}}{2} \int d^4x e^{ip\cdot x} \\ &\quad \left\{ \text{Tr} \left[\gamma_\mu C^{kk'}(x) \gamma_\alpha C U^{jj'T}(x) C \right] \text{Tr} \left[\gamma_\beta C^{m'n}(-x) \gamma_\nu C D^{m'mT}(-x) C \right] \right. \\ &\quad + \text{Tr} \left[\gamma_\nu C^{kk'}(x) \gamma_\beta C U^{jj'T}(x) C \right] \text{Tr} \left[\gamma_\alpha C^{m'n}(-x) \gamma_\mu C D^{m'mT}(-x) C \right] \\ &\quad \pm \text{Tr} \left[\gamma_\mu C^{kk'}(x) \gamma_\beta C U^{jj'T}(x) C \right] \text{Tr} \left[\gamma_\alpha C^{m'n}(-x) \gamma_\nu C D^{m'mT}(-x) C \right] \\ &\quad \left. \pm \text{Tr} \left[\gamma_\nu C^{kk'}(x) \gamma_\alpha C U^{jj'T}(x) C \right] \text{Tr} \left[\gamma_\beta C^{m'n}(-x) \gamma_\mu C D^{m'mT}(-x) C \right] \right\}, \\ \Pi(p) &= i\epsilon^{ijk}\epsilon^{imn}\epsilon^{i'j'k'}\epsilon^{i'm'n'} \int d^4x e^{ip\cdot x} \\ &\quad \text{Tr} \left[\gamma_\mu C^{kk'}(x) \gamma_\alpha C U^{jj'T}(x) C \right] \text{Tr} \left[\gamma^\alpha C^{m'n}(-x) \gamma^\mu C D^{m'mT}(-x) C \right],\end{aligned}\tag{15}$$

where the $\pm$ correspond to $t = \pm$ respectively, the $U_{ij}(x)$, $D_{ij}(x)$ and $C_{ij}(x)$ are the full u , d and c quark propagators respectively (the $U_{ij}(x)$ and $D_{ij}(x)$ can be written as $S_{ij}(x)$ for simplicity),

$$\begin{aligned}S_{ij}(x) &= \frac{i\delta_{ij}\not{x}}{2\pi^2 x^4} - \frac{\delta_{ij}\langle\bar{q}q\rangle}{12} - \frac{\delta_{ij}x^2\langle\bar{q}g_s\sigma Gq\rangle}{192} - \frac{ig_s G_{\alpha\beta}^a t_{ij}^a (\not{x}\sigma^{\alpha\beta} + \sigma^{\alpha\beta}\not{x})}{32\pi^2 x^2} - \frac{i\delta_{ij}x^2\not{x}g_s^2\langle\bar{q}q\rangle^2}{7776} \\ &\quad - \frac{\delta_{ij}x^4\langle\bar{q}q\rangle\langle g_s^2 GG\rangle}{27648} - \frac{1}{8}\langle\bar{q}_j\sigma^{\mu\nu}q_i\rangle\sigma_{\mu\nu} - \frac{1}{4}\langle\bar{q}_j\gamma^\mu q_i\rangle\gamma_\mu + \dots,\end{aligned}\tag{16}$$

$$\begin{aligned}C_{ij}(x) &= \frac{i}{(2\pi)^4} \int d^4k e^{-ik\cdot x} \left\{ \frac{\delta_{ij}}{k - m_c} - \frac{g_s G_{\alpha\beta}^n t_{ij}^n}{4} \frac{\sigma^{\alpha\beta}(\not{k} + m_c) + (\not{k} + m_c)\sigma^{\alpha\beta}}{(k^2 - m_c^2)^2} \right. \\ &\quad \left. + \frac{g_s D_{\alpha\beta} G_{\beta\lambda}^n t_{ij}^n (f^{\lambda\beta\alpha} + f^{\lambda\alpha\beta})}{3(k^2 - m_c^2)^4} - \frac{g_s^2 (t^a t^b)_{ij} G_{\alpha\beta}^a G_{\mu\nu}^b (f^{\alpha\beta\mu\nu} + f^{\alpha\mu\beta\nu} + f^{\alpha\mu\nu\beta})}{4(k^2 - m_c^2)^5} + \dots \right\}, \\ f^{\lambda\alpha\beta} &= (\not{k} + m_c)\gamma^\lambda(\not{k} + m_c)\gamma^\alpha(\not{k} + m_c)\gamma^\beta(\not{k} + m_c), \\ f^{\alpha\beta\mu\nu} &= (\not{k} + m_c)\gamma^\alpha(\not{k} + m_c)\gamma^\beta(\not{k} + m_c)\gamma^\mu(\not{k} + m_c)\gamma^\nu(\not{k} + m_c),\end{aligned}\tag{17}$$

and $t^n = \frac{\lambda^n}{2}$, the λ^n is the Gell-Mann matrix, $D_\alpha = \partial_\alpha - ig_s G_\alpha^n t^n$ [19], then compute the integrals both in the coordinate and momentum spaces to obtain the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ and $\Pi(p)$ therefore the QCD spectral densities. In Eq.(16), we retain the terms $\langle \bar{q}_j \sigma_{\mu\nu} q_i \rangle$ and $\langle \bar{q}_j \gamma_\mu q_i \rangle$ originate from the Fierz re-arrangement of the $\langle q_i \bar{q}_j \rangle$ to absorb the gluons emitted from the heavy quark lines to form $\langle \bar{q}_j g_s G_{\alpha\beta}^a t_{mn}^a \sigma_{\mu\nu} q_i \rangle$ and $\langle \bar{q}_j \gamma_\mu q_i g_s D_\nu G_{\alpha\beta}^a t_{mn}^a \rangle$ so as to extract the mixed condensate and four-quark condensates $\langle \bar{q} g_s \sigma G q \rangle$ and $g_s^2 \langle \bar{q} q \rangle^2$, respectively. One can consult Ref.[14] for some technical details in the operator product expansion.

Once the analytical expressions are obtained, we can take the quark-hadron duality below the continuum thresholds s_0 and perform Borel transform with respect to the variable $P^2 = -p^2$ to obtain the following QCD sum rules:

$$\lambda_Z^2 \exp\left(-\frac{M_Z^2}{T^2}\right) = \int_{4m_c^2}^{s_0} ds \rho(s) \exp\left(-\frac{s}{T^2}\right), \quad (18)$$

where

$$\rho(s) = \rho_0(s) + \rho_3(s) + \rho_4(s) + \rho_5(s) + \rho_6(s) + \rho_7(s) + \rho_8(s) + \rho_{10}(s), \quad (19)$$

the subscripts 0, 3, 4, 5, 6, 7, 8 and 10 denote the dimensions of the vacuum condensates in the operator product expansion, the T^2 denotes the Borel parameter. The explicit expressions of the QCD spectral densities $\rho_i(s)$ are given in the appendix. In this article, we carry out the operator product expansion to the vacuum condensates up to dimension-10 and discard the perturbative corrections. Furthermore, we assume vacuum saturation for the higher dimension vacuum condensates. The terms associate with $\frac{1}{T^2}$, $\frac{1}{T^4}$, $\frac{1}{T^6}$ in the QCD spectral densities ρ_i , $i = 7, 8, 10$, manifest themselves at small values of the T^2 , we have to choose large values of the T^2 to warrant convergence of the operator product expansion and appearance of the Borel platforms. In the Borel windows, the higher dimension vacuum condensates play a less important role. In summary, the higher dimension vacuum condensates play an important role in determining the Borel windows therefore the ground state masses and pole residues, though they maybe play a less important role in the Borel windows. We should take them into account consistently.

Differentiate Eq.(18) with respect to $\frac{1}{T^2}$, then eliminate the pole residues λ_Z , we obtain the QCD sum rules for the masses of the scalar, axial-vector and tensor tetraquark states,

$$M_Z^2 = \frac{\int_{4m_c^2}^{s_0} ds \frac{d}{d(-1/T^2)} \rho(s) \exp\left(-\frac{s}{T^2}\right)}{\int_{4m_c^2}^{s_0} ds \rho(s) \exp\left(-\frac{s}{T^2}\right)}. \quad (20)$$

3 Numerical results and discussions

The vacuum condensates are taken to be the standard values $\langle \bar{q} q \rangle = -(0.24 \pm 0.01 \text{ GeV})^3$, $\langle \bar{q} g_s \sigma G q \rangle = m_0^2 \langle \bar{q} q \rangle$, $m_0^2 = (0.8 \pm 0.1) \text{ GeV}^2$, $\langle \frac{\alpha_s G G}{\pi} \rangle = (0.33 \text{ GeV})^4$ at the energy scale $\mu = 1 \text{ GeV}$ [18, 19, 21]. The quark condensate and mixed quark condensate evolve with the renormalization group equation, $\langle \bar{q} q \rangle(\mu) = \langle \bar{q} q \rangle(Q) \left[\frac{\alpha_s(Q)}{\alpha_s(\mu)} \right]^{\frac{4}{3}}$ and $\langle \bar{q} g_s \sigma G q \rangle(\mu) = \langle \bar{q} g_s \sigma G q \rangle(Q) \left[\frac{\alpha_s(Q)}{\alpha_s(\mu)} \right]^{\frac{2}{27}}$.

In the article, we take the $\overline{MS}$ mass $m_c(m_c) = (1.275 \pm 0.025) \text{ GeV}$ from the Particle Data Group [22], and take into account the energy-scale dependence of the $\overline{MS}$ mass from the renormalization group equation,

$$\begin{aligned} m_c(\mu) &= m_c(m_c) \left[\frac{\alpha_s(\mu)}{\alpha_s(m_c)} \right]^{\frac{12}{25}}, \\ \alpha_s(\mu) &= \frac{1}{b_0 t} \left[1 - \frac{b_1 \log t}{b_0^2 t} + \frac{b_1^2 (\log^2 t - \log t - 1) + b_0 b_2}{b_0^4 t^2} \right], \end{aligned} \quad (21)$$

where $t = \log \frac{\mu^2}{\Lambda^2}$, $b_0 = \frac{33-2n_f}{12\pi}$, $b_1 = \frac{153-19n_f}{24\pi^2}$, $b_2 = \frac{2857-\frac{5033}{9}n_f+\frac{325}{27}n_f^2}{128\pi^3}$, $\Lambda = 213 \text{ MeV}$, 296 MeV and 339 MeV for the flavors $n_f = 5, 4$ and 3 , respectively [22].

In the conventional QCD sum rules [18, 19], there are two criteria (pole dominance and convergence of the operator product expansion) for choosing the Borel parameter T^2 and threshold parameter s_0 . The light tetraquark states cannot satisfy the two criteria, though it is not an indication of non-existence of the light tetraquark states (For detailed discussions about this subject, one can consult Ref.[23]). We impose the two criteria on the hidden charmed tetraquark states to choose the Borel parameter T^2 and threshold parameter s_0 . Furthermore, we take it for granted that the energy gap between the ground states and the first radial excited states is about 0.5 GeV .

In Fig.1, the masses of the scalar ($J = 0$), axial-vector ($J = 1$) and tensor ($J = 2$) tetraquark states are plotted with variations of the Borel parameters T^2 , energy scales μ , and continuum threshold parameters s_0 , where $s_0^{J=2} = s_0^{J=1} = 20.5 \text{ GeV}^2$, $s_0^{J=0} = 18.5 \text{ GeV}^2$, $s_0^{Z(4050)} = 21.0 \text{ GeV}^2$, and $s_0^{Z(4250)} = 22.5 \text{ GeV}^2$. From the figure, we can see that the masses decrease monotonously with increase of the energy scales. The parameters $s_0^{J=2} = s_0^{J=1} = 20.5 \text{ GeV}^2$ and $\mu = 1.8 \text{ GeV}$ can reproduce the experimental values of the masses of the $Z_c(4020)$ and $Z_c(4025)$ in the cases of $J^{PC} = 1^{+-}$ or 2^{++} assignments. In Ref.[12], we observe that the predictions based on the QCD sum rules disfavor assigning the $Z_c(4020)$ and $Z_c(4025)$ as the diquark-antidiquark type vector tetraquark states.

In Ref.[14], we observe that the energy scale $\mu = 1.5 \text{ GeV}$ is the lowest energy scale to reproduce the experimental values of the masses of the $X(3872)$ and $Z_c(3900)$ or $Z_c(3885)$. In Ref.[12], we suggest a formula to estimate the energy scales of the QCD spectral densities in the QCD sum rules for the hidden charmed tetraquark states, $\mu = \sqrt{M_{X/Y/Z}^2 - (2\mathbb{M}_c)^2}$, with the effective c -quark mass $\mathbb{M}_c = 1.8 \text{ GeV}$. The heavy tetraquark system could be described by a double-well potential with two light quarks $q'\bar{q}$ lying in the two wells respectively. In the heavy quark limit, the c (and b) quark can be taken as a static well potential, which binds the light quark q' to form a diquark in the color antitriplet channel or binds the light antiquark $\bar{q}$ to form a meson in the color singlet channel (or a meson-like state in the color octet channel). Then the heavy tetraquark states are characterized by the effective heavy quark masses $\mathbb{M}_Q$ (or constituent quark masses) and the virtuality $\sqrt{M_{X/Y/Z}^2 - (2\mathbb{M}_Q)^2}$ (or bound energy not as robust). It is natural to take the energy scale $\mu = \text{virtuality}$. The resulting energy scales are $\mu = 1.5 \text{ GeV}$ for the $Z_c(3900)$ and $X(3872)$ [14], $\mu = 3.0 \text{ GeV}$ for the $Y(4660)$ [12], $\mu = 1.8 \text{ GeV}$ for the $Z_c(4020)$ and $Z_c(4025)$ in the present case.

If the $Z(4050)$ and $Z(4250)$ are scalar tetraquark states, the energy scale $\mu = 1.8 \text{ GeV}$ is too large, see Fig.1.

- If the $Z(4050)$ is a scalar tetraquark state, the energy scale $\mu = 1.2 \text{ GeV}$ is the optimal energy scale.
- If the $Z(4250)$ is a scalar tetraquark state, the energy scale $\mu = 1.0 \text{ GeV}$ is the optimal energy scale.

In the two cases, the energy scales $\mu \ll 1.8 \text{ GeV}$, it is very odd that the energy scales of the QCD spectral densities of the $Z(4050)$ and $Z(4250)$ are much smaller than that of the $Z_c(4020)$ and $Z_c(4025)$. We can draw the conclusion tentatively that the predictions based on the QCD sum rules disfavor assigning the $Z(4050)$ and $Z(4250)$ as the $C\gamma_\mu - C\gamma^\mu$ type scalar tetraquark states.

In Refs.[2, 3], we obtain the masses $(4.36 \pm 0.18) \text{ GeV}$, $(4.37 \pm 0.18) \text{ GeV}$ and $(4.56 \pm 0.14) \text{ GeV}$ for the $C\gamma_\mu - C\gamma^\mu$, $C\gamma_5 - C\gamma_5$ and $C\gamma_\mu\gamma_5 - C\gamma^\mu\gamma_5$ type scalar $c\bar{c}d\bar{d}$ tetraquark states respectively using the QCD sum rules. In calculations, we take the mass $m_c(\mu = 1 \text{ GeV}) = (1.35 \pm 0.10) \text{ GeV}$, the energy scale $\mu = 1 \text{ GeV}$ is much smaller than that determined by the formula $\mu = \sqrt{M_{X/Y/Z}^2 - (2\mathbb{M}_c)^2}$. The masses of the $C\gamma_\mu\gamma_5 - C\gamma^\mu\gamma_5$ type scalar tetraquark states are much larger than that of the $C\gamma_\mu - C\gamma^\mu$ and $C\gamma_5 - C\gamma_5$ type scalar tetraquark states [2, 3]. If the $Z(4050)$ and $Z(4250)$ are scalar tetraquark states, they may be the $C\gamma_\mu\gamma_5 - C\gamma^\mu\gamma_5$ or $C - C$ type.

Under the Fierz re-arrangement, we can rearrange the diquark-antidiquark type currents into




Figure 1: The masses with variations of the Borel parameters T^2 and energy scales μ for the $J = 2, 1, 0$ tetraquark states. The horizontal lines denote the experimental values of the masses of the $Z_c(4020)$ and $Z_c(4025)$ in the case of $J = 1$ and $J = 2$; the experimental values of the masses of the $Z(4050)$ and $Z(4250)$ in the case of $Z(4050, J = 0)$ and $Z(4250, J = 0)$. In the figures denoted by the $Z(4050, J = 0)$ and $Z(4250, J = 0)$, we take the $Z(4050)$ and $Z(4250)$ as the $J = 0$ tetraquark states, choose the continuum threshold parameters $s_0^{Z(4050)} = (M_{Z(4050)} + 0.5)^2 \approx 21.0 \text{ GeV}^2$ and $s_0^{Z(4250)} = (M_{Z(4250)} + 0.5)^2 \approx 22.5 \text{ GeV}^2$.

J^{PC}	$T^2(\text{GeV}^2)$	$s_0(\text{GeV}^2)$	pole	$M_Z(\text{GeV})$	λ_Z
2^{++}	$2.6 - 3.0$	20.5 ± 1.0	$(50 - 73)\%$	$4.02^{+0.09}_{-0.09}$	$3.97^{+0.58}_{-0.52} \times 10^{-2} \text{GeV}^5$
1^{+-}	$2.8 - 3.2$	20.5 ± 1.0	$(51 - 72)\%$	$4.02^{+0.07}_{-0.08}$	$0.80^{+0.11}_{-0.11} \times 10^{-2} \text{GeV}^4$
0^{++}	$2.0 - 2.4$	18.5 ± 1.0	$(50 - 78)\%$	$3.85^{+0.15}_{-0.09}$	$3.15^{+0.75}_{-0.53} \times 10^{-2} \text{GeV}^5$

Table 1: The Borel parameters, continuum threshold parameters, pole contributions, masses and pole residues of the scalar, axial-vector and tensor tetraquark states.

the color singlet-singlet type + color octet-octet type currents [24], the energy scale formula can be applied to study the molecular states in the QCD sum rules [25].

We take the energy scales as $\mu = 1.8 \text{ GeV}$, 1.8 GeV and 1.4 GeV for the $J = 2, 1$ and 0 tetraquark states, respectively. In Fig.2, the contributions of the pole terms are plotted with variations of the threshold parameters s_0 and Borel parameters T^2 . In calculations, we observe that the values $s_0^{J=2} \leq 18.5 \text{ GeV}^2$, $s_0^{J=1} \leq 18.5 \text{ GeV}^2$ and $s_0^{J=0} \leq 16.5 \text{ GeV}^2$ are too small to satisfy the pole dominance condition and result in reasonable Borel windows. In Fig.3, the contributions of different terms in the operator product expansion are plotted with variations of the Borel parameters T^2 for the threshold parameters $s_0^{J=2} = s_0^{J=1} = 20.5 \text{ GeV}^2$, $s_0^{J=0} = 18.5 \text{ GeV}^2$. The contributions of the vacuum condensates of dimensions 3, 5, 6, 8, 10 change quickly with variations of the Borel parameters at the regions $T^2 < 2.4 \text{ GeV}^2$ and 2.0 GeV^2 for the $J = 2$ and 0 tetraquark states, respectively, the contributions of the vacuum condensates of dimensions 3, 5 change quickly with variations of the Borel parameter at the region $T^2 < 2.4 \text{ GeV}^2$ for the $J = 1$ tetraquark state, which does not warrant platforms for the masses.

In this article, the Borel parameters are chosen as $T^2 = (2.6 - 2.8) \text{ GeV}^2$, $(2.8 - 3.2) \text{ GeV}^2$ and $(2.0 - 2.4) \text{ GeV}^2$ for the $J = 2, 1$ and 0 tetraquark states, respectively, the continuum threshold parameters are chosen as $s_0 = (20.5 \pm 1.0) \text{ GeV}^2$, $(20.5 \pm 1.0) \text{ GeV}^2$ and $(18.5 \pm 1.0) \text{ GeV}^2$ for the $J = 2, 1$ and 0 tetraquark states, respectively. Then the convergent behavior in the operator product expansion is very good. Such Borel parameters and threshold parameters can also lead to analogous pole contributions. The Borel parameters, continuum threshold parameters and the pole contributions are shown explicitly in Table 1. The two criteria (pole dominance and convergence of the operator product expansion) of the QCD sum rules are fully satisfied, so we expect to make reasonable predictions.

Taking into account all uncertainties of the input parameters, finally we obtain the values of the masses and pole residues of the scalar, axial-vector and tensor tetraquark states, which are shown explicitly in Figs.4-5 and Table 1. The predictions $M_{J=2} = (4.02^{+0.09}_{-0.09}) \text{ GeV}$, $M_{J=1} = (4.02^{+0.07}_{-0.08}) \text{ GeV}$ are consistent with the experimental values $M_{Z_c(4025)} = (4026.3 \pm 2.6 \pm 3.7) \text{ MeV}$, $M_{Z_c(4020)} = (4022.9 \pm 0.8 \pm 2.7) \text{ MeV}$ from the BESIII collaboration [7, 8]. The present predictions favor assigning the $Z_c(4020)$ and $Z_c(4025)$ as the $J^{PC} = 1^{+-}$ and 2^{++} diquark-antidiquark type tetraquark states. More experimental data on the spin and parity are still needed to identify the $Z_c(4020)$ and $Z_c(4025)$, while the $Z(4050)$ and $Z(4250)$ still need confirmation.

In Table 1, the pole residues $\lambda_{Z(1^{+-})} = (0.80^{+0.11}_{-0.11}) \times 10^{-2} \text{ GeV}^4$ and $\lambda_{Z(2^{++})} = (3.97^{+0.58}_{-0.52}) \times 10^{-2} \text{ GeV}^5$ are quite different due to the definitions, see Eq.(13), although the interpolating currents $\eta_{\mu\nu}^{t=\pm}$ are similar. The $\lambda_{Z(1^{+-})}$ has smaller dimension of mass compared to the $\lambda_{Z(2^{++})}$, the $\lambda_{Z(0^{++})}$, $\lambda_{Z(1^{+-})} M_{Z(1^{+-})}$, $\lambda_{Z(2^{++})}$ are of the same magnitude. The correlation function $\Pi_{\mu\nu\alpha\beta}^{J=2}(p)$ ($\Pi_{\mu\nu\alpha\beta}^{J=1}(p)$) is symmetric (anti-symmetric) in exchanging the Lorentz indexes $\mu \leftrightarrow \nu$ or $\alpha \leftrightarrow \beta$, the vacuum condensates of the dimensions 8 and 10 and some other vacuum condensates disappear at the QCD side of the correlation function $\Pi_{\mu\nu\alpha\beta}^{J=1}(p)$ due to the anti-symmetry property of the Lorentz indexes, which leads to weaker current-meson coupling, $\lambda_{Z(1^{+-})} M_{Z(1^{+-})} < \lambda_{Z(2^{++})}$. If we study the strong decays of the 1^{+-} and 2^{++} tetraquark states with the three-point QCD sum rules by taking the pole residues $\lambda_{Z(1^{+-})}$ and $\lambda_{Z(2^{++})}$ as input parameters, the difference in the




Figure 2: The pole contributions with variations of the Borel parameters T^2 and threshold parameters s_0 , where the A , B , C , D , E and F denote the threshold parameters $s_0 = 18.5, 19.5, 20.5, 21.5, 22.5$ and 23.5 GeV^2 respectively for the $J = 2, 1$ tetraquark states; $16.5, 17.5, 18.5, 19.5, 20.5$ and 21.5 GeV^2 respectively for the $J = 0$ tetraquark state.




Figure 3: The contributions of different terms in the operator product expansion with variations of the Borel parameters T^2 , where the 0, 3, 4, 5, 6, 7, 8 and 10 denote the dimensions of the vacuum condensates.




Figure 4: The masses with variations of the Borel parameters T^2 , where the horizontal lines denote the experimental values of the masses of the $Z_c(4020)$ and $Z_c(4025)$.




Figure 5: The pole residues with variations of the Borel parameters T^2 .

pole residues at the hadronic side is expected to be compensated by the difference in the spectral densities at the QCD side, so the difference in the pole residues will not influence the decay widths significantly.

The BESIII collaboration observed the $Z_c^\pm(4025)$ and $Z_c^\pm(4020)$ in the following processes [7, 8],

$$\begin{aligned} e^+e^- &\rightarrow Z_c^\pm(4025)\pi^\mp \rightarrow (D^*\bar{D}^*)^\pm(0^{++}, 1^{+-}, 2^{++}, 0^{-+}, 1^{--}, 2^{-+}, 3^{--})\pi^\mp, \\ e^+e^- &\rightarrow Z_c^\pm(4020)\pi^\mp \rightarrow (h_c\pi)^\pm(1^{--}, 0^{++}, 1^{+-}, 2^{++})\pi^\mp, \end{aligned} \quad (22)$$

where we present the possible quantum numbers J^{PC} of the $(D^*\bar{D}^*)^\pm$ and $(h_c\pi)^\pm$ systems in the brackets. If the $Z_c^\pm(4025)$ and $Z_c^\pm(4020)$ are the same particle, the quantum numbers are $J^{PC} = 1^{--}, 0^{++}, 1^{+-}, 2^{++}$. On the other hand, the $Z_c^\pm(4025)\pi^\mp$ and $Z_c^\pm(4020)\pi^\mp$ systems have the quantum numbers $J^{PC} = 1^{--}$, then the survived quantum numbers of the $Z_c^\pm(4025)$ and $Z_c^\pm(4020)$ are $J^{PC} = 1^{--}, 1^{+-}$ and 2^{++} . The predictions based on the QCD sum rules reduce the possible quantum numbers of the $Z_c(4025)$ and $Z_c(4020)$ to $J^{PC} = 1^{+-}$ and 2^{++} [12]. The strong decays

$$Y(4260)/\gamma^*(4260) \rightarrow Z_c^\pm(4025/4020)(2^{++})\pi^\mp, \quad (23)$$

take place through relative D-wave, and are kinematically suppressed in the phase-space compared to the strong decays

$$Y(4260)/\gamma^*(4260) \rightarrow Z_c^\pm(4025/4020)(1^{+-})\pi^\mp. \quad (24)$$

The 2^{++} assignment is disfavored, but not excluded.

In the following, we list out the possible strong decays of the $Z_c^\pm(4025)$ and $Z_c^\pm(3900)$ in the case of the $J^{PC} = 1^{+-}$ assignment.

$$\begin{aligned} Z_c^\pm(4025)(1^{+-}) &\rightarrow h_c(1P)\pi^\pm, J/\psi\pi^\pm, \eta_c\rho^\pm, \eta_c(\pi\pi)_P^\pm, \chi_{c1}(\pi\pi)_P^\pm, (D\bar{D}^*)^\pm, (D^*\bar{D})^\pm, \\ Z_c^\pm(3900)(1^{+-}) &\rightarrow h_c(1P)\pi^\pm, J/\psi\pi^\pm, \eta_c\rho^\pm, \eta_c(\pi\pi)_P^\pm, \chi_{c1}(\pi\pi)_P^\pm, \end{aligned} \quad (25)$$

where the $(\pi\pi)_P$ denotes the P-wave $\pi\pi$ systems have the same quantum numbers of the ρ . We take the $Z_c(4025)$ and $Z_c(4020)$ as the same particle in the $J^{PC} = 1^{+-}$ assignment, and will denote them as $Z_c(4025)$. In Ref.[14], we observe that the $Z_c(3900)$ couples to the axial-vector current η_{1+-}^μ . In the following, we perform Fierz re-arrangement both in the color and Dirac-spinor spaces to obtain the result,

$$\begin{aligned} \eta_{1+-}^\mu &= \frac{\epsilon^{ijk}\epsilon^{imn}}{\sqrt{2}} \{u^j C\gamma_5 c^k \bar{d}^m \gamma^\mu C \bar{c}^n - u^j C\gamma^\mu c^k \bar{d}^m \gamma_5 C \bar{c}^n\}, \\ &= \frac{1}{2\sqrt{2}} \{i\bar{c}\bar{i}\gamma_5 c \bar{d}\gamma^\mu u - i\bar{c}\gamma^\mu c \bar{d}i\gamma_5 u + \bar{c}u \bar{d}\gamma^\mu \gamma_5 c - \bar{c}\gamma^\mu \gamma_5 u \bar{d}c \\ &\quad - i\bar{c}\gamma_\nu \gamma_5 c \bar{d}\sigma^{\mu\nu} u + i\bar{c}\sigma^{\mu\nu} c \bar{d}\gamma_\nu \gamma_5 u - i\bar{c}\sigma^{\mu\nu} \gamma_5 u \bar{d}\gamma_\nu c + i\bar{c}\gamma_\nu u \bar{d}\sigma^{\mu\nu} \gamma_5 c\}, \end{aligned} \quad (26)$$

the components $\bar{c}\sigma^{\mu\nu} c \bar{d}\gamma_\nu \gamma_5 u$, $\bar{c}\gamma^\mu c \bar{d}i\gamma_5 u$, $\bar{c}i\gamma_5 c \bar{d}\gamma^\mu u$, $\bar{c}i\gamma_5 c \bar{d}\gamma^\mu u$ couple to the $h_c(1P)\pi^\pm$, $J/\psi\pi^\pm$, $\eta_c\rho^\pm$, $\eta_c(\pi\pi)_P^\pm$, respectively. The strong decays

$$Z_c^\pm(3900)(1^{+-}) \rightarrow h_c(1P)\pi^\pm, J/\psi\pi^\pm, \eta_c\rho^\pm, \eta_c(\pi\pi)_P^\pm, \quad (27)$$

are Okubo-Zweig-Iizuka super-allowed, we take the decays to the $(\pi\pi)_P^\pm$ final states as Okubo-Zweig-Iizuka super-allowed according to the decays $\rho \rightarrow \pi\pi$. The BESIII collaboration observed no evidence of the $Z_c(3900)$ in the process $e^+e^- \rightarrow \pi^+\pi^-h_c$ at center-of-mass energies (3.90 – 4.42) GeV [8]. We expect to observe the $Z_c^\pm(3900)$ in the $h_c(1P)\pi^\pm$ final states in the futures when a large amount of events are accumulated. The components $\bar{c}\sigma^{\mu\nu} \gamma_5 u \bar{d}\gamma_\nu c$ and $\bar{c}\gamma_\nu u \bar{d}\sigma^{\mu\nu} \gamma_5 c$

couples to the scattering state $(D^* \bar{D}^*)^+$. In the nonrelativistic and heavy quark limit, the current $\bar{c}\sigma^{\mu\nu}\gamma_5 u \bar{d}\gamma_\nu c$ is reduced to the following form,

$$\begin{aligned}\bar{c}\sigma^{0j}\gamma_5 u \bar{d}\gamma_j c &\propto \xi_c^\dagger \sigma^j \zeta_u \chi_d^\dagger \vec{\sigma} \cdot \vec{k}_d \sigma^j \xi_c \propto \xi_c^\dagger \frac{\sigma^j}{2} \zeta_u \chi_d^\dagger \frac{\sigma^j}{2} \xi_c = \vec{S}_{\bar{D}^*} \cdot \vec{S}_{D^*}, \\ \bar{c}\sigma^{ij}\gamma_5 u \bar{d}\gamma_j c &\propto \epsilon^{ijk} \xi_c^\dagger \sigma^k \vec{\sigma} \cdot \vec{k}_u \zeta_u \chi_d^\dagger \vec{\sigma} \cdot \vec{k}_d \sigma^j \xi_c \propto \epsilon^{ijk} \xi_c^\dagger \frac{\sigma^k}{2} \zeta_u \chi_d^\dagger \frac{\sigma^j}{2} \xi_c = \vec{S}_{D^*} \times \vec{S}_{\bar{D}^*},\end{aligned}\quad (28)$$

where the ξ, ζ, χ are the two-component spinors of the quark fields, the $\vec{k}$ are the three-vectors of the quark fields, the σ^i are the pauli matrixes, and the $\vec{S}$ are the spin operators. It is obvious that the currents $\bar{c}\sigma^{\mu\nu}\gamma_5 u \bar{d}\gamma_\nu c$ and $\bar{c}\gamma_\nu u \bar{d}\sigma^{\mu\nu}\gamma_5 c$ couple to the $J^P = 0^+$ and 1^+ $(D^* \bar{D}^*)^+$ states. However, the strong decays $Z_c^\pm(3900)(1^{+-}) \rightarrow (D^* \bar{D}^*)^\pm$ are kinematically forbidden. The $Z_c(4025)$ and $Z_c(3900)$ have the same quantum numbers and analogous strong decays but different masses and quark configurations.

We can search for the $Z_c^\pm(4025)(1^{+-})$ in the final states $h_c(1P)\pi^\pm, \chi_{c1}\pi^\pm, J/\psi\pi^\pm, \eta_c\rho^\pm, \eta_c(\pi\pi)_P^\pm, \chi_{c1}(\pi\pi)_P^\pm$. Now we perform Fierz re-arrangement both in the color and Dirac-spinor spaces to obtain the following result,

$$\begin{aligned}\eta_{1^{+-}}^{\mu\nu} &= \frac{\epsilon^{ijk}\epsilon^{imn}}{\sqrt{2}} \{u^j C\gamma^\mu c^k \bar{d}^m \gamma^\nu C\bar{c}^n - u^j C\gamma^\nu c^k \bar{d}^m \gamma^\mu C\bar{c}^n\}, \\ &= \frac{1}{2\sqrt{2}} \{i\bar{d}u \bar{c}\sigma^{\mu\nu} c + i\bar{d}\sigma^{\mu\nu} u \bar{c}c + i\bar{d}c \bar{c}\sigma^{\mu\nu} u + i\bar{d}\sigma^{\mu\nu} c \bar{c}u \\ &\quad - \bar{c}\sigma^{\mu\nu}\gamma_5 c \bar{d}i\gamma_5 u - \bar{c}i\gamma_5 c \bar{d}\sigma^{\mu\nu}\gamma_5 u - \bar{c}\sigma^{\mu\nu}\gamma_5 u \bar{d}i\gamma_5 c - \bar{d}i\gamma_5 c \bar{c}\sigma^{\mu\nu}\gamma_5 u \\ &\quad + i\epsilon^{\mu\nu\alpha\beta} \bar{c}\gamma^\alpha \gamma_5 c \bar{d}\gamma^\beta u - i\epsilon^{\mu\nu\alpha\beta} \bar{c}\gamma^\alpha c \bar{d}\gamma^\beta \gamma_5 u \\ &\quad + i\epsilon^{\mu\nu\alpha\beta} \bar{c}\gamma^\alpha \gamma_5 u \bar{d}\gamma^\beta c - i\epsilon^{\mu\nu\alpha\beta} \bar{c}\gamma^\alpha u \bar{d}\gamma^\beta \gamma_5 c\}.\end{aligned}\quad (29)$$

The scattering states $J/\psi\pi^\pm, \eta_c\rho^\pm, \eta_c(\pi\pi)_P^\pm, \chi_{c1}(\pi\pi)_P^\pm, (DD^*)^+$ couple to the components $\bar{c}\sigma^{\mu\nu}\gamma_5 c \bar{d}i\gamma_5 u, \bar{c}i\gamma_5 c \bar{d}\sigma^{\mu\nu}\gamma_5 u, \bar{c}i\gamma_5 c \bar{d}\sigma^{\mu\nu}\gamma_5 u, \epsilon^{\mu\nu\alpha\beta} \bar{c}\gamma^\alpha \gamma_5 c \bar{d}\gamma^\beta u, \bar{c}\sigma^{\mu\nu}\gamma_5 u \bar{d}i\gamma_5 c$, respectively. The strong decays

$$Z_c^\pm(4025)(1^{+-}) \rightarrow J/\psi\pi^\pm, \eta_c\rho^\pm, \eta_c(\pi\pi)_P^\pm, \chi_{c1}(\pi\pi)_P^\pm, (DD^*)^\pm, \quad (30)$$

are Okubo-Zweig-Iizuka super-allowed. In this article, we take the decays to the $(\pi\pi)_P^\pm/(\pi\pi\pi)_P^0$ final states as Okubo-Zweig-Iizuka super-allowed according to the decays $\rho \rightarrow \pi\pi/\omega \rightarrow \pi\pi\pi$.

We can also search for the neutral partner $Z_c^0(4025)(1^{+-})$ in the following strong and electromagnetic decays,

$$\begin{aligned}Z_c^0(4025)(1^{+-}) &\rightarrow h_c(1P)\pi^0, J/\psi\pi^0, J/\psi\eta, \eta_c\rho^0, \eta_c\omega, \eta_c(\pi\pi)_P^0, \chi_{cj}(\pi\pi)_P^0, \\ &\quad \eta_c(\pi\pi\pi)_P^0, \chi_{cj}(\pi\pi\pi)_P^0, \eta_c\gamma, \chi_{cj}\gamma, (DD^*)^0,\end{aligned}\quad (31)$$

where the $(\pi\pi\pi)_P$ denotes the P-wave $\pi\pi\pi$ systems with the same quantum numbers of the ω .

On the other hand, if the $Z_c(4025)$ and $Z_c(4020)$ are different particles, we can search for the $Z_c^\pm(4025/4020)(0^{++})$ and $Z_c^\pm(4025/4020)(2^{++})$ in the following strong decays,

$$\begin{aligned}Z_c^\pm(4025/4020)(0^{++}) &\rightarrow \eta_c\pi^\pm, J/\psi\rho^\pm, J/\psi(\pi\pi)_P^\pm, \chi_{c1}\pi^\pm, D\bar{D}, D^*\bar{D}^*, \\ Z_c^\pm(4025/4020)(2^{++}) &\rightarrow \eta_c\pi^\pm, J/\psi\rho^\pm, J/\psi(\pi\pi)_P^\pm, \chi_{c1}\pi^\pm, D\bar{D}, D^*\bar{D}^*.\end{aligned}\quad (32)$$

The strong decays

$$Y(4260)/\gamma^*(4260) \rightarrow Z_c^\pm(4025/4020)(0^{++})\pi^\mp, \quad (33)$$

cannot take place. The 0^{++} assignment is excluded. If the $Z_c^\pm(4025/4020)(0^{++})$ states are observed one day, it is odd indeed.

In the following, we perform Fierz re-arrangement to the tensor and scalar currents $\eta_{2^{++}}^{\mu\nu}$ and $\eta_{0^{++}}$ both in the color and Dirac-spinor spaces to obtain the results,

$$\begin{aligned}
\eta_{2^{++}}^{\mu\nu} &= \frac{\epsilon^{ijk}\epsilon^{imn}}{\sqrt{2}} \{ u^j C \gamma^\mu c^k \bar{d}^m \gamma^\nu C \bar{c}^n + u^j C \gamma^\nu c^k \bar{d}^m \gamma^\mu C \bar{c}^n \} , \\
&= \frac{1}{2\sqrt{2}} \{ \bar{c} \gamma^\mu \gamma_5 c \bar{d} \gamma^\nu \gamma_5 u + \bar{c} \gamma^\nu \gamma_5 c \bar{d} \gamma^\mu \gamma_5 u - \bar{c} \gamma^\mu c \bar{d} \gamma^\nu u - \bar{c} \gamma^\nu c \bar{d} \gamma^\mu u \\
&\quad + \bar{c} \gamma^\mu \gamma_5 u \bar{d} \gamma^\nu \gamma_5 c + \bar{c} \gamma^\nu \gamma_5 u \bar{d} \gamma^\mu \gamma_5 c - \bar{c} \gamma^\mu u \bar{d} \gamma^\nu c - \bar{c} \gamma^\nu u \bar{d} \gamma^\mu c \\
&\quad + g_{\alpha\beta} (\bar{c} \sigma^{\mu\alpha} c \bar{d} \sigma^{\nu\beta} u + \bar{c} \sigma^{\nu\alpha} c \bar{d} \sigma^{\mu\beta} u + \bar{c} \sigma^{\mu\alpha} u \bar{d} \sigma^{\nu\beta} c + \bar{c} \sigma^{\nu\alpha} u \bar{d} \sigma^{\mu\beta} c) \\
&\quad + g^{\mu\nu} \left(\bar{c} c \bar{d} u + \bar{c} i \gamma_5 c \bar{d} i \gamma_5 u + \bar{c} \gamma_\alpha c \bar{d} \gamma^\alpha u - \bar{c} \gamma_\alpha \gamma_5 c \bar{d} \gamma^\alpha \gamma_5 u - \frac{1}{2} \bar{c} \sigma_{\alpha\beta} c \bar{d} \sigma^{\alpha\beta} u \right. \\
&\quad \left. + \bar{c} u \bar{d} c + \bar{c} i \gamma_5 u \bar{d} i \gamma_5 c + \bar{c} \gamma_\alpha u \bar{d} \gamma^\alpha c - \bar{c} \gamma_\alpha \gamma_5 u \bar{d} \gamma^\alpha \gamma_5 c - \frac{1}{2} \bar{c} \sigma_{\alpha\beta} u \bar{d} \sigma^{\alpha\beta} c \right) \} , \quad (34)
\end{aligned}$$

$$\begin{aligned}
\eta_{0^{++}} &= u^j C \gamma_\mu c^k \bar{d}^m \gamma^\mu C \bar{c}^n , \\
&= \bar{c} c \bar{d} u + \bar{c} i \gamma_5 c \bar{d} i \gamma_5 u + \frac{1}{2} \bar{c} \gamma_\alpha c \bar{d} \gamma^\alpha u - \frac{1}{2} \bar{c} \gamma_\alpha \gamma_5 c \bar{d} \gamma^\alpha \gamma_5 u \\
&\quad + \bar{c} u \bar{d} c + \bar{c} i \gamma_5 u \bar{d} i \gamma_5 c + \frac{1}{2} \bar{c} \gamma_\alpha u \bar{d} \gamma^\alpha c - \frac{1}{2} \bar{c} \gamma_\alpha \gamma_5 u \bar{d} \gamma^\alpha \gamma_5 c , \quad (35)
\end{aligned}$$

where we add the quantum numbers 2^{++} and 0^{++} as subscripts to show the J^{PC} explicitly. The currents $\eta_{2^{++}}^{\mu\nu}$ and $\eta_{0^{++}}$ couple to the $Z_c^\pm(4025/4020)(2^{++})$ and $Z_c^\pm(3850)(0^{++})$, respectively. We denote the scalar hidden charmed tetraquark states with the mass 3850 MeV as the $Z_c(3850)$, see Table 1. Then we obtain the Okubo-Zweig-Iizuka super-allowed decays by taking into account the couplings to the meson-meson pairs,

$$\begin{aligned}
Z_c^\pm(4025/4020)(2^{++}) &\rightarrow \eta_c \pi^\pm, J/\psi \rho^\pm, J/\psi(\pi\pi)_P^\pm, \chi_{c1} \pi^\pm, D\bar{D}, D^* \bar{D}^*, \\
Z_c^\pm(3850)(0^{++}) &\rightarrow \eta_c \pi^\pm, J/\psi(\pi\pi)_P^\pm, \chi_{c1} \pi^\pm, D\bar{D}. \quad (36)
\end{aligned}$$

We can search for the scalar and tensor tetraquark states in the futures at the BESIII, LHCb and Belle-II. The diquark-antidiquark type current with special quantum numbers couples to a special tetraquark state, while the current can be re-arranged both in the color and Dirac-spinor spaces, and changed to a current as a special superposition of color singlet-singlet type currents. The color singlet-singlet type currents couple to the meson-meson pairs. The diquark-antidiquark type tetraquark state can be taken as a special superposition of a series of meson-meson pairs, and embodies the net effects. The decays to its components (meson-meson pairs) are Okubo-Zweig-Iizuka super-allowed, the kinematically allowed decays take place easily.

The ground state masses of the scalar tetraquark states are $M_{J=0} = 3832$ MeV or 3723 MeV in the constituent diquark model [26], $M_{J=0} = 3852$ MeV in the relativistic quark model [27], $M_{J=0} = 3729$ MeV in the relativized quark model [28]. The present prediction $M_{J=0} = (3.85_{-0.09}^{+0.15})$ GeV is compatible with the values from Refs.[26, 27].

4 Conclusion

In this article, we distinguish the charge conjugations of the interpolating currents, calculate the contributions of the vacuum condensates up to dimension-10 in the operator product expansion, study the $C\gamma_\mu - C\gamma_\nu$ type scalar, axial-vector and tensor tetraquark states in details with the QCD sum rules. In calculations, we use the formula $\mu = \sqrt{M_{X/Y/Z}^2 - (2M_c)^2}$ suggested in our previous work to determine the energy scales of the QCD spectral densities, the μ can be interpreted as the virtuality (or bound energy not as robust) in the heavy quark limit. The predictions $M_{J=2} = (4.02_{-0.09}^{+0.09})$ GeV, $M_{J=1} = (4.02_{-0.08}^{+0.07})$ GeV are consistent with the experimental

values $M_{Z_c(4025)} = (4026.3 \pm 2.6 \pm 3.7) \text{ MeV}$, $M_{Z_c(4020)} = (4022.9 \pm 0.8 \pm 2.7) \text{ MeV}$ from the BESIII collaboration, which favor assigning the $Z_c(4020)$ and $Z_c(4025)$ as the $J^{PC} = 1^{+-}$ or 2^{++} diquark-antidiquark type tetraquark states. The prediction $M_{J=0} = (3.85^{+0.15}_{-0.09}) \text{ GeV}$ disfavors assigning the $Z(4050)$ and $Z(4250)$ as the $J^{PC} = 0^{++}$ diquark-antidiquark type tetraquark states. There is no candidate for the scalar hidden charmed tetraquark state, the prediction $M_{J=0} = (3.85^{+0.15}_{-0.09}) \text{ GeV}$ can be confronted with the experimental data in the futures at the BESIII, LHCb and Belle-II. More experimental data on the spin and parity are still needed to identify the $Z_c(4020)$ and $Z_c(4025)$, while the $Z(4050)$ and $Z(4250)$ still need confirmation. Furthermore, we discuss the strong decays of the 0^{++} , 1^{+-} , 2^{++} diquark-antidiquark type tetraquark states in details, which are of phenomenological interest. The pole residues can be taken as basic input parameters to study relevant processes of the scalar, axial-vector and tensor tetraquark states with the three-point QCD sum rules.

Appendix

The spectral densities at the level of the quark-gluon degrees of freedom,

$$\begin{aligned} \rho_0^{J=2}(s) &= \frac{1}{15360\pi^6} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z)^3 (s - \overline{m}_c^2)^2 (293s^2 - 190s\overline{m}_c^2 + 17\overline{m}_c^4) \\ &+ \frac{1}{5120\pi^6} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z)^2 (s - \overline{m}_c^2)^4, \end{aligned} \quad (37)$$

$$\rho_3^{J=2}(s) = -\frac{m_c \langle \bar{q}q \rangle}{16\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z)(1-y-z) (s - \overline{m}_c^2) (3s - \overline{m}_c^2), \quad (38)$$

$$\begin{aligned} \rho_4^{J=2}(s) &= -\frac{m_c^2}{11520\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{z}{y^2} + \frac{y}{z^2} \right) (1-y-z)^3 \\ &\quad \{56s - 17\overline{m}_c^2 + 10\overline{m}_c^4 \delta(s - \overline{m}_c^2)\} \\ &- \frac{m_c^2}{3840\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{z}{y^2} + \frac{y}{z^2} \right) (1-y-z)^2 (s - \overline{m}_c^2) \\ &- \frac{1}{15360\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z)(1-y-z)^2 (185s^2 - 208s\overline{m}_c^2 + 43\overline{m}_c^4) \\ &+ \frac{1}{7680\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z)(1-y-z) (s - \overline{m}_c^2)^2 \\ &- \frac{1}{2304\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z)(1-y-z)^2 (15s^2 - 16s\overline{m}_c^2 + 3\overline{m}_c^4) \\ &- \frac{1}{13824\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z)^3 (25s^2 - 24s\overline{m}_c^2 + 3\overline{m}_c^4) \\ &- \frac{1}{6912\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z) (25s^2 - 24s\overline{m}_c^2 + 3\overline{m}_c^4) \\ &- \frac{1}{4608\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z)^2 (s - \overline{m}_c^2)^2 \\ &- \frac{1}{6912\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (s - \overline{m}_c^2) (13s - 5\overline{m}_c^2), \end{aligned} \quad (39)$$

$$\begin{aligned}\rho_5^{J=2}(s) &= \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{32\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (2s - \bar{m}_c^2) \\ &\quad + \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{144\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) (2s - \bar{m}_c^2),\end{aligned}\quad (40)$$

$$\begin{aligned}\rho_6^{J=2}(s) &= \frac{m_c^2 \langle \bar{q} q \rangle^2}{6\pi^2} \int_{y_i}^{y_f} dy + \frac{g_s^2 \langle \bar{q} q \rangle^2}{3240\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz y z \{56s - 17\bar{m}_c^2 + 10\bar{m}_c^4 \delta(s - \bar{m}_c^2)\} \\ &\quad + \frac{g_s^2 \langle \bar{q} q \rangle^2}{3240\pi^4} \int_{y_i}^{y_f} dy y(1-y) (s - \tilde{m}_c^2) \\ &\quad - \frac{g_s^2 \langle \bar{q} q \rangle^2}{9720\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) \left\{ 45 \left(\frac{z}{y} + \frac{y}{z} \right) (2s - \bar{m}_c^2) + \left(\frac{z}{y^2} + \frac{y}{z^2} \right) \right. \\ &\quad \left. m_c^2 [19 + 20\bar{m}_c^2 \delta(s - \bar{m}_c^2)] + (y+z) [18(3s - \bar{m}_c^2) + 10\bar{m}_c^4 \delta(s - \bar{m}_c^2)] \right\} \\ &\quad - \frac{g_s^2 \langle \bar{q} q \rangle^2}{9720\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) \left\{ 15 \left(\frac{z}{y} + \frac{y}{z} \right) (2s - \bar{m}_c^2) + \left(\frac{z}{y^2} + \frac{y}{z^2} \right) \right. \\ &\quad \left. m_c^2 [6 + 5\bar{m}_c^2 \delta(s - \bar{m}_c^2)] + (y+z) [56s - 17\bar{m}_c^2 + 10\bar{m}_c^4 \delta(s - \bar{m}_c^2)] \right\},\end{aligned}\quad (41)$$

$$\begin{aligned}\rho_7^{J=2}(s) &= \frac{m_c^3 \langle \bar{q} q \rangle}{144\pi^2 T^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{y}{z^3} + \frac{z}{y^3} + \frac{1}{y^2} + \frac{1}{z^2} \right) (1-y-z) \bar{m}_c^2 \delta(s - \bar{m}_c^2) \\ &\quad - \frac{m_c \langle \bar{q} q \rangle}{48\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{y}{z^2} + \frac{z}{y^2} \right) (1-y-z) \{1 + \bar{m}_c^2 \delta(s - \bar{m}_c^2)\} \\ &\quad + \frac{m_c \langle \bar{q} q \rangle}{48\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left\{ 1 + \frac{\bar{m}_c^2}{3} \delta(s - \bar{m}_c^2) \right\} \\ &\quad + \frac{m_c \langle \bar{q} q \rangle}{432\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{1-y}{y} + \frac{1-z}{z} \right) \{1 + \bar{m}_c^2 \delta(s - \bar{m}_c^2)\} \\ &\quad - \frac{m_c \langle \bar{q} q \rangle}{288\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \{1 + \tilde{m}_c^2 \delta(s - \tilde{m}_c^2)\},\end{aligned}\quad (42)$$

$$\begin{aligned}\rho_8^{J=2}(s) &= -\frac{m_c^2 \langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle}{12\pi^2} \int_0^1 dy \left(1 + \frac{\tilde{m}_c^2}{T^2} \right) \delta(s - \tilde{m}_c^2) \\ &\quad - \frac{m_c^2 \langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle}{216\pi^2} \int_0^1 dy \frac{1}{y(1-y)} \delta(s - \tilde{m}_c^2),\end{aligned}\quad (43)$$

$$\begin{aligned}
\rho_{10}^{J=2}(s) = & \frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle^2}{96\pi^2 T^6} \int_0^1 dy \tilde{m}_c^4 \delta(s - \tilde{m}_c^2) \\
& - \frac{m_c^4 \langle \bar{q} q \rangle^2}{108 T^4} \langle \frac{\alpha_s G G}{\pi} \rangle \int_0^1 dy \left\{ \frac{1}{y^3} + \frac{1}{(1-y)^3} \right\} \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} q \rangle^2}{36 T^2} \langle \frac{\alpha_s G G}{\pi} \rangle \int_0^1 dy \left\{ \frac{1}{y^2} + \frac{1}{(1-y)^2} \right\} \delta(s - \tilde{m}_c^2) \\
& - \frac{m_c^2 \langle \bar{q} q \rangle^2}{324 T^2} \langle \frac{\alpha_s G G}{\pi} \rangle \int_0^1 dy \frac{1}{y(1-y)} \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle^2}{864 \pi^2 T^4} \int_0^1 dy \frac{1}{y(1-y)} \tilde{m}_c^2 \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle^2}{576 \pi^2 T^2} \int_0^1 dy \frac{1}{y(1-y)} \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} q \rangle^2}{108 T^6} \langle \frac{\alpha_s G G}{\pi} \rangle \int_0^1 dy \tilde{m}_c^4 \delta(s - \tilde{m}_c^2) , \tag{44}
\end{aligned}$$

$$\begin{aligned}
\rho_0^{J=1}(s) = & \frac{1}{3072\pi^6 s} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z)^3 (s - \overline{m}_c^2)^2 (49s^2 - 30s\overline{m}_c^2 + \overline{m}_c^4) \\
& + \frac{1}{3072\pi^6 s} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z)^2 (s - \overline{m}_c^2)^3 (3s + \overline{m}_c^2) , \tag{45}
\end{aligned}$$

$$\rho_3^{J=1}(s) = -\frac{m_c \langle \bar{q} q \rangle}{16\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z)(1-y-z) (s - \overline{m}_c^2) , \tag{46}$$

$$\begin{aligned}
\rho_4^{J=1}(s) = & -\frac{m_c^2}{2304\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{z}{y^2} + \frac{y}{z^2} \right) (1-y-z)^3 \\
& \left\{ 8s - \overline{m}_c^2 + \frac{5\overline{m}_c^4}{3} \delta(s - \overline{m}_c^2) \right\} \\
& -\frac{m_c^2}{2304\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{z}{y^2} + \frac{y}{z^2} \right) (1-y-z)^2 \overline{m}_c^2 \\
& -\frac{1}{9216\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (1-y-z)^2 (5s^2 - 3\overline{m}_c^4) \\
& +\frac{1}{4608\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (1-y-z) (s^2 - \overline{m}_c^4) \\
& +\frac{1}{2304\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (1-y-z)^2 (5s - 4\overline{m}_c^2) \\
& +\frac{1}{41472\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z)^3 (55s^2 - 48s\overline{m}_c^2 + 3\overline{m}_c^4) \\
& +\frac{1}{6912\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z) (5s^2 - 3\overline{m}_c^4) \\
& -\frac{1}{3456\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z)^2 (s - \overline{m}_c^2) (2s - \overline{m}_c^2) \\
& +\frac{1}{1728\pi^4 s} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (s - \overline{m}_c^2) (2s - \overline{m}_c^2) , \tag{47}
\end{aligned}$$

$$\begin{aligned}
\rho_5^{J=1}(s) = & \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{64\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) \\
& -\frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{288\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) , \tag{48}
\end{aligned}$$

$$\begin{aligned}
\rho_6^{J=1}(s) = & \frac{g_s^2 \langle \bar{q} q \rangle^2}{648\pi^4 s} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz \left\{ 8s - \overline{m}_c^2 + \frac{5\overline{m}_c^4}{3} \delta(s - \overline{m}_c^2) \right\} \\
& +\frac{g_s^2 \langle \bar{q} q \rangle^2}{1944\pi^4 s} \int_{y_i}^{y_f} dy y(1-y) \tilde{m}_c^2 \\
& -\frac{g_s^2 \langle \bar{q} q \rangle^2}{1296\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) \left\{ 3 \left(\frac{z}{y} + \frac{y}{z} \right) + \left(\frac{z}{y^2} + \frac{y}{z^2} \right) m_c^2 \delta(s - \overline{m}_c^2) \right. \\
& \left. + (y+z) [8 + 2\overline{m}_c^2 \delta(s - \overline{m}_c^2)] \right\} \\
& -\frac{g_s^2 \langle \bar{q} q \rangle^2}{11664\pi^4 s} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) \left\{ 27 \left(\frac{z}{y} + \frac{y}{z} \right) s + 11 \left(\frac{z}{y^2} + \frac{y}{z^2} \right) \right. \\
& \left. m_c^2 \overline{m}_c^2 \delta(s - \overline{m}_c^2) + (y+z) [6(8s - \overline{m}_c^2) + 10\overline{m}_c^4 \delta(s - \overline{m}_c^2)] \right\} , \tag{49}
\end{aligned}$$

$$\begin{aligned}
\rho_7^{J=1}(s) = & \frac{m_c^3 \langle \bar{q}q \rangle}{288\pi^2 T^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{y}{z^3} + \frac{z}{y^3} + \frac{1}{y^2} + \frac{1}{z^2} \right) (1-y-z) \delta(s - \bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle}{96\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{y}{z^2} + \frac{z}{y^2} \right) (1-y-z) \delta(s - \bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle}{288\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \delta(s - \bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle}{864\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{1-y}{y} + \frac{1-z}{z} \right) \delta(s - \bar{m}_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle}{576\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \delta(s - \tilde{m}_c^2), \tag{50}
\end{aligned}$$

$$\begin{aligned}
\rho_0^{J=0}(s) = & \frac{1}{256\pi^6} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z)^3 (s - \bar{m}_c^2)^2 (7s^2 - 6s\bar{m}_c^2 + \bar{m}_c^4) \\
& + \frac{1}{256\pi^6} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z)^2 (s - \bar{m}_c^2)^3 (3s - \bar{m}_c^2), \tag{51}
\end{aligned}$$

$$\rho_3^{J=0}(s) = -\frac{m_c \langle \bar{q}q \rangle}{8\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z)(1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2), \tag{52}$$

$$\begin{aligned}
\rho_4^{J=0}(s) = & -\frac{m_c^2}{192\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{z}{y^2} + \frac{y}{z^2} \right) (1-y-z)^3 \\
& \left\{ 2s - \bar{m}_c^2 + \frac{\bar{m}_c^4}{6} \delta(s - \bar{m}_c^2) \right\} \\
& - \frac{m_c^2}{384\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{z}{y^2} + \frac{y}{z^2} \right) (1-y-z)^2 (3s - 2\bar{m}_c^2) \\
& - \frac{1}{768\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (1-y-z)^2 (10s^2 - 12s\bar{m}_c^2 + 3\bar{m}_c^4) \\
& + \frac{1}{384\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (1-y-z) (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{1}{384\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (1-y-z)^2 (10s^2 - 12s\bar{m}_c^2 + 3\bar{m}_c^4) \\
& + \frac{1}{3456\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z)^3 (10s^2 - 12s\bar{m}_c^2 + 3\bar{m}_c^4) \\
& + \frac{1}{576\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (1-y-z) (10s^2 - 12s\bar{m}_c^2 + 3\bar{m}_c^4) \\
& + \frac{1}{576\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z)^2 (s - \bar{m}_c^2) (2s - \bar{m}_c^2) \\
& + \frac{1}{288\pi^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz (s - \bar{m}_c^2) (2s - \bar{m}_c^2), \tag{53}
\end{aligned}$$

$$\begin{aligned}\rho_5^{J=0}(s) &= \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{32\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (y+z) (3s - 2\bar{m}_c^2) \\ &\quad - \frac{m_c \langle \bar{q} g_s \sigma G q \rangle}{48\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) (3s - 2\bar{m}_c^2) ,\end{aligned}\quad (54)$$

$$\begin{aligned}\rho_6^{J=0}(s) &= \frac{m_c^2 \langle \bar{q} q \rangle^2}{3\pi^2} \int_{y_i}^{y_f} dy + \frac{g_s^2 \langle \bar{q} q \rangle^2}{54\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz yz \left\{ 2s - \bar{m}_c^2 + \frac{\bar{m}_c^4}{6} \delta(s - \bar{m}_c^2) \right\} \\ &\quad + \frac{g_s^2 \langle \bar{q} q \rangle^2}{324\pi^4} \int_{y_i}^{y_f} dy y(1-y) (3s - 2\tilde{m}_c^2) \\ &\quad - \frac{g_s^2 \langle \bar{q} q \rangle^2}{648\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) \left\{ 3 \left(\frac{z}{y} + \frac{y}{z} \right) (3s - 2\bar{m}_c^2) + \left(\frac{z}{y^2} + \frac{y}{z^2} \right) \right. \\ &\quad \left. m_c^2 [2 + \bar{m}_c^2 \delta(s - \bar{m}_c^2)] + (y+z) [12(2s - \bar{m}_c^2) + 2\bar{m}_c^4 \delta(s - \bar{m}_c^2)] \right\} \\ &\quad - \frac{g_s^2 \langle \bar{q} q \rangle^2}{1944\pi^4} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz (1-y-z) \left\{ 15 \left(\frac{z}{y} + \frac{y}{z} \right) (3s - 2\bar{m}_c^2) + 7 \left(\frac{z}{y^2} + \frac{y}{z^2} \right) \right. \\ &\quad \left. m_c^2 [2 + \bar{m}_c^2 \delta(s - \bar{m}_c^2)] + (y+z) [12(2s - \bar{m}_c^2) + 2\bar{m}_c^4 \delta(s - \bar{m}_c^2)] \right\} ,\end{aligned}\quad (55)$$

$$\begin{aligned}\rho_7^{J=0}(s) &= \frac{m_c^3 \langle \bar{q} q \rangle \langle \frac{\alpha_s GG}{\pi} \rangle}{144\pi^2} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{y}{z^3} + \frac{z}{y^3} + \frac{1}{y^2} + \frac{1}{z^2} \right) (1-y-z) \\ &\quad \left(1 + \frac{\bar{m}_c^2}{T^2} \right) \delta(s - \bar{m}_c^2) \\ &\quad - \frac{m_c \langle \bar{q} q \rangle \langle \frac{\alpha_s GG}{\pi} \rangle}{48\pi^2} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{y}{z^2} + \frac{z}{y^2} \right) (1-y-z) \{ 2 + \bar{m}_c^2 \delta(s - \bar{m}_c^2) \} \\ &\quad + \frac{m_c \langle \bar{q} q \rangle \langle \frac{\alpha_s GG}{\pi} \rangle}{48\pi^2} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \{ 2 + \bar{m}_c^2 \delta(s - \bar{m}_c^2) \} \\ &\quad - \frac{m_c \langle \bar{q} q \rangle \langle \frac{\alpha_s GG}{\pi} \rangle}{144\pi^2} \int_{y_i}^{y_f} dy \int_{z_i}^{1-y} dz \left(\frac{1-y}{y} + \frac{1-z}{z} \right) \{ 2 + \bar{m}_c^2 \delta(s - \bar{m}_c^2) \} \\ &\quad - \frac{m_c \langle \bar{q} q \rangle \langle \frac{\alpha_s GG}{\pi} \rangle}{288\pi^2} \int_{y_i}^{y_f} dy \{ 2 + \tilde{m}_c^2 \delta(s - \tilde{m}_c^2) \} ,\end{aligned}\quad (56)$$

$$\begin{aligned}\rho_8^{J=0}(s) &= - \frac{m_c^2 \langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle}{6\pi^2} \int_0^1 dy \left(1 + \frac{\tilde{m}_c^2}{T^2} \right) \delta(s - \tilde{m}_c^2) \\ &\quad + \frac{m_c^2 \langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle}{36\pi^2} \int_0^1 dy \frac{1}{y(1-y)} \delta(s - \tilde{m}_c^2) ,\end{aligned}\quad (57)$$

$$\begin{aligned}
\rho_{10}^{J=0}(s) = & \frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle^2}{48\pi^2 T^6} \int_0^1 dy \tilde{m}_c^4 \delta(s - \tilde{m}_c^2) \\
& - \frac{m_c^4 \langle \bar{q} q \rangle^2}{54T^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dy \left\{ \frac{1}{y^3} + \frac{1}{(1-y)^3} \right\} \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} q \rangle^2}{18T^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dy \left\{ \frac{1}{y^2} + \frac{1}{(1-y)^2} \right\} \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} q \rangle^2}{54T^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dy \frac{1}{y(1-y)} \delta(s - \tilde{m}_c^2) \\
& - \frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle^2}{144\pi^2 T^4} \int_0^1 dy \frac{1}{y(1-y)} \tilde{m}_c^2 \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} g_s \sigma G q \rangle^2}{32\pi^2 T^2} \int_0^1 dy \frac{1}{y(1-y)} \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^2 \langle \bar{q} q \rangle^2}{54T^6} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dy \tilde{m}_c^4 \delta(s - \tilde{m}_c^2), \tag{58}
\end{aligned}$$

the subscripts 0, 3, 4, 5, 6, 7, 8, 10 denote the dimensions of the vacuum condensates, $y_f = \frac{1+\sqrt{1-4m_c^2/s}}{2}$, $y_i = \frac{1-\sqrt{1-4m_c^2/s}}{2}$, $z_i = \frac{ym_c^2}{ys-\tilde{m}_c^2}$, $\bar{m}_c^2 = \frac{(y+z)m_c^2}{yz}$, $\tilde{m}_c^2 = \frac{m_c^2}{y(1-y)}$, $\int_{y_i}^{y_f} dy \rightarrow \int_0^1 dy$, $\int_{z_i}^{1-y} dz \rightarrow \int_0^{1-y} dz$ when the δ functions $\delta(s - \bar{m}_c^2)$ and $\delta(s - \tilde{m}_c^2)$ appear. The condensates $\langle \frac{\alpha_s}{\pi} GG \rangle$, $\langle \bar{q} q \rangle \langle \frac{\alpha_s}{\pi} GG \rangle$, $\langle \bar{q} q \rangle^2 \langle \frac{\alpha_s}{\pi} GG \rangle$, $\langle \bar{q} g_s \sigma G q \rangle^2$ and $g_s^2 \langle \bar{q} q \rangle^2$ are the vacuum expectations of the operators of the order $\mathcal{O}(\alpha_s)$. The four-quark condensate $g_s^2 \langle \bar{q} q \rangle^2$ comes from the terms $\langle \bar{q} \gamma_\mu t^a q g_s D_\eta G_{\lambda\tau}^a \rangle$, $\langle \bar{q}_j D_\mu^\dagger D_\nu^\dagger D_\alpha^\dagger q_i \rangle$ and $\langle \bar{q}_j D_\mu D_\nu D_\alpha q_i \rangle$, rather than comes from the perturbative corrections of $\langle \bar{q} q \rangle^2$. The condensates $\langle g_s^3 GGG \rangle$, $\langle \frac{\alpha_s GG}{\pi} \rangle^2$, $\langle \frac{\alpha_s GG}{\pi} \rangle \langle \bar{q} g_s \sigma G q \rangle$ have the dimensions 6, 8, 9 respectively, but they are the vacuum expectations of the operators of the order $\mathcal{O}(\alpha_s^{3/2})$, $\mathcal{O}(\alpha_s^2)$, $\mathcal{O}(\alpha_s^{3/2})$ respectively, and discarded. We take the truncations $n \leq 10$ and $k \leq 1$ in a consistent way, the operators of the orders $\mathcal{O}(\alpha_s^k)$ with $k > 1$ are discarded. Furthermore, the values of the condensates $\langle g_s^3 GGG \rangle$, $\langle \frac{\alpha_s GG}{\pi} \rangle^2$, $\langle \frac{\alpha_s GG}{\pi} \rangle \langle \bar{q} g_s \sigma G q \rangle$ are very small, and they can be neglected safely.

Acknowledgements

This work is supported by National Natural Science Foundation, Grant Numbers 11375063, and Natural Science Foundation of Hebei province, Grant Number A2014502017. The author would like to thank Prof. T. Huang for suggesting this subject.

References

- [1] R. Mizuk et al, Phys. Rev. **D78** (2008) 072004.
- [2] Z. G. Wang, Eur. Phys. J. **C59** (2009) 675; Z. G. Wang, Eur. Phys. J. **C62** (2009) 375; Z. G. Wang, Phys. Rev. **D79** (2009) 094027.
- [3] Z. G. Wang, Eur. Phys. J. **C67** (2010) 411.
- [4] D. Ebert, R. N. Faustov and V. O. Galkin, Eur. Phys. J. **C58** (2008) 399.
- [5] S. H. Lee, K. Morita and M. Nielsen, Nucl. Phys. **A815** (2009) 29; S. H. Lee, K. Morita and M. Nielsen, Phys. Rev. **D78** (2008) 076001; G. J. Ding, Phys. Rev. **D79** (2009) 014001; Y. R. Liu and Z. Y. Zhang, Phys. Rev. **C80** (2009) 015208; X. Liu, Z. G. Luo, Y. R. Liu and

- S. L. Zhu, Eur. Phys. J. **C61** (2009) 411; T. F. Carames, A. Valcarce and J. Vijande, Phys. Rev. **D82** (2010) 054032.
- [6] J. P. Lees et al, Phys. Rev. **D85** (2012) 052003.
- [7] M. Ablikim et al, Phys. Rev. Lett. **112** (2014) 132001.
- [8] M. Ablikim et al, Phys. Rev. Lett. **111** (2013) 242001.
- [9] G. Li, Eur. Phys. J. **C73** (2013) 2621; X. Wang, Y. Sun, D. Y. Chen, X. Liu and T. Matsuki, Eur. Phys. J. **C74** (2014) 2761.
- [10] F. K. Guo, C. Hidalgo-Duque, J. Nieves and M. P. Valderrama, Phys. Rev. **D88** (2013) 054007; J. He, X. Liu, Z. F. Sun and S. L. Zhu, Eur. Phys. J. **C73** (2013) 2635; C. Y. Cui, Y. L. Liu and M. Q. Huang, Eur. Phys. J. **C73** (2013) 2661; W. Chen, T. G. Steele, M. L. Du and S. L. Zhu, Eur. Phys. J. **C74** (2014) 2773; K. P. Khemchandani, A. Martinez Torres, M. Nielsen and F. S. Navarra, Phys. Rev. **D89** (2014) 014029; A. Martinez Torres, K. P. Khemchandani, F. S. Navarra, M. Nielsen and E. Oset, Phys. Rev. **D89** (2014) 014025.
- [11] C. F. Qiao and L. Tang, Eur. Phys. J. **C74** (2014) 2810.
- [12] Z. G. Wang, Eur. Phys. J. **C74** (2014) 2874. .
- [13] E. S. Swanson, Phys. Rept. **429** (2006) 243; S. Godfrey and S. L. Olsen, Ann. Rev. Nucl. Part. Sci. **58** (2008) 51; M. B. Voloshin, Prog. Part. Nucl. Phys. **61** (2008) 455; N. Drenska, R. Faccini, F. Piccinini, A. Polosa, F. Renga and C. Sabelli, Riv. Nuovo Cim. **033** (2010) 633; N. Brambilla et al, Eur. Phys. J. **C71** (2011) 1534.
- [14] Z. G. Wang and T. Huang, Phys. Rev. **D89** (2014) 054019.
- [15] A. De Rujula, H. Georgi and S. L. Glashow, Phys. Rev. **D12** (1975) 147; T. DeGrand, R. L. Jaffe, K. Johnson and J. E. Kiskis, Phys. Rev. **D12** (1975) 2060.
- [16] R. L. Jaffe and F. Wilczek, Phys. Rev. Lett. **91** (2003) 232003; R. L. Jaffe, Phys. Rept. **409** (2005) 1.
- [17] Z. G. Wang, Eur. Phys. J. **C71** (2011) 1524; R. T. Kleiv, T. G. Steele and A. Zhang, Phys. Rev. **D87** (2013) 125018.
- [18] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147** (1979) 385; Nucl. Phys. **B147** (1979) 448.
- [19] L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127** (1985) 1.
- [20] Z. G. Wang, Eur. Phys. J. **C73** (2013) 2533.
- [21] P. Colangelo and A. Khodjamirian, hep-ph/0010175; B. L. Ioffe, Prog. Part. Nucl. Phys. **56** (2006) 232.
- [22] J. Beringer et al, Phys. Rev. **D86** (2012) 010001.
- [23] Z. G. Wang, Nucl. Phys. **A791** (2007) 106.
- [24] M. Nielsen, F. S. Navarra and S. H. Lee, Phys. Rept. **497** (2010) 4.
- [25] Z. G. Wang, Eur. Phys. J. **C63** (2009) 115; Z. G. Wang, Z. C. Liu and X. H. Zhang, Eur. Phys. J. **C64** (2009) 373; Z. G. Wang, Phys. Lett. **B690** (2010) 403; Z. G. Wang and X. H. Zhang, Commun. Theor. Phys. **54** (2010) 323; Z. G. Wang and X. H. Zhang, Eur. Phys. J. **C66** (2010) 419.

- [26] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. **D71** (2005) 014028.
- [27] D. Ebert, R. N. Faustov and V. O. Galkin, Phys. Lett. **B634** (2006) 214.
- [28] M. V. Carlucci, F. Giannuzzi, G. Nardulli, M. Pellicoro and S. Stramaglia, Eur. Phys. J. **C57** (2008) 569.