

International Journal of Modern Physics: Conference Series
 © World Scientific Publishing Company

Lattice Input on the Inclusive τ Decay V_{us} Puzzle

P. A. BOYLE, L. DEL DEBBIO and R.J. HUDSPITH

*Physics and Astronomy, University of Edinburgh
 Edinburgh EH9 3JZ, UK*

N. GARRON

*School of Mathematics, Trinity College
 Dublin 2, Ireland*

E. KERRANE

*Instituto de Física Teórica UAM/CSIC, Universidad Autónoma de Madrid
 Cantoblanco E-28049 Madrid, Spain*

K. MALTMAN*and J.M. ZANOTTI

*CSSM, University of Adelaide,
 Adelaide, SA 5005, Australia*

Received Day Month Year

Revised Day Month Year

Recent analyses of flavor-breaking hadronic- τ -decay-based sum rules produce values of $|V_{us}| \sim 3\sigma$ low compared to 3-family unitarity expectations. An unresolved systematic issue is the significant variation in $|V_{us}|$ produced by different prescriptions for treating the slowly converging $D = 2$ OPE series. We investigate the reliability of these prescriptions using lattice data for various flavor-breaking correlators and show the fixed-scale prescription is clearly preferred. Preliminary updates of the conventional τ -based, and related mixed τ -electroproduction-data-based, sum rule analyses incorporating B-factory results for low-multiplicity strange τ decay mode distributions are then performed. Use of the preferred FOPT $D = 2$ OPE prescription is shown to significantly reduce the discrepancy between 3-family unitarity expectations and the sum rule results.

The conventional inclusive hadronic τ decay determination of $|V_{us}|$ ¹ is obtained by applying the finite energy sum rule (FESR) relation, involving polynomial weight $w(s)$ and kinematic-singularity-free correlator $\Pi(s)$ with spectral function $\rho(s)$,

$$\int_0^{s_0} w(s) \rho(s) ds = -\frac{1}{2\pi i} \oint_{|s|=s_0} w(s) \Pi(s) ds, \quad (1)$$

to the flavor-breaking (FB) difference $\Delta\Pi_\tau \equiv [\Pi_{V+A;ud}^{(0+1)} - \Pi_{V+A;us}^{(0+1)}]$, where

*Permanent address: Math and Statistics, York University, Toronto, ON Canada M3J 1P3

2 *K. MALTMAN*

$\Pi_{V/A;ij}^{(J)}(s)$ are the spin $J = 0, 1$ components of the flavor ij , vector (V) or axial vector (A) current-current 2-point functions. The spectral functions, $\rho_{V/A;ij}^{(0+1)}$, hence also $\Delta\rho_\tau$, are related to the normalized differential decay distributions, $dR_{V/A;ij}/ds$, of flavor ij V- or A-current-induced τ decay widths, $R_{V/A;ij} \equiv \Gamma[\tau^- \rightarrow \nu_\tau \text{ hadrons}_{V/A;ij}(\gamma)]/\Gamma[\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e(\gamma)]$, by

$$dR_{V/A;ij}/ds = 12\pi^2 |V_{ij}|^2 S_{EW} \left[w_\tau(y_\tau) \rho_{V/A;ij}^{(0+1)}(s) - w_L(y_\tau) \rho^{(0)}(s) \right] / m_\tau^2, \quad (2)$$

with $y_\tau = s/m_\tau^2$, V_{ij} the ij CKM matrix element, $w_\tau(y) = (1-y)^2(1+2y)$, $w_L(y) = y(1-y)^2$, and S_{EW} a short-distance electroweak correction factor. The $J = 0$ (longitudinal) contributions in (2) are well known phenomenologically and, due to problems with the corresponding $D = 2$ OPE series, usually subtracted from dR/ds ^{1,2}. The subtracted result, $dR_{V/A;ij}^{(0+1)}/ds$, allows the construction of $J = 0 + 1$ reweighted analogues, $R_{V+A;ij}^w(s_0) = \int_0^{s_0} ds [w(s)/w_\tau(y_\tau)] dR_{V+A;ij}^{(0+1)}/ds$, for any $w(s)$ and $s_0 < m_\tau^2$. Defining $\delta R_{V+A}^w(s_0) = [R_{V+A;ud}^w(s_0)/|V_{ud}|^2] - [R_{V+A;us}^w(s_0)/|V_{us}|^2]$, one has, for s_0 large enough to allow use of the OPE on the RHS of (1), ¹

$$|V_{us}| = \sqrt{R_{V+A;us}^w(s_0) / \left[\frac{R_{V+A;ud}^w(s_0)}{|V_{ud}|^2} - \delta R_{V+A}^{w,OPE}(s_0) \right]}. \quad (3)$$

This relation has usually been employed in un-reweighted form, with $w = w_\tau$, and the single value $s_0 = m_\tau^2$ ¹. This has the advantage that $R_{V+A;ud,us}^{w_\tau}(m_\tau^2)$ is determinable from branching fraction information alone, but the disadvantage of precluding tests of the s_0 - and $w(s)$ -independence of the analysis, which could otherwise be used to investigate potential systematic uncertainties (in particular, those associated with the treatment of OPE contributions). Such self-consistency tests were carried out in Refs. 2, 3, 4, and non-trivial $w(s)$ - and s_0 -dependences observed, suggesting shortcomings in the experimental data and/or OPE representation.

The most obvious potential OPE problem lies in the rather slow convergence of the $D = 2$ OPE series. In terms of the running \overline{MS} quantities $m_s(Q^2)$ and $\bar{a} \equiv \alpha_s(Q^2)/\pi$, the $D = 2$ series, which is known to 4-loops, is given by

$$[\Delta\Pi_\tau(Q^2)]_{D=2}^{OPE} = \frac{3}{2\pi^2} \frac{m_s^2(Q^2)}{Q^2} \sum_{k=0} c_k^\tau \bar{a}^k \quad (4)$$

with $c_k^\tau = 1, 7/3, 19.93, 208.75$ for $k = 0 \cdots 3$ ⁵. Since $\bar{a}(m_\tau^2) \simeq 0.10$, $c_3^\tau \bar{a}^3 > c_2^\tau \bar{a}^2$ at the spacelike point on the contour for all $s_0 \leq m_\tau^2$. The problematic convergence complicates the assessment of $D = 2$ truncation errors, and manifests itself, e.g., in the ~ 0.0020 difference in $|V_{us}|$ values obtained using two alternate (CIPT or FOPT) versions of the 4-loop-truncated, w_τ -weighted series.

An alternate determination employs the FB combination $\Delta\Pi_{\tau-EM} \equiv 9\Pi_{EM} - 5\Pi_{ud;V}^{(0+1)} + \Pi_{ud;A}^{(0+1)} - \Pi_{us;V+A}^{(0+1)}$ in place of $\Delta\Pi_\tau$ ⁶. Inclusive electroproduction cross-sections fix the electromagnetic (EM) spectral function. By construction, the $\Delta\Pi_{\tau-EM}$ $D = 2$ series is strongly suppressed, having the form (4), with $c_k^\tau \rightarrow$

$c_k^{\tau-EM} = 0, -1/3, -4.384, -44.943$ for $k = 0 \cdots 3$. The $D = 4$ series is also strongly suppressed. OPE contributions to $\Delta\Pi_{\tau-EM}$ FESRs, hence also estimated OPE errors, are thus very small⁶, and the resulting $|V_{us}|$ errors essentially entirely experimental. A check of this predicted suppression is thus of interest.

We investigate the relative merits of the fixed-scale (FOPT-like) and local-scale ($\mu^2 = Q^2$, i.e., CIPT-like) treatments of the $\Delta\Pi_\tau$ $D = 2$ series, and the level of $\Delta\Pi_{\tau-EM}$ suppression, by comparing OPE expectations and lattice data for the two correlator combinations over a range of Euclidean Q^2 . Five RBC/UKQCD domain wall fermion ensembles are employed, three, with $m_\pi = 293, 349, 399$ MeV, having $1/a = 2.31$ GeV⁸, and two, with $m_\pi = 171, 248$ MeV, having $1/a = 1.37$ GeV⁷. For technical reasons, conserved-local versions of the flavor us 2-point functions are numerically challenging and hence, for $\Delta\Pi_\tau$, local-local versions are used. To check that this does not produce residual lattice artifacts which would impact our conclusions, we have also performed the OPE-lattice comparison, using conserved-local data, for the alternate flavor-diagonal FB combination $\Delta\Pi_{diag} \equiv \Pi_{V;\ell\ell} - \Pi_{V;ss}$, whose $D = 2$ series is very similar to that of $\Delta\Pi_\tau$ ($c_k^\tau \rightarrow c_k^{diag} = 1, 8/3, 24.32, 253.69$ for $k = 0 \cdots 3$ in (4)). The results confirm those of the local-local study.

Representative OPE-lattice data comparisons for $\Delta\Pi_\tau$ are shown, for the $1/a = 2.13$ GeV, $m_\pi = 293$ MeV ensemble, in Fig. 1. The left (right) panel comparison employs the fixed-scale (local-scale) prescription for the $D = 2$ OPE series. The fixed-scale versions match much better the Q^2 dependence of the lattice results, with the 3-loop-truncated version thereof best matching the overall normalization.





Fig. 1. OPE and lattice $\Delta\Pi_\tau$ data, $1/a = 2.31$ GeV, $m_\pi = 293$ MeV ensemble, $O(\bar{a}^{1,2,3})$ $D = 2$ OPE truncation, fixed-scale (left panel) or local-scale (right panel) $D = 2$ prescription

The comparison of lattice data for $\Delta\Pi_\tau$ and $\Delta\Pi_{\tau-EM}$ confirms the very strong suppression of $\Delta\Pi_{\tau-EM}$ ⁴ (see Ref. 4 for the relevant figure).

4 *K. MALTMAN*Fig. 2. $|V_{us}|$ from preliminary updates of the FB $\Delta\Pi_\tau$ and $\Delta\Pi_{\tau-EM}$ FESRs

We turn to preliminary updates of the $|V_{us}|$ analyses. For the $D = 2$ OPE series, we employ the 3-loop-truncated FOPT prescription favored by lattice data, and for the ud spectral integrals, OPAL data⁹, as updated in Ref. 10. For the us spectral integrals, recent B-factory results are used for the $K\pi$ ¹¹, $K^-\pi^-\pi^+$ ¹² and $K_s\pi^-\pi^0$ ¹³ exclusive mode distributions, and ALEPH results¹⁴, updated for current branching fractions (BFs), for all other modes. Contributions from the latter lie higher in the spectrum, and have much larger errors. The B-factory distributions are unit normalized, and also require current BF's for their overall scales. We work with BF's obtained in a $\pi_{\mu 2}$, $K_{\mu 2}$ -constrained HFAG fit, supplemented by the update to $B[\tau^- \rightarrow K_s^0 \pi^- \pi^0 \nu_\tau]$ produced by the recent Belle result¹³. Other non-trivial shifts in the us BF's also remain possible. To illustrate the changes to $|V_{us}|$ that could result, we consider also an alternate set of us BF's with the recent larger, but not yet finalized, BaBar results¹⁵ for $B[\tau^- \rightarrow K^- n \pi^0 \nu_\tau]$, $n \leq 3$, used in place of those of the HFAG fit. The first set of us BF's is labelled “ us BF set #1” below, the second, alternate set “ us BF set #2”. Changes to the us BF's alter the inclusive us spectral distribution, and hence can affect both the magnitude of $|V_{us}|$ and the s_0 -dependence of the results. The significantly larger preliminary BaBar $K^-\pi^0$ BF is particularly relevant for the FB FESRs considered here, which weight more strongly the low- s part of the spectrum. We consider FESRs employing the weights w_τ and $w_2(y) = (1 - y)^2$. w_2 weights less strongly the higher- s , large-error region of the us spectral distribution. Differences between results obtained using the two different weights can thus point to issues with the us spectral distribution.

$|V_{us}|$ results obtained from the w_τ and w_2 versions of the $\Delta\Pi_\tau$ FESR are shown, as a function of s_0 , and also the choice of the input us BF set, in the left panel of Fig. 2. Similar results for the $\Delta\Pi_{\tau-EM}$ FESR are shown in the right panel. w_2 results, which are less sensitive to the large-error high- s region, show better s_0 -stability in both cases. For w_τ , s_0 -stability is also better for the $\Delta\Pi_{\tau-EM}$ case,

where OPE contributions are suppressed. The convergence of w_τ results to the more stable w_2 ones as $s_0 \rightarrow m_\tau^2$, seen for both the $\Delta\Pi_\tau$ and $\Delta\Pi_{\tau-EM}$ FESRs, suggests the possibility of residual OPE problems in the w_τ case, where cancellations on the contour play a larger role. Finally we note that results obtained using the FOPT prescription preferred by the lattice data agree better with 3-family unitarity expectations than do those (not shown here) obtained using CIPT, as do those obtained using us BF set #2 in place of us BF set #1. More details of these analyses will be presented elsewhere.

We close by stressing the preference for FOPT over CIPT for the $D = 2$ OPE series. The prescription which underlies CIPT (of summing logarithmic terms to all orders while truncating the series of non-logarithmic terms), though plausible, is motivated by heuristic arguments not generally valid for divergent series¹⁶, and performs poorly when tested against lattice data for the FB correlators.

Acknowledgments

Computations were performed using the STFC's DiRAC facilities at Swansea and Edinburgh. PAB, LDD, and RJH were supported by an STFC Consolidated Grant, and by the EU under Grant Agreement PITN-GA-2009-238353 (ITN STRONGnet); EK by the Comunidad Autònoma de Madrid under the program HEPHACOS S2009/ESP-1473 and the EU under Grant Agreement PITN-GA-2009-238353 (ITN STRONGnet); KM by NSERC (Canada); and JMZ by the Australian Research Council grant FT100100005.

References

1. E. Gamiz *et al.*, *JHEP* **0301**, 060 (2003); *Phys. Rev. Lett.* **94**, 011803 (2005); *PoS KAON (2008)*, 008 (2008); E. Gamiz, arXiv:1303.3971 [hep-ph].
2. K. Maltman, C.E. Wolfe, *Phys. Lett.* **B639**, 283 (2006), *ibid.* *B650*, 27 (2007); K. Maltman *et al.*, *Int. J. Mod. Phys.* **A23**, 3191 (2008), *Nucl. Phys. Proc. Suppl.* **189**, 175 (2009).
3. K. Maltman, *Nucl. Phys. Proc. Suppl.* **218**, 146 (2011).
4. P.A. Boyle *et al.*, *PoS Confinement X (2012)*, 100 (2012).
5. K.G. Chetyrkin and A. Kwiatkowski, *Z. Phys.* **C59**, 525 (1993) and hep-ph/9805232; P.A. Baikov, K.G. Chetyrkin and J.H. Kuhn, *Phys. Rev. Lett.* **95**, 012003 (2005).
6. K. Maltman, *Phys. Lett.* **B672**, 257 (2009).
7. R. Arthur *et al.*, *Phys. Rev.* **D87**, 094514 (2013).
8. Y. Aoki *et al.*, *Phys. Rev.* **D83**, 074508 (2011).
9. K. Ackerstaff *et al.* (OPAL Collab.), *Eur. Phys. J.* **C7**, 571 (1999).
10. D. Boito *et al.*, *Phys. Rev.* **D85**, 093015 (2012).
11. B. Aubert, *et al.* (BaBar Collab.), *Phys. Rev.* **D76**, 051104 (2007) 051104; D. Epifanov, *et al.* (Belle Collab.), *Phys. Lett.* **B654**, 65 (2007).
12. I.M. Nugent (for the BaBar Collaboration), arXiv:1301.7105 [hep-ex].
13. S. Ryu (for the Belle Collaboration), arXiv:1302.4565 [hep-ex].
14. R. Barate *et al.* (ALEPH Collab.), *Eur. Phys. J.* **C11**, 599 (1999).
15. A. Adametz, U. Heidelberg PhD thesis, July 2011 and the BaBar Collab., in progress.
16. M. Beneke and M. Jamin, *JHEP* **0809**, 044 (2008).