

Kaon Freeze-out Dynamics in $\sqrt{s_{NN}}=200$ GeV Au+Au Collisions at RHIC

MICHAL ŠUMBERA FOR THE STAR COLLABORATION

Nuclear Physics Institute ASCR, 250 68 Řež, Czech Republic

Measurements of three-dimensional correlation functions of like-sign low transverse momentum kaon pairs from Au+Au collisions at top RHIC energy $\sqrt{s_{NN}}=200$ GeV are presented. The extracted kaon source function is narrower than the pion one and does not have the long tail along the pair transverse momentum direction. This indicates a much smaller role of long-lived resonance decays and/or of the emission duration on kaon emission. Three-dimensional Gaussian shape of the kaon source function can be adequately reproduced by Therminator simulations with resonance contributions taken into account. Comparison to pion data at the same energy reveals that the kaon Gaussian radii in the outward and sideward directions scale with the transverse mass m_T . In the longitudinal direction, unlike at lower SPS energies, the Gaussian radii do not seem to follow the exact m_T scaling between kaons and pions.

1. Introduction

The momentum correlations of particles at small relative momenta in their center-of-mass system contain an important information about space-time characteristics of the production process on a femtometer scale, so serving as a correlation femtoscopy tool (see, e.g., [1, 2] and references therein). For non-interacting identical particles, like photons, these correlations result from the interference of the production amplitudes due to symmetrization requirement of quantum statistics (QS). Additional important source of femtoscopic correlations comes from Coulomb and strong final state interaction (FSI). It provides an important information on coalescence femtoscopy and correlation femtoscopy with unlike particles, including the studies of space-time asymmetries in particle production and strong interaction between specific particles [1].

In heavy ion experiments two-particle interferometry with identical charged hadrons has been for a long time used as a reliable technique to extract the space-time characteristics of the hot expanding fireball [1, 3]. Object of study is three-dimensional (3D) correlation function $C(\mathbf{q})$, where $2\mathbf{q}$ is the

difference between the momenta of the two particles in the pair center-of-mass system (PCMS). The correlation function is defined as the ratio of the 3D relative momentum distribution of particle pairs from the same event to pairs constructed from mixed events. In the following a right-handed Cartesian coordinate system with z (*long*) parallel to the beam direction, x (*out*) pointing in the direction of the pair total transverse momentum and y (*side*) perpendicular to x and z will be used. $C(\mathbf{q})$ is related to the probability $S(\mathbf{r})$ to emit a pair of particles with a pair separation vector $\mathbf{r}$ in the PCMS via Koonin-Pratt equation [3]:

$$C(\mathbf{q}) - 1 \equiv R(\mathbf{q}) = \int d\mathbf{r} K(\mathbf{q}, \mathbf{r}) S(\mathbf{r}), \quad K(\mathbf{q}, \mathbf{r}) \equiv |\phi(\mathbf{q}, \mathbf{r})|^2 - 1. \quad (1)$$

The wave function of relative motion of two particle $\phi(\mathbf{q}, \mathbf{r})$ incorporates both QS and FSI effects. The bias arising from the frequently used Gaussian assumption on the source shape [1, 3] can be avoided if we directly extract $S(\mathbf{r})$ inverting Eq. (1) numerically [4, 5]. No assumption on the source shape is thus needed.

In [6] data on femtoscopic correlations of like-sign charged pions from 20% most central Au+Au collisions at $\sqrt{s_{NN}}=200$ GeV were analysed exploiting expansion of the source function $S(\mathbf{r})$ into a Cartesian harmonic basis [5, 7]:

$$S(\mathbf{r}) = \sum_{l, \alpha_1 \dots \alpha_l} S_{\alpha_1 \dots \alpha_l}^l(r) A_{\alpha_1 \dots \alpha_l}^l(\Omega_{\mathbf{r}}), \quad (2)$$

where $l = 0, 1, 2, \dots$, $\alpha_i = x, y$ or z , $A_{\alpha_1 \dots \alpha_l}^l(\Omega_{\mathbf{q}})$ are Cartesian harmonic basis elements and $\Omega_{\mathbf{q}}$ is the solid angle in $\mathbf{q}$ space. Significant non-Gaussian features including a long range tail in *out* and *long* directions in the pion source function were found. Model comparisons were used to study lifetime and emission duration of expanding fireball. Sizeable emission time differences between emitted pions were required to allow models to be successfully matched to these tails. However, an interpretation of pion correlations in terms of pure hydrodynamic evolution is complicated by significant contributions from later stages of the reaction, such as decays of long-lived resonances and anomalous diffusion from rescattering [8].

A purer probe of the fireball decay could be obtained with kaons which have less contribution from long life-time resonances and suffer less rescattering than pions. Their lower yields, however, make it difficult to carry out a detailed 3D source shape analysis. A one-dimensional kaon source image measurement for the same colliding system as in Ref. [6] was reported by the PHENIX collaboration [10]. The measurement corresponds to a fairly broad range of the pair transverse momentum $2k_T$, which makes the interpretation more ambiguous. In particular, information about the transverse expansion of the system contained in the k_T -dependence of the emission radii is lost. The one-dimensional nature of the measurement also less constrains the model predictions than would be available from a 3D

measurement. A different aspect of the fireball expansion can be addressed by studying the k_T -dependence of the source size. Data at SPS energies [9], as well as at RHIC [10] at relatively higher k_T values, showed a scaling behaviour between pions and kaons, as expected from perfect hydrodynamics [11].

In this contribution we present recent STAR 3D analyses [12] of the shape and k_T -dependent size of the kaon source at mid-rapidity using low transverse momentum like-sign kaon pairs produced in $\sqrt{s_{NN}}=200$ GeV central Au+Au collisions.

2. Data analysis

The kaon source shape was analyzed using 4.6 million 0–20% central events from 2004, and 16 million 0–20% central events from 2007. The k_T -dependent analysis was carried out using 6.6 million 0–30% central events from 2004. The STAR Time Projection Chamber was used to select charged kaons with rapidity $|y| < 0.5$ and transverse momenta $0.1 < p_T < 1.0$ GeV/c. Only pairs with $0.2 < k_T < 0.36$ GeV/c were accepted. In the k_T -dependent analysis, kaon pairs were collected in two bins: $0.2 < k_T < 0.36$ GeV/c and $0.36 < k_T < 0.48$ GeV/c. The 3D correlation function $C(\mathbf{q})$ defined via Eq. (1) was constructed as a ratio of the $N_{\text{same}}(\mathbf{q})$, for K^+K^+ and K^-K^- pairs in the same event to $N_{\text{mixed}}(\mathbf{q})$. $C(\mathbf{q})$ is flat and normalized to unity over $60 < |\mathbf{q}| < 100$ MeV/c. For further experimental details see [12].

From the measured 3D correlation function correlation moments

$$R_{\alpha_1 \dots \alpha_l}^l(q) = \frac{(2l+1)!!}{l!} \int \frac{d\Omega_{\mathbf{q}}}{4\pi} A_{\alpha_1 \dots \alpha_l}^l(\Omega_{\mathbf{q}}) R(\mathbf{q}) \quad (3)$$

were extracted. The lowest correlation moment R^0 agrees with the direction independent correlation function $R(q)$ within statistical errors. Even moments with $l > 4$ were found to be consistent with zero within statistical uncertainty. As expected from symmetry considerations, the same was also found for odd moments. Therefore in this analysis, the sum in Eq. (2) is truncated at $l = 4$ and expressed in terms of independent moments only. Up to order 4, there are 6 independent moments: R^0 , R_{xx}^2 , R_{yy}^2 , R_{xxx}^4 , R_{yyy}^4 and R_{xxyy}^4 . Dependent moments are obtained from independent ones [5, 7].

Fitting the truncated series to the measured 3D correlation function with a 3D Gaussian,

$$S^G(r_x, r_y, r_z) = \frac{\lambda}{(2\sqrt{\pi})^3 R_x R_y R_z} \exp\left[-\left(\frac{r_x^2}{4R_x^2} + \frac{r_y^2}{4R_y^2} + \frac{r_z^2}{4R_z^2}\right)\right], \quad (4)$$

has yielded the independent moments as a function of q . In Eq.(4) R_x , R_y and R_z are the characteristic radii of the source in the *out*, *side* and *long* directions, and λ represents the overall correlation strength. While λ may be

sensitive to feed-down from long-lived resonance decays, remainder sample contamination or track splitting/merging not removed by purity and track quality cuts, the radii are virtually independent of these effects. Technically, the fit is carried out as a simultaneous fit on the even independent moments up to $l=4$, yielding $\chi^2/ndf=1.7$ in the source shape analysis, $\chi^2/ndf=1.1$ and $\chi^2/ndf=1.3$ in the $0.2 < k_T < 0.36$ GeV/ c and $0.36 < k_T < 0.48$ GeV/ c bins in the k_T -dependent analysis, respectively. The three Gaussian radii and the amplitude obtained from this fit are listed in Table 1.

Table 1. Parameters obtained from the 3D Gaussian source function fits for the different datasets. The first errors are statistical, the second errors are systematic.

Year	2004+2007	2004	
Centrality	0%–20%	0%–30%	
k_T [GeV/ c]	0.2–0.36	0.2–0.36	0.36–0.48
R_x [fm]	$4.8 \pm 0.1 \pm 0.2$	$4.3 \pm 0.1 \pm 0.4$	$4.5 \pm 0.2 \pm 0.3$
R_y [fm]	$4.3 \pm 0.1 \pm 0.1$	$4.0 \pm 0.1 \pm 0.3$	$3.7 \pm 0.1 \pm 0.1$
R_z [fm]	$4.7 \pm 0.1 \pm 0.2$	$4.3 \pm 0.2 \pm 0.4$	$3.6 \pm 0.2 \pm 0.3$
λ	$0.49 \pm 0.02 \pm 0.05$	$0.39 \pm 0.01 \pm 0.09$	$0.27 \pm 0.01 \pm 0.04$

The shape assumption was tested using a double Gaussian trial function and by pushing the fit parameters to the edges of their errors. Other systematic errors were obtained under varying conditions including magnetic field, data collection periods, charge and various sample purity selections. The systematic errors are largely governed by the limited statistics available.

3. Results

The source function profiles in the x , y and z directions are shown on the left panel of Figure 1 (circles). The two solid curves around the Gaussian source function profiles represent the error band arising from the statistical and systematic errors on the 3D Gaussian fit. Note that the latter becomes important for large r values only. The 3D pion source functions from PHENIX [6] are shown for comparison purposes (squares). While the Gaussian radii are similar, there is a striking difference between the source shapes of the two particle species, especially in the *out* direction. Note that the PHENIX and STAR pion measurements are fully consistent [14]. We have used the STAR tune of the Therminator Blast-Wave model (solid triangles) [15, 16] to gain a better understanding of this difference. The simulation reproduces the source function profiles with emission duration $\Delta\tau=0$ (solid upward-pointing triangles). However, with resonance contribution switched off, Therminator gives a distribution that is narrower than the measurement (empty triangles). Also note that the pion source func-

tion is reproduced with Therminator only when non-zero emission duration is assumed [6]. Recent simulations of the kaon source function [17] with the hydrokinetic model (HKM) [18] show a good agreement in the *side* direction, although it is slightly over the measurements at larger radii in *out* and *long* (downward-pointing triangles).




Fig. 1. (Color online) Left: Kaon source function profiles extracted from the data (solid circles) compared to 3D pion source function (squares) from PHENIX [6], and to the Therminator (triangles pointing upwards) and HKM (triangles pointing downwards) models. Right: Transverse mass dependence of Gaussian radii and the λ for the 30% most central Au+Au collisions (solid stars). PHENIX data are also plotted (dots). Squares are HKM, solid curves are Buda-Lund model calculations. The 20% most central data are also shown for comparison (open stars).

The right panel of Figure 1 shows the dependence of the Gaussian radii in the longitudinally co-moving system (LCMS) ($R_{out}=R_x/\gamma$, $R_{side}=R_y$ and $R_{long}=R_z$; γ is the Lorentz boost in the outward direction from the LCMS to the PCMS frame) as a function of the transverse mass $m_T = (m^2 + k_T^2)^{1/2}$. PHENIX kaon data [10] are also shown. The perfect fluid hydrodynamics calculations from the Buda-Lund model [11] and the hydrokinetic model (HKM) [18] with Glauber initial conditions are plotted for comparison purposes. While pions are well described by the Buda Lund model in the whole interval shown [11], low- k_T kaons in the *long* direction seem to favour HKM over the Buda-Lund model, suggesting that that contrary to lower SPS energies [9] the m_T -scaling in the *long* direction is broken at RHIC.

4. Summary

We have presented the first model-independent extraction of the 3D kaon source by the STAR Collaboration [12], at mid-rapidity in $\sqrt{s_{NN}}=200$ GeV central Au+Au collisions, using low- k_T kaon pair correlations and the Cartesian surface-spherical harmonic decomposition technique. No significant non-Gaussian tail has been observed. Comparison with the Thermanator model calculations indicates that, although the transverse extent of the source is similar to pions, the shape and the size in the longitudinal direction is very different. This can be attributed to resonance decays, but also indicates that kaons and pions may be subject to different freeze-out dynamics. Although the Gaussian radii follow m_T -scaling in the outward and sideward directions, the scaling appears to be broken in the longitudinal direction. Thus the hydro-kinetic predictions [18] are favoured over pure hydrodynamical model calculations.

Acknowledgments

This research was supported by the grant LA09013 of the Ministry of Education of the Czech Republic.

References

- [1] R. Lednicky, Nucl. Phys. **A774**, 189 (2006).
- [2] R. Lednicky, J. Phys. **G35**, 125109 (2008); Phys. Part. Nucl. **40**, 307 (2009).
- [3] M. A. Lisa *et al.*, Ann. Rev. Nucl. Part. Sci. **55**, 357 (2005).
- [4] D. A. Brown and P. Danielewicz, Phys. Lett. B **398**, 252 (1997).
- [5] P. Danielewicz and S. Pratt, Phys. Lett. **B 618**, 60 (2005).
- [6] S. Afanasiev *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **100**, 232301 (2008).
- [7] P. Danielewicz and S. Pratt, Phys. Rev. C **75** (2007) 034907.
- [8] M. Csanád, T. Csörgő and M. Nagy, Braz. J. Phys. **37**, 1002 (2007).
- [9] S. V. Afanasiev *et al.* (NA49 Collaboration), Phys. Lett. B **557**, 157 (2003).
- [10] S. Afanasiev *et al.* (PHENIX Collaboration) Phys. Rev. Lett. **103**, 142301 (2009).
- [11] M. Csanád and T. Csörgő, Acta Phys. Polon. Supp. **1**, 521 (2008).
- [12] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. C **88**, 034906 (2013).
- [13] B. I. Abelev *et al.* (STAR Collaboration), Phys. Rev. C **74**, 054902 (2006).
- [14] P. Chung (STAR Collaboration), Phys. Part. Nucl. Lett. **8**, 1019 (2011).
- [15] A. Kisiel *et al.*, Comput. Phys. Commun. **174**, 669 (2006).
- [16] A. Kisiel *et al.*, Phys. Rev. C **73**, 064902 (2006).
- [17] V. M. Shapoval, Y. M. Sinyukov and I. A. Karpenko, arXiv:1308.6272.
- [18] I. A. Karpenko and Y. M. Sinyukov, Phys. Rev. C **81**, 054903 (2010).

Title of article capitalized only at beginning of line

AUTHOR 1, AUTHOR 2
Institution address 1

AUTHOR 3
Institution address 2

This is the abtract of my presentation at ISMD 2013.

1. Introduction

Section is here. For sample Figures and Table entries see the LaTeX.
Volume will be printed in 6-1/8”x9-1/8” format.

Table 1. Caption for the Table here.

Cat1	Cat2	Cat3	Cat4
val1	val2	val3	val4

Acknowledgments

This work is supported by everyone.

References

[1] John Smith, Journ. Impossible Results **999**, 1234 (2025).

Fig. 1. Figure caption here.