

LINKING STAROBINSKY-TYPE INFLATION IN NO-SCALE SUPERGRAVITY TO MSSM

C. PALLIS

*Departament de Física Teòrica and IFIC,
Universitat de València-CSIC,
E-46100 Burjassot, SPAIN*

*Department of Physics, University of Cyprus,
P.O. Box 20537, Nicosia 1678, CYPRUS*

ABSTRACT

A novel realization of the Starobinsky inflationary model within a moderate extension of the Minimal Supersymmetric Standard Model (MSSM) is presented. The proposed superpotential is uniquely determined by applying a continuous R and a \mathbb{Z}_2 discrete symmetry, whereas the Kähler potential is associated with a no-scale-type $SU(54, 1)/SU(54) \times U(1)_R \times \mathbb{Z}_2$ Kähler manifold. The inflaton is identified with a Higgs-like modulus whose the vacuum expectation value controls the gravitational strength. Thanks to a strong enough coupling (with a parameter c_T involved) between the inflaton and the Ricci scalar curvature, inflation can be attained even for subplanckian values of the inflaton with $c_T \geq 76$ and the corresponding effective theory being valid up to the Planck scale. The inflationary observables turn out to be in agreement with the current data and the inflaton mass is predicted to be $3 \cdot 10^{13}$ GeV. At the cost of a relatively small superpotential coupling constant, the model offers also a resolution of the μ problem of MSSM. Supplementing MSSM by three right-handed neutrinos we show that spontaneously arising couplings between the inflaton and the particle content of MSSM not only ensure a sufficiently low reheating temperature but also support a scenario of non-thermal leptogenesis consistently with the neutrino oscillation parameters for gravitino heavier than about 10^4 GeV.

Keywords: Cosmology, Supersymmetric models, Supergravity, Modified Gravity;

PACS codes: 98.80.Cq, 11.30.Qc, 12.60.Jv, 04.65.+e, 04.50.Kd

Published in *J. Cosmol. Astropart. Phys.* **04**, 024 (2014); *J. Cosmol. Astropart. Phys.* **07**, 01E (2017)

CONTENTS

1	INTRODUCTION	1
2	MODEL DESCRIPTION	2
2.1	THE SUPERPOTENTIAL	3
2.2	THE KÄHLER POTENTIAL	5
3	THE INFLATIONARY SCENARIO	6
3.1	THE INFLATIONARY POTENTIAL	6
3.2	THE INFLATIONARY OBSERVABLES – REQUIREMENTS	9
3.3	THE EFFECTIVE CUT-OFF SCALE	10
3.4	NUMERICAL RESULTS	12
4	THE R SYMMETRY AND THE μ PROBLEM OF MSSM	14
4.1	GENERATION OF THE μ TERM OF MSSM	14
4.2	CONNECTION WITH THE MSSM PHENOMENOLOGY	15
5	NON-THERMAL LEPTOGENESIS AND NEUTRINO MASSES	16
5.1	THE INFLATON DECAY	16
5.2	LEPTON-NUMBER AND GRAVITINO ABUNDANCES	18
5.3	LEPTON-NUMBER ASYMMETRY AND NEUTRINO MASSES	18
5.4	POST-INFLATIONARY REQUIREMENTS	19
5.5	NUMERICAL RESULTS	20
6	CONCLUSIONS	23
	REFERENCES	24

1 INTRODUCTION

After the announcement of the recent PLANCK results [1, 2], inflation based on the potential of the Starobinsky model [3] has gained a lot of momentum [4–11] since it predicts [3, 12] a (scalar) spectral index very close to the one favored by the fitting of the observations by the standard power-law cosmological model with *cold dark matter* (CDM) and a cosmological constant (Λ CDM). In particular, it has been shown that Starobinsky-type inflation can be realized within extensions of the *Standard Model* (SM) [13] or *Minimal SUSY SM* (MSSM) [14]. However, the realization of this type of inflation within *Supergravity* (SUGRA) is not unique. Different super- and Kähler potentials are proposed [5–7] which result to the same scalar potential. Prominent, however, is the idea [4, 5] of implementing this type of inflation using a Kähler potential, K , corresponding to a $SU(N, 1)/SU(N) \times U(1)$ Kähler manifold inspired by the no-scale models [15, 16]. Such a symmetry fixes beautifully the form of K up to an holomorphic function f_K which exclusively depends on a modulus-like field and plays the role of a varying gravitational coupling. The stabilization of the non-inflaton accompanying field can to be conveniently arranged by higher order terms in K . In this context, a variety of models are proposed in which inflaton can be identified with either a matter-like [4, 5, 14] or a modulus-like [5, 6] inflaton. The former option seems to offer a more suitable framework [14] for connecting the inflationary physics with a low-energy theory, such as the MSSM endowed with right handed neutrinos, N_i^c , since the non-inflaton modulus is involved in the no-scale mechanism of *soft SUSY breaking* (SSB). On the other hand, the inflationary superpotential, W_{MI} , is arbitrarily chosen and not protected by any symmetry. Given that, the inflaton takes transplanckian values during inflation, higher order corrections – e.g., by non-renormalizable terms in W_{MI} – with not carefully tuned coefficients may easily invalidate or strongly affect [8, 17] the predictions of an otherwise successful inflationary scenario.

It would be interesting, therefore, to investigate if the shortcoming above can be avoided in the presence of a strong enough coupling of the inflaton to gravity [18, 19], as done [20–25] in the models of *non-minimal Inflation* (nMI). This idea can be implemented keeping the no-scale structure of K , since the involved f_K can be an analytic function, selected conveniently. In view of the fact that f_K depends only on a modulus-like field, we here focus on this kind of inflaton – contrary to Ref. [14]. As a consequence, the direct connection of the inflationary model with the mechanism of the SSB is lost. Note, in passing, that despite their attractive features, the no-scale models [14] of SSB enface difficulties – e.g., viable SUSY spectra are obtained only when the boundary conditions for the SSB terms are imposed beyond the *Grand Unified Theory* (GUT) scale and so the low energy observables depend on the specific GUT.

Focusing on a modulus-like inflaton, the link to MSSM can be established through the adopted W_{MI} . Its form in our work is fixed by imposing a continuous R symmetry, which reduces to the well-known R -parity of MSSM, and a \mathbb{Z}_2 discrete symmetry. As a consequence, W_{MI} resembles the one used in the widely employed models [26, 27] of standard *F-term Hybrid Inflation* (FHI) – with singlet waterfall field though. As a bonus, a dynamical generation of the reduced Planck scale arises in *Jordan Frame* (JF) through the *vacuum expectation value* (v.e.v) of the inflaton. Therefore the inflaton acquires a higgs-character as in the theories of induced gravity [28, 29]. To produce an inflationary plateau with the selected W_{MI} , f_K is to be taken quadratic, in accordance with the adopted symmetries. This is to be contrasted with the so-called modified Cecotti model [5–8, 30] where the inflaton appears linearly in the super- and Kähler potentials. The inclusion of two extra parameters compared to the original model – cf. [5, 6, 8] – allows us to attain inflationary solutions for subplanckian values of the inflaton with the successful inflationary predictions of the model being remained intact. As a bonus, the *ultraviolet* (UV) cut-off scale [9, 31, 32] of the theory can be identified with the Planck scale and so, concerns regarding the naturalness of the model can be safely eluded.

Our inflationary model – let name it for short *no-scale modular inflation* (nSMI) – has ramifications to other fundamental open problems of the MSSM and post-inflationary cosmological evolution. As a consequence of the adopted $U(1)_R$ symmetry, the generation [27, 33] of the mixing term between the two electroweak Higgses is explained via the v.e.v of the non-inflaton accompanying field, provided that a coupling constant in W_{MI} is rather suppressed. Finally, the observed [34] *baryon asymmetry of the universe* (BAU) can be explained via spontaneous [35, 36] *non-thermal leptogenesis* (nTL) [37] consistently with the \tilde{G} constraint [38–40], the data [41, 42] on the neutrino oscillation parameters as long as the masses of the gravitino (\tilde{G}) lie in the multi-TeV region – as dictated in many versions [43–45] of MSSM after the recent LHC [46, 47] results on the Higgs boson mass.

The basic ingredients – particle content and structure of the super- and Kähler potentials – of our model are presented in Sec. 2. In Sec. 3 we describe the inflationary potential, derive the inflationary observables and confront them with observations. Sec. 4 is devoted to the resolution of the μ problem of MSSM. In Sec. 5 we outline the scenario of nTL, exhibit the relevant imposed constraints and describe our predictions for neutrino masses. Our conclusions are summarized in Sec. 6. Throughout the text, the subscript of type $_{,\chi}$ denotes derivation *with respect to* (w.r.t) the field χ (e.g., $_{,\chi\chi} = \partial^2/\partial\chi^2$) and charge conjugation is denoted by a star.

2 MODEL DESCRIPTION

We focus on a moderated extension of MSSM with three N_i^c 's augmented by two superfields, a matter-like S and a modulus-like T , which are singlets under $G_{\text{SM}} = SU(3)_c \times SU(2)_L \times U(1)_Y$. Besides the local symmetry of MSSM, G_{SM} , the model possesses also the baryon number symmetry $U(1)_B$, a nonanomalous R symmetry $U(1)_R$ and a discrete \mathbb{Z}_2 . Note that global continuous symmetries can effectively arise [48] in many compactified string theories. The charge assignments under the

	S	T	H_u	H_d	L_i	N_i^c	e_i^c	Q_i	u_i^c	d_i^c
$U(1)_B$	0	0	0	0	0	0	0	1/3	-1/3	-1/3
$U(1)_R$	2	0	0	0	1	1	1	1	1	1
\mathbb{Z}_2	0	1	0	0	0	0	0	0	0	0

Table 1: The global charges of the superfields of our model.

global symmetries of the various matter and Higgs superfields are listed in Table 1. We below present the structure of the superpotential (Sec. 2.1) and the Kähler potential (Sec. 2.2) of our model.

2.1 THE SUPERPOTENTIAL

The superpotential of our model naturally splits into two parts:

$$W = W_{\text{MSSM}} + W_{\text{MI}}, \quad (2.1a)$$

where W_{MSSM} is the part of W which contains the usual terms – except for the μ term – of MSSM, supplemented by Yukawa interactions among the left-handed leptons and N_i^c

$$W_{\text{MSSM}} = h_{ijE} e_i^c L_j H_d + h_{ijD} d_i^c Q_j H_d + h_{ijU} u_i^c Q_j H_u + h_{ijN} N_i^c L_j H_u. \quad (2.1b)$$

Here the i th generation $SU(2)_L$ doublet left-handed quark and lepton superfields are denoted by Q_i and L_i respectively, whereas the $SU(2)_L$ singlet antiquark [antilepton] superfields by u_i^c and d_i^c [e_i^c and N_i^c] respectively. The electroweak Higgs superfields which couple to the up [down] quark superfields are denoted by H_u [H_d].

On the other hand, W_{MI} is the part of W which is relevant for nSMI, the generation of the μ term of MSSM and the Majorana masses for N_i^c 's. It takes the form

$$W_{\text{MI}} = \lambda S (T^2 - M^2/2) + \lambda_\mu S H_u H_d + \frac{1}{2} M_{iN^c} N_i^c{}^2 + \lambda_{ijN^c} T^2 N_i^c N_j^c / 2m_P, \quad (2.1c)$$

where $m_P = 2.44 \cdot 10^{18}$ GeV is the reduced Planck mass. The imposed $U(1)_R$ symmetry ensures the linearity of W_{MI} w.r.t S . This fact allows us to isolate easily via its derivative the contribution of the inflaton T into the F-term SUGRA scalar potential, placing S at the origin. The imposed \mathbb{Z}_2 prohibits the existence of the term ST which, although does not drastically modifies our proposal, it complicates the determination of SUSY vacuum and the inflationary dynamics. On the other hand, the imposed symmetries do not forbid non-renormalizable terms of the form T^{2n+2} where $n \geq 1$ is an integer. For this reason we are obliged to restrict ourselves to subplanckian values of T .

The second term in the *right-hand side* (r.h.s) of Eq. (2.1c) provides the μ term of MSSM along the lines of Ref. [27, 33] – see Sec. 4. The third term is the Majorana mass term for the N_i^c 's and we assume that it overshadows (for sufficiently low λ_{ijN^c} 's) the last non-renormalizable term which is neglected henceforth. Here we work in the so-called *right-handed neutrino basis*, where M_{iN^c} is diagonal, real and positive. These masses together with the Dirac neutrino masses in Eq. (2.1b) lead to the light neutrino masses via the seesaw mechanism. The same term is important for the decay [35, 36] of the inflaton after the end of nSMI to \tilde{N}_i^c , whose subsequent decay can activate nTL. As a result of the imposed \mathbb{Z}_2 , a term of the form $T N_i^c{}^2$ is prohibited and so the decay of T into N_i^c is processed by suppressed SUGRA-induced interactions [35], guaranteing thereby a sufficiently low reheate temperature compatible with the \tilde{G} constraint and successful nTL – see Sec. 5.1.

Since the no-scale SUGRA adopted here leads to the non-renormalizable F-term (scalar) potential in Eq. (3.1), we expect that it yields an effective SUSY theory which depends not only on the superpotential, W_{MI} in Eq. (2.1c), but also on the Kähler potential, K in Eq. (2.6a) – see Sec. 2.2. To trace out this behavior we apply the generic formula for the SUSY F-term potential [49]:

$$V_{\text{SUSY}} = K^{\alpha\bar{\beta}} W_{\text{MI}\alpha} W_{\text{MI}\bar{\beta}}^*, \quad (2.2)$$

which is obtained from the SUGRA potential in Eq. (3.1) – see Sec. 3.1 below – if we perform an expansion in powers $1/m_{\text{P}}$ and take the limit $m_{\text{P}} \rightarrow \infty$. The Kähler potential, K , employed here can not be expanded in powers of $1/m_{\text{P}}$, since unity is not included in the argument of the logarithm – in contrast to the K 's used in Ref. [50,51]. In Eq. (2.2) $K^{\alpha\bar{\beta}}$ is the inverse of the Kähler metric $K_{\alpha\bar{\beta}}$ with $z^\alpha = T, S, H_u, H_d$ and \tilde{N}_i^c where the complex scalar components of the superfields T, S, H_u and H_d are denoted by the same symbol whereas this of N_i^c by \tilde{N}_i^c . We find that $K_{\alpha\bar{\beta}}$ reads

$$(K_{\alpha\bar{\beta}}) = \left(\frac{3}{\Omega}\right)^2 \begin{pmatrix} \frac{12c_T^2|T|^2}{m_{\text{P}}^2} & -\frac{2c_T ST}{m_{\text{P}}^2} & -\frac{2c_T H_u T}{m_{\text{P}}^2} & -\frac{2c_T H_d T}{m_{\text{P}}^2} & -\frac{2c_T \tilde{N}_i^c T}{m_{\text{P}}^2} \\ -\frac{2c_T S^* T^*}{m_{\text{P}}^2} & -\frac{\Omega}{3} & \frac{S^* H_u}{3m_{\text{P}}^2} & \frac{S^* H_d}{3m_{\text{P}}^2} & \frac{S^* \tilde{N}_i^c}{3m_{\text{P}}^2} \\ -\frac{2c_T H_u^* T^*}{m_{\text{P}}^2} & \frac{H_u^* S}{3m_{\text{P}}^2} & -\frac{\Omega}{3} & \frac{H_u^* H_d}{3m_{\text{P}}^2} & \frac{H_u^* \tilde{N}_i^c}{3m_{\text{P}}^2} \\ -\frac{2c_T H_d^* T^*}{m_{\text{P}}^2} & \frac{H_d^* S}{3m_{\text{P}}^2} & \frac{H_d^* H_u}{3m_{\text{P}}^2} & -\frac{\Omega}{3} & \frac{H_d^* \tilde{N}_i^c}{3m_{\text{P}}^2} \\ -\frac{2c_T \tilde{N}_i^{c*} T^*}{m_{\text{P}}^2} & \frac{\tilde{N}_i^{c*} S}{3m_{\text{P}}^2} & \frac{\tilde{N}_i^{c*} H_u}{3m_{\text{P}}^2} & \frac{c_T \tilde{N}_i^{c*} H_d}{3m_{\text{P}}^2} & -\frac{\Omega}{3} \end{pmatrix}, \quad (2.3)$$

where Ω is given in Eq. (2.7c) and we neglect the fourth order terms since we expect that these are not relevant for the low energy effective theory. The inverse of the matrix above is

$$(K^{\alpha\bar{\beta}}) = -\frac{\Omega}{3} \begin{pmatrix} (T^2 + T^{*2})/12c_T|T|^2 & S/6c_T T^* & H_u/6c_T T^* & H_d/6c_T T^* & \tilde{N}_i^c/6c_T T^* \\ S^*/6c_T T & 1 & 0 & 0 & 0 \\ H_u^*/6c_T T & 0 & 1 & 0 & 0 \\ H_d^*/6c_T T & 0 & 0 & 1 & 0 \\ \tilde{N}_i^{c*}/6c_T T & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (2.4)$$

Substituting this in Eq. (2.2), we end up with the following expression

$$\begin{aligned} V_{\text{SUSY}} = & -\frac{\Omega}{3} \left(\lambda^2 |T^2 + \lambda_\mu H_u H_d / \lambda - M^2/2|^2 + \lambda_\mu^2 (|H_u|^2 + |H_d|^2) |S|^2 + M_{iN^c}^2 |\tilde{N}_i^c|^2 \right. \\ & + \frac{2\lambda^2}{3c_T} |S|^2 (T^2 + T^{*2} - M^2/2) + \frac{\lambda\lambda_\mu}{c_T} (H_u H_d + H_u^* H_d^*) |S|^2 \\ & \left. + \frac{\lambda}{3c_T} M_{iN^c} (S^* \tilde{N}_i^{c2} + S \tilde{N}_i^{c*2}) \right). \end{aligned} \quad (2.5a)$$

The three first terms in the r.h.s of the expression above come from the terms $K^{\alpha\bar{\alpha}} W_{\text{MI}\alpha} W_{\text{MI}\bar{\alpha}}^*$ of Eq. (2.2) for $z^\alpha = S, H_u, H_d, \tilde{N}_i^c$. The fourth one comes from the terms

$$K^{\alpha\bar{\alpha}} W_{\text{MI}\alpha} W_{\text{MI}\bar{\alpha}}^* + K^{\alpha\bar{\beta}} W_{\text{MI}\alpha} W_{\text{MI}\bar{\beta}}^* + K^{\beta\bar{\alpha}} W_{\text{MI}\beta} W_{\text{MI}\bar{\alpha}}^*$$

for $z^\alpha = T$ and $z^\beta = S$; the residual terms arise from terms of the form $K^{\alpha\bar{\beta}} W_{\text{MI}\alpha} W_{\text{MI}\bar{\beta}}^* + K^{\beta\bar{\alpha}} W_{\text{MI}\beta} W_{\text{MI}\bar{\alpha}}^*$; for $z^\alpha = T$ and $z^\beta = H_u, H_d$ the fifth one and $z^\beta = \tilde{N}_i^c$ the last one.

From the potential in Eq. (2.5a), we find that the SUSY vacuum lies at

$$\langle H_u \rangle = \langle H_d \rangle = \langle \tilde{N}_i^c \rangle = 0, \quad \langle S \rangle \simeq 0 \quad \text{and} \quad \sqrt{2}|\langle T \rangle| = M. \quad (2.5b)$$

Contrary to the Cecotti model [5, 6, 30] our modulus T can take values $M \leq m_P$ at the SUSY vacuum. Also, $\langle T \rangle$ breaks spontaneously the imposed \mathbb{Z}_2 and so, it can comfortably decay via SUGRA-inspired decay channels – see Sec. 5.1 – reheating the universe and rendering [36] spontaneous nTL possible. No domain walls are produced due to the spontaneous breaking of \mathbb{Z}_2 at the SUSY vacuum, since this is broken already during nSMI.

With the addition of SSB terms, as required in a realistic model, the position of the vacuum shifts [27, 33] to non-zero $\langle S \rangle$ and an effective μ term is generated from the second term in the r.h.s of Eq. (2.1c) – see Sec. 4. Let us emphasize that SSB effects explicitly break $U(1)_R$ to the \mathbb{Z}_2^R matter parity, under which all the matter (quark and lepton) superfields change sign. Combining \mathbb{Z}_2^R with the \mathbb{Z}_2^f fermion parity, under which all fermions change sign, yields the well-known R -parity. Recall that this residual symmetry prevents the rapid proton decay, guarantees the stability of the *lightest SUSY particle* (LSP) and therefore it provides a well-motivated CDM candidate. Needless to say, finally, that such a connection of the Starobinsky-type inflation with this vital for MSSM R -symmetry can not be established within the modified Cecotti model [6, 7, 30], since no symmetry can prohibit a quadratic term for the modulus-like field in conjunction with the tadpole term in W_{MI} .

2.2 THE KÄHLER POTENTIAL

According to the general discussion of Ref. [15], the Kähler manifold which corresponds to a Kähler potential of the form

$$K = -3m_P^2 \ln \left(f_K(T) + f_K^*(T^*) - \frac{\Phi_A \Phi^{*\bar{A}}}{3m_P^2} + k_{S\Phi^A} \frac{|S|^2 |\Phi_A|^2}{3m_P^4} + \dots \right), \quad (2.6a)$$

with f_K being an holomorphic function of T , exhibits a $SU(N, 1)/SU(N) \times U(1)_R \times \mathbb{Z}_2$ global symmetry. Here $N - 1 = 53$ is the number of scalar components of S , N_i^c and the MSSM superfields which are collectively denoted as

$$\Phi^A = \tilde{e}_i^c, \tilde{u}_i^c, \tilde{d}_i^c, \tilde{N}_i^c, \tilde{L}_i, \tilde{Q}_i, H_u, H_d \text{ and } S. \quad (2.6b)$$

Note that summation over the repeated (small or capital) Greek indices is implied. The third term in the r.h.s of Eq. (2.6a) – with coefficients $k_{S\Phi^A}$ being taken, for simplicity, real – is included since it has an impact on the scalar mass spectrum along the inflationary track – see Sec. 3.1. In particular, the term with coefficient $k_{SS} = k_S \simeq 1$ assists us to avoid the tachyonic instabilities encountered in similar models [5–7, 18, 19] – see Sec. 3.1. The ellipsis represents higher order terms which are irrelevant for the inflationary dynamics since they do not mix the inflaton T with the matter fields. This is, in practice, a great simplification compared to the models of nMI – cf. Ref. [25]. Contrary to other realizations of the Starobinsky model – cf. Ref. [5–7] –, we choose f_K to be quadratic and not linear with respect to T , i.e.,

$$f_K(T) = c_T T^2 / m_P^2 \quad (2.6c)$$

in accordance with the imposed \mathbb{Z}_2 symmetry which forbids a linear term – the coefficient c_T is taken real too. As in the case of Eq. (2.1c), non-renormalizable terms of the form T^{2n+2} , with integer $n \geq 1$, are allowed but we can safely ignore them restricting ourselves to $T \leq m_P$.

The interpretation of the adopted K in Eq. (2.6a) can be given in the “physical” frame by writing the JF action for the scalar fields $\Phi^\alpha = \Phi^A, T$. To extract it, we start with the corresponding EF action

within SUGRA [18, 21, 25] which can be written as

$$S = \int d^4x \sqrt{-\hat{\mathbf{g}}} \left(-\frac{1}{2} m_{\text{P}}^2 \hat{\mathcal{R}} + K_{\alpha\bar{\beta}} \dot{\Phi}^\alpha \dot{\Phi}^{*\bar{\beta}} - \hat{V}_{\text{MI0}} + \dots \right), \quad (2.7a)$$

where $K_{\alpha\bar{\beta}} = \hat{K}_{\Phi^\alpha \Phi^{*\bar{\beta}}}$ with $K^{\bar{\beta}\alpha} K_{\alpha\bar{\gamma}} = \delta_{\bar{\gamma}}^{\bar{\beta}}$, $\hat{\mathbf{g}}$ is the determinant of the EF metric $\hat{g}_{\mu\nu}$, $\hat{\mathcal{R}}$ is the EF Ricci scalar curvature, \hat{V}_{MI0} is defined in Sec. 3.1, the dot denotes derivation w.r.t the JF cosmic time and the ellipsis represents terms irrelevant for our analysis. Performing then a suitable conformal transformation, along the lines of Ref. [21, 25] we end up with the following action in the JF

$$S = \int d^4x \sqrt{-\mathbf{g}} \left(-\frac{m_{\text{P}}^2}{2} \left(-\frac{\Omega}{3} \right) \mathcal{R} + m_{\text{P}}^2 \Omega_{\alpha\bar{\beta}} \dot{\Phi}^\alpha \dot{\Phi}^{*\bar{\beta}} - V_{\text{SUSY}} + \dots \right), \quad (2.7b)$$

where $g_{\mu\nu} = -(3/\Omega) \hat{g}_{\mu\nu}$ is the JF metric with determinant \mathbf{g} , \mathcal{R} is the JF Ricci scalar curvature, and we use the shorthand notation $\Omega_\alpha = \Omega_{,\Phi^\alpha}$ and $\Omega_{\bar{\alpha}} = \Omega_{,\Phi^{*\bar{\alpha}}}$. The corresponding frame function can be found from the relation

$$-\frac{\Omega}{3} = e^{-K/3m_{\text{P}}^2} = f_K(T) + f_K^*(T^*) - \frac{\Phi_A \Phi^{*\bar{A}}}{3m_{\text{P}}^2} + k_{S\Phi^A} \frac{|S|^2 |\Phi_A|^2}{3m_{\text{P}}^4} + \dots \quad (2.7c)$$

The last result reveals that T has no kinetic term, since $\Omega_{,TT^*} = 0$. This is a crucial difference between the Starobinsky-type models and those [25] of nMI, with interest consequences [9] to the derivation of the ultraviolet cutoff scale of the theory – see Sec. 3.2. Furthermore, given that $\langle \Phi^A \rangle \simeq 0$, recovering the conventional Einstein gravity at the SUSY vacuum, Eq. (2.5b), dictates

$$f_K(\langle T \rangle) + f_K^*(\langle T^* \rangle) = 1 \Rightarrow M = m_{\text{P}} / \sqrt{c_T}. \quad (2.8)$$

Given that the analysis of inflation in both frames yields equivalent results [12, 52], we below – see Sec. 3.1 and 3.2 – carry out the derivation of the inflationary observables exclusively in the EF.

3 THE INFLATIONARY SCENARIO

In this section we outline the salient features of our inflationary scenario (Sec. 3.1) and then, we present its predictions in Sec. 3.4, calculating a number of observable quantities introduced in Sec. 3.2. We also provide a detailed analysis of the UV behavior of the model in Sec. 3.3.

3.1 THE INFLATIONARY POTENTIAL

The EF F-term (tree level) SUGRA scalar potential, \hat{V}_{MI0} , of our model – see Eq. (2.7a) – is obtained from W_{MI} in Eq. (2.1c) and K in Eq. (2.6a) by applying the standard formula:

$$\hat{V}_{\text{MI0}} = e^{K/m_{\text{P}}^2} \left(K^{\alpha\bar{\beta}} F_\alpha F_{\bar{\beta}}^* - 3 \frac{|W_{\text{MI}}|^2}{m_{\text{P}}^2} \right), \quad (3.1)$$

where $F_\alpha = W_{\text{MI},\Phi^\alpha} + K_{\Phi^\alpha} W_{\text{MI}}/m_{\text{P}}^2$. Setting the fields $\Phi^\alpha = S, \tilde{N}_i^c, H_u$ and H_d at the origin the only surviving term of \hat{V}_{MI0} is

$$\hat{V}_{\text{MI0}} = e^{K/m_{\text{P}}^2} K^{SS^*} W_{\text{MI},S} W_{\text{MI},S}^* = \frac{\lambda^2 |2T^2 - M^2|^2}{4(f_K + f_K^*)^2}. \quad (3.2)$$

It is obvious from the result above that a form of f_K as the one proposed in Eq. (2.6c) can flatten \hat{V}_{MI0} sufficiently so that it can drive nSMI. Employing the dimensionless variables

$$x_\phi = \phi/m_{\text{P}}, \quad f_T = 1 - c_T x_\phi^2 \quad \text{and} \quad x_M = M/m_{\text{P}} \quad \text{with} \quad \phi = |T|/\sqrt{2} \quad (3.3)$$

and setting $\arg T = 0$, \widehat{V}_{MIO} and the corresponding Hubble parameter \widehat{H}_{MI} read

$$\widehat{V}_{\text{MIO}} = \frac{\lambda^2 m_{\text{P}}^4 (x_\phi^2 - x_M^2)^2}{4c_T^2 x_\phi^4} = \frac{\lambda^2 m_{\text{P}}^4 f_T^2}{4c_T^4 x_\phi^4} \quad \text{and} \quad \widehat{H}_{\text{MI}} = \frac{\widehat{V}_{\text{MIO}}^{1/2}}{\sqrt{3} m_{\text{P}}} \simeq \frac{\lambda m_{\text{P}}}{2\sqrt{3} c_T^2}, \quad (3.4)$$

where we put $x_M = 1/\sqrt{c_T}$ – by virtue of Eq. (2.8) – in the final expressions.

Expanding T and Φ^α in real and imaginary parts as follows

$$T = \frac{\phi}{\sqrt{2}} e^{i\theta/m_{\text{P}}} \quad \text{and} \quad X^\alpha = \frac{x^\alpha + i\bar{x}^\alpha}{\sqrt{2}} \quad \text{with} \quad X^\alpha = S, H_u, H_d, \tilde{N}_i^c \quad (3.5)$$

we can check the stability of the inflationary direction

$$\theta = x^\alpha = \bar{x}^\alpha = 0 \quad \text{where} \quad x^\alpha = s, h_u, h_d, \tilde{\nu}_i^c, \quad (3.6)$$

w.r.t the fluctuations of the various fields. In particular, we examine the validity of the extremum and minimum conditions, i.e.,

$$\left. \frac{\partial \widehat{V}_{\text{MIO}}}{\partial \widehat{\chi}^\alpha} \right|_{\text{Eq. (3.6)}} = 0 \quad \text{and} \quad \widehat{m}_{\chi^\alpha}^2 > 0 \quad \text{with} \quad \chi^\alpha = \theta, x^\alpha, \bar{x}^\alpha. \quad (3.7a)$$

Here $\widehat{m}_{\chi^\alpha}^2$ are the eigenvalues of the mass matrix with elements

$$\widehat{M}_{\alpha\beta}^2 = \left. \frac{\partial^2 \widehat{V}_{\text{MIO}}}{\partial \widehat{\chi}^\alpha \partial \widehat{\chi}^\beta} \right|_{\text{Eq. (3.6)}} \quad \text{with} \quad \chi^\alpha = \theta, x^\alpha, \bar{x}^\alpha \quad (3.7b)$$

and hat denotes the EF canonically normalized fields. Taking into account that along the configuration of Eq. (3.6) $K_{\alpha\bar{\beta}}$ defined below Eq. (2.7a) takes the form

$$(K_{\alpha\bar{\beta}}) = \text{diag} \left(6/x_\phi^2, \underbrace{1/c_T x_\phi^2, \dots, 1/c_T x_\phi^2}_{8 \text{ elements}} \right) \quad (3.8)$$

– here we take into account that H_u and H_d are $SU(2)_L$ doublets –, the kinetic terms of the various scalars in Eq. (2.7a) can be brought into the following form

$$K_{\alpha\bar{\beta}} \dot{\Phi}^\alpha \dot{\Phi}^{*\bar{\beta}} = \frac{1}{2} \left(\dot{\phi}^2 + \dot{\theta}^2 \right) + \frac{1}{2} \left(\dot{\hat{x}}_\alpha \dot{\hat{x}}^\alpha + \dot{\hat{x}}_\alpha \dot{\hat{x}}^\alpha \right), \quad (3.9a)$$

where the hatted fields are defined as follows

$$d\widehat{\phi}/d\phi = J = \sqrt{6}/x_\phi, \quad \widehat{\theta} = \sqrt{6}\theta, \quad \widehat{x}^\alpha = x^\alpha/\sqrt{c_T}x_\phi \quad \text{and} \quad \widehat{\bar{x}}^\alpha = \bar{x}^\alpha/\sqrt{c_T}x_\phi. \quad (3.9b)$$

Upon diagonalization of the relevant sub-matrices of $\widehat{M}_{\alpha\beta}^2$, Eq. (3.7b), we construct the scalar mass spectrum of the theory along the direction in Eq. (3.6). Our results are summarized in Table 2, assuming $k_{SH_u} \simeq k_{SH_d} = k_{SH}$ in order to avoid very lengthy formulas for the masses of \widehat{h}_\pm and $\widehat{\bar{h}}_\pm$. The various unspecified there eigenvalues are defined as follows:

$$\widehat{h}_\pm = (\widehat{h}_u \pm \widehat{h}_d)/\sqrt{2}, \quad \widehat{\bar{h}}_\pm = (\widehat{\bar{h}}_u \pm \widehat{\bar{h}}_d)/\sqrt{2} \quad \text{and} \quad \widehat{\psi}_\pm = (\widehat{\psi}_T \pm \widehat{\psi}_S)/\sqrt{2}, \quad (3.10a)$$

where the spinors ψ_T, ψ_S and N_i^c associated with the superfields S, T and N_i^c are related to the normalized ones in Table 2 as follows:

$$\widehat{\psi}_S = \sqrt{6}\psi_S/x_\phi, \quad \widehat{\psi}_T = \psi_T/\sqrt{c_T}x_\phi \quad \text{and} \quad \widehat{N}_i^c = N_i^c/\sqrt{c_T}x_\phi. \quad (3.10b)$$

FIELDS	EINGESTATES	MASSES SQUARED
1 real scalar	$\hat{\theta}$	$\hat{m}_\theta^2 = \lambda^2 m_P^2 (f_T + 2c_T^2 x_\phi^2) / 3c_T^4 x_\phi^4 \simeq 4\hat{H}_{\text{MI}}^2$
2 real scalars	$\hat{s}, \hat{\bar{s}}$	$\hat{m}_s^2 = \lambda^2 m_P^2 (1 + c_T x_\phi^2 (2 - c_T x_\phi^2 + 6k_S f_T^2)) / 6c_T^4 x_\phi^4$
4 real scalars	\hat{h}_+, \hat{h}_+	$\hat{m}_{h+}^2 = \lambda m_P^2 f_T (\lambda f_T f_{SH} + 6\lambda_\mu c_T^2 x_\phi^2) / 12c_T^4 x_\phi^4$
4 real scalars	\hat{h}_-, \hat{h}_-	$\hat{m}_{h-}^2 = \lambda m_P^2 f_T (\lambda f_T f_{SH} - 6\lambda_\mu c_T^2 x_\phi^2) / 12c_T^4 x_\phi^4$
6 real scalars	$\hat{\nu}_i^c, \hat{\bar{\nu}}_i^c$	$\hat{m}_{i\nu^c}^2 = (\lambda^2 m_P^2 f_T^2 f_{S\tilde{N}_i^c} + 12M_{iN^c}^2 c_T^3 x_\phi^2) / 12c_T^4 x_\phi^4$
2 Weyl spinors	$\hat{\psi}_\pm$	$\hat{m}_{\psi\pm}^2 \simeq \lambda^2 m_P^2 / 3c_T^4 x_\phi^4$
3 Weyl spinors	\hat{N}_i^c	$\hat{m}_{iN^c}^2 = M_{iN^c}^2 / c_T x_\phi^2$

Table 2: The mass spectrum of our model along the inflationary trajectory of Eq. (3.6).

We also use the shorthand notation:

$$f_{SH} = 2 + 3k_{SH} c_T x_\phi^2 \quad \text{and} \quad f_{S\tilde{N}_i^c} = 2 + 3k_{S\tilde{N}_i^c} c_T x_\phi^2. \quad (3.11)$$

Note that, due to the large effective masses that the χ 's in Eq. (3.7b) acquire during nSMI, they enter a phase of oscillations about $\chi = 0$ with reducing amplitude. As a consequence – see Eq. (3.9b) –, $\dot{\hat{\chi}} \simeq \dot{\chi} / \sqrt{f_K}$ since the quantity $\dot{f}_K / 2f_K^{3/2} \chi$, involved in relating $\dot{\hat{\chi}}$ to $\dot{\chi}$, turns out to be negligibly small compared with $\dot{\chi} / \sqrt{f_K}$ – cf. Ref. [24]. Moreover, we have numerically verified that the various masses remain greater than \hat{H}_{MI} during the last 50 e-foldings of nSMI, and so any inflationary perturbations of the fields other than the inflaton are safely eliminated – see also Sec. 3.4.

From Table 2 it is evident that $k_S \gtrsim 1$ assists us to achieve $\hat{m}_s^2 > 0$ – in accordance with the results of Ref. [5–7]. On the other hand, given that $f_T \leq 0$, $\hat{m}_{h-}^2 > 0$ requires

$$\lambda f_T f_{SH} + 6\lambda_\mu c_T^2 x_\phi^2 < 0 \Rightarrow \lambda_\mu < -\frac{\lambda f_T f_{SH}}{6c_T^2 x_\phi^2} \simeq \frac{\lambda}{3c_T} + \frac{1}{2} \lambda k_{SH} x_\phi^2 \simeq 2 \cdot 10^{-5} - 10^{-6}, \quad (3.12)$$

as k_{SH} decreases from 3 to 0.5. Here we have made use of Eqs. (3.16a) and (3.20b) – see Sec. 3.2. We do not consider such a condition on λ_μ as unnatural, given that h_{11U} in Eq. (2.1b) is of the same order of magnitude too – cf. Ref. [53]. In Table 2 we also present the masses squared of chiral fermions along the trajectory of Eq. (3.6), which can be served for the calculation of the one-loop radiative corrections. Employing the well-known Coleman-Weinberg formula [54], we find that the one-loop corrected inflationary potential is

$$\begin{aligned} \hat{V}_{\text{MI}} = \hat{V}_{\text{MI}0} &+ \frac{1}{64\pi^2} \left(\hat{m}_\theta^4 \ln \frac{\hat{m}_\theta^2}{\Lambda^2} + 2\hat{m}_s^4 \ln \frac{\hat{m}_s^2}{\Lambda^2} + 4\hat{m}_{h+}^4 \ln \frac{\hat{m}_{h+}^2}{\Lambda^2} + 4\hat{m}_{h-}^4 \ln \frac{\hat{m}_{h-}^2}{\Lambda^2} \right. \\ &+ \left. 2 \sum_{i=1}^3 \left(\hat{m}_{i\nu^c}^4 \ln \frac{\hat{m}_{i\nu^c}^2}{\Lambda^2} - \hat{m}_{iN^c}^4 \ln \frac{\hat{m}_{iN^c}^2}{\Lambda^2} \right) - 4\hat{m}_{\psi\pm}^4 \ln \frac{\hat{m}_{\psi\pm}^2}{\Lambda^2} \right), \end{aligned} \quad (3.13)$$

where Λ is a *renormalization group* (RG) mass scale. As we numerically verify the one-loop corrections have no impact on our results. The absence of gauge interactions and of a direct renormalizable coupling between T and N_i^c assists to that direction – cf. Ref. [25, 55]. Based on \hat{V}_{MI} , we can proceed to the analysis of nSMI in the EF, employing the standard slow-roll approximation [56, 57]. It can be shown [28] that the results calculated this way are the same as if we had calculated them using the non-minimally coupled scalar field in the JF.

3.2 THE INFLATIONARY OBSERVABLES – REQUIREMENTS

A successful inflationary scenario has to be compatible with a number of observational requirements which are outlined in the following.

3.2.1 The number of e-foldings, \hat{N}_* , that the scale $k_* = 0.05/\text{Mpc}$ suffers during nSMI has to be adequate to resolve the horizon and flatness problems of the standard Big Bag cosmology. Assuming that nSMI is followed in turn by a decaying-particle, radiation and matter domination and employing standard methods [20], we can easily derive the required \hat{N}_* for our model, with the result:

$$\hat{N}_* \simeq 19.4 + 2 \ln \frac{\hat{V}_{\text{MI}}(\phi_*)^{1/4}}{1 \text{ GeV}} - \frac{4}{3} \ln \frac{\hat{V}_{\text{MI}}(\phi_f)^{1/4}}{1 \text{ GeV}} + \frac{1}{3} \ln \frac{T_{\text{rh}}}{1 \text{ GeV}} + \frac{1}{2} \ln \frac{f_K(\phi_*)}{f_K(\phi_f)^{1/3}}, \quad (3.14)$$

where $\phi_* [\hat{\phi}_*]$ is the value of $\phi [\hat{\phi}]$ when k_* crosses the inflationary horizon. Also $\phi_f [\hat{\phi}_f]$ is the value of $\phi [\hat{\phi}]$ at the end of nSMI determined, in the slow-roll approximation, by the condition:

$$\max\{\hat{\epsilon}(\phi_f), |\hat{\eta}(\phi_f)|\} = 1, \quad (3.15a)$$

where the slow-roll parameters read

$$\hat{\epsilon} = \frac{m_{\text{P}}^2}{2} \left(\frac{\hat{V}_{\text{MI},\hat{\phi}}}{\hat{V}_{\text{MI}}} \right)^2 = \frac{m_{\text{P}}^2}{2J^2} \left(\frac{\hat{V}_{\text{MI},\phi}}{\hat{V}_{\text{MI}}} \right)^2 \simeq \frac{4}{3f_T^2} \quad (3.15b)$$

and

$$\hat{\eta} = m_{\text{P}}^2 \frac{\hat{V}_{\text{MI},\hat{\phi}\hat{\phi}}}{\hat{V}_{\text{MI}}} = \frac{m_{\text{P}}^2}{J^2} \left(\frac{\hat{V}_{\text{MI},\phi\phi}}{\hat{V}_{\text{MI}}} - \frac{\hat{V}_{\text{MI},\phi}}{\hat{V}_{\text{MI}}} \frac{J_{,\phi}}{J} \right) \simeq \frac{4(1+f_T)}{3f_T^2}. \quad (3.15c)$$

The termination of nSMI is triggered by the violation of the ϵ criterion at a value of ϕ equal to ϕ_f , which is calculated to be

$$\hat{\epsilon}(\phi_f) = 1 \Rightarrow \phi_f = m_{\text{P}} \left((1 + 2/\sqrt{3})/c_T \right)^{1/2}, \quad (3.16a)$$

since the violation of the η criterion occurs at $\phi = \tilde{\phi}_f$ such that

$$\hat{\eta}(\tilde{\phi}_f) = 1 \Rightarrow \tilde{\phi}_f = m_{\text{P}} (5/3c_T)^{1/2} < \phi_f. \quad (3.16b)$$

On the other hand, \hat{N}_* can be calculated via the relation

$$\hat{N}_* = \frac{1}{m_{\text{P}}^2} \int_{\hat{\phi}_f}^{\hat{\phi}_*} d\hat{\phi} \frac{\hat{V}_{\text{MI}}}{\hat{V}_{\text{MI},\hat{\phi}}} = \frac{1}{m_{\text{P}}^2} \int_{\phi_f}^{\phi_*} d\phi J^2 \frac{\hat{V}_{\text{MI}}}{\hat{V}_{\text{MI},\phi}}. \quad (3.17)$$

Given that $\phi_f \ll \phi_*$, we can find a relation between ϕ_* and \hat{N}_* as follows

$$\hat{N}_* \simeq \frac{3c_T}{4m_{\text{P}}^2} (\phi_*^2 - \phi_f^2) \Rightarrow \phi_* \simeq 2m_{\text{P}} \sqrt{\hat{N}_*/3c_T}. \quad (3.18a)$$

Obviously, nSMI with subplanckian ϕ 's can be achieved if

$$\phi_* \leq m_{\text{P}} \Rightarrow c_T \geq 4\hat{N}_*/3 \simeq 76 \quad (3.18b)$$

for $\hat{N}_* \simeq 52$. Therefore we need relatively large c_T 's.

3.2.2 The amplitude A_s of the power spectrum of the curvature perturbation generated by ϕ at the pivot scale k_* is to be confronted with the data [1, 2], i.e.

$$A_s^{1/2} = \frac{1}{2\sqrt{3}\pi m_P^3} \frac{\widehat{V}_{\text{MI}}(\widehat{\phi}_*)^{3/2}}{|\widehat{V}_{\text{MI},\phi}(\widehat{\phi}_*)|} = \frac{1}{2\pi m_P^2} \sqrt{\frac{\widehat{V}_{\text{MI}}(\phi_*)}{6\widehat{\epsilon}(\phi_*)}} \simeq 4.685 \cdot 10^{-5}. \quad (3.19)$$

Since the scalars listed in Table 2 are massive enough during nSMI, the curvature perturbations generated by ϕ are solely responsible for A_s . Substituting Eqs. (3.15b) and (3.18a) into the relation above, we obtain

$$\sqrt{A_s} = \frac{\lambda m_P^2 f_T(\phi_*)^2}{8\sqrt{2}\pi c_T^2 \phi_*^2} \Rightarrow \lambda \simeq 6\pi \sqrt{2A_s} c_T / \widehat{N}_*. \quad (3.20a)$$

Combining the last equality with Eq. (3.19), we find that λ is to be proportional to c_T , for almost constant \widehat{N}_* . Indeed, we obtain

$$\lambda \simeq 3.97 \cdot 10^{-4} \pi c_T / \widehat{N}_* \Rightarrow c_T \simeq 41637 \lambda \text{ for } \widehat{N}_* \simeq 52. \quad (3.20b)$$

3.2.3 The (scalar) spectral index n_s , its running a_s , and the scalar-to-tensor ratio r must be consistent with the fitting [1, 2] of the observational data, i.e.,

$$(a) \ n_s = 0.96 \pm 0.014, \quad (b) \ -0.0314 \leq a_s \leq 0.0046 \quad \text{and} \quad (c) \ r < 0.11 \quad (3.21)$$

at 95% confidence level (c.l.). The observable quantities above can be estimated through the relations:

$$n_s = 1 - 6\widehat{\epsilon}_* + 2\widehat{\eta}_* \simeq 1 - 2/\widehat{N}_* - 9/2\widehat{N}_*^2, \quad (3.22a)$$

$$a_s = \frac{2}{3} (4\widehat{\eta}_*^2 - (n_s - 1)^2) - 2\widehat{\xi}_* \simeq -2\widehat{\xi}_* \simeq -2/\widehat{N}_*^2 + 3/2\widehat{N}_*^3, \quad (3.22b)$$

$$r = 16\widehat{\epsilon}_* \simeq 12/\widehat{N}_*^2, \quad (3.22c)$$

where $\widehat{\xi} = m_P^4 \widehat{V}_{\text{MI},\phi} \widehat{V}_{\text{MI},\phi\phi} / \widehat{V}_{\text{MI}}^2 = m_P \sqrt{2\widehat{\epsilon}} \widehat{\eta}_{,\phi} / J + 2\widehat{\eta}\widehat{\epsilon}$. The variables with subscript $*$ are evaluated at $\phi = \phi_*$ and Eqs. (3.15b) and (3.15c) have been employed.

3.3 THE EFFECTIVE CUT-OFF SCALE

As anticipated in Eq. (3.18b), the realization of nSMI with subplanckian ϕ 's requires relatively large c_T 's. This fact may [31] jeopardize the validity of the classical approximation, on which the analysis of the inflationary behavior is based. To see if this problem – which is rather questionable [18, 32] though – insists here, we have to extract the UV cut-off scale, Λ_{UV} , of the effective theory.

We first determine Λ_{UV} analyzing the small-field behavior of the model in EF along the lines of Ref. [9]. The EF action S in Eq. (2.7a) along the path of Eq. (3.6) is written as

$$S = \int d^4x \sqrt{-\widehat{\mathbf{g}}} \left(-\frac{1}{2} m_P^2 \widehat{\mathcal{R}} + \frac{1}{2} J^2 \dot{\phi}^2 - \widehat{V}_{\text{MI}0} + \dots \right). \quad (3.23a)$$

Given the form of J in Eq. (3.9b) an expansion of the kinetic term in Eq. (3.23a) about zero is not doable. Therefore we expand it about $\langle \phi \rangle = m_P / \sqrt{c_T}$ – see Eqs. (2.5b) and (2.8) – and we find

$$J^2 \dot{\phi}^2 = 6c_T \left(1 - \frac{2\sqrt{c_T} \delta\phi}{m_P} + \frac{3c_T \delta\phi^2}{m_P^2} - \frac{4c_T \sqrt{c_T} \delta\phi^3}{m_P^3} + \frac{5c_T^2 \delta\phi^4}{m_P^4} - \dots \right) \dot{\phi}^2, \quad (3.23b)$$

where $\delta\phi = (\phi - M)$. Since there is no canonically normalized leading kinetic term, we define the canonically normalized inflaton at the SUSY vacuum $\widehat{\delta\phi} = \sqrt{6c_T} \delta\phi$ – see also Sec. 5.1 – and we reexpress Eq. (3.23b) in terms of $\widehat{\delta\phi}$, with result

$$J^2 \dot{\phi}^2 = \left(1 - \sqrt{\frac{2}{3}} \frac{\widehat{\delta\phi}}{m_P} + \frac{1}{2} \frac{\widehat{\delta\phi}^2}{m_P^2} - \frac{\sqrt{2}}{3\sqrt{3}} \frac{\widehat{\delta\phi}^3}{m_P^3} + \frac{5}{36} \frac{\widehat{\delta\phi}^4}{m_P^4} - \dots \right) \dot{\widehat{\delta\phi}}^2. \quad (3.23c)$$

On the other hand, $\widehat{V}_{\text{MI}0}$ in Eq. (3.4) can be expanded also in terms of $\widehat{\delta\phi}$ as follows

$$\widehat{V}_{\text{MI}0} = \frac{\lambda^2 m_{\text{P}}^2}{6c_T^2} \widehat{\delta\phi}^2 \left(1 - \sqrt{\frac{3}{2}} \frac{\widehat{\delta\phi}}{m_{\text{P}}} + \frac{25}{24} \frac{\widehat{\delta\phi}^2}{m_{\text{P}}^2} - \dots \right). \quad (3.23d)$$

From the derived expressions in Eqs. (3.23c) and (3.23d) we conclude that $\Lambda_{\text{UV}} = m_{\text{P}}$ and therefore our model is valid up to m_{P} as the original Starobinsky model [9].

The resulting Λ_{UV} represents essentially the unitarity-violation scale [31] of the $\delta\phi - \delta\phi$ scattering process via s -channel graviton, $h^{\mu\nu}$, exchange in the JF. The relevant vertex is $c_T \delta\phi^2 \square h / m_{\text{P}}$ – with $h = h_{\mu}^{\mu}$ – can be derived from the first term in the r.h.s of Eq. (2.7b) expanding the JF metric $g_{\mu\nu}$ about the flat spacetime metric $\eta_{\mu\nu}$ and the inflaton ϕ about its v.e.v as follows:

$$g_{\mu\nu} \simeq \eta_{\mu\nu} + h_{\mu\nu}/m_{\text{P}} \quad \text{and} \quad \phi = \langle\phi\rangle + \delta\phi. \quad (3.24)$$

Retaining only the terms with two derivatives of the excitations, the part of the lagrangian corresponding to the two first terms in the r.h.s of Eq. (2.7b) takes the form

$$\begin{aligned} \delta\mathcal{L} &= -\frac{\langle f_K \rangle}{4} F_{\text{EH}}(h^{\mu\nu}) + \left(m_{\text{P}} \langle f_{K,\phi} \rangle + \frac{c_T \delta\phi}{2m_{\text{P}}} \right) (\square h - \partial_{\mu} \partial_{\nu} h^{\mu\nu}) \delta\phi + \dots \\ &= -\frac{1}{8} F_{\text{EH}}(\bar{h}^{\mu\nu}) + \frac{1}{2} \partial_{\mu} \bar{\delta\phi} \partial^{\mu} \bar{\delta\phi} + \frac{1}{2\sqrt{2}} \frac{c_T}{m_{\text{P}}} \frac{\sqrt{\langle f_K \rangle}}{\langle f_K \rangle} \bar{\delta\phi}^2 \square \bar{h} + \dots, \end{aligned} \quad (3.25a)$$

where the function F_{EH} , related to the the linearized Einstein-Hilbert part of the lagrangian, reads

$$F_{\text{EH}}(h^{\mu\nu}) = h^{\mu\nu} \square h_{\mu\nu} - h \square h + 2\partial_{\rho} h^{\mu\rho} \partial^{\nu} h_{\mu\nu} - 2\partial_{\nu} h^{\mu\nu} \partial_{\mu} h \quad (3.25b)$$

and the JF canonically normalized fields $\bar{h}_{\mu\nu}$ and $\bar{\delta\phi}$ are defined by the relations

$$\bar{\delta\phi} = \sqrt{\frac{\langle \bar{f}_K \rangle}{\langle f_K \rangle}} \delta\phi \quad \text{and} \quad \bar{h}_{\mu\nu} = \sqrt{\langle f_K \rangle} h_{\mu\nu} + \frac{m_{\text{P}} \langle f_{K,\phi} \rangle}{\sqrt{\langle f_K \rangle}} \eta_{\mu\nu} \delta\phi \quad \text{with} \quad \bar{f}_K = 3m_{\text{P}}^2 f_{K,\phi}^2. \quad (3.25c)$$

The interaction originating from the last term in the r.h.s of Eq. (3.25a) gives rise to a scattering amplitude which is written in terms of the center-of-mass energy E as follows

$$\mathcal{A} \sim \left(\frac{E}{\Lambda_{\text{UV}}} \right)^2 \quad \text{with} \quad \Lambda_{\text{UV}} = \frac{m_{\text{P}}}{3\sqrt{2}c_T} \frac{\langle \bar{f}_K \rangle}{\sqrt{\langle f_K \rangle}} = m_{\text{P}}, \quad (3.26)$$

where $\langle f_K \rangle = 1/2$ and $\langle \bar{f}_K \rangle = 3c_T$ and Λ_{UV} is identified as the UV cut-off scale in the JF, since \mathcal{A} remains within the validity of the perturbation theory provided that $E < \Lambda_{\text{UV}}$.

Although the expansions in Eqs. (3.23d) and (3.25a) are obtained for $\phi \simeq \langle\phi\rangle$ and are not valid [32] during nSMI, we consider Λ_{UV} as the overall UV cut-off scale of the model since reheating is an unavoidable stage of the inflationary dynamics [9]. Therefore, the validity of the effective theory implies [31]

$$\widehat{V}_{\text{MI}}(\phi_{\star})^{1/4} \ll \Lambda_{\text{UV}} \quad \text{with} \quad \Lambda_{\text{UV}} = m_{\text{P}}, \quad (3.27)$$

which is much less restrictive than the corresponding condition applied in the models of nMI with quartic scalar potential, where Λ_{UV} turns out to be equal to m_{P} divided by the strength of the non-minimal coupling to gravity – cf. Ref. [9, 21, 22, 25].




Figure 1: The inflationary scale $\hat{V}_{\text{MI}}^{1/4}$ as a function of ϕ for $\lambda = 2.26 \cdot 10^{-5}$ and $M = m_{\text{P}}$ ($c_T = 1$) or $\lambda = 1.7 \cdot 10^{-3}$ and $M/m_{\text{P}} = 0.115$ ($c_T = 76$) or $\lambda = 0.1$ and $M/m_{\text{P}} = 0.015$ ($c_T = 4500$). The values corresponding to ϕ_* and ϕ_{f} are also depicted.

3.4 NUMERICAL RESULTS

As can be easily seen from the relevant expressions above, the inflationary dynamics of our model depends on the following parameters:

$$\lambda, c_T, \lambda_\mu, k_{SS} = k_S, k_{SH}, k_{S\tilde{N}^c}, M_{iN^c} \text{ and } T_{\text{rh}}.$$

Recall that M is related to c_T via Eq. (2.8). Our results are essentially independent of λ_μ and k 's, provided that we choose them so as \hat{m}_{h-}^2 and \hat{m}_s^2 in Table 2 are positive for every allowed λ . We therefore set $\lambda_\mu = 10^{-6}$, $k_S = k_{S\tilde{N}^c} = 1$ and $k_{SH} = 1.5$ throughout our calculation. Moreover we take into account the contribution to \hat{V}_{MI} , Eq. (3.13), only from the heaviest N_i^c which is taken to be $M_{3N^c} = 10^{14}$ GeV – cf. Sec. 5.5. We also choose $\Lambda \simeq 10^{13}$ GeV so as the one-loop corrections in Eq. (3.13) vanish at the SUSY vacuum, Eqs. (2.5b) and (2.8). Finally T_{rh} can be calculated self-consistently in our model as a function of the inflaton mass, $\hat{m}_{\delta\phi}$ and the strength of the various inflaton decays – see Sec. 5.1. However, since the inflationary predictions depend very weakly on T_{rh} – see Eq. (3.14) – we prefer to take here a constant $T_{\text{rh}} = 6 \cdot 10^8$ GeV as suggested by our results on post-inflationary evolution – see Sec. 5.5. Upon substitution of \hat{V}_{MI} from Eq. (3.13) in Eqs. (3.15a), (3.17) and (3.19) we extract the inflationary observables as functions of c_T , λ and ϕ_* . The two latter parameters can be determined by enforcing the fulfilment of Eq. (3.14) and (3.19), for every chosen c_T . Our numerical findings are quite close to the analytic ones listed in Sec. 3.2 for the sake of presentation.

The importance of the two extra variables (M and c_T) – in Eqs. (2.1c), (2.6a) and (2.6c) – compared to the Cecotti model [5–8] in reducing ϕ_* below m_{P} can be easily inferred from Fig. 1. We there depict $\hat{V}_{\text{MI}}^{1/4}$ as a function of ϕ (both normalized to m_{P}) for $\lambda = 2.26 \cdot 10^{-5}$ and $c_T = 1$ or $\lambda = 0.0017$ and $c_T = 76$ or $\lambda = 0.1$ and $c_T = 4500$ – the last value saturates an upper bound on c_T derived in Sec. 4. Note that for $c_T = 1$ (or $x_M = 1$) our result matches that of the original Starobinsky model [4, 52] – with the mass scale appearing in that model being replaced by $\lambda m_{\text{P}} \simeq 2.2 \cdot 10^{13}$ GeV. Increasing c_T , λ increases too, whereas ϕ_* and M decrease and for $c_T \geq 76$, ϕ_* becomes subplanckian. On the other hand, we have to clarify that the corresponding values of the inflaton in the EF remain transplanckian, since integrating the first equation in Eq. (3.9b) and using Eqs. (3.18a) and (3.16a) we find:

$$\hat{\phi} = \hat{\phi}_c + \sqrt{6}m_{\text{P}} \ln(\phi/M) \Rightarrow \begin{cases} \hat{\phi}_* - \hat{\phi}_c \simeq \sqrt{6}m_{\text{P}} \ln 2(\hat{N}_*/3)^{1/2} \\ \hat{\phi}_{\text{f}} - \hat{\phi}_c \simeq \sqrt{6}m_{\text{P}} \ln(1 + 2/\sqrt{3})^{1/2}. \end{cases} \quad (3.28)$$




Figure 2: The allowed by Eqs. (3.14), (3.19) and (3.27) values of c_T (solid line), $10^3 x_M$ (dashed line) and $10^3 \hat{V}_{\text{MI}}(\phi_\star)^{1/4}/\Lambda_{\text{UV}}$ (dotted line) [ϕ_f (solid line) and ϕ_\star (dashed line)] versus λ (a) [(b)] for $k_S = 1$, $\lambda_\mu = 10^{-6}$, $M_{3N^c} = 10^{14}$ GeV and $T_{\text{rh}} = 6 \cdot 10^8$ GeV.

where $\hat{\phi}_c$ is a constant of integration. E.g., setting $\hat{\phi}_c = 0$, we obtain $\hat{\phi}_\star = 5.3 m_P$ and $\hat{\phi}_f = 0.94 m_P$ for any c_T – with constant \hat{N}_\star . We do not consider this result as an upset of our proposal, since the inflaton field defined in the JF enters W_{MI} and K . Therefore, possible corrections from non-renormalizable terms, which may be avoided for subplanckian values of inflaton, are applied in this frame, which is mostly considered as the physical frame.

From Fig. 1 we also infer that $\hat{V}_{\text{MI}}^{1/4}/m_P$ remains almost constant during nSMI. Indeed, if we plug Eqs. (3.18a) and (3.20b) into Eq. (3.4), we obtain

$$\hat{V}_{\text{MI}0}(\phi_\star)^{1/4}/m_P \simeq \left(3\pi\sqrt{2A_s}/\hat{N}_\star\right)^{1/2} \simeq 0.0033 \ll 1. \quad (3.29)$$

This result is more explicitly displayed in Fig. 2 too, where we draw the allowed values of c_T (solid line), $10^3 x_M$ (dashed line) and $10^3 \hat{V}_{\text{MI}}(\phi_\star)^{1/4}/m_P$ (dotted line) [ϕ_f (solid line) and ϕ_\star (dashed line)] versus λ (a) [(b)]. The lower bound of the depicted lines comes from the saturation of Eq. (3.18b) whereas the upper bound originates from the perturbative bound on λ , $\lambda \leq \sqrt{4\pi} \simeq 3.54$. In Fig. 2-(a) we see that Eq. (3.27) is readily satisfied along the various curves and we can verify our analytic estimation in Eq. (3.20b). Moreover, the variation of ϕ_f and ϕ_\star as a function of λ – drawn in Fig. 2-(b) – is consistent with Eqs. (3.16a) and (3.18a). The overall allowed parameter space of our model is

$$76 \lesssim c_T \lesssim 1.5 \cdot 10^5, \quad 0.11 \gtrsim x_M \gtrsim 0.002 \quad \text{and} \quad 1.7 \cdot 10^{-3} \lesssim \lambda \lesssim 3.54 \quad \text{for} \quad \hat{N}_\star \simeq 52. \quad (3.30a)$$

Letting λ or c_T vary within its allowed region in Eq. (3.30a), we obtain

$$0.961 \lesssim n_s \lesssim 0.963, \quad -7.4 \lesssim a_s/10^{-4} \lesssim -6.7 \quad \text{and} \quad 4.2 \gtrsim r/10^{-3} \gtrsim 3.8, \quad (3.30b)$$

whereas the masses of the various scalars in Table 2 remain well above \hat{H}_{MI} both during and after nSMI for the selected k_S , λ_μ and M_{3N^c} . E.g., for $\phi = \phi_\star$ and $c_T = 150$, we obtain

$$(\hat{m}_\theta^2, \hat{m}_s^2, \hat{m}_{h^-}^2, \hat{m}_{h^+}^2, \hat{m}_{3\nu^c}^2) / \hat{H}_{\text{MI}}^2 \simeq (4, 905, 342, 342, 282). \quad (3.30c)$$

Clearly, the predicted a_s and r lie within the allowed ranges given in Eq. (3.21b) and Eq. (3.21c) respectively, whereas n_s turns out to be impressively close to its central observationally favored value – see Eq. (3.21a). Therefore, the inclusion of extra parameters, compared to the Cecotti model [5–8], does not affect the successful predictions on the inflationary observables.

4 THE R SYMMETRY AND THE μ PROBLEM OF MSSM

A byproduct of the R symmetry associated with our model is that it assists us to understand the origin of μ term of MSSM. To see how this works, in Sec. 4.1, we estimate the μ parameter and, in Sec. 4.2, we control its compatibility with phenomenologically acceptable values obtained in the context of the *Constrained MSSM* (CMSSM) [47].

4.1 GENERATION OF THE μ TERM OF MSSM

First we write the part of the scalar potential which includes the SSB terms corresponding to W_{MI} in Eq. (2.1c). We have

$$V_{\text{soft}} = \left(\lambda A_\lambda S T^2 + \lambda_\mu A_\mu S H_u H_d + B_{iN^c} M_{iN^c} \tilde{N}_i^c \tilde{N}_i^c - a_S S \lambda M^2 + \text{h.c.} \right) + m_\alpha^2 |\Phi^\alpha|^2, \quad (4.1)$$

where m_α , A_λ , A_μ , B_{iN^c} and a_S are SSB mass parameters. Rotating S in the real axis by an appropriate R -transformation, choosing conveniently the phases of A_λ and a_S so as the total low energy potential $V_{\text{tot}} = V_{\text{SUSY}} + V_{\text{soft}}$ to be minimized – see Eq. (2.5a) – and substituting in V_{soft} the SUSY v.e.s of T , H_u , H_d and \tilde{N}_i^c from Eq. (2.5b) we get

$$\langle V_{\text{tot}}(S) \rangle = \frac{\lambda^2 M^2}{3c_T m_P^2} S^2 \left(c_T M^2 - \frac{1}{3} S^2 \right) - 2\lambda a_{3/2} m_{3/2} M^2 S, \quad (4.2)$$

where we set $|A_\lambda| + |a_S| = 2a_{3/2}m_{3/2}$ with $m_{3/2}$ being the \tilde{G} mass and $a_{3/2} > 0$ a parameter of order unity which parameterizes our ignorance for the dependence of $|A_\lambda|$ and $|a_S|$ on $m_{3/2}$. Making use of the induced-gravity condition in Eq. (6), we can write the extremum condition for $\langle V_{\text{tot}}(S) \rangle$ as follows

$$\frac{d}{dS} \langle V_{\text{tot}}(S) \rangle = 0 \Rightarrow -\frac{2\lambda^2 m_P^2}{3c_T^2} \left(\frac{2}{3} \frac{S^3}{m_P^2} - S + \frac{3}{\lambda} c_T a_{3/2} m_{3/2} \right) = 0. \quad (4.3a)$$

For $\langle S \rangle \ll m_P$, one of the solutions of the last equation is

$$\langle S \rangle \simeq 3c_T a_{3/2} m_{3/2} / \lambda \simeq 1.25 \cdot 10^5 a_{3/2} m_{3/2}, \quad (4.3b)$$

where we employ Eq. (3.20b) which yields λ as a function of c_T . At this S value, $\langle V_{\text{tot}}(S) \rangle$ develops a minimum since

$$\frac{d^2}{dS^2} \langle V_{\text{tot}}(S) \rangle = \frac{2\lambda^2}{3c_T^2} (m_P^2 - 2S^2) \quad (4.4a)$$

becomes positive for $m_{3/2} \ll m_P$. Indeed, inserting Eq. (4.3b) into Eq. (4.4a) we obtain the constraint

$$3\sqrt{2}c_T \frac{a_{3/2}m_{3/2}}{\lambda m_P} \leq 1 \Rightarrow m_{3/2} \leq \frac{\lambda m_P}{3\sqrt{2}a_{3/2}c_T} \simeq 1.38 \cdot 10^{13} \text{ GeV}, \quad (4.4b)$$

which is comfortably fulfilled in the case of the low scale SUSY. The other two solutions of Eq. (4.3a) violate this bound. The generated μ parameter from the second term in the r.h.s of Eq. (2.1c) is

$$\mu = \lambda_\mu \langle S \rangle \simeq 3\lambda_\mu c_T a_{3/2} m_{3/2} / \lambda \simeq 1.25 \cdot 10^5 \lambda_\mu a_{3/2} m_{3/2}. \quad (4.5)$$

Note that λ_μ (and so μ) may have either sign without any essential alteration in the stability analysis of the inflationary system – see Table 2. Thanks to the magnitude of the proportionality constant, any ratio $|\mu|/a_{3/2}m_{3/2} \lesssim 2.5$ is accessible for the λ_μ values allowed by Eq. (3.12) with k_{SH} of order unity. Ergo, the resulting μ is λ independent, in sharp contrast to the originally proposed scheme in Ref. [27], and no hierarchy of the type $\mu \ll m_{3/2}$ is required – see e.g. the second row of Table 3 below where $\mu/a_{3/2}m_{3/2} \simeq 2.13$.

Obviously the proposed resolution of the μ problem of MSSM relies on the existence of non-zero A_λ and/or a_S . These issues depend on the adopted model of SSB. We single out the following cases:

- (i) If we wish to be fully consistent the no-scale structure of K and suppose that the modulus, z , which is responsible for the SSB, is contained (somehow) in the logarithm of Eq. (2.6a), K is of the “sequestered-sector” form [58] and has the property that it generates no tree-level SSB scalar masses for the visible-sector fields and vanishing trilinear coupling constants. In this case the anomaly-mediated SSB [33, 58] is the dominant mechanism for obtaining $A_\lambda \neq 0$ and/or $a_S \neq 0$. Since the involved superfields T and S are G_{SM} singlets, we expect $A_\lambda = 0$. However, according to the superconformal formalism, M^2 can be rescaled as $M^2\varphi^2$ (where φ is a superconformal compensator) and, in the presence of SSB, a non vanishing $a_S = 2m_{3/2}$ comes out.
- (ii) If we decide to deviate from the no-scale form of K in Eq. (2.6a), we can suppose that z is not contained in the logarithm, and has an almost canonical Kähler potential [10, 59]. In a such circumstance, both A_λ and a_S are expected to be non-zero, as in the gravity-mediated SSB [59], giving rise again to $\langle S \rangle \neq 0$.

In both cases above, our superpotential in Eq. (2.1a) has to be extended by a SSB sector which should ensure the successful stabilization of z – cf. Ref. [10, 58, 60]. We expect that these terms do not disturb the inflationary dynamics. Alternatively, the μ problem can be resolved [61] by imposing a Peccei-Quinn symmetry which is broken spontaneously at an intermediate scale by the v.e.v.s of two G_{SM} singlets which enter the superpotential via non-renormalizable terms. This scheme, already adopted, e.g., in Ref. [22, 62], can be applied as first realized in Ref. [61] in the case (ii) above and somehow modified in the case (i).

Let us clarify, finally, that the due hierarchy in Eq. (3.12) between λ_μ and λ , is the inverse to that imposed in the models [27] of FHI, where S plays the role of inflaton and T , H_u and H_d are confined at zero – playing the role of the waterfall fields. This is because, at the end of FHI, the mass squared of T becomes negative for $S < M/\sqrt{2}$ and the mass matrix squared of the scalars $H_u - H_d$ develop a negative eigenvalue for $S < M\sqrt{\lambda/2\lambda_\mu}$. Consequently, the correct cosmological scenario can be attained if we ensure that, at the end of FHI, T acquires its v.e.v, while H_u and H_d remain equal to zero. To this end we demand [27] $\lambda_\mu > \lambda$ so as the tachyonic instability in the T direction occurs first, and T start evolving towards its v.e.v, whereas H_u and H_d continue to be confined to zero. In our case, though, $|T|$ is the inflaton while S and the $H_u - H_d$ system are safely stabilized at the origin both during and after the end of nSMI. Therefore, $|T|$ is led at its vacuum whereas H_u and H_d take their non-vanishing v.e.v.s during the electroweak phase transition triggered by radiative corrections.

4.2 CONNECTION WITH THE MSSM PHENOMENOLOGY

Taking advantage from the updated investigation of the parameter space of CMSSM in Ref. [47] we can easily verify that the μ and $m_{3/2}$ values satisfying Eq. (4.5) are consistent with the values required by the analyses of the low energy observables of MSSM. We concentrate on CMSSM which is the most predictive, restrictive and well-motivated version of MSSM, employing the free parameters:

$$\text{sign}\mu, \tan\beta = \langle H_u \rangle / \langle H_d \rangle, M_{1/2}, m_0, \text{ and } A_0.$$

Here $\text{sign}\mu$ is the sign of μ and the three last mass parameters denote the common gaugino mass, scalar mass, and trilinear coupling constant, respectively, defined at a high scale which is determined by the unification of the gauge coupling constants. The parameter $|\mu|$ is not free, since it is computed at low scale enforcing the conditions for the electroweak symmetry breaking. The values of these parameters can be tightly restricted imposing a number of cosmo-phenomenological constraints. Some updated results are recently presented in Ref. [47], where we can also find the best-fit values of $|A_0|$, m_0 and $|\mu|$ listed in Table 3. We see that there are four allowed regions characterized by the specific mechanism for suppressing the relic density of the lightest sparticle which can act as dark matter.

CMSSM REGION	$ A_0 (\text{TeV})$	$m_0(\text{TeV})$	$ \mu (\text{TeV})$	$a_{3/2}$	$\lambda_\mu(10^{-6})$
A/H Funnel	9.9244	9.136	1.409	1.086	1.071
$\tilde{\tau}$ Coannihilation	1.2271	1.476	2.621	0.831	16.11
\tilde{t} Coannihilation	9.965	4.269	4.073	2.33	3.081
$\tilde{\chi}_1^\pm$ Coannihilation	9.2061	9.000	0.983	1.023	0.805

Table 3: The required λ_μ values for $m_0 = m_{3/2}$, $|A_\lambda| = |a_S| = |A_0|$ and $k_{SH} = 1.5$ which render our model compatible with the best-fit points of the CMSSM as found in Ref. [47].

If we identify m_0 with $m_{3/2}$ and $|A_0|$ with $|A_\lambda| = |a_S|$ we can derive first $a_{3/2}$ and then the λ_μ values which yield the phenomenologically desired $|\mu|$ – see the two rightmost columns in Table 3. Here we assume that renormalization effects are negligible. Since the required λ_μ 's are compatible with Eq. (3.12) for $k_{SH} = 1.5$, we conclude that the whole inflationary scenario can be successfully combined with CMSSM. On the other hand, only the regions which become consistent with the dark matter requirements thanks to A/H funnel and $\tilde{\chi}_1^\pm$ coannihilation can be consistent with the gravitino limit on T_{rh} – see Sec. 5.4. Indeed, in these cases $m_{3/2} \simeq 9$ TeV and so, the unstable \tilde{G} becomes cosmologically safe with the presented in Table 4 T_{rh} values, necessitated for satisfactory leptogenesis.

5 NON-THERMAL LEPTOGENESIS AND NEUTRINO MASSES

We below specify how our inflationary scenario makes a transition to the radiation dominated era (Sec. 5.1) and give an explanation of the observed BAU (Sec. 5.2) consistently with the \tilde{G} constraint and the low energy neutrino data (Sec. 5.3). Our results are summarized in Sec. 5.5.

5.1 THE INFLATON DECAY

When nSMI is over, the inflaton continues to roll down towards the SUSY vacuum, Eq. (2.5b). Soon after, it settles into a phase of damped oscillations around the minimum of \hat{V}_{MIO} – note that θ is stabilized during and after nSMI at the origin and so, it does not participate neither into inflationary nor to post-inflationary dynamics. The (canonically normalized) inflaton, $\hat{\delta\phi} = \sqrt{6c_T}\delta\phi$ – see, also, Sec. 3.3 –, acquires mass which is given by

$$\hat{m}_{\delta\phi} = \left\langle \hat{V}_{\text{MIO},\hat{\phi}\hat{\phi}} \right\rangle^{1/2} = \left\langle \hat{V}_{\text{MIO},\phi\phi}/J^2 \right\rangle^{1/2} = \lambda m_{\text{P}}/\sqrt{3}c_T \simeq 3 \cdot 10^{13} \text{ GeV}, \quad (5.1)$$

where we make use of Eq. (3.20b) in the last step. Since Eq. (3.8) implies $\langle K_{A\bar{A}} \rangle = 1$ for $\langle x_\phi \rangle = 1/\sqrt{c_T}$ – see Eqs. (3.3), (2.5b) and (2.8) –, the EF canonically normalized fields Φ^A in Eq. (2.6b) are not distinguished from the JF ones at the SUSY vacuum.

The decay of $\hat{\delta\phi}$ is processed through the following decay channels:

5.1.1 Decay channel into N_i^c 's. The lagrangian which describes these decay channels arises from the part of the SUGRA langrangian [59] containing two fermions. In particular,

$$\begin{aligned} \mathcal{L}_{\hat{\delta\phi} \rightarrow N_i^c} &= -\frac{1}{2}e^{K/2m_{\text{P}}^2} W_{,N_i^c N_i^c} N_i^c N_i^c + \text{h.c.} = \frac{3}{2} \frac{M}{m_{\text{P}}} c_T^{1/2} \delta\phi N_i^c N_i^c + \dots \\ &= \lambda_{iN^c} \hat{\delta\phi} N_i^c N_i^c + \dots \quad \text{with} \quad \lambda_{iN^c} = \sqrt{3} M_{iN^c} / 2\sqrt{2} m_{\text{P}}, \end{aligned} \quad (5.2a)$$

where an expansion around $\langle \phi \rangle$ is performed in order to extract the result above. We observe that although there is not direct coupling between T and N_i^c in W_{MI} – recall that we assume that the third

term in the r.h.s of Eq. (2.1c) prevails over the last one –, an adequately efficient decay channel arises, which gives rise to the following decay width

$$\widehat{\Gamma}_{\delta\phi \rightarrow N_i^c} = \frac{1}{16\pi} \lambda_{iN^c}^2 \widehat{m}_{\delta\phi} (1 - 4M_{iN^c}^2/\widehat{m}_{\delta\phi}^2)^{3/2}, \quad (5.2b)$$

where we take into account that $\widehat{\delta\phi}$ decays to identical particles.

5.1.2. Decay channel into H_u and H_d . The lagrangian term which describes the relevant interaction comes from the F-term SUGRA scalar potential in Eq. (3.1). Namely, we obtain

$$\begin{aligned} \mathcal{L}_{\widehat{\delta\phi} \rightarrow H_u H_d} &= -\frac{1}{2} e^{K/m_P^2} K^{SS^*} |W_S|^2 = -\frac{1}{2} \lambda \lambda_\mu (\phi^2 - M^2) H_u^* H_d^* + \dots \\ &= -\lambda_H \widehat{m}_{\delta\phi} \widehat{\delta\phi} H_u^* H_d^* + \dots \quad \text{with } \lambda_H = \lambda_\mu/\sqrt{2}. \end{aligned} \quad (5.3a)$$

This interaction gives rise to the following decay width

$$\widehat{\Gamma}_{\delta\phi \rightarrow H} = \frac{2}{8\pi} \lambda_H^2 \widehat{m}_{\delta\phi}, \quad (5.3b)$$

where we take into account that H_u and H_d are $SU(2)_L$ doublets. Eq. (3.12) facilitates the reduction of $\widehat{\Gamma}_{\delta\phi \rightarrow H}$ to a level which allows for the decay mode into N_i^c 's playing its important role for nTL.

5.1.3. Three-particle decay channels. Focusing on the same part of the SUGRA langrangian [59] as in the paragraph 5.1.1, for a typical trilinear superpotential term of the form $W_y = yXYZ$ – cf. Eq. (2.1b) –, where y is a Yukawa coupling constant, we obtain the interactions described by

$$\begin{aligned} \mathcal{L}_{\widehat{\delta\phi} \rightarrow XYZ} &= -\frac{1}{2} e^{K/2m_P^2} (W_{y,YZ} \psi_Y \psi_Z + W_{y,XZ} \psi_X \psi_Z + W_{y,XY} \psi_X \psi_Y) + \text{h.c.} \\ &= \lambda_y \frac{\widehat{\delta\phi}}{m_P} (X \psi_Y \psi_Z + Y \psi_X \psi_Z + Z \psi_X \psi_Y) + \text{h.c.} \quad \text{with } \lambda_y = \sqrt{3/2} (y/2), \end{aligned} \quad (5.4a)$$

where ψ_X, ψ_Y and ψ_Z are the chiral fermions associated with the superfields X, Y and Z whose the scalar components are denoted with the superfield symbol. Working in the large $\tan \beta$ regime which yields similar y 's for the 3rd generation, we conclude that the interaction above gives rise to the following 3-body decay width

$$\widehat{\Gamma}_{\delta\phi \rightarrow XYZ} = \frac{14n_f}{512\pi^3} \lambda_y^2 \frac{\widehat{m}_{\delta\phi}^3}{m_P^2}, \quad (5.4b)$$

where for the third generation we take $y \simeq (0.4 - 0.6)$, computed at the $\widehat{m}_{\delta\phi}$ scale, and $n_f = 14$ [$n_f = 16$] for $\widehat{m}_{\delta\phi} < M_{3N^c}$ [$\widehat{m}_{\delta\phi} > M_{3N^c}$] – summation is taken over $SU(3)_c$ and $SU(2)_L$ indices.

Since the decay width of the produced N_i^c is much larger than $\widehat{\Gamma}_{\delta\phi}$ the reheating temperature, T_{rh} , is exclusively determined by the inflaton decay and is given by [63]

$$T_{\text{rh}} = \left(\frac{72}{5\pi^2 g_*} \right)^{1/4} \sqrt{\widehat{\Gamma}_{\delta\phi} m_P} \quad \text{with } \widehat{\Gamma}_{\delta\phi} = \widehat{\Gamma}_{\delta\phi \rightarrow N_i^c} + \widehat{\Gamma}_{\delta\phi \rightarrow H} + \widehat{\Gamma}_{\delta\phi \rightarrow XYZ}, \quad (5.5)$$

where $g_* \simeq 228.75$ counts the effective number of relativistic degrees of freedom of the MSSM spectrum at the temperature $T \simeq T_{\text{rh}}$. Let us clarify here that in our models there is no decay of a scalaron as in the original (non-SUSY) [3, 13] Starobinsky inflation and some [11] of its SUGRA realizations; thus, T_{rh} in our case is slightly lower than that obtained there. Indeed, spontaneous decay of the inflaton to scalars takes place only via three-body interactions which are suppressed compared to the two-body decays of scalaron. On the other hand, we here get also $\widehat{\Gamma}_{\delta\phi \rightarrow H}$ in Eq. (5.3b), due to explicit coupling of $\widehat{\delta\phi}$ into H_u and H_d , which can be kept at the same level with $\widehat{\Gamma}_{\delta\phi \rightarrow XYZ}$ due to the rather low λ_μ 's required here – see Eq. (3.12).

5.2 LEPTON-NUMBER AND GRAVITINO ABUNDANCES

The mechanism of nTL [37] can be activated by the out-of-equilibrium decay of the N_i^c 's produced by the $\widehat{\delta\phi}$ decay, via the interactions in Eq. (5.2a). If $T_{\text{rh}} \ll M_{iN^c}$, the out-of-equilibrium condition [64] is automatically satisfied. Namely, N_i^c decay into (fermionic and bosonic components of) H_u and L_i via the tree-level couplings derived from the last term in the r.h.s of Eq. (2.1b). The resulting – see Sec. 5.3 – lepton-number asymmetry ε_i (per N_i^c decay) after reheating can be partially converted via sphaleron effects into baryon-number asymmetry. In particular, the B yield can be computed as

$$(a) \ Y_B = -0.35Y_L \quad \text{with} \quad (b) \ Y_L = 2\frac{5}{4}\frac{T_{\text{rh}}}{\widehat{m}_{\delta\phi}} \sum_{i=1}^3 \frac{\widehat{\Gamma}_{\delta\phi \rightarrow N_i^c}}{\widehat{\Gamma}_{\delta\phi}} \varepsilon_i. \quad (5.6)$$

The numerical factor in the r.h.s of Eq. (5.6a) comes from the sphaleron effects, whereas the one (5/4) in the r.h.s of Eq. (5.6b) is due to the slightly different calculation [63] of T_{rh} – cf. Ref. [64].

The required for successful nTL T_{rh} must be compatible with constraints on the \widetilde{G} abundance, $Y_{\widetilde{G}}$, at the onset of *nucleosynthesis* (BBN). This is estimated to be [39, 40]:

$$Y_{\widetilde{G}} \simeq 1.9 \cdot 10^{-22} T_{\text{rh}}/\text{GeV}, \quad (5.7)$$

where we assume that \widetilde{G} is much heavier than the gauginos of MSSM. Let us note that non-thermal \widetilde{G} production within SUGRA is [35] also possible but strongly dependent on the mechanism of SSB. It can be easily suppressed [36, 65] when a tiny mixing arises between the inflaton and the field responsible for SSB provided that the mass of the latter is much lower than the inflationary scale. Therefore, we here prefer to adopt the conservative estimation of $Y_{\widetilde{G}}$ in Eq. (5.7).

Both Eqs. (5.6) and (5.7) calculate the correct values of the B and \widetilde{G} abundances provided that no entropy production occurs for $T < T_{\text{rh}}$. This fact can be achieved if the Polonyi-like field z decays early enough without provoking a late episode of secondary reheating. In both cases of Sec. 4, z is expected to be displaced from its true minimum to lower values due to large mass that it acquires during nSMI. In the course of the decaying-inflaton period which follows nSMI, z adiabatically tracks an instantaneous minimum [66] until the Hubble parameter becomes of the order of its mass. Successively it starts to oscillate about the true SUSY breaking minimum and may or may not dominate the Universe, depending on the initial amplitude of the coherent oscillations. The domination may be eluded in a very promising scenario [10, 60] which can be constructed assuming that z is strongly stabilized through a large enough coupling in a higher order term of Kähler potential, similar to that used for the stabilization of S – see Eq. (2.6a). A subsequent difficulty is the possible over-abundance of the LSPs which are produced by the z decay. From that perspective, it seems that the case (ii) – cf. Ref. [60, 66] – is more tolerable than the case (i) – see Ref. [67].

5.3 LEPTON-NUMBER ASYMMETRY AND NEUTRINO MASSES

As mentioned in Sec. 5.2, the decay of N_i^c emerging from the $\widehat{\delta\phi}$ decay, can generate [68] a lepton asymmetry ε_i caused by the interference between the tree and one-loop decay diagrams, provided that a CP-violation occurs in h_{ijN} 's – see Eq. (2.1b). The produced ε_i can be expressed in terms of the Dirac mass matrix of ν_i , m_D , defined in the N_i^c -basis, as follows:

$$\varepsilon_i = \sum_{j \neq i} \frac{\text{Im} \left[(m_D^\dagger m_D)_{ij}^2 \right]}{8\pi \langle H_u \rangle^2 (m_D^\dagger m_D)_{ii}} \left(F_S(x_{ij}) + F_V(x_{ij}) \right), \quad (5.8a)$$

where $x_{ij} := M_{jN^c}/M_{iN^c}$, $\langle H_u \rangle \simeq 174 \text{ GeV}$, for large $\tan \beta$ and the functions $F_{V,S}$ read [68]

$$F_V(x) = -x \ln(1 + x^{-2}) \quad \text{and} \quad F_S(x) = \frac{-2x}{x^2 - 1}. \quad (5.8b)$$

Also m_D is the Dirac mass matrix of ν_i 's and $m_D^\dagger m_D$ in Eq. (5.8a) can be written as follows:

$$m_D^\dagger m_D = U^{c\dagger} d_D^\dagger d_D U^c. \quad (5.8c)$$

where U^c are the 3×3 unitary matrix which relates N_i^c in the N_i^c -basis with the corresponding in the weak basis. With the help of the seesaw formula, m_{iD} and M_{iN^c} involved in Eq. (5.8a) can be related to the light-neutrino mass matrix m_ν . Working in the N_i^c -basis, we have

$$m_\nu = -m_D d_{N^c}^{-1} m_D^\dagger \quad \text{where} \quad d_{N^c} = \text{diag}(M_{1N^c}, M_{2N^c}, M_{3N^c}) \quad (5.9)$$

with $M_{1N^c} \leq M_{2N^c} \leq M_{3N^c}$ real and positive. Based on the analysis of Ref. [24, 70], we find \bar{m}_ν via

$$\bar{m}_\nu = U_\nu^* d_\nu U_\nu^\dagger \quad \text{where} \quad d_\nu = \text{diag}(m_{1\nu}, m_{2\nu}, m_{3\nu}) \quad (5.10)$$

with $m_{1\nu}$, $m_{2\nu}$ and $m_{3\nu}$ being the real and positive light neutrino mass eigenvalues. These can be found assuming *normal* [*inverted*] *ordered* (NO [IO]) $m_{i\nu}$'s and using a reference neutrino mass and the observed [41, 42] low energy neutrino mass-squared differences. Also U_ν is the PMNS matrix which is a function of the mixing angles θ_{ij} and the CP-violating Majorana (φ_1 and φ_2) and Dirac (δ) phases. Taking also m_{iD} as input parameters we can construct the complex symmetric matrix

$$W = -d_D^{-1} \bar{m}_\nu d_D^{-1} \quad (5.11)$$

from which we can extract d_{N^c} as follows [24, 70]:

$$d_{N^c}^{-2} = U^{c\dagger} W W^\dagger U^c. \quad (5.12)$$

Acting this way – see Sec. 5.5 –, we can determine the elements of U^c and the M_{iN^c} 's, compute $m_D^\dagger m_D$ through Eq. (5.8c) and finally obtain the ε_i 's via Eq. (5.8a).

5.4 POST-INFLATIONARY REQUIREMENTS

The success of our post-inflationary scenario can be judged, if, in addition to the constraints of Sec. 3.2, it is consistent with the following requirements:

5.4.1 The bounds on M_{1N^c} :

$$(a) \ M_{1N^c} \gtrsim 10T_{\text{rh}} \quad \text{and} \quad (b) \ \widehat{m}_{\delta\phi} \geq 2M_{1N^c}. \quad (5.13)$$

The first inequality is applied to avoid any erasure of the produced Y_L due to ν_1^c mediated inverse decays and $\Delta L = 1$ scatterings [70]. The second bound ensures that the decay of $\widehat{\delta\phi}$ into a pair of N_i^c 's is kinematically allowed for at least one species of the N_i^c 's.

5.4.2 Constraints from neutrino physics. We take as inputs the best-fit values [41] – see also Ref. [42] – on the neutrino mass-squared differences, $\Delta m_{21}^2 = 7.62 \cdot 10^{-3} \text{ eV}^2$ and $\Delta m_{31}^2 = (2.55 [-2.43]) \cdot 10^{-3} \text{ eV}^2$, on the mixing angles, $\sin^2 \theta_{12} = 0.32$, $\sin^2 \theta_{13} = 0.0246 [0.025]$, and $\sin^2 \theta_{23} = 0.613 [0.6]$ and the Dirac phase $\delta = 0.8\pi [-0.03\pi]$ for NO [IO] $m_{i\nu}$'s. Moreover, the sum of $m_{i\nu}$'s is bounded from above by the current data [1, 34], as follows

$$\sum_i m_{i\nu} \leq 0.28 \text{ eV at } 95\% \text{ c.l.} \quad (5.14)$$

5.4.3 The observational results on Y_B [1, 34]

$$Y_B \simeq (8.55 \pm 0.217) \cdot 10^{-11} \text{ at } 95\% \text{ c.l.} \quad (5.15)$$

5.4.4 The bounds on $Y_{3/2}$ imposed [40] by successful BBN:

$$Y_{3/2} \lesssim \begin{cases} 10^{-14} \\ 10^{-13} \\ 10^{-12} \end{cases} \quad \text{for } m_{3/2} \simeq \begin{cases} 0.69 \text{ TeV}, \\ 10.6 \text{ TeV}, \\ 13.5 \text{ TeV}. \end{cases} \quad (5.16)$$

Here we consider the conservative case where \tilde{G} decays with a tiny hadronic branching ratio.

5.5 NUMERICAL RESULTS

As shown in Sec. 5.1, nSMI predicts a constant value of $\hat{m}_{\delta\phi}$. Consequently, T_{rh} and Y_B – see Eqs. (5.5) and (5.6) – are largely independent of the precise value of c_T and λ in the range of Eq. (3.30a) – contrary to the case of FHI [26, 62]. Just for definiteness we specify that throughout this section we take $c_T = 150$ which corresponds to $\lambda = 0.0034$, $n_s = 0.963$ and $\hat{m}_{\delta\phi} = 3 \cdot 10^{13}$ GeV. On the other hand, T_{rh} and Y_B depend on λ_μ , y and the masses of the N_i^c 's into which $\hat{\delta\phi}$ decays. Throughout our computation we take $y = 0.5$, which is a typical value encountered [53] into various MSSM settings with large $\tan\beta$, and so the corresponding decay width via Eq. (5.4b) is confined to $\hat{\Gamma}_{\delta\phi \rightarrow XYZ} = 0.45$ GeV. Note that varying y in its plausible [53] range (0.4 – 0.6), $\hat{\Gamma}_{\delta\phi \rightarrow XYZ}$ ranges from 0.28 to 0.64 GeV causing minor changes to our results.

Following the bottom-up approach described in Sec. 5.3, we find the M_{iN^c} 's by using as inputs the m_{iD} 's, a reference mass of the ν_i 's – $m_{1\nu}$ for NO $m_{i\nu}$'s, or $m_{3\nu}$ for IO $m_{i\nu}$'s –, the two Majorana phases φ_1 and φ_2 of the PMNS matrix, and the best-fit values, mentioned in Sec. 5.4, for the low energy parameters of neutrino physics. In our numerical code, we also estimate, following Ref. [69], the RG evolved values of the latter parameters at the scale of nTL, $\Lambda_L = \hat{m}_{\delta\phi}$, by considering the MSSM with $\tan\beta \simeq 50$ as an effective theory between Λ_L and the SSB scale, $M_{\text{SUSY}} = 1.5$ TeV. We evaluate the M_{iN^c} 's at Λ_L , and we neglect any possible running of the m_{iD} 's and M_{iN^c} 's. Therefore, we present their values at Λ_L .

Fixing λ_μ at an intermediate value in its allowed region – see Eq. (3.12) – $\lambda_\mu = 10^{-6}$ which results, via Eq. (5.3b) in $\hat{\Gamma}_{\delta\phi \rightarrow H} = 1.3$ GeV we can get a first picture for the parameters which yield Y_B and $Y_{3/2}$ compatible with Eqs. (5.15) and (5.16), respectively in Table 4. We consider strongly NO (cases A and B), almost degenerate (cases C, D and E) and strongly IO (cases F and G) $m_{i\nu}$'s. In all cases the current limit of Eq. (5.14) is safely met – in the case D this limit is almost saturated. We observe that with NO or IO $m_{i\nu}$'s, the resulting M_{iN^c} 's are also hierarchical. With degenerate $m_{i\nu}$'s, the resulting M_{iN^c} 's are closer to one another. Consequently, in the latter case more $\hat{\delta\phi}$ -decay channels are available, whereas for the case A only a single decay channel is open. In all other cases – even in the case C where the decay channel $\hat{\delta\phi} \rightarrow N_3^c N_3^c$ is kinematically permitted –, the dominant contributions to Y_B arise from ε_2 . Therefore, the branching ratios, which are also presented in Table 4, $\hat{\Gamma}_{\delta\phi \rightarrow N_i^c} / \hat{\Gamma}_{\delta\phi}$ with $i = 1$ for the case A and $i = 2$ for the other cases are crucial for the calculation Y_B from Eq. (5.6). We notice that these ratios introduce a considerable reduction in the derivation of Y_B , given that $\hat{\Gamma}_{\delta\phi \rightarrow N_i^c} < \hat{\Gamma}_{\delta\phi \rightarrow XYZ} < \hat{\Gamma}_{\delta\phi \rightarrow H}$. This reduction can be eluded if we adopt – as in Ref. [22, 26, 62] – the resolution of the μ problem proposed in Ref. [61] since then the decay mode in Eq. (5.3a) disappears. In Table 4 shown also are the values of T_{rh} , the majority of which are close to $6 \cdot 10^8$ GeV, and the corresponding $Y_{3/2}$'s, which are consistent with Eq. (5.16) mostly for $m_{3/2} \gtrsim 11$ TeV. These large values are in nice agreement with the ones needed for the solution of the μ problem of MSSM, as explained in Sec. 4.

Since we do not consider any particular GUT here, the m_{iD} 's are free parameters. For the sake of comparison, however, we mention that the simplest realization of a SUSY Left-Right [Pati-Salam] GUT predicts [62, 71] $h_{iN} = h_{iE}$ [$m_{iD} = m_{iU}$], where m_{iU} are the masses of the up-type quarks and we ignore any possible mixing between generations. Taking into account the SUSY threshold

PARAMETERS	CASES						
	A	B	C	D	E	F	G
	NORMAL HIERARCHY		ALMOST DEGENERACY			INVERTED HIERARCHY	
LOW SCALE PARAMETERS							
$m_{1\nu}/0.1\text{ eV}$	0.01	0.1	0.5	0.7	0.7	0.5	0.49
$m_{2\nu}/0.1\text{ eV}$	0.09	0.1	0.51	0.7	0.7	0.51	0.5
$m_{3\nu}/0.1\text{ eV}$	0.5	0.5	0.71	0.86	0.5	0.1	0.05
$\sum_i m_{i\nu}/0.1\text{ eV}$	0.6	0.7	1.7	2.3	1.9	1.1	1
φ_1	0	$2\pi/3$	$\pi/2$	$\pi/2$	0	$-3\pi/4$	$\pi/4$
φ_2	$\pi/2$	$\pi/2$	$\pi/3$	$2\pi/3$	$-2\pi/3$	$5\pi/4$	$-2\pi/3$
LEPTOGENESIS-SCALE PARAMETERS							
$m_{1D}/0.1\text{ GeV}$	16	15	9	20	7	20	5
m_{2D}/GeV	40	8.3	10.5	10.3	7.5	5.3	11.8
$m_{3D}/10\text{ GeV}$	10	10	3.56	10	10	10	4
$M_{1N^c}/10^{11}\text{ GeV}$	12.3	2.2	0.16	0.58	0.11	0.7	0.12
$M_{2N^c}/10^{12}\text{ GeV}$	22.2	1.8	1.8	1.75	1	1.6	2.2
$M_{3N^c}/10^{14}\text{ GeV}$	25	4	0.15	0.73	0.74	2.7	1.2
OPEN DECAY CHANNELS OF THE INFLATON, $\widehat{\delta\phi}$, INTO N_i^c							
$\widehat{\delta\phi} \rightarrow$	N_1^c	$N_{1,2}^c$	$N_{1,2,3}^c$	$N_{1,2}^c$	$N_{1,2}^c$	$N_{1,2}^c$	$N_{1,2}^c$
$\widehat{\Gamma}_{\delta\phi\rightarrow N_i^c}/\widehat{\Gamma}_{\delta\phi}\text{ (\%)} $	3	7	7	6.5	2.3	5	9.8
RESULTING B -YIELD							
$10^{11}Y_B$	8.54	8.7	8.7	8.5	8.4	8.4	8.5
RESULTING T_{rh} AND \widetilde{G} -YIELD							
$T_{\text{rh}}/10^8\text{ GeV}$	5.9	5.9	6.3	5.9	5.8	5.9	6.1
$10^{13}Y_{3/2}$	1.1	1.1	1.2	1.13	1.11	1.1	1.15

Table 4: Parameters yielding the correct BAU for various neutrino mass schemes, $k_{SH} = 1.5$, $\lambda_\mu = 10^{-6}$ and $y = 0.5$. Shown also are the branching ratios of the $\widehat{\delta\phi}$ decay into N_i^c with $i = 2$ except for the case A where $i = 1$. Recall that these results are independent of the variables λ, c_T, k_S and $k_{S\widetilde{N}^c}$.

corrections [53] in the context of MSSM with universal gaugino masses and $\tan \beta \simeq 50$ – favored by the recent LHC results [46] – these predictions are translated as follows:

$$(m_{1D}^0, m_{2D}^0, m_{3D}^0) \simeq \begin{cases} (0.023, 4.9, 100) \text{ GeV} & \text{for a Left-Right GUT,} \\ (0.0005, 0.24, 100) \text{ GeV} & \text{for a Pati-Salam GUT.} \end{cases} \quad (5.17)$$

Comparing these values with those listed in Table 4, we remark that our model is not compatible with any GUT-inspired pattern of large hierarchy between the m_{iD} 's, especially in the two lighter




Figure 3: Contours in the $\lambda_\mu - m_{2D}$ plane yielding the central Y_B in Eq. (5.15) consistently with the inflationary requirements for $k_{SH} = 1.5$, $y = 0.5$ and the values of $m_{i\nu}$, m_{1D} , m_{3D} , φ_1 , and φ_2 which correspond to the cases B (solid line), D (dashed line), and F (dot-dashed line) of Table 4.

generations, since $m_{1D} \gg m_{1D}^0$ and $m_{2D} > m_{2D}^0$. On the other hand, in the cases A, B, D, E and F we are able to place $m_{3D} \simeq m_{3D}^0$. This arrangement can be understood if we take into account that m_{1D} and m_{2D} separately influences the derivation of M_{1N^c} and M_{2N^c} respectively – see, e.g., Ref. [22, 70]. Consequently, the displayed $m_{2D} \sim 10$ GeV assists us to obtain the ε_2 's required by Eq. (5.15) – note that in the case A $m_{2D} \simeq 40$ GeV kinematically blocks the channel $\widehat{\delta\phi} \rightarrow N_2^c N_2^c$. On the other hand, $m_{1D} \gtrsim 0.5$ GeV is necessitated in order to obtain the observationally favored ε_1 in the case A and fulfill Eq. (5.13a) in the other cases. Note that the phases φ_1 and φ_2 in Table 4 are selected in each case, so that the required m_{iD} and M_{iN^c} , which dominate the Y_B calculation, and the resulting T_{rh} are almost minimized.

In order to extend the conclusions inferred from Table 4 to the case of a variable λ_μ , we can examine how the central value of Y_B in Eq. (5.15) can be achieved by varying m_{2D} as a function of λ_μ . The resulting contours in the $\kappa - m_{2D}$ plane are presented in Fig. 3 – since the range of Y_B in Eq. (5.15) is very narrow, the 95% c.l. width of these contours is negligible. The convention adopted for these lines is also described in the figure. In particular, we use solid, dashed, or dot-dashed line for $m_{i\nu}$, m_{1D} , m_{3D} , φ_1 , and φ_2 corresponding to the cases B, D, or F of Table 4 respectively. Since increasing λ_μ , the resulting T_{rh} is expected to get larger than that shown in Table 4 – see Eqs. (5.3b) and (5.5) – we select for the plot in Fig. 3 one case from every low-energy mass scheme of $m_{i\nu}$'s with M_{1N^c} large enough, such that Eq. (5.13a) is comfortably satisfied for every λ_μ within the range of Eq. (3.12) with $k_{SH} = 1.5$. This equation sets, actually, the limits on the contours depicted in Fig. 3. For $\lambda_\mu \gtrsim 6 \cdot 10^{-7}$ we get $\widehat{\Gamma}_{\delta\phi \rightarrow H} > \widehat{\Gamma}_{\delta\phi \rightarrow XYZ}$ and so, increasing λ_μ the branching fraction in Eq. (5.6b) drops and larger m_{2D} 's are required to obtain Y_B compatible with Eq. (5.15). On the other hand, for $\lambda_\mu \lesssim 6 \cdot 10^{-7}$, $\widehat{\Gamma}_{\delta\phi \rightarrow XYZ}$ gets larger than $\widehat{\Gamma}_{\delta\phi \rightarrow H}$ and so, the branching fraction in Eq. (5.6b) remains almost constant and no sizable variation of m_{2D} is required. At the upper termination points of the contours, we obtain $T_{rh} \simeq 5 \cdot 10^9$ GeV or $Y_{\widetilde{G}} \simeq 9.4 \cdot 10^{-13}$. The constraint of Eq. (5.16), therefore, will cut any possible extension of the curves would be available for possible larger λ_μ 's. Along the depicted contours, the resulting M_{2N^c} 's vary in the range $(1.4 - 4) \cdot 10^{12}$ GeV whereas M_{1N^c} and M_{3N^c} remain close to their values presented in the corresponding cases of Table 4.

In conclusion, nTL is a realistic possibility within our model, thanks to the spontaneously arising couplings in SUGRA, even without direct couplings of the inflaton to N_i^c 's in W .

6 CONCLUSIONS

We investigated a variant of the Starobinsky inflation, which can be embedded in a moderate extension of MSSM supplemented by three N_i^c 's and two more superfields, the inflaton and an accompanied field. Key role in our proposal plays a continuous R symmetry, which is reduced to the well-known R -parity of MSSM, a \mathbb{Z}_2 discrete symmetry and a no-scale-type symmetry imposed on the Kähler manifold. The adopted symmetries have a number of ensuing consequences: (i) The inflaton appears quadratically in the super- and Kähler potentials; (ii) it couples to N_i^c via SUGRA-induced interactions ensuring low T_{rh} and no important contributions to the one-loop radiative corrections; (iii) the μ problem of MSSM can be elegantly resolved provided that a related parameter in superpotential is somehow suppressed. The last issue can be naturally incorporated in various schemes of SSB with relatively large – of the order $(10^4 - 10^6)$ GeV – $m_{3/2}$'s which facilitate the explanation of the recently observed mass of the electroweak Higgs and the satisfaction of the \tilde{G} constraint.

The next important modification of our set-up compared to other incarnations – cf. Ref. [5–7] – of the Starobinsky inflation in SUGRA is the introduction of a variable scale (M) – besides the existing one in Ref. [5–7] – in the superpotential and a parameter (c_T) in the Kähler potential which was ultimately confined in the range $76 \leq c_T \leq 1.5 \cdot 10^5$. One of these parameters (M and c_T) can be eliminated demanding that the gravitational strength takes its conventional value at the SUSY vacuum of the theory. Actually our inflationary model interpolates between the Starobinsky [3] and the induced-gravity [28, 29] inflation. Variation of the free model parameters (λ and c_T) gives us the necessary flexibility in order to obtain inflation for subplanckian values of the inflaton. Consequently, our proposal is stable against possible corrections from higher order terms in the super- and/or Kähler potentials. Moreover, we showed that the one-loop radiative corrections remain subdominant during inflation and the corresponding effective theory is trustable up to m_P .

Despite the addition of the extra parameters, our scheme remains very predictive since all the possible sets (λ, c_T) which are compatible with the two inflationary requirements, concerning the number of the e-foldings and the normalization of the curvature perturbation, yield almost constant values of r and n_s and a unique inflaton mass, $\hat{m}_{\delta\phi}$. In particular, we find $n_s \simeq 0.963$, $a_s \simeq -0.00068$ and $r \simeq 0.0038$, which are in excellent agreement with the current data, and $\hat{m}_{\delta\phi} = 3 \cdot 10^{13}$ GeV. Moreover, the post-inflationary evolution within our model remains intact from the variation of the inflationary parameters (λ and c_T). Implementing the (type I) seesaw mechanism for the generation of the light neutrino masses, we restricted their Dirac masses, m_{iD} , and the masses of N_i^c 's, M_{iN^c} , fulfilling a number of requirements, which originate from the BAU, the (unstable) \tilde{G} abundance and the neutrino oscillation parameters. Namely, we found $m_{1D} \geq 0.5$ GeV and $m_{2D} \simeq 10$ GeV resulting mostly to $M_{1N^c} \simeq 10^{11}$ GeV and $M_{2N^c} \simeq 10^{12}$ GeV.

As a bottom line, we would like to emphasize that the Starobinsky-type inflation in no-scale SUGRA can be linked to the phenomenology of MSSM, even if it is not realized by a matter-like inflaton as in Ref. [14]. In our framework, this type of inflation, driven by a modulus-like field, suggests a resolution of the μ problem of MSSM, compatible with large values of $m_{3/2}$ and it is followed by a robust cosmological scenario – already applied in many inflationary settings [22, 24, 26, 36, 37, 62] – ensuring spontaneous nTL reconcilable with the \tilde{G} constraint and the neutrino oscillation parameters.

ACKNOWLEDGEMENTS

The author would like to cordially thank G. Lazarides, A. Mazumdar, Q. Shafi and O. Vives for helpful discussions. This research was supported by the Generalitat Valenciana under contract PROMETEOII/2013/017.

REFERENCES

- [1] G. Hinshaw *et al.* [WMAP Collaboration], [arXiv:1212.5226](#).
- [2] P.A.R. Ade *et al.* [Planck Collaboration], [arXiv:1303.5082](#).
- [3] A.A. Starobinsky, *Phys. Lett. B* **91**, 99 (1980);
V.F. Mukhanov and G.V. Chibisov, *JETP Lett.* **33**, 532 (1981);
A.A. Starobinsky, *Sov. Astron. Lett.* **9**, 302 (1983).
- [4] J. Ellis, D.V. Nanopoulos and K. A. Olive, *Phys. Rev. Lett.* **111**, 111301 (2013) [[arXiv:1305.1247](#)].
- [5] J. Ellis, D.V. Nanopoulos and K.A. Olive, *J. Cosmol. Astropart. Phys.* **10**, 009 (2013) [[arXiv:1307.3537](#)].
- [6] R. Kallosh and A. Linde, *J. Cosmol. Astropart. Phys.* **06**, 028 (2013) [[arXiv:1306.3214](#)].
- [7] D. Roest, M. Scalisi and I. Zavala, *J. Cosmol. Astropart. Phys.* **11**, 007 (2013) [[arXiv:1307.4343](#)].
- [8] F. Farakos, A. Kehagias and A. Riotto, *Nucl. Phys.* **B876** (2013) 187 [[arXiv:1307.1137](#)].
- [9] A. Kehagias, A.M. Dizgah and A. Riotto, *Phys. Rev. D* **89**, 043527 (2014) [[arXiv:1312.1155](#)].
- [10] W. Buchmüller, V. Domcke and K. Kamada, *Phys. Lett. B* **726**, 467 (2013) [[arXiv:1306.3471](#)];
W. Buchmüller, V. Domcke and C. Wieck, [arXiv:1309.3122](#).
- [11] S.V. Ketov and A.A. Starobinsky, *Phys. Rev. D* **83**, 063512 (2011) [[arXiv:1011.0240](#)];
S. V. Ketov and N. Watanabe, *J. Cosmol. Astropart. Phys.* **03**, 011 (2011) [[arXiv:1101.0450](#)];
S.V. Ketov and A.A. Starobinsky, *J. Cosmol. Astropart. Phys.* **08**, 022 (2012) [[arXiv:1203.0805](#)];
S.V. Ketov and S. Tsujikawa, *Phys. Rev. D* **86**, 023529 (2012) [[arXiv:1205.2918](#)];
F. Brisce, L. Modesto and S. Tsujikawa, [arXiv:1308.1413](#);
E.J. Copeland, C. Rahmede and I.D. Saltas, [arXiv:1311.0881](#);
L. Sebastiani, G. Cognola, R. Myrzakulov, S.D. Odintsov and S. Zerbini, [arXiv:1311.0744](#).
- [12] S. Tsujikawa, J. Ohashi, S. Kuroyanagi and A. De Felice, *Phys. Rev. D* **88**, 023529 (2013) [[arXiv:1305.3044](#)].
- [13] D.S. Gorbunov and A.G. Panin, *Phys. Lett. B* **700**, 157 (2011) [[arXiv:1009.2448](#)];
D. Gorbunov and A. Tokareva, [arXiv:1212.4466](#).
- [14] J. Ellis, D.V. Nanopoulos and K.A. Olive, *Phys. Rev. D* **89**, 043502 (2014) [[arXiv:1310.4770](#)];
J. Ellis, A. Mustafayev and K.A. Olive, *Eur. Phys. J. C* **69**, 219 (2010) [[arXiv:1004.5399](#)].
- [15] E. Cremmer, S. Ferrara, C. Kounnas and D.V. Nanopoulos, *Phys. Lett. B* **133**, 61 (1983);
J.R. Ellis, C. Kounnas and D.V. Nanopoulos, *Nucl. Phys. B* **247**, 373 (1984).
- [16] For a review, see A.B. Lahanas and D.V. Nanopoulos, *Phys. Rept.* **145**, 198 (1987).
- [17] K. Nakayama, F. Takahashi and T.T. Yanagida, *J. Cosmol. Astropart. Phys.* **08**, 038 (2013) [[arXiv:1305.5099](#)];
J. Ellis and N.E. Mavromatos, *Phys. Rev. D* **88**, 085029 (2013) [[arXiv:1308.1906](#)];
K. Nakayama, F. Takahashi and T.T. Yanagida, [arXiv:1311.4253](#).
- [18] M.B. Einhorn and D.R.T. Jones, *J. High Energy Phys.* **03**, 026 (2010) [[arXiv:0912.2718](#)];
H.M. Lee, *J. Cosmol. Astropart. Phys.* **08**, 003 (2010) [[arXiv:1005.2735](#)];
S. Ferrara *et al.*, *Phys. Rev. D* **83**, 025008 (2011) [[arXiv:1008.2942](#)].
- [19] R. Kallosh and A. Linde, *J. Cosmol. Astropart. Phys.* **11**, 011 (2010) [[arXiv:1008.3375](#)];
R. Kallosh, A. Linde and D. Roest, [arXiv:1310.3950](#).
- [20] C. Pallis, *Phys. Lett. B* **692**, 287 (2010) [[arXiv:1002.4765](#)].
- [21] C. Pallis and N. Toumbas, *J. Cosmol. Astropart. Phys.* **02**, 019 (2011) [[arXiv:1101.0325](#)].
- [22] C. Pallis and N. Toumbas, *J. Cosmol. Astropart. Phys.* **12**, 002 (2011) [[arXiv:1108.1771](#)];
C. Pallis and N. Toumbas, “Open Questions in Cosmology” (InTech, 2012) [[arXiv:1207.3730](#)].
- [23] M. Arai, S. Kawai and N. Okada, *Phys. Rev. D* **84**, 123515 (2011) [[arXiv:1107.4767](#)];
M.B. Einhorn and D.R.T. Jones, *J. Cosmol. Astropart. Phys.* **11**, 049 (2012) [[arXiv:1207.1710](#)].
- [24] C. Pallis and Q. Shafi, *Phys. Rev. D* **86**, 023523 (2012) [[arXiv:1204.0252](#)].

- [25] C. Pallis, PoS **CORFU2012**, 061 (2013) [arXiv:1307.7815].
- [26] G.R. Dvali, Q. Shafi and R.K. Schaefer, *Phys. Rev. Lett.* **73**, 1886 (1994) [hep-ph/9406319];
C. Pallis and Q. Shafi, *Phys. Lett. B* **725**, 327 (2013) [arXiv:1304.5202].
- [27] G.R. Dvali, G. Lazarides and Q. Shafi, *Phys. Lett. B* **424**, 259 (1998) [hep-ph/9710314].
- [28] A. Zee, *Phys. Rev. Lett.* **42**, 417 (1979);
F.S. Accetta, D. J. Zoller, and M. S. Turner, *Phys. Rev. D* **31**, 3046 (1985);
D. S. Salopek, J. R. Bond and J.M. Bardeen, *Phys. Rev. D* **40**, 1753 (1989);
R. Fakir and W. G. Unruh, *Phys. Rev. D* **41**, 1792 (1990).
- [29] J.L. Cervantes-Cota and H. Dehnen, *Phys. Rev. D* **51**, 395 (1995) [astro-ph/9412032];
N. Kaloper, L. Sorbo and J. Yokoyama, *Phys. Rev. D* **78**, 043527 (2008) [arXiv:0803.3809];
A. Cerioni, F. Finelli, A. Tronconi and G. Venturi, *Phys. Rev. D* **81**, 123505 (2010) [arXiv:1005.0935].
- [30] S. Cecotti, *Phys. Lett. B* **190**, 86 (1987);
S. Ferrara, R. Kallosh and A. Van Proeyen, arXiv:1309.4052.
- [31] C.P. Burgess, H.M. Lee and M. Trott, *J. High Energy Phys.* **09**, 103 (2009) [arXiv:0902.4465];
J.L.F. Barbon and J.R. Espinosa, *Phys. Rev. D* **79**, 081302 (2009) [arXiv:0903.0355];
C.P. Burgess, H.M. Lee and M. Trott, *J. High Energy Phys.* **07**, 007 (2010) [arXiv:1002.2730];
M.P. Hertzberg, *J. High Energy Phys.* **11**, 023 (2010) [arXiv:1002.2995].
- [32] R.N. Lerner and J. McDonald, *J. Cosmol. Astropart. Phys.* **04**, 015 (2010) [arXiv:0912.5463];
F. Bezrukov *et al.*, *J. High Energy Phys.* **016**, 01 (2011) [arXiv:1008.5157];
X. Calmet and R. Casadio, arXiv:1310.7410.
- [33] G.F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, *J. High Energy Phys.* **12**, 027 (1998) [hep-ph/9810442];
E. Katz, Y. Shadmi and Y. Shirman, *J. High Energy Phys.* **08**, 015 (1999) [hep-ph/9906296];
N. Setzer and S. Spinner, *J. High Energy Phys.* **06**, 137 (2011) [arXiv:1008.3774];
Q. Shafi and Z. Tavartkiladze, hep-ph/0408156.
- [34] P.A.R. Ade *et al.* [Planck Collaboration], arXiv:1303.5076.
- [35] M. Endo, F. Takahashi and T.T. Yanagida, *Phys. Rev. D* **76**, 083509 (2007) [arXiv:0706.0986].
- [36] M. Endo, F. Takahashi and T.T. Yanagida, *Phys. Rev. D* **74**, 123523 (2006) [hep-ph/0611055].
- [37] G. Lazarides and Q. Shafi, *Phys. Lett. B* **258**, 305 (1991);
K. Kumeke, T. Moroi and T. Yanagida, *Prog. Theor. Phys.* **92**, 437 (1994) [hep-ph/9405337];
G. Lazarides, R.K. Schaefer and Q. Shafi, *Phys. Rev. D* **56**, 1324 (1997) [hep-ph/9608256].
- [38] M.Yu. Khlopov and A.D. Linde, *Phys. Lett. B* **138**, 265 (1984);
J. Ellis, J.E. Kim, and D.V. Nanopoulos, *Phys. Lett. B* **145**, 181 (1984).
- [39] M. Bolz, A. Brandenburg and W. Buchmüller, *Nucl. Phys.* **B606**, 518 (2001);
M. Bolz, A. Brandenburg and W. Buchmüller, *Nucl. Phys.* **B790**, 336 (2008) (E) [hep-ph/0012052];
J. Pradler and F.D. Steffen, *Phys. Rev. D* **75**, 023509 (2007) [hep-ph/0608344].
- [40] R.H. Cyburt *et al.*, *Phys. Rev. D* **67**, 103521 (2003) [astro-ph/0211258];
M.Kawasaki, K.Kohri and T.Moroi, *Phys. Lett. B* **625**, 7 (2005) [astro-ph/0402490];
M. Kawasaki, K. Kohri and T. Moroi, *Phys. Rev. D* **71**, 083502 (2005) [astro-ph/0408426];
J.R. Ellis, K.A. Olive and E. Vangioni, *Phys. Lett. B* **619**, 30 (2005) [astro-ph/0503023].
- [41] D.V. Forero, M. Tortola and J.W.F. Valle, *Phys. Rev. D* **86**, 073012 (2012) [arXiv:1205.4018].
- [42] G.L. Fogli *et al.*, *Phys. Rev. D* **86**, 013012 (2012) [arXiv:1205.5254].
- [43] M. Hindmarsh and D.R.T. Jones, *Phys. Rev. D* **87**, 075022 (2013) [arXiv:1203.6838];
A. Arbey, A. Deandrea, F. Mahmoudi and A. Tarhini, *Phys. Rev. D* **87**, 115020 (2013) [arXiv:1304.0381];
- [44] J.L. Feng, K.T. Matchev and T. Moroi, *Phys. Rev. Lett.* **84**, 2322 (2000) [hep-ph/9908309];
J.L. Feng and D. Sanford, *Phys. Rev. D* **86**, 055015 (2012) [arXiv:1205.2372];
V.E. Mayes, *Int. J. Mod. Phys. A* **28**, 1350061 (2013) [arXiv:1302.4394].

- [45] K. Harigaya, M. Ibe and T.T. Yanagida, [arXiv:1310.0643](#);
J.L. Evans, M. Ibe, K.A. Olive and T.T. Yanagida, *Eur. Phys. J. C* **73**, 2468 (2013) [[arXiv:1302.5346](#)];
J.L. Evans, K.A. Olive, M. Ibe and T.T. Yanagida, *Eur. Phys. J. C* **73**, 2611 (2013) [[arXiv:1305.7461](#)].
- [46] G. Aad *et al.* [ATLAS Collaboration], *Phys. Lett. B* **716**, 1 (2012) [[arXiv:1207.7214](#)];
S. Chatrchyan *et al.* [CMS Collaboration], *Phys. Lett. B* **716**, 30 (2012) [[arXiv:1207.7235](#)].
- [47] P. Athron *et al.* [GAMBIT Collaboration], [arXiv:1705.07935](#).
- [48] G. Lazarides, C. Panagiotakopoulos, and Q. Shafi, *Phys. Rev. Lett.* **56**, 432 (1986).
- [49] S.P. Martin, *Adv. Ser. Direct. High Energy Phys.* **21**, 1 (2010) [[hep-ph/9709356](#)].
- [50] G. Lazarides and C. Pallis, *J. High Energy Phys.* **11**, 114 (2015) [[arXiv:1508.06682](#)].
- [51] C. Pallis, *J. Cosmol. Astropart. Phys.* **10**, no. 10, 037 (2016) [[arXiv:1606.09607](#)].
- [52] S. Tsujikawa and B. Gumjudpai, *Phys. Rev. D* **69**, 123523 (2004) [[astro-ph/0402185](#)];
A. De Felice *et al.*, *J. Cosmol. Astropart. Phys.* **08**, 021 (2011) [[arXiv:1105.4685](#)];
K. Kamada *et al.*, *Phys. Rev. D* **86**, 023504 (2012) [[arXiv:1203.4059](#)].
- [53] S. Antusch and M. Spinrath, *Phys. Rev. D* **78**, 075020 (2008) [[arXiv:0804.0717](#)].
- [54] S.R. Coleman and E.J. Weinberg, *Phys. Rev. D* **7**, 1888 (1973).
- [55] V.N. Şenoğuz and Q. Shafi, *Phys. Lett. B* **668**, 6 (2008) [[arXiv:0806.2798](#)].
- [56] G. Lazarides, *Lect. Notes Phys.* **592**, 351 (2002) [[hep-ph/0111328](#)];
G. Lazarides, *J. Phys. Conf. Ser.* **53**, 528 (2006) [[hep-ph/0607032](#)].
- [57] D.H. Lyth and A. Riotto, *Phys. Rept.* **314**, 1 (1999) [[hep-ph/9807278](#)];
A. Mazumdar and J. Rocher, *Phys. Rept.* **497**, 85 (2011) [[arXiv:1001.0993](#)].
- [58] L. Randall and R. Sundrum, *Nucl. Phys.* **B557**, 79 (1999) [[hep-th/9810155](#)];
J.A. Bagger, T. Moroi, and E. Poppitz, *J. High Energy Phys.* **04**, 009 (2000) [[hep-th/9911029](#)];
P. Binetruy, M.K. Gaillard and B.D. Nelson, *Nucl. Phys.* **B604**, 32 (2001) [[hep-ph/0011081](#)].
- [59] H.P. Nilles, *Phys. Rept.* **110**, 1 (1984).
- [60] M. Dine, R. Kitano, A. Morisse and Y. Shirman, *Phys. Rev. D* **73**, 123518 (2006) [[hep-ph/0604140](#)];
R. Kitano, *Phys. Lett. B* **641**, 203 (2006) [[hep-ph/0607090](#)];
J.L. Evans, M.A.G. Garcia and K.A. Olive, [arXiv:1311.0052](#).
- [61] G. Lazarides and Q. Shafi, *Phys. Rev. D* **58**, 071702 (1998) [[hep-ph/9803397](#)].
- [62] R. Armillius, G. Lazarides and C. Pallis, *Phys. Rev. D* **89**, 065032 (2014) [[arXiv:1309.6986](#)].
- [63] C. Pallis, *Nucl. Phys.* **B751**, 129 (2006) [[hep-ph/0510234](#)].
- [64] K. Hamaguchi, *Phd Thesis* [[hep-ph/0212305](#)];
W. Buchmuller, R.D. Peccei and T. Yanagida, *Ann. Rev. Nucl. Part. Sci.* **55**, 311 (2005) [[hep-ph/0502169](#)].
- [65] M. Dine, R. Kitano, A. Morisse and Y. Shirman, *Phys. Rev. D* **73**, 123518 (2006) [[hep-ph/0604140](#)];
K. Nakayama, F. Takahashi and T.T. Yanagida, *Phys. Lett. B* **718**, 526 (2012) [[arXiv:1209.2583](#)].
- [66] A.D. Linde, *Phys. Rev. D* **53**, 4129 (1996) [[hep-th/9601083](#)];
K. Nakayama, F. Takahashi and T.T. Yanagida, *Phys. Rev. D* **84**, 123523 (2011) [[arXiv:1109.2073](#)];
K. Nakayama, F. Takahashi and T.T. Yanagida, *Phys. Lett. B* **714**, 256 (2012) [[arXiv:1203.2085](#)].
- [67] M. Endo *et al.*, *J. Cosmol. Astropart. Phys.* **02**, 018 (2007) [[hep-ph/0612263](#)];
S. Nakamura and M. Yamaguchi *Phys. Lett. B* **655**, 167 (2007) [[arXiv:0707.4538](#)].
- [68] M. Flanz, E.A. Paschos and U. Sarkar, *Phys. Lett. B* **345**, 248 (1995); (*E*) *ibid.* **382**, 447 (1996) [[hep-ph/9411366](#)];
M. Flanz, E.A. Paschos, U. Sarkar and J. Weiss, *Phys. Lett. B* **389**, 693 (1996) [[hep-ph/9607310](#)];
L. Covi, E. Roulet and F. Vissani, *Phys. Lett. B* **384**, 169 (1996) [[hep-ph/9605319](#)].
- [69] S. Antusch, J. Kersten, M. Lindner and M. Ratz, *Nucl. Phys.* **B674**, 401 (2003) [[hep-ph/0305273](#)].
- [70] V.N. Şenoğuz, *Phys. Rev. D* **76**, 013005 (2007) [[arXiv:0704.3048](#)].
- [71] N. Karagiannakis, G. Lazarides and C. Pallis, *Int. J. Mod. Phys. A* **28**, 1330048 (2013) [[arXiv:1305.2574](#)].