

December 17, 2013

Status of hadronic light-by-light scattering in the muon $g - 2$ *

Andreas Nyffeler[†]

*Regional Centre for Accelerator-based Particle Physics, Harish-Chandra Research
Institute, Chhatnag Road, Jhusi, Allahabad - 211 019, India*

*Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics,
University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland*

Abstract

We give an update on the status of the hadronic light-by-light scattering contribution to the muon $g - 2$. We review recent work by various groups, list some of the open problems and give an outlook on how to better control the uncertainty of this contribution. This is necessary in order to fully profit from planned future muon $g - 2$ experiments to test the Standard Model. Despite some recent developments, we think that the estimate $a_{\mu}^{\text{HLbL}} = (116 \pm 40) \times 10^{-11}$ still gives a fair description of the current situation.

PACS numbers: 14.60.Ef, 13.40.Em, 12.38.Lg

*Based on presentations at the *International Workshop on e^+e^- collisions from Φ to Ψ 2013 (PHIPSI13)*, September 9-12, 2013, Rome, Italy, the *14th Meeting of the Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies*, September 13, 2013, Frascati, Italy, and the *LC13 Workshop: Exploring QCD from the infrared regime to heavy flavour scales at B-factories, the LHC and a Linear Collider*, September 16-20, 2013, Trento, Italy. Some material added compared to the text submitted to the proceedings of the LC13 Workshop.

[†]nyffeler@hri.res.in

1 Introduction

The anomalous magnetic moment of the muon has served over many years as an important test of the Standard Model, see the reviews [1, 2]. It is also sensitive to potential contributions from New Physics. The current status of the muon $g - 2$ is summarized in Table 1 where we list the different contributions in theory (QED, weak, hadronic) from various recent sources and compare with the experimental value. More references to earlier work can be found in the quoted papers and in Refs. [1, 2]. The experimental world average is dominated by the final result of the Brookhaven muon $g - 2$ experiment [7], corrected for a small shift in the ratio of the magnetic moments of the muon and the proton [8]. We observe a difference between experiment and theory of more than three standard deviations

$$a_\mu^{\text{exp}} - a_\mu^{\text{th}} = (293 \pm 88) \times 10^{-11} \quad (3.3\sigma). \quad (1)$$

Unfortunately, the theoretical uncertainties [1, 2, 9] from hadronic vacuum polarization (HVP) and hadronic light-by-light scattering (HLbL) make it difficult to interpret this discrepancy as a clear sign of New Physics. Most recent evaluations [1, 5, 10, 2, 9], which differ slightly in the treatment of the hadronic contributions, obtain deviations of $3 - 4\sigma$. In Ref. [11] the hadronic cross-section data below 1 GeV was fitted to a (broken) Hidden Local Symmetry (HLS) model and a discrepancy in the muon $g - 2$ between $3.7 - 4.9\sigma$ was observed, depending on the selected data.

Table 1: Standard Model contributions to $a_\mu \times 10^{11}$ and comparison of theory and experiment.

Contribution	Value	Error	Reference
QED	116 584 718.853	0.036	[3]
Weak	153.6	1.0	[4]
Leading order HVP	6 907.5	47.2	[5]
Higher order HVP	-100.3	2.2	[5]
HLbL	116	40	[6, 1]
Theory (total)	116 591 796	62	-
Experiment	116 592 089	63	[7]
Experiment - Theory (3.3σ)	293	88	-

The HLbL contribution to the muon $g - 2$ involves the Green function of four electromagnetic currents, connected to off-shell photons, see Figure 1 and Ref. [1] for details and references. The relevant scales for the off-shell photons in HLbL are about $0 - 2$ GeV, i.e. larger than the muon mass, and therefore a pure low-energy effective field theory approach with muons, photons and pions fails [12]. In contrast to the HVP contribution, HLbL cannot be directly related to experimental data and therefore various models have




Figure 1: The different contributions to HLbL scattering and their chiral and large- N_c counting.

been employed to estimate HLbL. One uses some hadronic model with exchanges and loops of resonances at low energies and some form of (dressed, constituent) quark-loop at high energies as short-distance complement of the low-energy hadronic models. The dependence on several momenta leads, however, to a mixing of long and short distances and makes it difficult to avoid a double counting of quark-gluon and hadronic contributions. In Ref. [13] a classification of the different contributions to HLbL based on the chiral and large- N_c counting was proposed, see Figure 1. In general, all the interactions of the hadrons or the quarks with the photons are dressed by some form factors, e.g. via $\rho - \gamma$ mixing. Note that in the Feynman diagrams in Fig. 1 form factors with off-shell photons and off-shell hadrons enter [14]. Constraints on the models can be obtained from experimental data, e.g. on the various form factors, and from theory, e.g. chiral perturbation theory at low energies and short-distance constraints from perturbative QCD and the Operator Product Expansion (OPE) at high momenta.

2 Current status of HLbL and recent developments

A selection of estimates for HLbL is presented in Table 2. Note that the Refs. [15, 16] are the only full calculations of HLbL to date, using, as much as possible, one model for all the contributions (HLS model in Ref. [15], Extended Nambu-Jona-Lasinio (ENJL) model in Ref. [16]). Both calculations showed that the exchanges of the lightest pseudoscalar states, π^0, η, η' , dominate numerically, which can be understood from the large- N_c counting. The contributions from the (dressed) pion-loop and the (dressed) quark-loop are subdominant, but not negligible, and they happen to largely cancel each other numerically. The final results for the total HLbL contribution were rather close in both models. In Ref. [27] an ansatz for the pion-photon transition form factor with a minimal number of narrow vector resonances in large- N_c QCD (lowest meson dominance (LMD, LMD+V)) was matched to short-distance constraints from the OPE. The reevaluation of the pion-

Table 2: Summary of selected estimates for the different contributions to $a_\mu^{\text{HLbL}} \times 10^{11}$. For comparison, the last line shows some results when no form factors are used (undressed).

π, K -loops	π^0, η, η'	axial-vect.	scalars	quark-loop	Total	Ref.
-4.5(8.1)	82.7(6.4)	1.7(1.7)	-	9.7(11.1)	89.6(15.4)	[15]
-19(13)	85(13)	2.5(1.0)	-6.8(2.0)	21(3)	83(32)	[16]
-	83(12)	-	-	-	80(40)	[17]
0(10)	114(10)	22(5)	-	0	136(25)	[18]
-	-	-	-	-	110(40)	[19]
-19(19)	114(13)	15(10)	-7(7)	2.3 [c-quark]	105(26)	[20]
-19(13)	99(16)	22(5)	-7(2)	21(3)	116(40)	[6, 1]
-	81(2)	-	-	107(2)	188(4)	[21]
-	-	-	-	-	118-148	[22]
-	68(3) [π^0]	-	-	82(6)	150(3)	[23]
-	-	-	-	-	76(4)-125(7)	[24]
-(11 - 71)	-	-	-	-	-	[25]
-20(5)	-	-	-	-	-	[26]
-45	$+\infty$	-	-	60	-	no FF

pole contribution to HLbL in Ref. [17] with the ansatz from Ref. [27] then revealed a sign error in the earlier calculations [15, 16]. Furthermore, for the neutral pion contribution, a 2-dimensional integral representation was derived in Ref. [17] for a certain class of vector meson dominance (VMD)-like form factors. In the end, one integrates over the lengths of the two independent loop-momenta $|Q_1|, |Q_2|$ in Euclidean space. Schematically,

$$a_\mu^{\text{HLbL};\pi^0} = \int_0^\infty dQ_1 \int_0^\infty dQ_2 \sum_i w_i(Q_1, Q_2) f_i(Q_1, Q_2), \quad (2)$$

with universal weight functions w_i and where the (model) dependence on the form factors resides in the functions f_i , see Ref. [17] for details. Note that the expressions in Ref. [17] with on-shell form factors are in general not valid as they stand. As pointed out in Ref. [18], one needs to set the form factor at the external vertex to a constant to obtain the pion-*pole* contribution. But the expressions are valid for the constant Wess-Zumino-Witten (WZW) form factor and the off-shell VMD form factor. The weight functions w_i are plotted in Figure 2. One can see that the relevant momentum regions are about 0 – 1.25 GeV, see also the numerical analysis in Ref. [28]. As long as the form factors in different models lead to a damping, we expect comparable results for $a_\mu^{\text{HLbL};\pi^0}$ at the level of 20%, which is indeed the case, see the papers quoted in Table 2 and Ref. [29]. For the general case with off-shell form factors, a 3-dimensional integral representation was derived in Ref. [1], where the integrations run over Q_1, Q_2 and $\cos \theta$, where $Q_1 \cdot Q_2 = |Q_1||Q_2| \cos \theta$.

Later, Ref. [18] derived new short-distance constraints from the four-point function on the pion-pole and axial-vector-pole contributions, which do not allow for any form




Figure 2: The different weight functions w_i in Eq. (2) from Ref. [17].

factors at the external vertex. Reference [18] also included the mixing of two axial-vector nonets and studied the pion-loop within the HLS model in more detail. All this lead to a substantial enhancement of these contributions to HLbL. More recently, a new short-distance constraint on the off-shell form factor at the external vertex in pion-exchange was derived in Ref. [6], which yielded, again in the framework of the lowest meson dominance approximation to large- N_c QCD, a value for this contribution about half-way between the results in Refs. [15, 16, 17] and those in Ref. [18]. Note that the compilations [19, 20, 1] and Ref. [6] are largely based on the full calculations [15, 16], with revised or newly calculated values for some of the contributions. More recent estimates, mostly for the pseudoscalar contribution, can be found in Ref. [29]. While most of these evaluations agree at the level of 15%, if one takes the extreme values, there is a spread of $a_\mu^{\text{HLbL;PS}} = (59 - 107) \times 10^{-11}$.

Until 2010, a consensus had been reached about the central value $a_\mu^{\text{HLbL}} \approx 110 \times 10^{-11}$, but there was a discussion on how to estimate the error, more progressively, $\pm 26 \times 10^{-11}$, in Ref. [20] and more conservatively, $\pm 40 \times 10^{-11}$, in Refs. [6, 1]. In view of the precision goal of future $g - 2$ experiments at Fermilab and J-PARC [30] with $\delta a_\mu = 16 \times 10^{-11}$ and the continued progress in improving the error in HVP, the HLbL contribution might soon be the main uncertainty in the theory prediction, if it cannot be brought under better control [1, 2, 9].

In the last few years, several works have appeared which yield much larger (absolute)

values for some of the contributions, see Table 2. In Ref. [21] the quark-loop was studied using a Dyson-Schwinger equation approach. In contrast to Refs. [15, 16], no damping compared to the bare constituent quark-loop result was seen, when a dressing was included. Note that this calculation of the quark-loop is not yet complete and that earlier, very large results for the quark-loop seem to have been affected by some errors in the numerics in certain parts. The large size of the quark-loop contribution in Ref. [21] was questioned in the papers [22, 23], using different quark-models and approaches, see also the ballpark prediction for HLbL in Ref. [24]. The pion-loop contribution was analyzed in Ref. [25]. The authors stressed the importance of the pion-polarizability effect and the role of the axial-vector resonance a_1 , which are not included in the models used in Refs. [15, 16]. Depending on the value of the pion-polarizability and the model for the a_1 resonance used, a large variation was seen. The issue was taken up in Ref. [26] where different models for the pion-loop were studied. The inclusion of the a_1 resonance was attempted, but no finite result for $g - 2$ could be achieved. With a cutoff of 1 GeV, a result close to the earlier estimate in Ref. [16] was obtained. Reference [26] also pointed out that the very small (absolute) value for the pion-loop in Ref. [15] could be due to the fact that the HLS model used in Ref. [15] has a wrong high-energy behavior and that there is some cancellation between positive and negative contributions in the pion-loop in HLbL.

3 Outlook

Concerning the future, maybe lattice QCD will provide a reliable calculation of HLbL at some point, see Ref. [31] for some promising recent results. In the meantime, only a close collaboration between theory and experiment can lead to a better controlled estimate for HLbL. On the theory side, the hadronic models can be improved by short-distance constraints from perturbative QCD to have a better matching at high momenta. One can also use dispersion relations to connect the theory with experimental data, e.g. in $\gamma\gamma \rightarrow \pi\pi$ [32]. Also the issue about whether the dressing of the bare constituent quark-loop leads to a suppression or an enhancement needs to be studied further. This problem is also related to the question whether there is any double counting involved.

On the experimental side, the information on various processes (decays, form factors, cross-sections) of hadrons interacting with photons at low and intermediate momenta, $|q| \leq 2$ GeV, can help to constrain the models. Important experiments which should be pursued include more precise measurements of the (transition) form factors of light pseudoscalars with possibly two off-shell photons in the process $e^+e^- \rightarrow e^+e^-P$ ($P = \pi^0, \eta, \eta'$) and the two-photon decay width and the (double) Dalitz decays of these mesons. This could further reduce the error of the dominant pseudoscalar exchange contribution [33]. Concerning the pion-loop contribution, in addition to studying $\gamma\gamma \rightarrow \pi\pi$, measurements of the pion-polarizability in various processes, e.g. in radiative pion decay $\pi^+ \rightarrow e^+\nu_e\gamma$, in radiative pion photoproduction $\gamma p \rightarrow \gamma'\pi^+n$ or with the hadronic Primakoff effect $\pi A \rightarrow \pi'\gamma A$ or $\gamma A \rightarrow \pi^+\pi^-A$ (with some nucleus A), can help to improve the mod-

els [25]. For the development of models with the axial-vector resonance a_1 and estimates of the sizable axial-vector contribution, information about the decays $a_1 \rightarrow \rho\pi, \pi\gamma$ would be useful as well. Finally, to extract the needed quantities from experiment will also require the development of dedicated Monte Carlo programs for the relevant processes [34].

4 Conclusions

If the recent results for the quark-loop and pion-loop are taken at face value, one obtains the range $a_\mu^{\text{HLbL}} = (64 - 202) \times 10^{-11}$. While the new approaches raise some important issues and point to potential shortcomings in the previously used models, these calculations are also still preliminary and further studies are needed. Therefore, the estimate

$$a_\mu^{\text{HLbL}} = (116 \pm 40) \times 10^{-11} \quad (3)$$

from Refs. [6, 1] still seems to give a fair description of the current situation.

Acknowledgments

I would like to thank the organizers of the *International Workshop on e^+e^- collisions from Φ to Ψ 2013 (PHIPSI13)*, of the *Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies* and of the *LC13 Workshop: Exploring QCD from the infrared regime to heavy flavour scales at B-factories, the LHC and a Linear Collider* for the opportunity to present reviews on the HLbL in the muon $g-2$. Generous support from the organizers, the INFN, the ECT*, Trento, Italy, and the Heinrich-Greinacher-Stiftung, Physics Institute, University of Bern, Switzerland, is gratefully acknowledged. The AEC is supported by the “Innovations- und Kooperationsprojekt C-13” of the “Schweizerische Universitätskonferenz SUK/CRUS.”

References

- [1] F. Jegerlehner, A. Nyffeler, Phys. Rept. **477**, 1 (2009).
- [2] J.P. Miller *et al.*, Ann. Rev. Nucl. Part. Sci. **62**, 237 (2012); M. Knecht, talk at PHIPSI13, Rome, September 2013.
- [3] T. Aoyama *et al.*, Phys. Rev. Lett. **109**, 111808 (2012).
- [4] C. Gnendiger, D. Stöckinger, H. Stöckinger-Kim, Phys. Rev. D **88**, 053005 (2013); H.G. Fargnoli *et al.*, arXiv:1311.4890.
- [5] F. Jegerlehner, R. Szafron, Eur. Phys. J. C **71**, 1632 (2011).
- [6] A. Nyffeler, Phys. Rev. D **79**, 073012 (2009).

- [7] G.W. Bennett *et al.* [Muon $g - 2$ Collaboration], Phys. Rev. D **73**, 072003 (2006).
- [8] P.J. Mohr, B.N. Taylor, D.B. Newell, Rev. Mod. Phys. **80**, 633 (2008).
- [9] G. Venanzoni, talk at LC13, Trento, September 2013; T. Blum *et al.*, arXiv:1311.2198.
- [10] M. Davier *et al.*, Eur. Phys. J. C **71**, 1515 (2011) [Erratum-ibid. C **72**, 1874 (2012)]; K. Hagiwara *et al.*, J. Phys. G **38**, 085003 (2011); T. Teubner, talks at PHIPSI13, Rome and at Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies, Frascati, September 2013.
- [11] M. Benayoun *et al.*, Eur. Phys. J. C **72**, 1848 (2012); **73**, 2453 (2013); M. Benayoun, talk at PHIPSI13, Rome, September 2013; F. Jegerlehner, Acta Phys. Polon. B **44**, 2257 (2013).
- [12] M. Knecht *et al.*, Phys. Rev. Lett. **88**, 071802 (2002).
- [13] E. de Rafael, Phys. Lett. B **322**, 239 (1994).
- [14] F. Jegerlehner, Acta Phys. Polon. B **38**, 3021 (2007).
- [15] M. Hayakawa, T. Kinoshita, A.I. Sanda, Phys. Rev. Lett. **75**, 790 (1995); Phys. Rev. D **54**, 3137 (1996); M. Hayakawa, T. Kinoshita, Phys. Rev. D **57**, 465 (1998) [Erratum-ibid. D **66**, 019902 (2002)].
- [16] J. Bijnens, E. Pallante, J. Prades, Phys. Rev. Lett. **75**, 1447 (1995) [Erratum-ibid. **75**, 3781 (1995)]; Nucl. Phys. B **474**, 379 (1996); **626**, 410 (2002).
- [17] M. Knecht, A. Nyffeler, Phys. Rev. D **65**, 073034 (2002).
- [18] K. Melnikov, A. Vainshtein, Phys. Rev. D **70**, 113006 (2004).
- [19] J. Bijnens, J. Prades, Mod. Phys. Lett. A **22**, 767 (2007); J.P. Miller, E. de Rafael, B.L. Roberts, Rep. Prog. Phys. **70**, 795 (2007).
- [20] J. Prades, E. de Rafael, A. Vainshtein, in *Advanced Series on Directions in High Energy Physics - Vol. 20 "Lepton Dipole Moments"*, edited by B.L Roberts, W.J. Marciano (World Scientific, Singapore, 2010), p.303; and arXiv:0901.0306.
- [21] T. Goecke, C.S. Fischer, R. Williams, Phys. Rev. D **83**, 094006 (2011) [Erratum-ibid. D **86**, 099901 (2012)]; Phys. Rev. D **87**, 034013 (2013).
- [22] R. Boughezal, K. Melnikov, Phys. Lett. B **704**, 193 (2011).
- [23] D. Greynat, E. de Rafael, JHEP **07** (2012) 020.
- [24] P. Masjuan, M. Vanderhaeghen, arXiv:1212.0357.

- [25] K.T. Engel, H.H. Patel, M.J. Ramsey-Musolf, Phys. Rev. D **86**, 037502 (2012); K.T. Engel, Ph.D. thesis, May 2013; K.T. Engel, M.J. Ramsey-Musolf, arXiv:1309.2225.
- [26] M. Zahiri Abyaneh, arXiv:1208.2554; J. Bijnens, M. Zahiri Abyaneh, EPJ Web Conf. **37**, 01007 (2012) [arXiv:1208.3548]; J. Bijnens, talk at MesonNet 2013 International Workshop, Prague, June 2013 [see: arXiv:1308.2575].
- [27] M. Knecht, A. Nyffeler, Eur. Phys. J. C **21**, 659 (2001).
- [28] A. Nyffeler, PoS (CD12) 045 (2013) [arXiv:1306.5987].
- [29] A.E. Dorokhov, W. Broniowski, Phys. Rev. D **78**, 073011 (2008); A.E. Dorokhov, A.E. Radzhabov, A.S. Zhevlakov, Eur. Phys. J. C **71**, 1702 (2011); **72**, 2227 (2012); A.E. Radzhabov, talk at PHIPSI13, Rome, September 2013; D.K. Hong, D. Kim, Phys. Lett. B **680**, 480 (2009); L. Cappiello, O. Cata, G. D'Ambrosio, Phys. Rev. D **83**, 093006 (2011); K. Kampf, J. Novotny, Phys. Rev. D **84**, 014036 (2011); P. Masjuan, Phys. Rev. D **86**, 094021 (2012); R. Escribano, P. Masjuan, P. Sanchez-Puertas, arXiv:1307.2061.
- [30] Talks by S. Maxfield and K. Ishida at PHIPSI13, Rome, September 2013.
- [31] T. Blum, M. Hayakawa, T. Izubuchi, PoS (Lattice 2012) 022 (2012) [arXiv:1301.2607] and references therein; updated at Lattice 2013 (private communication by T. Blum).
- [32] B. Moussallam, arXiv:1305.3143 and arXiv:1312.0395; M. Hoferichter *et al.*, arXiv:1309.6877.
- [33] D. Babusci *et al.*, Eur. Phys. J. C **72**, 1917 (2012).
- [34] S. Actis *et al.* [Working Group on Radiative Corrections and Monte Carlo Generators for Low Energies], Eur. Phys. J. C **66**, 585 (2010); H. Czyż *et al.*, arXiv:1312.0454.