PROCEEDINGS OF SCIENCE

arXiv:1312.5076v1 [hep-ph] 18 Dec 2013

# Natural Seesaw Realization at the TeV Scale

**Emiliano MOLINARO***
*Technische Universität München*
*James-Franck-Straße, 85748 Garching, Germany*
*E-mail:* emiliano.molinaro@tum.de

We summarize the phenomenological constraints on seesaw scenarios defined at the TeV scale and provide a simple extension of the Standard Model which naturally leads to a testable mechanism of neutrino mass generation.

*The European Physical Society Conference on High Energy Physics*
*18-24 July, 2013*
*Stockholm, Sweden*

*Speaker.

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike Licence. 

We review the most important phenomenological constraints on type I/inverse seesaw scenarios defined at the electroweak scale. We also consider a simple extension of the Standard Model (SM) which realizes a "testable" seesaw scenario, without imposing any fine-tuning of the neutrino Yukawa couplings in order to generate light active neutrino masses.

## 1. Phenomenological constraints on the TeV scale seesaw parameter space

We consider a phenomenological seesaw [1] extension of the SM with two heavy Majorana fermion singlets $\mathcal{N}_{1,2}$, which in principle can be tested in low energy and collider experiments for masses $M_{1,2} \sim (100 - 1000)$ GeV.

Following [2], $\mathcal{N}_{1,2}$ have charged-current (CC) and neutral-current (NC) interactions with the SM leptons and interact with the Higgs boson. They are given by

$$\mathscr{L}^{\mathcal{N}}_{CC} = -\frac{g}{2\sqrt{2}}\,\overline{\ell}\,\gamma_\alpha\,(RV)_{\ell k}(1-\gamma_5)\,\mathcal{N}_k W^\alpha + \text{h.c.}\,, \tag{1.1}$$

$$\mathscr{L}^{\mathcal{N}}_{NC} = -\frac{g}{4c_w}\,\overline{\nu_{\ell L}}\,\gamma_\alpha\,(RV)_{\ell k}(1-\gamma_5)\,\mathcal{N}_k Z^\alpha + \text{h.c.}\,, \tag{1.2}$$

$$\mathscr{L}^{\mathcal{N}}_{H} = -\frac{g M_k}{4M_W}\,\overline{\nu_{\ell L}}\,(RV)_{\ell k}(1+\gamma_5)\,\mathcal{N}_k h + \text{h.c.} \tag{1.3}$$

The couplings $(RV)_{\ell k}$ ($\ell = e,\mu,\tau$ and $k = 1,2$) arise from the mixing between heavy and light Majorana neutrinos and, therefore, are suppressed by the seesaw scale. They can be conveniently parametrized as follows [2]:

$$|(RV)_{\ell 1}|^2 = \frac{1}{2}\frac{y^2v^2}{M_1^2}\frac{m_3}{m_2+m_3}\left|U_{\ell 3} + i\sqrt{m_2/m_3}U_{\ell 2}\right|^2, \quad \text{for normal hierarchy}\,, \tag{1.4}$$

$$|(RV)_{\ell 1}|^2 = \frac{1}{2}\frac{y^2v^2}{M_1^2}\frac{m_2}{m_1+m_2}\left|U_{\ell 2} + i\sqrt{m_1/m_2}U_{\ell 1}\right|^2, \quad \text{for inverted hierarchy}\,, \tag{1.5}$$

$$(RV)_{\ell 2} = \pm i\,(RV)_{\ell 1}\sqrt{\frac{M_1}{M_2}}\,,\ \ell = e,\mu,\tau\,, \tag{1.6}$$

where $U$ denotes the PMNS neutrino mixing matrix and $v \simeq 174$ GeV. The relative mass splitting of the two heavy Majorana neutrinos must be very small, $|M_1 - M_2|/M_1 \ll 1$, due to the current upper limit on the effective Majorana mass probed in neutrinoless double beta decay experiments [2]. In this case, the flavour structure of the neutrino Yukawa couplings is fixed by the neutrino oscillation parameters [2, 3] and the two heavy Majorana neutrinos form a pseudo-Dirac fermion. The parameter $y$ in the expressions above represents the largest eigenvalue of the matrix of the neutrino Yukawa couplings: $y^2v^2 = 2M_1^2\,(|(RV)_{e1}|^2 + \left|(RV)_{\mu 1}\right|^2 + |(RV)_{\tau 1}|^2)$.

Electroweak precision data (EWPD) provide an upper bound of the size of the Yukawa coupling $y$ for a given seesaw scale, that is $y \leq 0.06\,(M_1/100\ \text{GeV})$ [2]. However, the most stringent constraint on $y$ comes from lepton flavour violating observables, in particular from the present upper limit on $\mu^+ \to e^+\gamma$ branching ratio reported by the MEG experiment [4]: $\text{BR}(\mu \to e\gamma) < 5.7 \times 10^{-13}$ at 90% confidence level. Indeed, in this case, taking the best fit values of the neutrino oscillation parameters [5], we get the bound: $y \lesssim 0.026$ for $M_1 = 100$ GeV.

We show in Fig. 1 all the relevant constraints on the effective couplings $(RV)_{\mu 1}$ and $(RV)_{e1}$ which come from the requirement of reproducing neutrino oscillation data, from the EWPD bounds





**Figure 1:** Correlation between $|(RV)_{e1}|$ and $|(RV)_{\mu 1}|$ for $M_1 = 100$ GeV in the case of normal (left panel) and inverted (right panel) light neutrino mass spectrum (see the text for details).

and the current upper limit on $\mu \to e\gamma$. The seesaw mass scale is fixed at benchmark value $M_1 =$ 100 GeV. From Fig. 1, it is manifest that the allowed ranges of the right-handed (RH) neutrino couplings $|(RV)_{\mu 1}|$ and $|(RV)_{e1}|$, in the case of normal (left panel) and inverted (right panel) light neutrino mass spectrum, are confined in a small strip of the overall representative plane. This corresponds to the scatter plot of the points which are consistent with the $3\sigma$ allowed ranges of the neutrino oscillation parameters [5]. The region of the parameter space which is allowed by the EWPD is marked with solid lines. The region allowed by the current bound on the $\mu \to e\gamma$ decay rate is indicated with a dashed line, while the dot-dashed line shows the exclusion limit for $\mathrm{BR}(\mu \to e\gamma) < 10^{-14}$. The scatter points correspond to different values of the maximum neutrino Yukawa coupling $y$: $y = 0.001$ (blue $\circ$), *ii*) $y = 0.01$ (green $+$), *iii*) $y = 0.1$ (red $\times$), *iv*) $y = 1$ (orange $\Diamond$) and *v*) an arbitrary value of the Yukawa coupling $y \leq 1$ (cyan points).

As depicted in Fig. 1, in the case of a light neutrino mass spectrum with inverted hierarchy a strong suppression of the $\mu \to e\gamma$ decay rate is possible for specific values of the measured neutrino oscillation parameters. This is due to cancellations in the $\mu - e$ transition amplitude proportional to $|U_{\mu 2} + iU_{\mu 1}|$ which are possible in the case of neutrino mass spectrum with inverted hierarchy for $\sin\theta_{13} \gtrsim 0.13$ and CP conserving phases of the neutrino mixing matrix (see [2] for a detailed discussion).

## 2. A natural model realization of the TeV scale seesaw scenario

We discuss a simple model that naturally provides a testable seesaw scenario where the RH neutrino interactions in (1.1-1.3) are "sizable" and give rise to observable effects. We extend the scalar sector of the SM with an additional Higgs doublet, $H_2$, and a complex singlet, $\phi$. The fermion sector, instead, consists of the two RH neutrinos $N_{1,2}$, which allow to implement the seesaw mechanism in order to explain active neutrino masses and mixing. The resulting model Lagrangian is invariant under a global $U(1)_X$ symmetry, where $X$ corresponds to a generalization of the lepton

| Field | $L_\alpha$ | $e_{R\alpha}$ | $N_1$ | $N_2$ | $H_1$ | $H_2$ | $\phi$ |
|---|---|---|---|---|---|---|---|
| $\mathrm{SU(2)_L}$ | **2** | **1** | **1** | **1** | **2** | **2** | **1** |
| $\mathrm{U(1)_Y}$ | -1/2 | -1 | 0 | 0 | 1/2 | 1/2 | 0 |
| $\mathrm{U(1)_X}$ | -1 | -1 | -1 | +1 | 0 | 2 | -2 |

**Table 1:** Charge assignment of the fields.

number. The particle content of the model and the charge assignment of the fields are reported in Table 1.

In this scenario, the presence of $H_2$ and $\phi$ is motivated by the requirement of generating light neutrino masses through the type I/inverse/linear seesaw mechanism at the TeV scale. The most general scalar potential $\mathscr{V}_{\rm SB}$, invariant under the $\mathrm{SU(2)_L}\times\mathrm{U(1)_Y}\times[\mathrm{U(1)_X}]$ symmetry, is derived in [6]. Given the fields in Table 1, we get

$$\begin{aligned}\mathscr{V}_{\rm SB} &= -\mu_1^2\,(H_1^\dagger H_1)+\lambda_1\,(H_1^\dagger H_1)^2-\mu_2^2\,(H_2^\dagger H_2)+\lambda_2\,(H_2^\dagger H_2)^2-\mu_3^2\,\phi^*\phi+\lambda_3\,(\phi^*\phi)^2\\ &+\ \kappa_{12}\,(H_1^\dagger H_1)\,(H_2^\dagger H_2)+\kappa'_{12}\,(H_1^\dagger H_2)\,(H_2^\dagger H_1)+\kappa_{13}\,(H_1^\dagger H_1)\phi^*\phi+\kappa_{23}\,(H_2^\dagger H_2)\phi^*\phi\\ &-\ \frac{\mu'}{\sqrt{2}}\left((H_1^\dagger H_2)\phi+(H_2^\dagger H_1)\phi^*\right). \end{aligned} \tag{2.1}$$

The two $\mathrm{SU(2)_L}$ doublets $H_{1,2}$ and the singlet $\phi$ [6] take a non-zero vacuum expectation value (vev) $v_{1,2}$ and $v_\phi$, respectively. In this case, the global $\mathrm{U(1)_X}$ is spontaneously broken down to a $Z_2$ discrete symmetry. The scalar mass spectrum of the model consists of: 1 charged scalar $H^\pm$, 3 CP-even neutral scalars $h^0$, $H^0$ and $h_A$, 2 pseudo-scalars $A^0$ and $J$. The latter is the Goldstone boson associated with the breaking of the global $\mathrm{U(1)_X}$ symmetry and is usually dubbed Majoron in theories with spontaneously broken lepton charge. Since it is a massless particle, strong constraints apply on its couplings to the SM fermions: a hierarchical pattern for the vevs of the scalar fields, namely $v_2 \ll v_1, v_\phi$, is required in order to satisfy the astrophysical constraints on the Majoron phenomenology. As discussed in detail in [6], a suppressed value of $v_2 \propto \mu'$ is naturally realized from the minimization of the potential (2.1), due to the residual symmetries of the model.

In the limit of negligible $v_2$, the longitudinal gauge boson components are $W_L^\pm \sim H_1^\pm$ and $Z_L \sim \sqrt{2}\,\mathrm{Im}\,(H_1^0)$, while the scalar mass eigenstates are to a good approximation: $H^\pm \sim H_2^\pm$, $h_A \sim \sqrt{2}\,\mathrm{Re}\,(H_2^0)$, $A_0 \sim \sqrt{2}\,\mathrm{Im}\,(H_2^0)$ and $J \sim \sqrt{2}\,\mathrm{Im}\,(\phi)$. Moreover, the two neutral scalars $h^0$ and $H^0$ arise from the mixing of $\sqrt{2}\,\mathrm{Re}\,(H_1^0)$ and $\sqrt{2}\,\mathrm{Re}\,(\phi)$. Typically, we have $v_2 \lesssim 10$ MeV [6], $v_1 \simeq 246.2$ GeV, while $v_\phi$ is free. Recalling that only $H_1$ has Yukawa couplings to SM fermions (cf. Table 1), $h_A$, $A^0$ and $H^\pm$ couple to the SM sector only through gauge interactions and via the scalar quartic couplings, while $h^0$ and $H^0$ can have *a priori* sizable Yukawa couplings to SM fermions (see [6] for a discussion of the collider constraints on the scalar sector of the model).

**Neutrino mass generation**

We introduce for convenience a Dirac fermion field, $N_D \equiv P_R N_1 + P_L N_2^C$, where $P_{L,R}$ are the usual chiral projectors and $N_2^C \equiv C\overline{N_2}^T$ is the conjugate of the $N_2$ RH neutrino field. The most

general interaction Lagrangian of $N_D$ invariant under the global $U(1)_X$ symmetry is

$$\mathscr{L} \supset -m_N \overline{N_D} N_D - \left( Y_{\nu 1}^{\beta} \overline{N_D} \widetilde{H}_1^* L_\beta + Y_{\nu 2}^{\gamma} \overline{N_D}^C \widetilde{H}_2^* L_\gamma + \frac{\delta_N}{\sqrt{2}} \phi \overline{N_D} N_D^C + \text{h.c.} \right) \tag{2.2}$$

where $N_D^C \equiv C\overline{N}_D^T$ and $\widetilde{H}_k \equiv -i\sigma_2 H_k^*$ $(k = 1,2)$. The parameter $\delta_N$ is made real through a phase transformation. The Yukawa interactions $\propto Y_{\nu 1}$ ($Y_{\nu 2}$) couple $N_1$ ($N_2$) to the SM leptons. Therefore, after the Higgs doublets acquire a nonzero vev, the SM lepton number (i.e. the generalized $X$ charge) is explicitly violated by $Y_{\nu 2}$ mediated interactions. Furthermore, while the Dirac type mass $m_N$ conserves the lepton number, the term proportional to $\delta_N$ provides, after $\phi$ takes a nonzero vev, a Majorana mass term for both $N_1$ and $N_2$. In the case $m_N \gg \delta_N v_\phi$ we have a low energy realization of the type I/inverse seesaw scenario [6].

In the chiral basis $\left(\nu_{\mathbf{L}}\,, (N_1^C)_{\mathbf{L}}, (N_2^C)_{\mathbf{L}}\right)$, the $5\times 5$ symmetric neutrino mass matrix reads:

$$\mathscr{M}_\nu = \begin{pmatrix} \mathbf{0_{3\times 3}} & \mathbf{y_1} v_1 & \mathbf{y_2} v_2 \\ \mathbf{y_1}^{\mathrm{T}} v_1 & \delta_N v_\phi & m_N \\ \mathbf{y_2}^{\mathrm{T}} v_2 & m_N & \delta_N v_\phi \end{pmatrix} . \tag{2.3}$$

In the previous expression $\mathbf{0_{3\times 3}}$ denotes the null matrix of dimension 3 and we introduce the shorthand notation: $\mathbf{y_k} \equiv \left(Y_{\nu k}^e\,, Y_{\nu k}^\mu\,, Y_{\nu k}^\tau\right)^{\mathrm{T}}/\sqrt{2}$. The neutrino sector, therefore, consists of one massless neutrino, two massive light Majorana neutrinos and two heavy Majorana neutrinos $\mathscr{N}_{1,2}$. The latter are quasidegenerate, with masses $M_{1,2} = m_N \mp \delta_N v_\phi$, and form a pseudo-Dirac pair for $m_N \gg \delta_N v_\phi$ [6]. They have, therefore, naturally "sizable" CC and NC interactions with the SM leptons, eqs. (1.1) and (1.2), where in this case the mixing matrix elements $(RV)_{\ell 1}$ are proportional to the lepton number conserving Yukawa couplings $Y_{\nu k}^\ell$. The resulting effective light neutrino mass matrix is

$$(M_\nu)^{\alpha\beta} \simeq -\frac{v_1 v_2}{m_N} \left( \mathbf{y_1}^\alpha \mathbf{y_2}^\beta + \mathbf{y_2}^\alpha \mathbf{y_1}^\beta \right) + v_\phi\, \delta_N \frac{v_1^2}{m_N^2} \left( \mathbf{y_1}^\alpha \mathbf{y_1}^\beta + \mathbf{y_2}^\alpha \mathbf{y_2}^\beta \frac{v_2^2}{v_1^2} \right) . \tag{2.4}$$

The first term in (2.4) acts as a linear seesaw contribution and its suppression originates from the small vev $v_2$. The second term is typical of inverse seesaw scenarios, where the small ratio $v_\phi\, \delta_N / m_N$ suppresses the neutrino mass scale. Notice that, with only two RH neutrinos the linear seesaw contribution alone (i.e. neglecting $v_\phi$ in (2.4)) allows to fit all current neutrino oscillation data, while if $v_2 = 0$ and $v_\phi \neq 0$ the inverse seesaw scenario can only account for one massive light neutrino. Therefore, the complex scalar field $\phi$, with vev $v_\phi \neq 0$, is not mandatory in order to obtain two massive light neutrinos through the (linear) seesaw mechanism. On the other hand, $v_\phi \neq 0$ is a necessary condition to set a hierarchy between the Higgs doublet vevs, $v_2 \ll v_1$, without fine-tuning of the parameters [6]. Taking $\mu_N \ll 1$, the neutrino masses have a simple expression:

$$\begin{aligned} m_\nu^\pm &\simeq \frac{1}{m_N} \left( \sqrt{y_1^2 y_2^2 - \mu_N \left(y_1^2 + y_2^2\right) \mathrm{Re}(y_{12})} \pm \sqrt{|y_{12}|^2 - \mu_N \left(y_1^2 + y_2^2\right) \mathrm{Re}(y_{12})} \right) \\ &\simeq \frac{1}{m_N} \left( y_1 y_2 \pm |y_{12}| \right) \times \left( 1 \mp \frac{\mu_N}{2} \frac{\left(y_1^2 + y_2^2\right) \mathrm{Re}(y_{12})}{y_1 y_2\, |y_{12}|} \right) , \end{aligned} \tag{2.5}$$

with $y_i \equiv \sqrt{\mathbf{y_i}^\dagger \cdot \mathbf{y_i}}\, v_i$, $y_{12} \equiv \mathbf{y_1}^\dagger \cdot \mathbf{y_2}\, v_1 v_2$, $\eta_{12} \equiv \mathbf{y_1} \times \mathbf{y_2}\, v_1 v_2$ and $\mu_N = (\delta_N v_\phi)/m_N$. Notice that if the neutrino Yukawa vectors $\mathbf{y_1}$ and $\mathbf{y_2}$ are proportional, $m_\nu^-$ is exactly zero. For a normal hierarchical

spectrum, $m_\nu^+ = \sqrt{|\Delta m^2_{\rm A}|}$ and $m_\nu^- = \sqrt{\Delta m^2_\odot}$, while in the case of inverted hierarchy we have $m_\nu^+ = \sqrt{|\Delta m^2_{\rm A}|}$ and $m_\nu^- = \sqrt{|\Delta m^2_{\rm A}| - \Delta m^2_\odot}$, $|\Delta m^2_{\rm A}|$ and $\Delta m^2_\odot$ being the atmospheric and solar neutrino mass square differences, respectively. It is easy to show that at leading order in $\mu_N$, the neutrino mass parameters satisfy the relation [6]: $|\mathbf{y_1} \times \mathbf{y_2}|\, v_1 v_2/m_N \cong (\Delta m^2_\odot\, |\Delta m^2_{\rm A}|)^{1/4}$. Hence, barring accidental cancellations, the size of the neutrino Yukawa couplings is typically

$$|\mathbf{y_1}||\mathbf{y_2}| \approx 10^{-4}\,(m_N/1\ \text{TeV})\,(1\ \text{KeV}/v_2)\,. \tag{2.6}$$

Finally we point out that with the addition of a scalar field odd under $U(1)_X$ it is possible to realize a viable dark matter candidate in the model, which results naturally stable due to the presence of the remnant $Z_2$ symmetry. A variation of leptogenesis in this case is possible at the TeV scale [6].

This work was supported by the ERC Advanced Grant project "FLAVOUR"(267104).

## References

[1] P. Minkowski, Phys. Lett. B **67** (1977) 421; M. Gell-Mann, P. Ramond and R. Slansky, *Proceedings of the Supergravity Stony Brook Workshop*, New York 1979, eds. P. Van Nieuwenhuizen and D. Freedman; T. Yanagida, *Proceedings of the Workshop on Unified Theories and Baryon Number in the Universe*, Tsukuba, Japan 1979, eds. A. Sawada and A. Sugamoto; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980) 912.

[2] A. Ibarra, E. Molinaro and S. T. Petcov, *TeV Scale See-Saw Mechanisms of Neutrino Mass Generation, the Majorana Nature of the Heavy Singlet Neutrinos and $(\beta\beta)_{0\nu}$-Decay*, JHEP **1009** (2010) 108 [arXiv:1007.2378 [hep-ph]]. A. Ibarra, E. Molinaro and S. T. Petcov, *Low Energy Signatures of the TeV Scale See-Saw Mechanism*, Phys. Rev. D **84** (2011) 013005 [arXiv:1103.6217 [hep-ph]]. D. N. Dinh, A. Ibarra, E. Molinaro and S. T. Petcov, *The mu - e Conversion in Nuclei, mu $\to$ e $\gamma$, mu $\to$ 3e Decays and TeV Scale See-Saw Scenarios of Neutrino Mass Generation*, JHEP **1208** (2012) 125 [arXiv:1205.4671 [hep-ph]]. C. G. Cely, A. Ibarra, E. Molinaro and S. T. Petcov, *Higgs Decays in the Low Scale Type I See-Saw Model*, Phys. Lett. B **718** (2013) 957 [arXiv:1208.3654 [hep-ph]].

[3] M. Raidal, A. Strumia and K. Turzynski, *Low-scale standard supersymmetric leptogenesis*, Phys. Lett. B **609** (2005) 351 [Erratum-ibid. B **632** (2006) 752] [hep-ph/0408015]; M. Shaposhnikov, *A Possible symmetry of the nuMSM*, Nucl. Phys. B **763** (2007) 49 [hep-ph/0605047]; J. Kersten and A. Y. .Smirnov, *Right-Handed Neutrinos at CERN LHC and the Mechanism of Neutrino Mass Generation*, Phys. Rev. D **76** (2007) 073005 [arXiv:0705.3221 [hep-ph]]; M. B. Gavela, T. Hambye, D. Hernandez and P. Hernandez, *Minimal Flavour Seesaw Models*, JHEP **0909** (2009) 038 [arXiv:0906.1461 [hep-ph]].

[4] J. Adam *et al.* [MEG Collaboration], *New constraint on the existence of the mu+-> e+ gamma decay*, Phys. Rev. Lett. **110** (2013) 201801 [arXiv:1303.0754 [hep-ex]].

[5] D. V. Forero, M. Tortola and J. W. F. Valle, *Global status of neutrino oscillation parameters after Neutrino-2012*, Phys. Rev. D **86** (2012) 073012 [arXiv:1205.4018 [hep-ph]].

[6] F.X. Josse-Michaux and E. Molinaro, *A Common Framework for Dark Matter, Leptogenesis and Neutrino Masses*, Phys. Rev. D **84** (2011) 125021 [arXiv:1108.0482 [hep-ph]]. F. -X. Josse-Michaux and E. Molinaro, *Triplet scalar dark matter and leptogenesis in an inverse seesaw model of neutrino mass generation*, Phys. Rev. D **87** (2013) 3, 036007 [arXiv:1210.7202 [hep-ph]]