

Bosonic fractional quantum Hall states on the torus from conformal field theory

Anne E B Nielsen¹ and Germán Sierra²

¹ Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Straße 1, D-85748 Garching, Germany

² Instituto de Física Teórica, UAM-CSIC, Madrid, Spain

Abstract. The Kalmeyer-Laughlin state, which is a lattice version of the bosonic Laughlin state at filling factor one half, has attracted much attention due to its topological and chiral spin liquid properties. Here we show that the Kalmeyer-Laughlin state on the torus can be expressed in terms of a correlator of conformal fields from the $SU(2)_1$ Wess-Zumino-Witten model. This reveals an interesting underlying mathematical structure and provides a natural way to generalize the Kalmeyer-Laughlin state to arbitrary lattices on the torus. We find that the many-body Chern number of the states is unity for more different lattices, which suggests that the topological properties of the states are preserved when the lattice is changed. Finally, we analyze the symmetry properties of the states on square lattices.

1. Introduction

In [1], Moore and Read showed that a number of fractional quantum Hall (FQH) states can be expressed as certain correlators of conformal fields. More recently it has been shown that such a connection also holds for FQH states in lattice systems [2, 3] and for the interpolation between lattice and continuum systems [4]. The construction is interesting because it shows that the states can be seen as special cases of a more general mathematical framework. It also means that the states fulfil certain mathematical relations inherited from conformal field theory (CFT), and these are useful for deriving various properties of the states. Examples include the possibility to derive parent Hamiltonians, to study quasi particle or edge excitations, to investigate entanglement spectra and to derive correlation functions [1–8].

We shall here focus on a lattice version of the bosonic Laughlin state at filling factor $1/2$, which is known as the Kalmeyer-Laughlin (KL) state [9, 10]. On the Riemann sphere, i.e., the complex plane combined with the point at infinity, the KL state takes the form

$$\psi_{\text{KL}}(Z_1, Z_2, \dots, Z_M) = \prod_i G(Z_i) \prod_{i < j} (Z_i - Z_j)^2 \prod_i e^{-\frac{1}{4}|Z_i|^2} \quad (\text{plane}). \quad (1)$$

This wavefunction describes a state of M bosonic particles on a two-dimensional square or triangular lattice, and Z_i is the position of the i th particle written as a complex

number. The factor $\prod_i G(Z_i)$ is a gauge factor that is included to ensure that the state is a singlet, $G(Z_i) \in \{-1, +1\}$ for all i and the rest of the expression on the right hand side of (1) is the bosonic Laughlin state at filling factor $1/2$, except that the possible values of the positions Z_i are restricted to the considered lattice. The KL state has also been studied on a square lattice on the torus [11], and we provide an expression for this state in section 4. The KL state and closely related states have been analyzed in several papers, see e.g. [11–17]. These analyses show that the KL state has the same topological properties as the bosonic Laughlin state at half filling in the continuum. The investigated properties include, e.g., correlation functions, the fractional statistics of quasiparticle excitations, the chiral edge states, and topological entanglement entropy. Parent Hamiltonians have been obtained in [3, 17–21].

The huge interest in the KL state is due to its topological and chiral spin liquid properties combined with a relatively simple analytical wavefunction defined on a lattice. To characterize the properties of topological quantum states a number of different measures have been proposed, but more of these requires the state under investigation to be defined with a particular set of boundary conditions, e.g. periodic boundary conditions that correspond to defining the state on a torus. It is therefore very relevant to study different geometries.

In the present paper, we demonstrate that the KL states on the torus can be expressed as chiral correlators of fields from the $SU(2)_1$ Wess-Zumino-Witten (WZW) model. It has previously been shown [3, 4, 14] that this construction gives a KL-like state when considered on the Riemann sphere, and the present study extends this result to the torus geometry, where the relation to the KL state turns out to be exact. The CFT correlators can be defined for arbitrary lattices in 2D and are ensured to be singlets by construction. The CFT states thus provide a generalization of the KL states on the torus to arbitrary lattices, and by investigating the topological properties of the states on different lattices, we provide evidence that the topology remains the same at least for a broad class of lattices. For the case of square lattices with $L_x \times L_y$ spins, we also find linear combinations of the CFT states that are eigenstates of various symmetry operators. In particular, this allows us in certain cases to identify linear combinations of the states that are guaranteed to be orthogonal.

The paper is structured as follows. In section 2, we explain how wavefunctions can be constructed from conformal fields. We also introduce the CFT states that we study in this work and provide analytical expressions for them on the Riemann sphere and torus geometries. In section 3, we derive the analytical expressions for the CFT states on the torus. In section 4, we show that the CFT states are proportional to the KL states on the torus when defined on a square lattice. In section 5, we study the symmetry properties of the CFT states for the case of a square lattice. In particular, we find linear combinations of the two states that are eigenstates of different transformation operators and their eigenvalues. In section 6, we discuss the topological properties of the states, and we find among other things that the many-body Chern number of the states is unity for several different lattices. Section 7 concludes the paper, Appendix A summarizes the

definition and some properties of Riemann theta functions and Appendix B provides an alternative derivation of the singlet property of the states for four spins. To provide readers who are more interested in the concepts than the derivations with a faster route through the paper, we start sections 3, 4 and 5 with a summary of the results derived in the section.

2. Wavefunctions from conformal blocks

We first explain how states of spin systems can be constructed from conformal fields. Our starting point is the correlator

$$\langle \phi_{j_1}(z_1, \bar{z}_1) \phi_{j_2}(z_2, \bar{z}_2) \cdots \phi_{j_N}(z_N, \bar{z}_N) \rangle \quad (2)$$

of N primary fields $\phi_{j_i}(z_i, \bar{z}_i)$ of a CFT. Here, $\langle \dots \rangle$ stands for the vacuum expectation value, z_i is a coordinate in the complex plane, and \bar{z}_i is the complex conjugate of z_i . The field $\phi_{j_i}(z_i, \bar{z}_i)$ has spin j_i , and it therefore has $2j_i + 1$ components $\phi_{j_i, s_i}(z_i, \bar{z}_i)$, where s_i labels the $2j_i + 1$ possible values of the third component of the spin. It follows that (2) is a vector with $\prod_{i=1}^N (2j_i + 1)$ components.

The correlator (2) can be broken up into a sum over terms corresponding to different ways of fusing the fields present in the correlator to the vacuum state. Utilizing also that the field $\phi_{j_i}(z_i, \bar{z}_i)$ separates into a holomorphic and an antiholomorphic part, i.e., $\phi_{j_i}(z_i, \bar{z}_i) = \phi_{j_i}(z_i) \otimes \bar{\phi}_{j_i}(\bar{z}_i)$, we can write the components of (2) as

$$\langle \phi_{j_1, s_1}(z_1, \bar{z}_1) \cdots \phi_{j_N, s_N}(z_N, \bar{z}_N) \rangle = \sum_k f_k(s_j, z_j) \overline{f_k(s_j, z_j)}, \quad (3)$$

where

$$f_k(s_j, z_j) \equiv \langle \phi_{j_1, s_1}(z_1) \phi_{j_2, s_2}(z_2) \cdots \phi_{j_N, s_N}(z_N) \rangle_k \quad (4)$$

is a conformal block (also known as a chiral correlator), k labels the different ways of fusing the fields to the vacuum state, and the bar means complex conjugation.

The next step is to regard $f_k(s_j, z_j)$ as the components of a wavefunction $|\psi_k\rangle$ describing the state of N spins with spin quantum numbers j_1, j_2, \dots, j_N . Explicitly,

$$|\psi_k\rangle = \mathcal{C}_k \sum_{s_1, \dots, s_N} \psi_k(s_1, \dots, s_N) |s_1, \dots, s_N\rangle, \quad (5)$$

where

$$\psi_k(s_1, \dots, s_N) \propto f_k(s_j, z_j) \quad (6)$$

and \mathcal{C}_k is a normalization constant that allows us to choose the normalization of $\psi_k(s_1, \dots, s_N)$ after convenience. The complex coordinates $z_j = x_j + iy_j$ are fixed parameters of the wavefunction and are naturally interpreted as the physical positions (x_j, y_j) of the spins in the two-dimensional plane. In other words, the choice of z_j defines the lattice under consideration. Note that arbitrary lattices can be considered, since the only restriction on z_j is that $z_j \neq z_l$ for $j \neq l$.

In the present paper, we shall take the CFT to be the WZW model based on the Kac-Moody algebra $SU(2)_1$. This CFT has central charge $c = 1$ and two primary fields




Figure 1. Definition of the torus through the periods ω_1 and ω_2 . The points z and $z + n\omega_1 + m\omega_2$, where n and m are integers, are identified. The coordinate system is chosen such that ω_1 is real and positive and $\text{Im}(\omega_2) > 0$. The modular parameter τ is defined as $\tau = \omega_2/\omega_1$.

$\phi_0(z)$ and $\phi_{1/2}(z)$ with spin 0 and spin 1/2, respectively. $\phi_0(z)$ has conformal weight $h_0 = 0$, and $\phi_{1/2}(z)$ has conformal weight $h_{1/2} = 1/4$. Since the spin-0 primary field is the identity, we shall take all the fields in the correlator to be spin-1/2 primary fields. The number of conformal blocks is dictated by the fusion rules

$$\phi_0 \times \phi_0 = \phi_0, \quad \phi_0 \times \phi_{1/2} = \phi_{1/2}, \quad \phi_{1/2} \times \phi_{1/2} = \phi_0. \quad (7)$$

The fields can only fuse to the identity if N is even, and we shall therefore assume this to be the case throughout. The number of conformal blocks on a Riemann surface with genus g is then 2^g . There is hence one state on the Riemann sphere ($g = 0$) and two states on the torus ($g = 1$).

The state on the Riemann sphere (complex plane) can be written as [2, 3]

$$\psi_k(s_1, \dots, s_N) = \delta_{\mathbf{s}} \chi_{\mathbf{s}} \prod_{i < j} (z_i - z_j)^{(s_i s_j + 1)/2} \quad (\text{plane}), \quad (8)$$

where $k = 0$ can take only one value and $s_i \in \{-1, +1\}$ is defined such that the z -component of the spin of the i th field is $s_i/2$. The factor

$$\delta_{\mathbf{s}} = \begin{cases} 1 & \text{for } \sum_{i=1}^N s_i = 0 \\ 0 & \text{otherwise} \end{cases} \quad (9)$$

is the delta function factor, and

$$\chi_{\mathbf{s}} = \prod_{j=1}^N (-1)^{(j-1)(s_j+1)/2} \quad (10)$$

is the Marshall sign factor. This formula is the one used to consider lattices with open boundary conditions.

We shall here mostly be interested in periodic boundary conditions, which corresponds to defining the theory on the torus. The torus is defined by specifying two complex numbers ω_1 and ω_2 , which are called the periods of the torus, and then identifying all points in the complex plane that differ by integer multiples of these periods. The z_i in (4) are then restricted to lie within one parallelogram with sides ω_1

and ω_2 as illustrated in figure 1. ω_1 and ω_2 must be nonzero and have different phases, and we shall choose the coordinate system such that ω_1 is real and positive and ω_2 has positive imaginary part. It is convenient to define the modular parameter $\tau = \omega_2/\omega_1$ and use the scaled coordinates $\zeta_i = z_i/\omega_1$. In terms of these, the states (6) take the form

$$\psi_k(s_1, \dots, s_N) = \delta_{\mathbf{s}} \chi_{\mathbf{s}} \underbrace{\theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\sum_{i=1}^N \zeta_i s_i, 2\tau \right)}_{\text{Centre of mass factor}} \underbrace{\prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{(s_i s_j + 1)/2}}_{\text{Jastrow factor}} \quad (\text{torus}), \quad (11)$$

as we shall derive in the next section. Here, $k \in \{0, 1/2\}$ and $s_i \in \{-1, +1\}$. The definition and some properties of the Riemann theta function $\theta \begin{bmatrix} a \\ b \end{bmatrix}(\zeta, \tau)$ can be found in Appendix A, and

$$E(\zeta_i - \zeta_j, \tau) \equiv \frac{\theta_1(\zeta_i - \zeta_j, \tau)}{\partial_\zeta \theta_1(\zeta, \tau)|_{\zeta=0}}, \quad \theta_1(\zeta, \tau) \equiv \theta \begin{bmatrix} 1/2 \\ 1/2 \end{bmatrix}(\zeta, \tau), \quad (12)$$

is the prime form. In the following, we shall refer to the four factors on the right hand side of (11) as the delta function factor, the Marshall sign factor, the centre of mass factor, and the Jastrow factor, respectively.

3. Conformal blocks of the $SU(2)_1$ WZW model on the torus

In this section, we derive (11). We also demonstrate that if the numbering of the spins is altered, then (8) and (11) stay the same except that they are multiplied by the sign of the permutation needed to go from one numbering to the other. The choice of numbering is thus not important.

3.1. CFT correlators

We first need an expression for the correlator (2). To get this, we utilize the fact that the $SU(2)_1$ WZW model can be bosonized in terms of a massless free scalar field $\varphi(\zeta, \bar{\zeta}) = \varphi(\zeta) + \bar{\varphi}(\bar{\zeta})$ compactified on a circle of radius $R = \sqrt{2}$ [22]. Specifically, $\phi_{j_i, s_i}(\zeta_i, \bar{\zeta}_i) = \phi_{j_i, s_i}(\zeta_i) \otimes \bar{\phi}_{j_i, s_i}(\bar{\zeta}_i)$ can be expressed as the vertex operator

$$\phi_{j_i, s_i}(\zeta_i, \bar{\zeta}_i) = : e^{i s_i \varphi(\zeta_i, \bar{\zeta}_i)/\sqrt{2}} : = : e^{i s_i \varphi(\zeta_i)/\sqrt{2} + i s_i \bar{\varphi}(\bar{\zeta}_i)/\sqrt{2}} : , \quad (13)$$

where $: \dots :$ denotes normal ordering. We use here the scaled coordinates ζ_i , but we note that this changes the correlator (2) only by a constant factor. Note that the conformal weight of $: e^{i s_i \varphi(\zeta_i)/\sqrt{2}} :$ is $s_i^2/4 = 1/4$, as it should be.

We can now use the expression

$$\left\langle \prod_{i=1}^N : e^{i \nu_i \varphi(\zeta_i) + i \bar{\nu}_i \bar{\varphi}(\bar{\zeta}_i)} : \right\rangle = \frac{\delta_\nu \delta_{\bar{\nu}}}{|\eta(\tau)|^2} \sum_{(p, \bar{p}) \in \Gamma} A_0^p(\zeta_i, \nu_i) \overline{A_0^{\bar{p}}(\zeta_i, \bar{\nu}_i)} \quad (14)$$

for the correlator of a product of generic vertex operators on the torus derived in [23] (see also [24]). Here, $(\nu_i, \bar{\nu}_i) \in \Gamma$, Γ is the lattice of momenta given by

$$p = \frac{n}{R} + \frac{1}{2} m R, \quad \bar{p} = \frac{n}{R} - \frac{1}{2} m R, \quad n, m \in \mathbb{Z}, \quad (15)$$

$\delta_\nu \delta_{\bar{\nu}}$ is one for $\sum_i \nu_i = \sum_i \bar{\nu}_i = 0$ and zero otherwise,

$$\eta(\tau) = e^{\pi i \tau / 12} \prod_{n=1}^{\infty} (1 - e^{2\pi i \tau n}) \quad (16)$$

is the Dedekind eta function,

$$A_0^p(\zeta_i, \nu_i) = e^{i\pi p^2 \tau + 2\pi i p \sum_i \zeta_i \nu_i} \prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{\nu_i \nu_j}, \quad (17)$$

and the bar means complex conjugation and transformation of ν_i and p into $\bar{\nu}_i$ and \bar{p} .

For $R = \sqrt{2}$, the momenta become

$$p = \frac{1}{\sqrt{2}}(n + m), \quad \bar{p} = \frac{1}{\sqrt{2}}(n - m), \quad n, m \in \mathbb{Z}. \quad (18)$$

Noting that $n + m$ always has the same parity as $n - m$, we can write $n + m = 2r + 2k$ and $n - m = 2s + 2k$ and replace the sums over n and m by a sum over $k \in \{0, 1/2\}$ and sums over $r \in \mathbb{Z}$ and $s \in \mathbb{Z}$. Therefore

$$\sum_{(p, \bar{p}) \in \Gamma} A_0^p(\zeta_i, \nu_i) \overline{A_0^{\bar{p}}(\zeta_i, \nu_i)} = \sum_{k \in \{0, 1/2\}} \sum_{r \in \mathbb{Z}} A_0^{\sqrt{2}(r+k)}(\zeta_i, \nu_i) \overline{A_0^{\sqrt{2}(s+k)}(\zeta_i, \nu_i)}. \quad (19)$$

Combining (3), (13), (14), (17), (19), and (A.1), we conclude that the conformal blocks are

$$f_k(\zeta_i, s_i) = \frac{\xi_k \delta_s}{\eta(\tau)} \theta \left[\begin{matrix} k \\ 0 \end{matrix} \right] \left(\sum_i \zeta_i s_i, 2\tau \right) \prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{s_i s_j / 2}, \quad (20)$$

where $k \in \{0, 1/2\}$ and ξ_k is a phase factor that cannot be determined from the above arguments. ξ_k can be written as $\xi_k = e^{i g_k}$, where g_k is a real-valued function. Since g_k must also be holomorphic in ζ_i to ensure that f_k is holomorphic, we conclude that ξ_k cannot depend on ζ_i . It may, however, depend on s_i .

3.1.1. Singlet property The dependence of the phase factors ξ_k on s_i can be determined from the $SU(2)$ symmetry of the CFT, which dictates that the wavefunction must be a spin-0 state. Since ξ_k does not depend on ζ_i , it is sufficient to determine its value for a convenient choice of ζ_i . The idea in the following is to utilize that the torus locally looks like the plane. We can therefore find the phases by putting all the spins close together and comparing the conformal blocks to (8). Let us specifically put all ζ_i close to the origin, such that $|\zeta_i| \leq \epsilon$ for all i , where ϵ is a small positive number. We shall choose ϵ to be small enough that $N^2 \epsilon^2 \ll 1$, $|\sum_i \zeta_i s_i| \ll \text{Im}(\tau)^{1/2}$, and $|\zeta_i - \zeta_j| \ll \text{Im}(\tau)^{1/2}$ for all i and j . Then

$$\theta \left[\begin{matrix} k \\ 0 \end{matrix} \right] \left(\sum_i \zeta_i s_i, 2\tau \right) = \sum_{n \in \mathbb{Z}} e^{2\pi i \tau (n+k)^2 + 2\pi i (n+k) \sum_i \zeta_i s_i} = \sum_{n \in \mathbb{Z}} e^{2\pi i \tau (n+k)^2} + \mathcal{O}(\epsilon^2) \quad (21)$$

and

$$E(\zeta_i - \zeta_j, \tau) = \frac{1}{\partial_\zeta \theta_1(\zeta, \tau)|_{\zeta=0}} \sum_{n \in \mathbb{Z}} e^{\pi i \tau (n + \frac{1}{2})^2 + 2\pi i (n + \frac{1}{2})(\zeta_i - \zeta_j + \frac{1}{2})}$$

$$\begin{aligned}
 &= \frac{1}{\partial_\zeta \theta_1(\zeta, \tau)|_{\zeta=0}} \sum_{n \in \mathbb{Z}} 2\pi i \left(n + \frac{1}{2} \right) (\zeta_i - \zeta_j) e^{\pi i \tau (n + \frac{1}{2})^2 + \pi i (n + \frac{1}{2})} + \mathcal{O}(\epsilon^3) \\
 &= \zeta_i - \zeta_j + \mathcal{O}(\epsilon^3).
 \end{aligned} \tag{22}$$

Inserting these expressions in (20), we get

$$f_k(\zeta_i, s_i) \propto \xi_k \delta_{\mathbf{s}} \prod_{i < j} (\zeta_i - \zeta_j)^{s_i s_j / 2} (1 + \mathcal{O}(\epsilon^2)). \tag{23}$$

Since $\sum_{i < j} s_i s_j = \frac{1}{2} \sum_{i \neq j} s_i s_j = -\frac{1}{2} \sum_{i=1}^N s_i s_i = -\frac{N}{2}$ is a constant for $\sum_{j=1}^N s_j = 0$, we can replace $(\zeta_i - \zeta_j)$ by $(z_i - z_j)$ in (23). Comparing (23) to the $SU(2)$ invariant state on the complex plane (8), we observe that ξ_k is the Marshall sign factor defined in (10), provided we choose the phase convention of the square roots present in the Jastrow factor such that $(\zeta_i - \zeta_j)^{-1/2} = ((\zeta_i - \zeta_j)^{1/2})^{-1}$. Up to a constant phase factor, we thus have

$$f_k(\zeta_i, s_i) = \frac{\delta_{\mathbf{s}} \chi_{\mathbf{s}}}{\eta(\tau)} \theta \left[\begin{matrix} k \\ 0 \end{matrix} \right] \left(\sum_i \zeta_i s_i, 2\tau \right) \prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{s_i s_j / 2}, \tag{24}$$

where $E(\zeta_i - \zeta_j, \tau)^{-1/2} = (E(\zeta_i - \zeta_j, \tau)^{1/2})^{-1}$ is assumed. As far as the wavefunction is concerned, we are free to multiply the conformal block by an s_i -independent factor. To get (11), we multiply (24) by $\eta(\tau) \prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{1/2}$. Note that the latter factor changes the exponent in the Jastrow factor from the half-integer $s_i s_j / 2$ to the integer $(s_i s_j + 1) / 2$ so that the need to take square roots is eliminated. As a final remark, we note that there is a connection between the singlet property of the wavefunctions and Fay's trisecant identity as is clear from the alternative derivation of the singlet property for four spins given in Appendix B.

3.2. Independence of the ordering of the spins

Let us also demonstrate that the state (11) is independent (up to an overall sign factor) of the chosen ordering of the spins. Let p be a bijective map from $\{1, 2, \dots, N\}$ to $\{1, 2, \dots, N\}$, which defines an alternative ordering. Using this ordering, the state (11) takes the form

$$\begin{aligned}
 \psi_k^{(p)}(s_1, \dots, s_N) &= \delta_{\mathbf{s}} \prod_{j=1}^N (-1)^{(p(j)-1)(s_j+1)/2} \theta \left[\begin{matrix} k \\ 0 \end{matrix} \right] \left(\sum_{i=1}^N \zeta_i s_i, 2\tau \right) \\
 &\times \prod_{\{i, j \in \{1, 2, \dots, N\} | p(i) < p(j)\}} E(\zeta_i - \zeta_j, \tau)^{(s_i s_j + 1)/2}.
 \end{aligned} \tag{25}$$

Let us specifically consider the transposition

$$p(j) = \begin{cases} j & \text{for } j \neq j_1, j_2 \\ j_2 & \text{for } j = j_1 \\ j_1 & \text{for } j = j_2 \end{cases}, \tag{26}$$

where j_1 and j_2 are two numbers in $\{1, 2, \dots, N\}$ satisfying $j_1 < j_2$. Since E is an odd function in the first argument according to (A.10), we get

$$\begin{aligned} \frac{\psi_k^{(p)}(s_1, \dots, s_N)}{\psi_k(s_1, \dots, s_N)} &= \frac{(-1)^{(j_2-1)(s_{j_1}+1)/2+(j_1-1)(s_{j_2}+1)/2}}{(-1)^{(j_2-1)(s_{j_2}+1)/2+(j_1-1)(s_{j_1}+1)/2}} (-1)^{(s_{j_1}s_{j_2}+1)/2+\sum_{j=j_1+1}^{j_2-1}(s_{j_1}s_j+s_{j_2}s_j+2)/2} \\ &= (-1)^{(j_2-j_1)(s_{j_1}-s_{j_2})/2} (-1)^{(s_{j_1}s_{j_2}+1)/2+\sum_{j=j_1+1}^{j_2-1}s_j(s_{j_1}+s_{j_2})/2+j_2-j_1-1} \\ &= (-1)^{(j_2-j_1)(s_{j_1}-s_{j_2})/2+(s_{j_1}s_{j_2}+1)/2+(j_2-j_1-1)(s_{j_1}+s_{j_2})/2+j_2-j_1-1} \\ &= (-1)^{(j_2-j_1)(s_{j_1}+1)+(s_{j_1}-1)(s_{j_2}-1)/2-1} = -1. \end{aligned} \quad (27)$$

Swapping two labels in the ordering sequence thus only changes the overall sign of the wavefunction. Since any permutation p can be decomposed into a product of transpositions (26), we conclude that the choice of ordering is unimportant. The exact same arguments apply to (8), because $z_i - z_j$ is an odd function of $z_i - z_j$.

4. Connection to the KL states on the torus

The CFT states can be transformed into states describing particles on a lattice by regarding all spins with $s_i = +1$ as occupied sites and all spins with $s_i = -1$ as empty sites. Doing so, we demonstrate in this section that the CFT states on the torus (11) coincide with the KL states on the torus if the spins sit on an $L_x \times L_y$ square lattice with lattice constant $\sqrt{2\pi}$ and $\sum_j \zeta_j = 0$. The CFT states thus provide a natural generalization of the KL states to arbitrary lattices.

4.1. Statement of the result

Expressed mathematically, the desired square lattice is obtained by choosing

$$\omega_1 = \sqrt{2\pi}L_x, \quad \omega_2 = i\sqrt{2\pi}L_y, \quad \tau = iL_y/L_x, \quad (28)$$

and

$$z_{n+mL_x+1} = \sqrt{2\pi}[n - l_x + i(m - l_y)], \quad (29)$$

where

$$l_x = (L_x - 1)/2, \quad n \in \{0, 1, \dots, L_x - 1\}, \quad (30)$$

$$l_y = (L_y - 1)/2, \quad m \in \{0, 1, \dots, L_y - 1\}. \quad (31)$$

The result of this section is that

$$\psi_k(s_1, \dots, s_N) = (-1)^{2k} c\psi_k(Z_1, \dots, Z_{N/2}) \prod_i G_k(Z_i), \quad k = 0, 1/2. \quad (32)$$

The right hand side of this expression is the KL state on the torus. Specifically, $\psi_k(Z_1, \dots, Z_{N/2})$ is the Laughlin state with Landau level filling factor $1/2$ of $N/2$ particles on the torus [11, 25–27]

$$\begin{aligned} \psi_k(Z_1, \dots, Z_{N/2}) &= \\ &\theta \left[\begin{matrix} N/2 - k \\ -(N - 1) \end{matrix} \right] \left(\frac{2\sum_i Z_i}{\sqrt{2\pi}L_x}, 2\tau \right) \prod_{i < j} \left[\theta \left[\begin{matrix} 1/2 \\ 1/2 \end{matrix} \right] \left(\frac{Z_i - Z_j}{\sqrt{2\pi}L_x}, \tau \right) \right]^2 e^{-\frac{1}{2}\sum_i Y_i^2} \end{aligned} \quad (33)$$




Figure 2. We use four different ways to label the lattice sites. (n, m) is the coordinates on the lattice with $n = 0, 1, \dots, L_x - 1$ and $m = 0, 1, \dots, L_y - 1$. $j = n + mL_x + 1$ is the site index, which numbers the sites from 1 to N as shown with the labels above each site. z_j defined in (29) is the position in the complex plane of the site with index j (in units of the magnetic length, when comparing to FQH states), and $\zeta_j = z_j/\omega_1$, where $\omega_1 = \sqrt{2\pi}L_x$ is the width of the lattice in the x -direction.

with the particle positions $Z_i = X_i + iY_i$ restricted to the considered square lattice, i.e., $Z_i \in \{z_1, \dots, z_N\}$. Particles are identified with spins in the ‘up’ state such that $\{Z_1, \dots, Z_{N/2}\} = \{z_i | i \in \{1, 2, \dots, N\} \wedge s_i = 1\}$. As in (1), $\prod_i G_k(Z_i)$ is a gauge factor, which can only take values in the set $\{-1, +1\}$, and the purpose of this factor is to ensure that the right hand side of (32) is a singlet. Finally, c is an overall constant that only depends on the choice of L_x and L_y and is unimportant, since it is absorbed in the normalization of the state anyway. We find specifically that

$$\prod_{i=1}^{N/2} G_k(Z_i) = \begin{cases} \prod_{i=1}^N (-1)^{m_i q_i} & \text{for } L_y \text{ even} \\ \prod_{i=1}^N (-1)^{(n_i + m_i) q_i} & \text{for } L_y \text{ odd} \end{cases} . \quad (34)$$

Here, $n_i = \text{Re}(z_i)/\sqrt{2\pi} + l_x$ and $m_i = \text{Im}(z_i)/\sqrt{2\pi} + l_y$ as in (29) and

$$q_i = (s_i + 1)/2 \quad (35)$$

such that only sites with $s_i = +1$ contribute to the products. Note that q_i is the occupation number on site i since $q_i = 1$ for spin up and $q_i = 0$ for spin down. In the following, we derive (32) by rewriting each of the factors in (11).

4.2. The delta function factor

The condition $\sum_{i=1}^N s_i = 0$ translates into $\sum_{i=1}^N q_i = N/2$, and the delta function factor thus ensures that the lattice is half filled when a spin up is considered as an occupied site and a spin down is considered as an empty site. In other words, the delta function factor ensures that there are $N/2$ of the Z_i coordinates.

4.3. The Marshall sign factor

The Marshall sign factor can be rewritten into

$$\chi_{\mathbf{s}} = \prod_{j=1}^N (-1)^{(j-1)(s_j+1)/2} = \prod_{j=1}^N (-1)^{(j-1)q_j}. \quad (36)$$

4.4. The centre of mass factor

Let us first note that

$$\sum_{i=1}^N s_i z_i = 2 \sum_{i=1}^N q_i z_i - \sum_{i=1}^N z_i = 2 \sum_{i=1}^N q_i z_i = 2 \sum_{i=1}^{N/2} Z_i. \quad (37)$$

Using (A.2), (A.3), and the assumption that N is even, it then follows that

$$\theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\sum_i z_i s_i}{\sqrt{2\pi} L_x}, 2\tau \right) = (-1)^{2k\theta} \begin{bmatrix} N/2 - k \\ -(N-1) \end{bmatrix} \left(\frac{2 \sum_i Z_i}{\sqrt{2\pi} L_x}, 2\tau \right). \quad (38)$$

4.5. The Jastrow factor

We write

$$\prod_{i<j} E(\zeta_i - \zeta_j, \tau)^{(s_i s_j + 1)/2} = \prod_{i<j} E(\zeta_i - \zeta_j, \tau) \prod_{i<j} E(\zeta_i - \zeta_j, \tau)^{2q_i q_j} \prod_{i<j} E(\zeta_i - \zeta_j, \tau)^{-q_i - q_j}. \quad (39)$$

The first factor on the right hand side is an overall unimportant factor, the second factor appears in (33), and the third factor is to be compared to the Gaussian factor in (33). Since E is an odd function in the first argument according to (A.10), we have

$$\begin{aligned} \prod_{i<j} E(\zeta_i - \zeta_j, \tau)^{-q_i - q_j} &= \prod_{i<j} E(\zeta_i - \zeta_j, \tau)^{-q_i} \prod_{j<i} E(\zeta_j - \zeta_i, \tau)^{-q_i} \\ &= \prod_{j<i} (-1)^{-q_i} \prod_{i \neq j} E(\zeta_i - \zeta_j, \tau)^{-q_i} = \prod_{j=1}^N (-1)^{-(j-1)q_j} \prod_{i=1}^N f(\zeta_i)^{q_i}, \end{aligned} \quad (40)$$

where

$$f(\zeta_i) \equiv \prod_{j(\neq i)} E(\zeta_i - \zeta_j, \tau)^{-1} \quad (41)$$

and the notation $\prod_{j(\neq i)}$ means the product over all j except i and no product over i .

We next evaluate $f(\zeta_i)$. For $m \neq L_y - 1$, we find

$$\begin{aligned} \frac{f(\zeta_{n+(m+1)L_x+1})}{f(\zeta_{n+mL_x+1})} &= \frac{\prod_{\substack{0 \leq n' \leq L_x-1 \\ 0 \leq m' \leq L_y-1 \\ n'+im' \neq n+im}} E\left(\frac{n+im-n'-im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq n' \leq L_x-1 \\ 0 \leq m' \leq L_y-1 \\ n'+im' \neq n+i(m+1)}} E\left(\frac{n+i(m+1)-n'-im'}{L_x}, \tau\right)} \\ &= \frac{\prod_{\substack{0 \leq n' \leq L_x-1 \\ 0 \leq m' \leq L_y-1 \\ n'+im' \neq n+im}} E\left(\frac{n+im-n'-im'}{L_x}, \tau\right)}{\prod_{0 \leq n' \leq L_x-1} E\left(\frac{n+im-n'-i(L_y-1)}{L_x}, \tau\right)} \\ &= \frac{\prod_{\substack{0 \leq n' \leq L_x-1 \\ -1 \leq m' \leq L_y-2 \\ n'+im' \neq n+im}} E\left(\frac{n+im-n'-im'}{L_x}, \tau\right)}{\prod_{0 \leq n' \leq L_x-1} E\left(\frac{n+im-n'+i}{L_x}, \tau\right)} \end{aligned}$$

$$\begin{aligned}
&= \prod_{0 \leq n' \leq L_x - 1} e^{i\pi - i\pi\tau} e^{2\pi i[n + im - n' + i]/L_x} = e^{i\pi L_x - i\pi\tau L_x} e^{2\pi i[n + im - (L_x - 1)/2 + i]} \\
&= -e^{\pi L_y} e^{-2\pi(m+1)} = -e^{-2\pi(m - l_y) - \pi}, \tag{42}
\end{aligned}$$

where we have used (A.12). For $m = L_y - 1$, the relevant quantity is

$$\begin{aligned}
\frac{f(\zeta_{n+1})}{f(\zeta_{n+(L_y-1)L_x+1})} &= \frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n + i(L_y - 1)}} E\left(\frac{n + i(L_y - 1) - n' - im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n}} E\left(\frac{n - n' - im'}{L_x}, \tau\right)} \\
&= \frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n + i(L_y - 1)}} E\left(\frac{n + i(L_y - 1) - n' - im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n}} \left[E\left(\frac{n + iL_y - n' - im'}{L_x}, \tau\right) e^{i\pi - i\pi\tau + 2\pi i(n + iL_y - n' - im')/L_x} \right]} \\
&= \frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n + i(L_y - 1)}} E\left(\frac{n + i(L_y - 1) - n' - im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ -1 \leq m' \leq L_y - 2 \\ n' + im' \neq n - i}} \left[E\left(\frac{n + i(L_y - 1) - n' - im'}{L_x}, \tau\right) e^{i\pi - i\pi\tau + 2\pi i(n + i(L_y - 1) - n' - im')/L_x} \right]} \\
&= \frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ n' \neq n}} E\left(\frac{n - n'}{L_x}, \tau\right) e^{-i\pi(N-1) + i\pi\tau(N+1)}}{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ n' \neq n}} E\left(\frac{n - n'}{L_x} + \tau, \tau\right) \prod_{\substack{0 \leq n' \leq L_x - 1 \\ -1 \leq m' \leq L_y - 2}} e^{2\pi i(n + i(L_y - 1) - n' - im')/L_x}} \\
&= -\frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ n' \neq n}} E\left(\frac{n - n'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ n' \neq n}} \left[E\left(\frac{n - n'}{L_x}, \tau\right) e^{-i\pi - i\pi\tau - 2\pi i(n - n')/L_x} \right]} \times \frac{e^{i\pi\tau(N+1)}}{e^{2\pi i(n - L_x + i(L_y + 1)/2)L_y}} \\
&= \prod_{0 \leq n' \leq L_x - 1} e^{i\pi + i\pi\tau + 2\pi i(n - n')/L_x} \times \frac{e^{i\pi\tau N}}{e^{2\pi i(n - L_x + i(L_y + 1)/2)L_y}} = -(-1)^{L_y}. \tag{43}
\end{aligned}$$

For $n \neq L_x - 1$, we get

$$\begin{aligned}
\frac{f(\zeta_{n+1+mL_x+1})}{f(\zeta_{n+mL_x+1})} &= \frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n + im}} E\left(\frac{n + im - n' - im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n + 1 + im}} E\left(\frac{n + 1 + im - n' - im'}{L_x}, \tau\right)} = \frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n + im}} E\left(\frac{n + im - n' - im'}{L_x}, \tau\right)}{\prod_{\substack{-1 \leq n' \leq L_x - 2 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq n + im}} E\left(\frac{n + im - n' - im'}{L_x}, \tau\right)} \\
&= \frac{\prod_{0 \leq m' \leq L_y - 1} E\left(\frac{n + im - (L_x - 1) - im'}{L_x}, \tau\right)}{\prod_{0 \leq m' \leq L_y - 1} E\left(\frac{n + im + 1 - im'}{L_x}, \tau\right)} = (-1)^{L_y}, \tag{44}
\end{aligned}$$

where we used (A.11) in the last line. For $n = L_x - 1$, the relevant quantity is

$$\frac{f(\zeta_{mL_x+1})}{f(\zeta_{L_x-1+mL_x+1})} = \frac{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq L_x - 1 + im}} E\left(\frac{L_x - 1 + im - n' - im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq im}} E\left(\frac{im - n' - im'}{L_x}, \tau\right)}$$

$$\begin{aligned}
 & \prod_{\substack{0 \leq n' \leq L_x - 1 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq L_x - 1 + im}} E\left(\frac{L_x - 1 + im - n' - im'}{L_x}, \tau\right) \\
 = & - \frac{\prod_{\substack{-1 \leq n' \leq L_x - 2 \\ 0 \leq m' \leq L_y - 1 \\ n' + im' \neq -1 + im}} E\left(\frac{L_x - 1 + im - n' - im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq m' \leq L_y - 1 \\ m' \neq m}} E\left(\frac{im - im'}{L_x}, \tau\right)} \\
 = & - \frac{\prod_{\substack{0 \leq m' \leq L_y - 1 \\ m' \neq m}} E\left(\frac{im - im'}{L_x}, \tau\right)}{\prod_{\substack{0 \leq m' \leq L_y - 1 \\ m' \neq m}} E\left(\frac{L_x + im - im'}{L_x}, \tau\right)} = (-1)^{L_y}. \tag{45}
 \end{aligned}$$

The solution of these equations is

$$f(\zeta_{n+mL_x+1}) \propto \begin{cases} (-1)^m e^{-\pi(m-l_y)^2} & \text{for } L_y \text{ even} \\ (-1)^{n+m} e^{-\pi(m-l_y)^2} & \text{for } L_y \text{ odd} \end{cases}. \tag{46}$$

The result (32) is then obtained by combining the delta function factor, (36), (38), (39), (40), and (46).

5. Transformation properties of the CFT states on a square lattice on the torus

In this section, we investigate how the CFT states on the torus transform under different operations. This reveals the symmetries of the states and has the additional advantage of enabling us in certain cases to find linear combinations of the two states on the torus that are necessarily orthogonal. We assume throughout that the spins sit on an $L_x \times L_y$ square lattice as defined in (29), but unless it is required for the considered transformation to make sense, we shall not put other restrictions on L_x and L_y than demanding $N = L_x L_y$ to be even. For the sake of generality, we shall allow for twisted boundary conditions in the following. If the twist angles are θ_x and θ_y , this means that the state acquires a phase of $e^{i\theta_x N/2}$ if the lattice is displaced by $\sqrt{2\pi}L_x$ in the x -direction and acquires a phase of $e^{i\theta_y N/2}$ if the lattice is displaced by $\sqrt{2\pi}L_y$ in the y -direction. It is already known how to take the twists into account for the Laughlin states on the torus (see e.g. (5.5) in [27]), and by comparing the CFT states to this expression, we conclude that

$$\psi_{a,b}(s_1, \dots, s_N) = \delta_{\mathbf{s}} \chi_{\mathbf{s}} \theta \begin{bmatrix} a \\ b \end{bmatrix} \left(\sum_{i=1}^N \zeta_i s_i, 2\tau \right) \prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{(s_i s_j + 1)/2} \quad (\text{twist}) \tag{47}$$

provides the desired generalization, where

$$a = k + \frac{\theta_x}{4\pi}, \quad b = -\frac{\theta_y}{2\pi}, \quad k \in \{0, 1/2\}, \quad \sum_{j=1}^N \zeta_j = 0. \tag{48}$$

In analogy to (5), we also define

$$|\psi_{a,b}\rangle = \mathcal{C}_{a,b} \sum_{s_1, \dots, s_N} \psi_{a,b}(s_1, \dots, s_N) |s_1, \dots, s_N\rangle, \tag{49}$$

where $\mathcal{C}_{a,b}$ is the normalization constant.

Table 1. Eigenstates and eigenvalues in the subspace spanned by $\psi_0 = \psi_{0,0}(s_1, \dots, s_N)$ and $\psi_{1/2} = \psi_{1/2,0}(s_1, \dots, s_N)$ of the following operators: spin flip (F), translation by one lattice constant in the x -direction (T_x), translation by one lattice constant in the y -direction (T_y), rotation by 90° (R_{90°), rotation by 180° (R_{180°), time reversal followed by reflection in the x -direction ($M_x\Theta$), and time reversal followed by reflection in the y -direction ($M_y\Theta$). We consider four different types of square lattices on the torus: ‘equal’ stands for $L_x = L_y$ with L_x even, ‘e×e’ means L_x even and L_y even, ‘e×o’ means L_x even and L_y odd, and ‘o×e’ means L_x odd and L_y even (odd-by-odd is not allowed since $N = L_x L_y$ must be even). Note that ψ_0 and $\psi_{1/2}$ are defined as in (11) and are hence not normalized.

Lattice	Eigenstates	F	T_x	T_y	R_{90°	R_{180°	$M_x\Theta$	$M_y\Theta$
equal	$\psi_0 - (1 - \sqrt{2})\psi_{1/2}$	1	1	1	+1	1	1	1
	$\psi_0 - (1 + \sqrt{2})\psi_{1/2}$	1	1	1	-1	1	1	1
e×e	ψ_0	1	1	1	-	1	1	1
	$\psi_{1/2}$	1	1	1	-	1	1	1
e×o	ψ_0	$(-1)^{\frac{N}{2}}$	$(-1)^{\frac{L_x}{2}}$	1	-	$(-1)^{\frac{N}{2}}$	1	1
	$\psi_{1/2}$	$(-1)^{\frac{N}{2}}$	$-(-1)^{\frac{L_x}{2}}$	1	-	$(-1)^{\frac{N}{2}}$	1	1
o×e	$\psi_0 + \psi_{1/2}$	$(-1)^{\frac{N}{2}}$	1	$(-1)^{\frac{L_y}{2}}$	-	$(-1)^{\frac{N}{2}}$	1	1
	$\psi_0 - \psi_{1/2}$	$(-1)^{\frac{N}{2}}$	1	$-(-1)^{\frac{L_y}{2}}$	-	$(-1)^{\frac{N}{2}}$	1	1

The results of this section obtained for $\theta_x = \theta_y = 0$ are summarized in table 1. From this table we conclude that $\psi_0 = \psi_{0,0}(s_1, \dots, s_N)$ and $\psi_{1/2} = \psi_{1/2,0}(s_1, \dots, s_N)$ are orthogonal for even-by-odd lattices, whereas $\psi_0 + \psi_{1/2}$ and $\psi_0 - \psi_{1/2}$ are orthogonal for odd-by-even lattices. For $L_x = L_y$, it is the combinations $\psi_0 - (1 - \sqrt{2})\psi_{1/2}$ and $\psi_0 - (1 + \sqrt{2})\psi_{1/2}$ that are orthogonal. We also note that the operators describing translation in the x -direction and in the y -direction can be diagonalized simultaneously, which is not true in the continuum case [25]. This is so because the area of each lattice site is 2π , and the magnetic flux penetrating an area of 2π in the fractional quantum Hall setting is one flux quantum. The operator $(T_y)^{-1}(T_x)^{-1}T_yT_x$ thus moves all the particles (or spin ups) around one flux quantum, and therefore the Aharonov-Bohm phase picked up is a multiple of 2π . The transformation properties for general θ_x and θ_y can be found in (56), (66), (77), (91), (95), (102), and (103). Note also (82) and (83) for the transformation properties of a transformed set of translation operators.

5.1. General remarks for rearrangement transformations

Most of the transformations that we shall study below involve rearranging the spins on the lattice. An operator \mathcal{O} describing such a transformation is unitary and acts as

$$\mathcal{O}\sigma_j^\alpha\mathcal{O}^\dagger = \sigma_{d^{-1}(j)}^\alpha, \quad j = \{1, 2, \dots, N\}, \quad \alpha = x, y, z, \quad (50)$$

where σ_j^α is the α Pauli operator acting on the spin sitting at the site with index j and $d^{-1}(j)$ is a bijective map from $\{1, 2, \dots, N\}$ to $\{1, 2, \dots, N\}$ describing the

rearrangement. Let us for notational convenience define

$$\psi_{a,b}(p(j), \zeta_j, s_j, \tau) = \psi_{a,b}(s_1, \dots, s_N), \quad (51)$$

where $p(j)$ specifies the choice of ordering of the labels as in (25). The action of \mathcal{O} on the state (49) is

$$\begin{aligned} \mathcal{O}|\psi_{a,b}\rangle &= \mathcal{C}_{a,b} \mathcal{O} \sum_{s_1, \dots, s_N} \psi_{a,b}(j, \zeta_j, s_j, \tau) |s_1, \dots, s_N\rangle \\ &= \mathcal{C}_{a,b} \mathcal{O} \psi_{a,b}(j, \zeta_j, \sigma_j^z, \tau) \sum_{s_1, \dots, s_N} |s_1, \dots, s_N\rangle \\ &= \mathcal{C}_{a,b} \mathcal{O} \psi_{a,b}(j, \zeta_j, \sigma_j^z, \tau) \mathcal{O}^\dagger \mathcal{O} \sum_{s_1, \dots, s_N} |s_1, \dots, s_N\rangle \\ &= \mathcal{C}_{a,b} \mathcal{O} \psi_{a,b}(j, \zeta_j, \sigma_j^z, \tau) \mathcal{O}^\dagger \sum_{s_1, \dots, s_N} |s_1, \dots, s_N\rangle \\ &= \mathcal{C}_{a,b} \psi_{a,b}(j, \zeta_j, \sigma_{d^{-1}(j)}^z, \tau) \sum_{s_1, \dots, s_N} |s_1, \dots, s_N\rangle \\ &= \mathcal{C}_{a,b} \sum_{s_1, \dots, s_N} \psi_{a,b}(j, \zeta_j, s_{d^{-1}(j)}, \tau) |s_1, \dots, s_N\rangle. \end{aligned} \quad (52)$$

We can rewrite

$$\psi_{a,b}(j, \zeta_j, s_{d^{-1}(j)}, \tau) = \psi_{a,b}(d(j), \zeta_{d(j)}, s_j, \tau) = (-1)^{\mathcal{S}} \psi_{a,b}(j, \zeta_{d(j)}, s_j, \tau), \quad (53)$$

where \mathcal{S} is the number of times one has to swap two labels to change the numbering from $d(j)$ to j . The first equality in (53) follows from a relabelling of indices, and the second equality follows from (27), which trivially generalizes to the case of twisted boundary conditions. To find the transformation properties of the wavefunction under \mathcal{O} , we thus only need to determine \mathcal{S} and $\psi_{a,b}(j, \zeta_{d(j)}, s_j, \tau)$. Since the delta function factor and the Marshall sign factor do not depend on ζ_j , it is sufficient to consider the transformation properties of the centre of mass factor and the Jastrow factor. For later convenience, let us also define κ_d through the relation

$$\prod_{i < j} E(\zeta_{d(i)} - \zeta_{d(j)}, \tau)^{(s_i s_j + 1)/2} = \kappa_d \prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{(s_i s_j + 1)/2}. \quad (54)$$

5.2. Spin flip

Let us first consider the spin flip operator $F = \prod_{j=1}^N \sigma_j^x$, which is not of the rearrangement type. It transforms s_j into $-s_j$. The delta function and the Jastrow factor are easily seen to be invariant under this operation. The Marshall sign factor transforms as

$$\begin{aligned} \chi_{-s} &= \prod_{n=1}^N (-1)^{(n-1)(-s_n+1)/2} = \prod_{n=1}^N (-1)^{(n-1)(s_n-1)/2} = \chi_s \prod_{n=1}^N (-1)^{-(n-1)} \\ &= \chi_s (-1)^{(N-1)N/2} = \chi_s (-1)^{N/2}, \end{aligned} \quad (55)$$

and the transformation of the centre of mass factor follows from (A.4). Altogether

$$FC_{a,b}^{-1} |\psi_{a,b}\rangle = (-1)^{N/2} \mathcal{C}_{-a,-b}^{-1} |\psi_{-a,-b}\rangle. \quad (56)$$




Figure 3. The sets A , B , C , and D for a 5×4 lattice.

Note that the factors $\mathcal{C}_{a,b}^{-1}$ and $\mathcal{C}_{-a,-b}^{-1}$ simply remove the normalization factors from $|\psi_{a,b}\rangle$ and $|\psi_{-a,-b}\rangle$, respectively. Some special cases of (56) are

$$\begin{aligned}
 F|\psi_{0,0}\rangle &= (-1)^{N/2}|\psi_{0,0}\rangle, \\
 F|\psi_{1/2,0}\rangle &= (-1)^{N/2} \frac{\mathcal{C}_{1/2,0}}{\mathcal{C}_{-1/2,0}} |\psi_{-1/2,0}\rangle = (-1)^{N/2} |\psi_{1/2,0}\rangle, \\
 F|\psi_{0,1/2}\rangle &= (-1)^{N/2} \frac{\mathcal{C}_{0,1/2}}{\mathcal{C}_{0,-1/2}} |\psi_{0,-1/2}\rangle = (-1)^{N/2} |\psi_{0,1/2}\rangle, \\
 F|\psi_{1/2,1/2}\rangle &= (-1)^{N/2} \frac{\mathcal{C}_{1/2,1/2}}{\mathcal{C}_{-1/2,-1/2}} |\psi_{-1/2,-1/2}\rangle = -(-1)^{N/2} |\psi_{1/2,1/2}\rangle,
 \end{aligned} \tag{57}$$

where we have used (47), (49), (A.2) and (A.3). In particular, $\psi_{0,0}$ and $\psi_{1/2,0}$ are both eigenstates with eigenvalue $(-1)^{N/2}$ as stated in table 1.

5.3. Translation in the x -direction

The action of the translation operator T_x is to move the spins one lattice constant to the right, except for the rightmost column of spins which is wrapped around to the opposite edge. T_x is of the rearrangement type and is defined through the map

$$d(j) = \begin{cases} j - 1 & \text{for } j \in A, \\ j + (L_x - 1) & \text{for } j \in B, \end{cases} \tag{58}$$

which moves the lattice one lattice constant to the left. Here,

$$\begin{aligned}
 A &= \{n + mL_x + 1 | n = 1, \dots, L_x - 1 \wedge m = 0, 1, \dots, L_y - 1\}, \\
 B &= \{mL_x + 1 | m = 0, \dots, L_y - 1\},
 \end{aligned} \tag{59}$$

i.e., A is the set containing the indices of the sites in the $L_x - 1$ rightmost columns of the lattice and B is the set containing the indices of the sites in the leftmost column of the lattice as illustrated for a 5×4 lattice in figure 3(a). This map has $\mathcal{S} = L_y(L_x - 1)$.

To determine the transformation properties of the wavefunction, we note that

$$\zeta_{d(j)} = \begin{cases} \zeta_j - 1/L_x & \text{for } j \in A, \\ \zeta_j + (L_x - 1)/L_x & \text{for } j \in B. \end{cases} \tag{60}$$

Therefore

$$\sum_{j=1}^N \zeta_{d(j)} s_j = \sum_{j \in A} \left(\zeta_j - \frac{1}{L_x} \right) s_j + \sum_{j \in B} \left(\zeta_j + \frac{L_x - 1}{L_x} \right) s_j = \sum_{j=1}^N \zeta_j s_j + \sum_{j \in B} s_j, \tag{61}$$

and from (A.5) it then follows that

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} \left(\sum_{j=1}^N \zeta_{d(j)} s_j, 2\tau \right) = \theta \begin{bmatrix} a \\ b \end{bmatrix} \left(\sum_{j=1}^N \zeta_j s_j, 2\tau \right) e^{2\pi i a \sum_{j \in B} s_j}. \quad (62)$$

Regarding the Jastrow factor, we observe that

$$E(\zeta_{d(i)} - \zeta_{d(j)}, \tau) = \begin{cases} E(\zeta_i - \zeta_j, \tau) & \text{for } (i \in A \wedge j \in A) \vee (i \in B \wedge j \in B), \\ -E(\zeta_i - \zeta_j, \tau) & \text{for } (i \in A \wedge j \in B) \vee (i \in B \wedge j \in A). \end{cases} \quad (63)$$

The case $i \in A$ and $j \in B$ follows from

$$\begin{aligned} E(\zeta_{d(i)} - \zeta_{d(j)}, \tau) &= E\left(\zeta_i - \frac{1}{L_x} - \zeta_j - \frac{L_x - 1}{L_x}, \tau\right) = E(\zeta_i - \zeta_j - 1, \tau) \\ &= -E(\zeta_i - \zeta_j, \tau), \end{aligned} \quad (64)$$

where we have used (A.11), and the case $i \in B$ and $j \in A$ then follows from (A.10). For κ_d (see (54)), we therefore get

$$\kappa_d = (-1)^{\sum_{i \in A, j \in B} (s_i s_j + 1)/2} = (-1)^{-(\sum_{j \in B} s_j)^2/2 + L_y^2(L_x - 1)/2} = (-1)^{L_y^2(L_x - 2)/2}. \quad (65)$$

Collecting all the factors, we arrive at

$$T_x |\psi_{a,b}\rangle = \begin{cases} (-1)^{L_x/2} e^{2\pi i a \sum_{j \in B} \sigma_j^z} |\psi_{a,b}\rangle & \text{for } L_y \text{ odd,} \\ e^{2\pi i a \sum_{j \in B} \sigma_j^z} |\psi_{a,b}\rangle & \text{for } L_y \text{ even.} \end{cases} \quad (66)$$

For $a \in \{0, 1/2\}$, i.e., for $\theta_x = 0$, we can replace $\sum_{j \in B} \sigma_j^z$ by L_y , and it follows that $\psi_{0,b}$ and $\psi_{1/2,b}$ are eigenstates of T_x with the eigenvalues given in table 1.

5.4. Translation in the y -direction

The translation operator T_y in the y -direction is described by the map

$$d(j) = \begin{cases} j - L_x & \text{for } j \in C, \\ j + (L_y - 1)L_x & \text{for } j \in D, \end{cases} \quad (67)$$

which moves the lattice one lattice constant in the negative y -direction. Here

$$\begin{aligned} C &= \{L_x + 1, L_x + 2, \dots, L_x L_y\}, \\ D &= \{1, 2, \dots, L_x\}, \end{aligned} \quad (68)$$

i.e., C is the set of indices of the $L_y - 1$ uppermost rows of the lattice and D is the set of indices of the lowermost row of the lattice as illustrated for a 5×4 lattice in figure 3(b). This map has $\mathcal{S} = L_x(L_y - 1)$.

We note that

$$\zeta_{d(j)} = \begin{cases} \zeta_j - i/L_x & \text{for } j \in C, \\ \zeta_j + i(L_y - 1)/L_x & \text{for } j \in D. \end{cases} \quad (69)$$

Therefore

$$\sum_{j=1}^N \zeta_{d(j)} s_j = \sum_{j \in C} \left(\zeta_j - \frac{i}{L_x} \right) s_j + \sum_{j \in D} \left(\zeta_j + i \frac{L_y - 1}{L_x} \right) s_j = \sum_{j=1}^N \zeta_j s_j + \tau \sum_{j \in D} s_j, \quad (70)$$

and, using (A.6) with $p \rightarrow p/2$ and $\tau \rightarrow 2\tau$, it follows that

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} \left(\sum_{j=1}^N \zeta_{d(j)} s_j, 2\tau \right) = \theta \begin{bmatrix} a + \frac{L_x}{2} \\ b \end{bmatrix} \left(\sum_{j=1}^N \zeta_j s_j, 2\tau \right) e^{-i\pi\tau(\sum_{j \in D} s_j)^2/2 - i\pi \sum_{j \in D} s_j (\sum_{j=1}^N \zeta_j s_j + b)}. \quad (71)$$

To derive the transformation of the Jastrow factor, we observe that

$$E(\zeta_{d(i)} - \zeta_{d(j)}, \tau) = \begin{cases} E(\zeta_i - \zeta_j, \tau) & \text{for } i \in C \wedge j \in C, \\ E(\zeta_i - \zeta_j, \tau) & \text{for } i \in D \wedge j \in D, \\ e^{-i\pi - i\pi\tau - 2\pi i(\zeta_i - \zeta_j)} E(\zeta_i - \zeta_j, \tau) & \text{for } i \in D \wedge j \in C. \end{cases} \quad (72)$$

The case $i \in C$ and $j \in D$ is not relevant, because this automatically implies $i > j$ according to (68). The case $i \in D$ and $j \in C$ follows from

$$\begin{aligned} E(\zeta_{d(i)} - \zeta_{d(j)}, \tau) &= E\left(\zeta_i + i\frac{L_y - 1}{L_x} - \zeta_j + i\frac{1}{L_x}, \tau\right) = E(\zeta_i - \zeta_j + \tau, \tau) \\ &= e^{-i\pi - i\pi\tau - 2\pi i(\zeta_i - \zeta_j)} E(\zeta_i - \zeta_j, \tau), \end{aligned} \quad (73)$$

where we have used (A.12). κ_d (see (54)) is therefore

$$\kappa_d = (-1)^{\sum_{i \in D, j \in C} (s_i s_j + 1)/2} e^{-i\pi \sum_{i \in D, j \in C} [\tau + 2(\zeta_i - \zeta_j)] (s_i s_j + 1)/2}. \quad (74)$$

Note that

$$\begin{aligned} \sum_{i \in D, j \in C} [\tau + 2(\zeta_i - \zeta_j)] &= i\frac{L_y}{L_x} L_x^2 (L_y - 1) + 2L_x (L_y - 1) \sum_{i \in D} \zeta_i - 2L_x \sum_{j \in C} \zeta_j \\ &= iL_y L_x (L_y - 1) + 2L_x L_y \sum_{i \in D} \zeta_i \\ &= iL_y L_x (L_y - 1) + 2L_x L_y (-i) \frac{L_y - 1}{2L_x} L_x = 0. \end{aligned} \quad (75)$$

Therefore

$$\begin{aligned} \kappa_d &= (-1)^{\sum_{i \in D, j \in C} (s_i s_j + 1)/2} e^{-i\pi \sum_{i \in D, j \in C} [\tau + 2(\zeta_i - \zeta_j)] s_i s_j / 2} \\ &= (-1)^{L_x^2 (L_y - 2)/2} e^{i\pi\tau (\sum_{i \in D} s_i)^2 / 2 + i\pi \sum_{i=1}^N \zeta_i s_i \sum_{j \in D} s_j}. \end{aligned} \quad (76)$$

Collecting all the factors, we conclude that

$$T_y \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle = \begin{cases} (-1)^{L_y/2} e^{-i\pi b \sum_{j \in D} \sigma_j^z} \mathcal{C}_{a+\frac{1}{2}, b}^{-1} |\psi_{a+\frac{1}{2}, b}\rangle & \text{for } L_x \text{ odd,} \\ e^{-i\pi b \sum_{j \in D} \sigma_j^z} \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle & \text{for } L_x \text{ even.} \end{cases} \quad (77)$$

It follows that T_y transforms states in the subspace spanned by $\psi_{a,0}$ and $\psi_{a+1/2,0}$ into states in the same subspace, and one can therefore easily diagonalize T_y in these subspaces. The result for $a = 0$ is given in table 1.

5.5. Transformed translation operators

As a side remark, we show that the action of the operators

$$\tilde{T}_x = UT_x U^\dagger, \quad \tilde{T}_y = UT_y U^\dagger, \quad (78)$$

on the wavefunctions, where U is the unitary operator

$$U = \exp \left[\frac{i}{2} \sum_{n=0}^{L_x-1} \sum_{m=0}^{L_y-1} \left(\frac{n\theta_x}{L_x} + \frac{m\theta_y}{L_y} \right) \sigma_{n+mL_x+1}^z \right], \quad (79)$$

is in fact simpler than the action of T_x and T_y . We first find explicit expressions for \tilde{T}_x and \tilde{T}_y by writing $\tilde{T}_d = U(T_d U^\dagger T_d^\dagger) T_d$, $d = x, y$, and evaluating the expression in brackets using (50) with $\mathcal{O} = T_d$. This gives

$$\tilde{T}_x = \exp \left(-\frac{i}{2} \theta_x \sum_{j \in B} \sigma_j^z + \frac{i}{2} \frac{\theta_x}{L_x} \sum_{j=1}^N \sigma_j^z \right) T_x, \quad (80)$$

$$\tilde{T}_y = \exp \left(-\frac{i}{2} \theta_y \sum_{j \in D} \sigma_j^z + \frac{i}{2} \frac{\theta_y}{L_y} \sum_{j=1}^N \sigma_j^z \right) T_y. \quad (81)$$

Combining this result with (66) and (77) and remembering (48), we get

$$\tilde{T}_x \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle = \begin{cases} (-1)^{2k} (-1)^{L_x/2} \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle & \text{for } L_y \text{ odd,} \\ \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle & \text{for } L_y \text{ even,} \end{cases} \quad (82)$$

$$\tilde{T}_y \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle = \begin{cases} (-1)^{L_y/2} \mathcal{C}_{a+1/2,b}^{-1} |\psi_{a+1/2,b}\rangle & \text{for } L_x \text{ odd,} \\ \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle & \text{for } L_x \text{ even,} \end{cases} \quad (83)$$

where $k = a - \theta_x/(4\pi) \in \{0, 1/2\}$. We thus conclude that $\psi_{a,b}$ is an eigenstate of \tilde{T}_x and \tilde{T}_y for L_x even, whereas $\psi_{a,b} \pm \psi_{a+1/2,b}$ are eigenstates for L_x odd.

5.6. Rotation by 90°

In this subsection, we assume $L_x = L_y$ and study the action of a rotation by 90° . A rotation of the lattice by -90° is described by the map

$$d(n + mL_x + 1) = m + (L_x - 1 - n)L_x + 1, \quad (84)$$

where $n = 0, 1, \dots, L_x - 1$ and $m = 0, 1, \dots, L_x - 1$. For this map $\mathcal{S} = 3 \times N/4$.

Let us note that

$$\zeta_{d(j)} = -i\zeta_j = \zeta_j/\tau, \quad \tau = i = -1/i = -1/\tau. \quad (85)$$

We thus need to compute $\psi_{a,b}(j, \zeta_j/\tau, s_j, -1/\tau)$. The transformation done on this wavefunction is precisely the modular S -transformation (see, e.g., [22]), which takes

$$\zeta_j \rightarrow \frac{\zeta_j}{\tau}, \quad \tau \rightarrow -\frac{1}{\tau}, \quad (86)$$

and is defined for general τ . We have

$$\begin{aligned} \theta \begin{bmatrix} a \\ b \end{bmatrix} \left(\frac{\zeta}{\tau}, -\frac{2}{\tau} \right) &= \sum_{n \in \mathbb{Z}} e^{-\pi i \frac{2}{\tau} (n+a)^2 + 2\pi i (n+a) (\frac{\zeta}{\tau} + b)} = \sum_{k \in \mathbb{Z}} \int_{-\infty}^{\infty} e^{2\pi i k x - \pi i \frac{2}{\tau} (x+a)^2 + 2\pi i (x+a) (\frac{\zeta}{\tau} + b)} dx \\ &= \sqrt{\frac{\tau}{2i}} \sum_{k \in \mathbb{Z}} e^{-i2\pi a k + i \frac{\pi \tau}{2} (k + \frac{\zeta}{\tau} + b)^2} = \sqrt{\frac{\tau}{2i}} \sum_{\mu=0}^1 \sum_{n \in \mathbb{Z}} e^{-i2\pi a (2n+\mu) + i \frac{\pi \tau}{2} (2n+\mu + \frac{\zeta}{\tau} + b)^2} \end{aligned}$$

$$\begin{aligned}
 &= \sqrt{\frac{\tau}{2i}} \sum_{\mu=0}^1 \sum_{n \in \mathbb{Z}} e^{2\pi i \tau (n + \frac{\mu}{2} + \frac{b}{2})^2 + 2\pi i (n + \frac{\mu}{2} + \frac{b}{2})(\zeta - 2a) + \frac{\pi i \zeta^2}{2\tau} + 2\pi i a b} \\
 &= \sqrt{\frac{\tau}{2i}} e^{\pi i \zeta^2 / (2\tau) + 2\pi i a b} \sum_{\mu=0}^1 \theta \left[\begin{matrix} \frac{b}{2} + \frac{\mu}{2} \\ -2a \end{matrix} \right] (\zeta, 2\tau). \tag{87}
 \end{aligned}$$

A similar computation gives

$$\theta_1 \left(\frac{\zeta}{\tau}, -\frac{1}{\tau} \right) = -i(-i\tau)^{1/2} e^{i\pi \zeta^2 / \tau} \theta_1(\zeta, \tau), \tag{88}$$

which, together with (12), implies

$$E \left(\frac{\zeta_i - \zeta_j}{\tau}, -\frac{1}{\tau} \right) = \tau^{-1} e^{i\pi(\zeta_i - \zeta_j)^2 / \tau} E(\zeta_i - \zeta_j, \tau). \tag{89}$$

For the $L_x \times L_x$ lattice, we have $\sum_j \zeta_j = 0$ and $\sum_j \zeta_j^2 = 0$, and therefore

$$\prod_{i < j} e^{i\pi(\zeta_i - \zeta_j)^2 (s_i s_j + 1) / (2\tau)} = e^{i\pi \sum_i \sum_j (\zeta_i - \zeta_j)^2 (s_i s_j + 1) / (4\tau)} = e^{-i\pi (\sum_{j=1}^N \zeta_j s_j)^2 / (2\tau)}. \tag{90}$$

It follows that $\kappa_d = \tau^{-N(N-2)/4} \exp[-i\pi (\sum_{j=1}^N \zeta_j s_j)^2 / (2\tau)]$. Collecting the factors and using the facts that $\tau = i$ and that $N/4$ is an integer in our case, we find

$$R_{90^\circ} \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle = \frac{e^{2\pi i a b}}{\sqrt{2}} \left(\mathcal{C}_{b/2, -2a}^{-1} |\psi_{b/2, -2a}\rangle + \mathcal{C}_{b/2+1/2, -2a}^{-1} |\psi_{b/2+1/2, -2a}\rangle \right). \tag{91}$$

Particular cases are

$$R_{90^\circ} \mathcal{C}_{0,0}^{-1} |\psi_{0,0}\rangle = \frac{1}{\sqrt{2}} \left(\mathcal{C}_{0,0}^{-1} |\psi_{0,0}\rangle + \mathcal{C}_{1/2,0}^{-1} |\psi_{1/2,0}\rangle \right), \tag{92}$$

$$R_{90^\circ} \mathcal{C}_{1/2,0}^{-1} |\psi_{1/2,0}\rangle = \frac{1}{\sqrt{2}} \left(\mathcal{C}_{0,0}^{-1} |\psi_{0,0}\rangle - \mathcal{C}_{1/2,0}^{-1} |\psi_{1/2,0}\rangle \right).$$

From this one easily derives the results in table 1. Note also that the combination

$$|\psi_{0,0}(s_1, \dots, s_N)|^2 + |\psi_{1/2,0}(s_1, \dots, s_N)|^2 \tag{93}$$

is invariant under the rotation by 90° . This is a consequence of the fact that the correlator (3) of the complete field is modular invariant and that the absolute value of the factor $\eta(\tau) \prod_{i < j} E(\zeta_i - \zeta_j, \tau)^{1/2}$ that we removed from the conformal blocks in defining $\psi_{0,0}$ and $\psi_{1/2,0}$ in section 3.1.1 is invariant under a 90° rotation for the considered lattice. Simple eigenstates can also be found for $a \in \{1/4, 3/4\}$ and $b = 1/2$.

5.7. Rotation by 180°

The rotation by 180° has

$$d(j) = N - j, \quad \zeta_{d(j)} = -\zeta_j, \quad \mathcal{S} = N/2. \tag{94}$$

The transformation of the centre of mass factor follows from (A.7), and the Jastrow factor gives the factor $\kappa_d = 1$. Altogether, we therefore get

$$R_{180^\circ} \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle = (-1)^{N/2} \mathcal{C}_{-a,-b}^{-1} |\psi_{-a,-b}\rangle, \tag{95}$$

which is the same result as for the spin flip operator.

5.8. Time reversal combined with reflection

In the FQH setting, time reversal corresponds to inverting the direction of the magnetic field. The action of the time reversal operator Θ on a general spin state

$$|\psi\rangle = \sum_{s_1, \dots, s_N} \psi(s_1, \dots, s_N) |s_1, \dots, s_N\rangle \quad (96)$$

is [28]

$$\Theta|\psi\rangle = i^N \sum_{s_1, \dots, s_N} (-1)^{\sum_{j=1}^N (s_j+1/2)} \psi(-s_1, \dots, -s_N)^* |s_1, \dots, s_N\rangle. \quad (97)$$

Therefore time reversal takes $\mathcal{C}_{a,b}\psi_{a,b}(j, \zeta_j, s_j, \tau)$ into $\mathcal{C}_{a,b}\psi_{a,b}^*(j, \zeta_j, -s_j, \tau)$, where we choose $\mathcal{C}_{a,b}$ to be real. The delta function factor is not affected by this transformation, and the Marshall sign factor is changed by the factor $(-1)^{N/2}$ as in (55). Since τ is imaginary in our case, we get from (A.1) that

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} \left(-\sum_{j=1}^N \zeta_j s_j, 2\tau \right)^* = \theta \begin{bmatrix} a \\ -b \end{bmatrix} \left(\sum_{j=1}^N \zeta_j^* s_j, 2\tau \right), \quad (98)$$

$$E(\zeta_i - \zeta_j, \tau)^* = E(\zeta_i^* - \zeta_j^*, \tau). \quad (99)$$

The state is thus not invariant under time reversal, which is a generic feature for FQH wave functions.

We can get a second complex conjugation of ζ_j by acting with a reflection operator. The reflection operator M_x in the x -direction has

$$d(n + mL_x + 1) = L_x - n + mL_x, \quad \zeta_{d(j)} = -\zeta_j^*, \quad \mathcal{S} = N/2, \quad (100)$$

and the reflection operator M_y in the y -direction has

$$d(n + mL_x + 1) = n + (L_y - 1 - m)L_x + 1, \quad \zeta_{d(j)} = \zeta_j^*, \quad \mathcal{S} = N/2. \quad (101)$$

Applying one of these operators and using (A.4) and (A.10), we get

$$M_x \Theta \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle = \mathcal{C}_{-a,b}^{-1} |\psi_{-a,b}\rangle, \quad (102)$$

$$M_y \Theta \mathcal{C}_{a,b}^{-1} |\psi_{a,b}\rangle = \mathcal{C}_{a,-b}^{-1} |\psi_{a,-b}\rangle, \quad (103)$$

which, together with (A.2), leads to the results in table 1.

6. Many-body Chern number and other topological properties

A main reason for studying FQH states is their nontrivial topological properties. For the KL states, the topological properties derive from the center of mass factor [25, 27]. By now a number of measures have been identified that can be used to describe topological states, and some of these have already been used in [3] and [17] to demonstrate the topological nature of the CFT state on an irregular lattice on the sphere and the CFT states on square lattices on the torus. For the generalization of the KL states on the torus to arbitrary lattices presented in the present paper to be useful, it is important that the topological properties are preserved when the lattice is deformed away from




Figure 4. The many-body Chern number of the two CFT states is one for the 4×4 square lattice in a), the randomly distorted 4×4 square lattice in b), the triangular lattice in c), and the kagome lattice in d).

the square lattice. To test this property, we compute the many-body Chern number for the lattices shown in figure 4, and in all cases we get the expected value 1. (The computation is done as described in detail in [29, 30], and we use the wavefunction (47) with twisted boundary conditions.)

The number of degenerate ground states on higher genus surfaces is an important quantity to partially characterize topological states. Degeneracies may, however, arise for other reasons than topology. To talk about topologically degenerate states, the local structure of the states must be the same, i.e., the expectation value of any local operator must be the same for all the states in the thermodynamic limit. Local indistinguishability has been demonstrated for the two states in (11) for the case of a square lattice in [17]. One can observe from (47) that the two states on the torus can be transformed into each other by changing the twist angles. Since the local structure is changed little by this operation when the number of spins is large, this is also a sign of topology. Specifically, if θ_x is changed from 0 to 2π , a increases by $1/2$, and from (A.2) it then follows that the states are transformed as

$$\psi_{0,0}(s_1, \dots, s_N) \rightarrow \psi_{1/2,0}(s_1, \dots, s_N), \quad (104)$$

$$\psi_{1/2,0}(s_1, \dots, s_N) \rightarrow \psi_{0,0}(s_1, \dots, s_N). \quad (105)$$

This is illustrated in figure 5. If instead we increase θ_y from 0 to 2π , it follows from (47) and (A.3) that the states are transformed as

$$\psi_{0,0}(s_1, \dots, s_N) + \psi_{1/2,0}(s_1, \dots, s_N) \rightarrow \psi_{0,0}(s_1, \dots, s_N) - \psi_{1/2,0}(s_1, \dots, s_N), \quad (106)$$

$$\psi_{0,0}(s_1, \dots, s_N) - \psi_{1/2,0}(s_1, \dots, s_N) \rightarrow \psi_{0,0}(s_1, \dots, s_N) + \psi_{1/2,0}(s_1, \dots, s_N). \quad (107)$$

7. Conclusion

In conclusion, we have found that the KL states on the torus can be written as conformal blocks of primary fields from the $SU(2)_1$ WZW CFT. This representation of the states reveals an interesting underlying mathematical structure and gives a natural generalization of the KL states on the torus to arbitrary lattices. It also provides an easy way to ensure that the states are singlet states. In addition, we have found that the many-body Chern number for different lattices is unity and that the two states are transformed into each other under a twist of the boundary conditions. These findings




Figure 5. When θ_x increases from 0 to 2π , a increases by $1/2$. Since the wavefunction only depends on a modulus 1 (see (A.2)), it follows that $\psi_{0,b}$ is transformed into $\psi_{1/2,b}$ and vice versa. If θ_x is increased to 4π , we are back to the starting point.

suggest that the topological properties are preserved when the lattice is transformed away from the square lattice. Finally, we have analyzed the symmetry properties of the CFT states on the torus for the case of an $L_x \times L_y$ square lattice, which allowed us in certain cases to construct linear combinations of the states that are guaranteed to be orthogonal because they have different symmetries. Our work also shows that the KL states in different geometries can be obtained by evaluating the same conformal correlator in different geometries, and we believe that this holds more generally. The CFT representation found in the present paper constitutes an interesting starting point for further analytical investigations of the states. In [4], CFT representations of lattice Laughlin states with general filling factor $1/q$, where q is an integer, have been found on the Riemann sphere, and using these results, we note that the present work could be straightforwardly generalized to obtain lattice Laughlin states with filling factor $1/q$ on the torus. Higher genus versions of the CFT states fulfilling the CFT fusion rules can also be obtained. Another interesting perspective is to make similar constructions for other FQH states.

Acknowledgments

The authors acknowledge discussions with J. Ignacio Cirac, Denis Bernard, and Hong-Hao Tu. This work has been partially funded by EU through SIQS grant (FP7 600645) and by FIS2012-33642, QUITEMAD (CAM), and the Severo Ochoa Program.

Appendix A. Properties of the Riemann theta function

The Riemann theta function is defined as

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} (\zeta, \tau) = \sum_{n \in \mathbb{Z}} e^{i\pi\tau(n+a)^2 + 2\pi i(n+a)(\zeta+b)}, \quad (\text{A.1})$$

where a and b are real numbers, ζ is complex, and τ is complex with $\text{Im}(\tau) > 0$. From this definition, one easily derives the following identities

$$\theta \begin{bmatrix} a+c \\ b \end{bmatrix}(\zeta, \tau) = \theta \begin{bmatrix} a \\ b \end{bmatrix}(\zeta, \tau), \quad c \in \mathbb{Z}, \quad (\text{A.2})$$

$$\theta \begin{bmatrix} a \\ b+c \end{bmatrix}(\zeta, \tau) = e^{2\pi i a c} \theta \begin{bmatrix} a \\ b \end{bmatrix}(\zeta, \tau), \quad c \in \mathbb{Z}, \quad (\text{A.3})$$

$$\theta \begin{bmatrix} a \\ b \end{bmatrix}(-\zeta, \tau) = \theta \begin{bmatrix} -a \\ -b \end{bmatrix}(\zeta, \tau), \quad (\text{A.4})$$

$$\theta \begin{bmatrix} a \\ b \end{bmatrix}(\zeta + c, \tau) = e^{2\pi i a c} \theta \begin{bmatrix} a \\ b \end{bmatrix}(\zeta, \tau), \quad c \in \mathbb{Z}, \quad (\text{A.5})$$

$$\theta \begin{bmatrix} a \\ b \end{bmatrix}(\zeta + p\tau, \tau) = e^{-i\pi\tau p^2 - i2\pi p(\zeta+b)} \theta \begin{bmatrix} a+p \\ b \end{bmatrix}(\zeta, \tau), \quad p \in \mathbb{R}. \quad (\text{A.6})$$

Particular cases that we shall need are

$$\theta \begin{bmatrix} k \\ 0 \end{bmatrix}(-\zeta, \tau) = \theta \begin{bmatrix} k \\ 0 \end{bmatrix}(\zeta, \tau), \quad k = 0, \frac{1}{2}, \quad (\text{A.7})$$

$$\theta \begin{bmatrix} k \\ 0 \end{bmatrix}(\zeta \pm 1, 2\tau) = e^{\pm 2\pi i k} \theta \begin{bmatrix} k \\ 0 \end{bmatrix}(\zeta, 2\tau), \quad k = 0, \frac{1}{2}, \quad (\text{A.8})$$

$$\theta \begin{bmatrix} k \\ 0 \end{bmatrix}(\zeta \pm \tau, 2\tau) = e^{-i\pi\tau/2 \mp \pi i \zeta} \theta \begin{bmatrix} 1/2 - k \\ 0 \end{bmatrix}(\zeta, 2\tau), \quad k = 0, \frac{1}{2}, \quad (\text{A.9})$$

and

$$E(-\zeta, \tau) = -E(\zeta, \tau), \quad (\text{A.10})$$

$$E(\zeta \pm 1, \tau) = e^{\pm i\pi} E(\zeta, \tau), \quad (\text{A.11})$$

$$E(\zeta \pm \tau, \tau) = e^{-i\pi\tau \mp i\pi \mp 2\pi i \zeta} E(\zeta, \tau), \quad (\text{A.12})$$

where $E(\zeta, \tau)$ is the prime form defined in (12).

Appendix B. Singlet property for four spins from Fay's trisecant identity

Consider the case of four spins. From (11), the wavefunctions on the torus read

$$|\psi_k\rangle = \mathcal{C}_k(\psi_{1,k}|+1, -1, +1, -1\rangle + \psi_{2,k}|+1, -1, -1, +1\rangle + \psi_{3,k}|+1, +1, -1, -1\rangle + \psi_{1,k}|-1, +1, -1, +1\rangle + \psi_{2,k}|-1, +1, +1, -1\rangle + \psi_{3,k}|-1, -1, +1, +1\rangle), \quad (\text{B.1})$$

where

$$\psi_{1,k} = E(\zeta_1 - \zeta_3, \tau) E(\zeta_2 - \zeta_4, \tau) \theta \begin{bmatrix} k \\ 0 \end{bmatrix}(\zeta_1 - \zeta_2 + \zeta_3 - \zeta_4, 2\tau), \quad (\text{B.2})$$

$$\psi_{2,k} = -E(\zeta_1 - \zeta_4, \tau) E(\zeta_2 - \zeta_3, \tau) \theta \begin{bmatrix} k \\ 0 \end{bmatrix}(\zeta_1 - \zeta_2 - \zeta_3 + \zeta_4, 2\tau),$$

$$\psi_{3,k} = -E(\zeta_1 - \zeta_2, \tau) E(\zeta_3 - \zeta_4, \tau) \theta \begin{bmatrix} k \\ 0 \end{bmatrix}(\zeta_1 + \zeta_2 - \zeta_3 - \zeta_4, 2\tau).$$

This state is a singlet if and only if

$$\psi_{1,k} + \psi_{2,k} + \psi_{3,k} = 0. \quad (\text{B.3})$$

Now we use (A3) in [27] to express the theta function with argument 2τ as the product of two theta functions with argument τ ,

$$\theta \begin{bmatrix} k \\ 0 \end{bmatrix} (\zeta, 2\tau) = \frac{\eta(2\tau)}{\eta^2(\tau)} \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta}{2} + \frac{1}{4}, \tau \right) \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta}{2} - \frac{1}{4}, \tau \right), \quad k = 0, \frac{1}{2}, \quad (\text{B.4})$$

where η is the Dedekind eta function defined in (16). We can therefore write (B.2) as

$$\psi_{1,k} = [\eta(2\tau)/\eta^2(\tau)] E(\zeta_1 - \zeta_3, \tau) E(\zeta_2 - \zeta_4, \tau) \quad (\text{B.5})$$

$$\times \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta_1 - \zeta_2 + \zeta_3 - \zeta_4}{2} + \frac{1}{4}, \tau \right) \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta_1 - \zeta_2 + \zeta_3 - \zeta_4}{2} - \frac{1}{4}, \tau \right),$$

$$\psi_{2,k} = -[\eta(2\tau)/\eta^2(\tau)] E(\zeta_1 - \zeta_4, \tau) E(\zeta_2 - \zeta_3, \tau)$$

$$\times \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta_1 - \zeta_2 - \zeta_3 + \zeta_4}{2} + \frac{1}{4}, \tau \right) \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta_1 - \zeta_2 - \zeta_3 + \zeta_4}{2} - \frac{1}{4}, \tau \right),$$

$$\psi_{3,k} = -[\eta(2\tau)/\eta^2(\tau)] E(\zeta_1 - \zeta_2, \tau) E(\zeta_3 - \zeta_4, \tau)$$

$$\times \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta_1 + \zeta_2 - \zeta_3 - \zeta_4}{2} + \frac{1}{4}, \tau \right) \theta \begin{bmatrix} k \\ 0 \end{bmatrix} \left(\frac{\zeta_1 + \zeta_2 - \zeta_3 - \zeta_4}{2} - \frac{1}{4}, \tau \right).$$

These quantities satisfy (B.3) as a consequence of Fay's trisecant identity [31, 32]

$$\begin{aligned} E(\zeta_1 - \zeta_3, \tau) E(\zeta_2 - \zeta_4, \tau) \theta \begin{bmatrix} a \\ b \end{bmatrix} (\zeta, \tau) \theta \begin{bmatrix} a \\ b \end{bmatrix} (\zeta + \zeta_1 + \zeta_3 - \zeta_2 - \zeta_4, \tau) = \\ E(\zeta_1 - \zeta_4, \tau) E(\zeta_2 - \zeta_3, \tau) \theta \begin{bmatrix} a \\ b \end{bmatrix} (\zeta + \zeta_1 - \zeta_2, \tau) \theta \begin{bmatrix} a \\ b \end{bmatrix} (\zeta + \zeta_3 - \zeta_4, \tau) \\ + E(\zeta_1 - \zeta_2, \tau) E(\zeta_3 - \zeta_4, \tau) \theta \begin{bmatrix} a \\ b \end{bmatrix} (\zeta + \zeta_1 - \zeta_4, \tau) \theta \begin{bmatrix} a \\ b \end{bmatrix} (\zeta + \zeta_3 - \zeta_2, \tau). \end{aligned} \quad (\text{B.6})$$

This identity has already been used in CFT in connection with the bosonization formulas on the torus [33]. To prove (B.3) we have to choose ζ in (B.6) as

$$\zeta = -\frac{\zeta_1 - \zeta_2 + \zeta_3 - \zeta_4}{2} + \frac{1}{4} \quad (\text{B.7})$$

and use (A.7).

References

- [1] Moore G and Read N, *Nonabelions in the fractional quantum Hall effect*, 1991 *Nucl. Phys. B* **360** 362
- [2] Cirac J I and Sierra G, *Infinite matrix product states, conformal field theory, and the Haldane-Shastry model*, 2010 *Phys. Rev. B* **81** 104431
- [3] Nielsen A E B, Cirac J I and Sierra G, *Laughlin spin-liquid states on lattices obtained from conformal field theory*, 2012 *Phys. Rev. Lett.* **108** 257206
- [4] Tu H-H, Nielsen A E B, Cirac J I and Sierra G, *Lattice Laughlin states of bosons and fermions at filling fractions $1/q$* , arXiv:1311.3958
- [5] Blok B and Wen X-G, *Many-body systems with non-abelian statistics*, 1992 *Nucl. Phys. B* **374** 615
- [6] Wen X-G, Wu Y-S and Hatsugai Y, *Chiral operator product algebra and edge excitations of a fractional quantum Hall droplet*, 1994 *Nucl. Phys. B* **422** 476
- [7] Nielsen A E B, Cirac J I and Sierra G, *Laughlin spin-liquid states on lattices obtained from conformal field theory*, 2011 *J. Stat. Mech.* P11014
- [8] Tu H-H, *Projected BCS states and spin Hamiltonians for the $SO(n)_1$ Wess-Zumino-Witten model*, 2013 *Phys. Rev. B* **87** 041103(R)

- [9] Kalmeyer V and Laughlin R B, *Equivalence of the resonating-valence-bond and fractional quantum Hall states*, 1987 *Phys. Rev. Lett.* **59** 2095
- [10] Kalmeyer V and Laughlin R B, *Theory of the spin liquid state of the Heisenberg antiferromagnet*, 1989 *Phys. Rev. B* **39** 11879
- [11] Laughlin R B, *Spin Hamiltonian for which quantum Hall wavefunction is exact*, 1989 *Ann. Phys.* **191** 163
- [12] Wen X G, Wilczek F and Zee A, *Chiral spin states and superconductivity*, 1989 *Phys. Rev. B* **39** 11413
- [13] Laughlin R B and Zou Z, *Properties of the chiral-spin-liquid state*, 1990 *Phys. Rev. B* **41** 664
- [14] Balatsky A V, *Relation between the chiral-spin-liquid state and the chiral $SU(2)$ Wess-Zumino-Witten model*, 1991 *Phys. Rev. B* **43** 1257
- [15] Wen X-G, *Gapless boundary excitations in the quantum Hall states and in the chiral spin states*, 1991 *Phys. Rev. B* **43** 11025
- [16] Barkeshli M, *Transitions between chiral spin liquids and Z_2 spin liquids*, arXiv:1307.8194
- [17] Nielsen A E B, Sierra G and Cirac J I, *Local models of fractional quantum Hall states in lattices and physical implementation*, 2013 *Nat. Commun.* **4** 2864
- [18] Schroeter D F, Kapit E, Thomale R and Greiter M, *Spin Hamiltonian for which the chiral spin liquid is the exact ground state*, 2007 *Phys. Rev. Lett.* **99** 097202
- [19] Thomale R, Kapit E, Schroeter D F and Greiter M, *Parent Hamiltonian for the chiral spin liquid*, 2009 *Phys. Rev. B* **80** 104406
- [20] Kapit E and Mueller E, *Exact parent Hamiltonian for the quantum Hall states in a lattice*, 2010 *Phys. Rev. Lett.* **105** 215303
- [21] Bauer B, Keller B P, Dolfi M, Trebst S and Ludwig A W W, *Gapped and gapless spin liquid phases on the Kagome lattice from chiral three-spin interactions*, arXiv:1303.6963
- [22] Di Francesco P, Mathieu P and Sénéchal D, *Conformal Field Theory*, 1997 Springer-Verlag, New York
- [23] Dijkgraaf R, Verlinde E and Verlinde H, *$C = 1$ Conformal Field Theories on Riemann Surfaces*, 1988 *Commun. Math. Phys.* **115** 649
- [24] Alvarez-Gaumé L, Bost J-B, Moore G, Nelson P and Vafa C, *Bosonization on higher genus Riemann surfaces*, 1987 *Comm. Math. Phys.* **112** 503
- [25] Haldane F D M and Rezayi E H, *Periodic Laughlin-Jastrow wave functions for the fractional quantized Hall effect*, 1985 *Phys. Rev. B* **31** 2529
- [26] Haldane F D M, *Many-particle translational symmetries of two-dimensional electrons at rational Landau-level filling*, 1985 *Phys. Rev. Lett.* **55** 2095
- [27] Read N and Rezayi E, *Quasiholes and fermionic zero modes of paired fractional quantum Hall states: The mechanism for non-Abelian statistics*, 1996 *Phys. Rev. B* **54** 16864
- [28] Sakurai J J, *Modern Quantum Mechanics*, 1994 Addison-Wesley Publishing Company
- [29] Hatsugai Y, *Characterization of topological insulators: Chern numbers for ground state multiplet*, 2005 *J. Phys. Soc. Jpn.* **74** 1374
- [30] Hafezi M, Sørensen A S, Demler E and Lukin M D, *Fractional quantum Hall effect in optical lattices*, 2007 *Phys. Rev. A* **76** 023613
- [31] Fay J D, *Theta functions on Riemann surfaces*, 1973 Springer Notes in Mathematics 352, Springer
- [32] Mumford D, *Tata Lectures on Theta*, 1983-1984 volumes I and II, Birkhäuser
- [33] Verlinde E and Verlinde H, *Chiral bosonization, determinants and the string partition function*, 1987 *Nucl. Phys. B* **288** 357