

Out-of-equilibrium Baryogenesis and SuperWIMP Dark Matter

Giorgio Arcadi and Laura Covi

Institut für theoretische Physik, Friedrich-Hund-Platz 1, D-37077 Göttingen, Germany

Marco Nardecchia

*DAMTP, CMS, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, United Kingdom
Cavendish Laboratory, University of Cambridge, JJ Thomson Avenue, Cambridge CB3 0HE, United Kingdom*

The experimental fact that the energy density in Dark Matter and in Baryons is of the same order is one of the most puzzling in cosmology. In this letter we suggest a new mechanism able to explain this coincidence in the context of out-of-equilibrium baryogenesis with DM production "à la" SuperWIMP starting from the same initial particle. We then discuss two simple implementations of this scenario within supersymmetric models with gravitino DM.

PACS numbers: 98.80.Cq

INTRODUCTION

It is an amazing coincidence that both Dark Matter and the baryonic energy densities are approximately within a factor five of each other [1]. Since normally the two densities are generated by very different mechanisms and at different scales, e.g. a WIMP mechanism below the electroweak scale and leptogenesis at high scale, it is usually not possible to understand theoretically why the numbers are not much wider apart. Since the pioneering work by Sakharov [2], it has been realized that obtaining a sufficiently large baryon number is usually a much more difficult task than just to produce Dark Matter: baryogenesis requires a sufficiently large violation of C, CP and baryon number violation and a departure from thermal equilibrium, while DM production can take place even in thermal equilibrium and without any quantum number violation. One would therefore expect the density of Dark Matter to be much less suppressed than the baryon asymmetry and indeed, in order to obtain the observed number density for Dark Matter, one usually has to rely on either a reduced number density, e.g. via a non-relativistic decoupling, or a very small mass for the Dark Matter particle, which is often in tension with being *Cold* Dark Matter.

A simple way to connect Dark and Baryonic Matter is to invoke for both types of matter an asymmetry, like it has been proposed in Asymmetric Dark Matter (ADM) models [3]. In that case, the asymmetries in the two species can be related, and then in the simplest realizations the ratio between the matter densities can be simply explained through a ratio of masses. Then it is expected that the Dark Matter has a mass not much heavier than the proton and must interact sufficiently strongly to erase the symmetric density component.

In a similar spirit, in this letter we would like to propose and explore another way to connect the baryon and DM generation relying on baryogenesis through out-of-equilibrium decay [4, 5] and the SuperWIMP mecha-

nism [6–8]. In such a case both matter densities are produced from an initial mother particle and they are naturally suppressed compared to its initial density, by the CP violation, needed to generate the baryon asymmetry, and by the branching ratio into DM respectively. These two suppression factors can be naturally of the same order of magnitude and explain why the baryon and Dark Matter densities turn out to be so similar. In general then the ratio between the DM and the baryonic energy densities is independent of the original mother particle density and given just by masses, the decay CP asymmetry and branching ratios.

THE BASIC MECHANISM

The mechanism we would like to propose is very simple and relies on the possibility of generating both the baryon asymmetry and Dark Matter via the out-of-equilibrium decay of the same particle X . Let us consider first baryogenesis. In general if a massive particle X decays out-of-equilibrium in two channels with different baryon number and with a non-vanishing C- and CP-violation, one can obtain from the decay the baryon number [4, 5]

$$\Omega_b = \xi_b \epsilon_{CP} \frac{m_p}{m_X} BR(X \rightarrow b, \bar{b}) \Omega_X \quad (1)$$

where ξ_b is a coefficient taking into account the possible effects of wash-out processes and baryon number dilution (e.g. via annihilation of massive particles into photons), m_p, m_X are the proton and decaying particle masses, Ω_X is the initial density of the X particle at departure of equilibrium and $BR(X \rightarrow b, \bar{b})$ gives the branching ratio for the decay into baryons and antibaryons. The CP-violation in the decay is taken into account by ϵ_{CP} given as:

$$\epsilon_{CP} = \frac{\Gamma(X \rightarrow b) - \Gamma(X \rightarrow \bar{b})}{\Gamma(X \rightarrow b) + \Gamma(X \rightarrow \bar{b})}. \quad (2)$$

In order to generate a sufficiently large Ω_b one needs a large initial number density of the X -particle since both

ξ and ϵ_{CP} tend to suppress the final baryon density. For the general case of WIMP-like decoupling, $\Omega_X \propto m_X^2$, this condition requires a heavy X -particle possibly above the TeV scale with suppressed annihilation channels.

In this setting, we consider also the decay of the X -particle into Dark Matter. The presence of this additional decay channel does not modify the mechanism of baryogenesis discussed above, as long as the branching ratio into DM is negligibly small. The decays of the particle X after its freeze-out produce a DM abundance through the SuperWIMP mechanism [6–9] as:

$$\Omega_{DM} = \xi_{DM} \frac{m_{DM}}{m_X} BR(X \rightarrow DM + \text{anything}) \Omega_X \quad (3)$$

where in this case ξ_{DM} just accounts for the possible dilution after DM production. Therefore due to the presence of the other decay channels, the DM density is suppressed by the corresponding branching ratio and is much smaller than the original X density. Note that the time-scale of the X -particle decay is set by the total decay rate Γ_{tot} avoiding problems with Big Bang Nucleosynthesis as long as $\Gamma_{tot}^{-1} < 1$ s.

We see that in this scenario we expect both Ω_b and Ω_{DM} to be suppressed by small numbers compared to Ω_X and we can obtain their ratio as

$$\frac{\Omega_b}{\Omega_{DM}} = \xi \epsilon_{CP} \frac{m_p}{m_{DM}} \frac{BR(X \rightarrow b, \bar{b})}{BR(X \rightarrow DM + \text{anything})}, \quad (4)$$

with $\xi = \xi_b/\xi_{DM}$, independently of the initial density of the particle X . So we can indeed obtain naturally $\Omega_b/\Omega_{DM} \sim 1/5$, if the masses of the DM and of the proton are of the same order and if the branching ratio of the decay of X into DM is strongly suppressed in comparison to the other channels and is of order ϵ_{CP} . Note that in order to explain the whole Dark Matter abundance as coming from the same particle that produces the baryon asymmetry, all the other mechanisms that could produce DM in the early universe have to be subdominant.

In the following we will discuss two different implementations of this mechanism in the context of supersymmetric models with gravitino DM. Indeed if the X particle is a superpartner, it has always a decay channel to gravitino LSP with Planck suppressed decay rate. We have indeed for a fermionic X :

$$\Gamma(X \rightarrow \tilde{G} + \text{anything}) = \frac{1}{48\pi} \frac{m_X^5}{m_{3/2}^2 M_{Pl}^2} \quad (5)$$

where M_{Pl} is the reduced Planck mass, $M_{Pl} = 2.43 \times 10^{18}$ GeV, M_X is the mother particle mass and $m_{3/2}$ the gravitino mass. It is in this case therefore easy to achieve a small branching ratio into gravitino of the order of ϵ_{CP} . Indeed the condition (4) allows for a gravitino lighter than the proton, but the requirement of proton stability against processes mediated by the X -particle, points towards a gravitino heavier than 1 GeV, corresponding to $\epsilon_{CP} \geq BR(X \rightarrow \tilde{G} + \text{anything})$.

A SIMPLE IMPLEMENTATION OF THE MECHANISM AT THE TEV SCALE

Let us consider one this mechanism at the TeV scale in the case when baryogenesis occurs in a supersymmetric model with R-parity violation [10–12]. The model is based on the MSSM extended with the baryon and R-parity violating superpotential

$$W_B = \lambda''_{ijk} U_i^c D_j^c D_k^c \quad (6)$$

where U^c, D^c represent the squark chiral multiplets and i, j, k are generation indices. Then a Bino-like neutralino, not the LSP, can decay into different R-parity conserving channels, in particular the dominant one into a gluino and a quark-antiquark pair and subdominant decays into photon/Z and gravitino, as well as into three quarks via the R-parity violating coupling. Then the Bino plays the role of the X particle with decay rates given by [11, 13]

$$\begin{aligned} \Gamma(\tilde{B} \rightarrow \tilde{g} q_i \bar{q}_i) &= \frac{Y_i^2 \alpha_1 \alpha_3}{16\pi} \frac{m_{\tilde{B}}^5}{m_0^4} \\ \Gamma(\tilde{B} \rightarrow u_i d_j d_k + \bar{u}_i \bar{d}_j \bar{d}_k) &= \frac{|\lambda''_{ijk}|^2 \alpha_1 (Y_u - Y_d)^2}{16\pi^2} \frac{m_{\tilde{B}}^5}{m_0^4} \\ \Gamma(\tilde{B} \rightarrow \tilde{G} + \text{anything}) &= \frac{1}{48\pi} \frac{m_{\tilde{B}}^5}{m_{3/2}^2 M_{Pl}^2} \end{aligned} \quad (7)$$

where α_1, α_3 are the gauge coupling strenghts of the $U(1)_Y$ and $SU(3)$ gauge groups, $Y_{u,d,q} = 2/3, -1/3, 1/6$ are the hypercharges of the quarks u_R, d_R, q_L , $m_{\tilde{B}}$ and m_0 denote the Bino's and a common scalar superpartner's masses. In the rate we have already summed over color states, while the sum over the final state flavours gives factors $N_{RPC} = 6Y_q^2 + 3Y_u^2 + 3Y_d^2 = 11/6$ instead of Y_i^2 in the R-parity conserving case and $N_{RPV} = 9(Y_u - Y_d)^2 = 9$ instead of $(Y_u - Y_d)^2$ for the RPV decay for flavour-universal λ'' . If some of the squarks are substantially heavier than the others, the rates get smaller and we can obtain the limiting cases with decoupled $\tilde{u}_R, \tilde{d}_R, \tilde{q}_L$ by setting Y_u, Y_d, Y_q equal to zero respectively. In order for the neutralino to decay out-of-equilibrium, the scalar superpartners have to be sufficiently heavy to satisfy the relation $\Gamma_{tot} < H(T = m_{\tilde{B}})$ but lighter than the SUSY breaking scale $\sqrt{m_{3/2} M_{Pl}}$ in order for the decay into gravitino to have a small branching ratio. For universal λ'' , we obtain from these two conditions the window of squark mass:

$$\sqrt{m_{3/2} M_{Pl}} \gg (3A)^{-1/4} m_0 > \frac{1}{2\sqrt{\pi}} \left(\frac{g_\star}{10}\right)^{-1/8} m_{\tilde{B}}^{3/4} M_{Pl}^{1/4} \quad (8)$$

with $A = \alpha_1(|\lambda''|^2/\pi N_{RPV} + \alpha_3 N_{RPC})$ and g_\star counting the effective relativistic degrees of freedom at the time when the neutralino becomes non-relativistic.

The RPV decay may have a non-vanishing CP asymmetry, due to the one-loop diagrams with on-shell gluino




FIG. 1: We show here in red (dark grey) the region where Ω_b is between 0.01-0.04 in the plane m_0 vs $m_{\tilde{B}}$ for $\lambda'' = 0.5$. The black line gives $\Omega_{DM} h^2 = 0.11$ for $m_{3/2} = 1$ GeV. We can see that we can obtain both quantities in the observed range for $m_0 \sim 5 - 6 \times 10^6$ GeV and $m_{\tilde{B}} \geq 6$ TeV. In the orange (light grey) region, the Bino annihilation and other wash-out processes are still active during the decay.

and quark. Assuming that the couplings contain different complex phases and that the intermediate down squarks dominate the process, the CP violation parameter can be estimated as [11, 14]

$$\epsilon_{CP} = \frac{1}{20} \text{Im} [e^{i\phi}] \alpha_3 \frac{m_{\tilde{B}}^2}{m_0^2} \quad (9)$$

where ϕ denotes an effective CP-violating phase. If the Bino density is sufficiently large, as it can be arranged for large μ and pure Bino, i.e. when the annihilation is mostly into Higgs bosons, a baryon asymmetry can be generated as [11]

$$\Omega_b \sim 10^{-2} \left(\frac{m_{\tilde{B}}}{1 \text{ TeV}} \right) \left(\frac{\mu}{10^{3/2} m_0} \right)^2 \left(\frac{\alpha_1 |\lambda''|^2 N_{RPV}}{\pi A} \right) \quad (10)$$

for $\text{Im} [e^{i\phi}] = 1$.

So we can obtain for the baryon to DM ratio simply

$$\frac{\Omega_b}{\Omega_{DM}} \sim \xi N_{RPV} \left(\frac{m_{3/2}}{1 \text{ GeV}} \right) \left(\frac{\lambda''}{0.1} \right)^2 \left(\frac{m_{\tilde{B}}}{1 \text{ TeV}} \right)^2 \left(\frac{m_0}{10^3 \text{ TeV}} \right)^{-6}. \quad (11)$$

From this expression it is clear that we need $m_0 \gtrsim 10^3 \text{ TeV}$ to match the observed baryon to DM ratio and that the gravitino mass cannot be far from the proton mass and, likely, values lower than 1 GeV are favored. Such a large scalar mass is also required to obtain the right baryon number [11, 14]. We show in Figure 1 the regions of parameter space with the correct baryon and




FIG. 2: We show here in the plane $m_{\tilde{B}}$ vs $m_{3/2}$ the lines where the ratio Ω_b/Ω_{DM} takes the observed value for different values of the scalar mass m_0 . The matching colored regions have also the observed value for the baryon number, assuming $\mu = 10^2 m_0$.

DM densities in the plane of m_0 vs $m_{\tilde{B}}$ for a fixed gravitino mass of 1 GeV. In Figure 2 we give the same regions in the plane $m_{\tilde{B}}$ vs $m_{3/2}$ for fixed values of m_0 and μ .

In order for this mechanism to work, one has to be sure that the contributions to the gravitino density from thermal scatterings [15] and FIMP [16, 17] mechanism from the heavy scalars are not larger than the Super-WIMP one. The easiest way to achieve that simultaneously is to require a sufficiently low reheating temperature that the scalar superpartners are not in thermal equilibrium $T_{RH} < m_0$, while the FIMP decay of the gauginos is less important. A detailed study of all the relevant processes will be the subject of a longer publication [14].

A SIMPLE IMPLEMENTATION OF THE MECHANISM AT A HIGH SCALE

Let us consider now another implementation of the mechanism, where instead baryogenesis happens via leptogenesis at a high scale [18]. In this case the decay of the lightest RH neutrino and its superpartner, the RH sneutrino, both produce a lepton asymmetry with similar values of ϵ_{CP} [19]. The lightest RH neutrino and sneutrino decay rates into leptons and antileptons and their superpartners are given by

$$\Gamma_{\tilde{\nu}, \tilde{\psi}} = \Gamma(N, \tilde{N} \rightarrow L + \bar{L}) = \frac{(\lambda^\dagger \lambda)_{11} M_{N, \tilde{N}}}{4\pi} \quad (12)$$

so that to have an out-of-equilibrium decay we need

$$M_{N, \tilde{N}} > \frac{3}{4\pi^2} \sqrt{\frac{10}{g_\star}} (\lambda^\dagger \lambda)_{11} M_{Pl} \geq 10^9 \text{ GeV}. \quad (13)$$

Note that barring cancellations the heaviest light neutrino mass is well approximate by $\tilde{m}_1 \sim (\lambda^\dagger \lambda)_{11} v_u^2 / M_N \leq 1 \text{ eV}$ [20]. The generated lepton asymmetry is given by

$$Y_\ell = \xi_l \left(\epsilon_{CP,N} Y_N + \epsilon_{CP,\tilde{N}} Y_{\tilde{N}} \right), \quad (14)$$

where again ξ_l takes into account possible effects of wash-out processes as well as the change in degrees of freedom from the leptogenesis epoch to today and $Y_i = n_i/s$ is the number density of the species i rescaled by the entropy density s . In the preferred region of strong wash-out, where the baryon asymmetry is independent of the initial conditions [20], we have $\xi_l \sim 10^{-4} - 10^{-5}$. This lepton asymmetry is finally transferred into a baryon asymmetry by sphaleron processes as

$$\Omega_b = c_{SF} m_p Y_\ell \quad (15)$$

where $c_{SF} = 8/23$ is the sphaleron factor relating the original $B - L$ number to the baryon number [21].

In this setting the production of gravitino Dark Matter can happen via RH sneutrino decay. If the degeneracy in mass between the RH sneutrino and neutrino is lifted, i.e. for $\Delta M_N^2 \equiv M_{\tilde{N}}^2 - M_N^2 > 0$ and $M_{\tilde{N}} - M_N \simeq \frac{\Delta M_N^2}{2M_N} > m_{3/2}$ the decay can proceed into RH neutrino and gravitino. This points towards large mass splitting ΔM_N^2 . The decay rate of the RH sneutrino into a RH neutrino and a gravitino is

$$\begin{aligned} \Gamma(\tilde{N} \rightarrow \tilde{G} + N) &= \frac{1}{48\pi} \frac{M_{\tilde{N}}^5}{m_{3/2}^2 M_{Pl}^2} \left(1 - \frac{M_N^2}{M_{\tilde{N}}^2} \right)^4 \\ &\sim \frac{1}{48\pi} \frac{\Delta M_N^8}{m_{3/2}^2 M_{Pl}^2 M_{\tilde{N}}^3} \end{aligned} \quad (16)$$

so that the decay is suppressed both by M_{Pl} and the available phase space. Indeed the decay is governed by the Goldstino coupling that vanishes in the limit of conserved SUSY. If the mass splitting is too small, the three-body decay into Higgs, lepton and gravitino can be more important. For such three-body decay, away from resonances in the intermediate RH neutrino or higgsino, taken into account in Eqs. (12), (16), we have

$$\begin{aligned} \Gamma(\tilde{N} \rightarrow \tilde{G} + \ell h) &= \frac{(\lambda^\dagger \lambda)_{11}}{1536\pi^3} \frac{M_{\tilde{N}} \Delta M_N^4}{m_{3/2}^2 M_{Pl}^2} \\ &\sim \frac{\Gamma_{\tilde{\ell}}}{384\pi^2} \frac{\Delta M_N^4}{m_{3/2}^2 M_{Pl}^2}. \end{aligned} \quad (17)$$

We see therefore that the 3-body decay dominates over the 2-body as soon as

$$\frac{\Delta M_N^4}{M_N^4} < \frac{(\lambda^\dagger \lambda)_{11}}{32\pi^2}. \quad (18)$$

The branching ratio of the RH sneutrino decay into

gravitino is then

$$BR(\tilde{N} \rightarrow \tilde{G}) = \frac{\Delta M_N^4}{384\pi^2 m_{3/2}^2 M_{Pl}^2} \left(1 + \frac{\Delta M_N^4}{M_N^4} \frac{32\pi^2}{(\lambda^\dagger \lambda)_{11}} \right) \quad (19)$$

and it can naturally be of the same order as $\epsilon_{CP} \simeq 10^{-6}$ even for small λ . Taking $0 \leq Y_N \sim Y_{\tilde{N}}$ and $\epsilon_{CP,N} \sim \epsilon_{CP,\tilde{N}} = \epsilon_{CP}$, we have then for the ratio between baryon and DM energy density the relation:

$$\begin{aligned} \frac{\Omega_b}{\Omega_{DM}} &= 768\pi^2 c_{SF} \xi \epsilon_{CP} \frac{m_p m_{3/2} M_{Pl}^2}{\Delta M_N^4} \\ &\times \left(1 + \frac{\Delta M_N^4}{M_N^4} \frac{32\pi^2}{(\lambda^\dagger \lambda)_{11}} \right)^{-1}. \end{aligned} \quad (20)$$

Assuming for the parameters the typical values from thermal leptogenesis [20], i.e. $M_{\tilde{N}} \sim 10^{10} \text{ GeV}$, $(\lambda^\dagger \lambda)_{11} \sim 10^{-5}$, $\epsilon_{CP} \sim 10^{-6}$, and the strong wash-out regime with $c_{sf} \xi \sim 10^{-3}$, we obtain the observed value of the ratio for intermediate values of the mass difference ΔM_N^2 , i.e.

$$\sqrt{\Delta M_N^2} \sim 1.2 \times 10^8 \text{ GeV} \left(\frac{m_{3/2}}{1 \text{ GeV}} \right)^{1/4}. \quad (21)$$

Note that the result is independent of the RH sneutrino mass since the 3-body decay into gravitino dominates and has the same dependence on that mass as the lepton violating decays. For larger gravitino masses instead the 2-body decay plays the dominant role and the relation changes to

$$\sqrt{\Delta M_N^2} \sim 3.3 \times 10^8 \text{ GeV} \left(\frac{m_{3/2}}{1 \text{ TeV}} \right)^{1/8} \left(\frac{M_{\tilde{N}}}{10^{10} \text{ GeV}} \right)^{1/2} \quad (22)$$

for $(\lambda^\dagger \lambda)_{11} \sim 10^{-5}$. We show in Figure 3 the curves for the observed value of Ω_b/Ω_{DM} in the plane ϵ_{CP} vs gravitino mass. For values $\sqrt{\Delta M_N^2} \sim 10^7 - 10^8 \text{ GeV}$ it is possible to match the observation for $\epsilon_{CP} \sim 10^{-6}$ as required by thermal leptogenesis. In case of non-thermal leptogenesis, the RH sneutrino and neutrino number densities can be larger and allow also for smaller values of ϵ_{CP} and therefore ΔM_N^2 .

In both cases we need a quite large mass splitting in the RH neutrino multiplet, larger than expected for the other SM superpartners, but still smaller than the RH sneutrino masses. Such a splitting could arise naturally if the RH sneutrino couples directly to the supersymmetry breaking sector.

In this case again, in order for the mechanism to work, we need to suppress all other production channels of gravitinos, which is not so simple at high temperature. Possibly the easiest way would be to consider a heavy gravitino that couples more weakly or to reduce the scale of leptogenesis and so also $M_{\tilde{N}}$ as in non-thermal leptogenesis [22, 23].




FIG. 3: Curves of correct values of the ratio Ω_b/Ω_{DM} in the plane ϵ vs $m_{3/2}$ for different values of the mass difference ΔM_N^2 . We take $M_{\tilde{N}} = M_N = 10^{10}$ GeV and $(\lambda^\dagger \lambda)_{11} \sim 10^{-5}$. A typical value $\epsilon_{CP} = 10^{-6}$ is given by the green (dashed) line.

CONCLUSIONS

We have considered the possibility of producing both DM and the baryon asymmetry from the out-of-equilibrium decay of a single particle. In the general case this allows to obtain densities of the same order since both the baryon and the DM densities are strongly suppressed, by the CP asymmetry or by the decay branching ratio respectively. We are in this way able to obtain the observed ratio just from fundamental parameters like masses and couplings, independently of the mother particle density.

The mechanism can in principle be embedded both in low and high scale baryogenesis models with gravitino DM, as we have discussed in sections 3 and 4, as long as the usual thermal gravitino production by scatterings or decays is sufficiently suppressed. In both the models we find that at least part of the supersymmetric spectrum must be quite heavy. In the first case the squarks are required to be much heavier than the gauginos in order to achieve the out-of-equilibrium condition and also the right branching ratio of the Bino into gravitino. In the case of leptogenesis, a sufficiently large mass splitting in the RH neutrino multiplet is needed, implying very heavy scalar superpartners if extended to the whole MSSM superpartners. Nevertheless the gravitino LSP and also other part of the spectrum can still reside below or at the TeV scale.

We would like to thank Yossi Nir, Riccardo Rattazzi,

Fabrizio Rompineve Sorbello and Damien George for useful discussions. G.A. thanks the CERN Theory Division and the LPTH Orsay for the warm hospitality during part of the completion of this work.

G.A. and L.C. acknowledge partial financial support by the EU FP7 ITN Invisibles (Marie Curie Actions, PITN-GA-2011-289442). L.C. acknowledges partial financial support by the German-Israeli Foundation for scientific research and development (GIF).

-
- [1] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5076 [astro-ph.CO].
 - [2] A. D. Sakharov, Pisma Zh. Eksp. Teor. Fiz. **5** (1967) 32 [JETP Lett. **5** (1967) 24] [Sov. Phys. Usp. **34** (1991) 392] [Usp. Fiz. Nauk **161** (1991) 61].
 - [3] For a review on asymmetric DM see K. M. Zurek, arXiv:1308.0338 [hep-ph].
 - [4] E. W. Kolb and S. Wolfram, Nucl. Phys. B **172** (1980) 224 [Erratum-ibid. B **195** (1982) 542].
 - [5] J. A. Harvey, E. W. Kolb, D. B. Reiss and S. Wolfram, Nucl. Phys. B **201** (1982) 16.
 - [6] L. Covi, J. E. Kim and L. Roszkowski, Phys. Rev. Lett. **82** (1999) 4180 [hep-ph/9905212].
 - [7] J. L. Feng, A. Rajaraman and F. Takayama, Phys. Rev. Lett. **91** (2003) 011302 [hep-ph/0302215].
 - [8] J. L. Feng, A. Rajaraman and F. Takayama, Phys. Rev. D **68** (2003) 063504 [hep-ph/0306024].
 - [9] G. Arcadi and L. Covi, JCAP **1308** (2013) 005 [arXiv:1305.6587 [hep-ph]].
 - [10] Y. Cui and R. Sundrum, Phys. Rev. D **87** (2013) 116013 [arXiv:1212.2973 [hep-ph]].
 - [11] Y. Cui, arXiv:1309.2952 [hep-ph].
 - [12] F. R. Sorbello, arXiv:1310.0840 [hep-ph].
 - [13] E. A. Baltz and P. Gondolo, Phys. Rev. D **57** (1998) 2969 [hep-ph/9709445].
 - [14] G. Arcadi, L. Covi and M. Nardecchia, work in progress.
 - [15] M. Bolz, A. Brandenburg and W. Buchmüller, Nucl. Phys. B **606** (2001) 518 [Erratum-ibid. B **790** (2008) 336] [hep-ph/0012052].
 - [16] L. J. Hall, K. Jedamzik, J. March-Russell and S. M. West, JHEP **1003** (2010) 080 [arXiv:0911.1120 [hep-ph]].
 - [17] C. Cheung, G. Elor and L. Hall, Phys. Rev. D **84** (2011) 115021 [arXiv:1103.4394 [hep-ph]].
 - [18] M. Fukugita and T. Yanagida, Phys. Lett. B **174** (1986) 45.
 - [19] L. Covi, E. Roulet and F. Vissani, Phys. Lett. B **384** (1996) 169 [hep-ph/9605319].
 - [20] W. Buchmüller, P. Di Bari and M. Plumacher, Annals Phys. **315** (2005) 305 [hep-ph/0401240].
 - [21] J. A. Harvey and M. S. Turner, Phys. Rev. D **42** (1990) 3344.
 - [22] G. F. Giudice, M. Peloso, A. Riotto and I. Tkachev, JHEP **9908** (1999) 014 [hep-ph/9905242].
 - [23] T. Asaka, H. B. Nielsen and Y. Takahashi, Nucl. Phys. B **647** (2002) 252 [hep-ph/0207023].