

ESTIMATES OF BANDS FOR LAPLACIANS ON PERIODIC EQUILATERAL METRIC GRAPHS

EVGENY KOROTYAEV AND NATALIA SABUROVA

ABSTRACT. We consider Laplacians on periodic equilateral metric graphs. The spectrum of the Laplacian consists of an absolutely continuous part (which is a union of an infinite number of non-degenerated spectral bands) plus an infinite number of flat bands, i.e., eigenvalues of infinite multiplicity. We estimate the Lebesgue measure of the bands on a finite interval in terms of geometric parameters of the graph. The proof is based on spectral properties of discrete Laplacians.

1. INTRODUCTION AND MAIN RESULTS

We consider metric Laplacians Δ_M on $\mathbb{Z}^d$ -periodic equilateral metric graphs (each edge has unit length). Such operators arise naturally as simplified models in mathematics, physics, chemistry, and engineering when one considers propagation of waves of various nature through a quasi-one-dimensional system that looks like a thin neighborhood of a graph.

It is well-known that the spectrum of the Laplacian Δ_M consists of an absolutely continuous part plus an infinite number of flat bands (i.e., eigenvalues with infinite multiplicity). These and other properties of Δ_M are discussed, e.g., in [BK12], [P12] and see references therein. The absolutely continuous spectrum consists of an infinite number of spectral bands separated by gaps. There is a known problem: to estimate the spectrum and gaps of the Laplacian on periodic metric graph. Note that in the case of the Schrödinger operators $-\Delta + Q$ with a periodic potential Q in $\mathbb{R}^d$ there are two-sided estimates of potentials in terms of gap lengths only at $d = 1$ in [K98], [K03]. We do not know other estimates. For the case of periodic graphs we know only two papers about estimates of spectrum and gaps:

(1) Lledó and Post [LP08] estimated the positions of spectral bands of Laplacians both on metric and discrete graphs in terms of eigenvalues of the operator on finite graphs (the so-called eigenvalue bracketing).

(2) Korotyaev and Saburova [KS14] considered Schrödinger operators on the discrete graphs and estimated the Lebesgue measure of their spectrum in terms of geometric parameters of the graph only.

Our main goal is to estimate the spectral bands and gaps for the Laplacian on the metric graph in terms of geometric parameters of the graph. Due to Cattaneo [C97], in order to study the spectrum of the Laplacian Δ_M on the equilateral metric graph it is enough to consider only the spectral interval $[0, \pi^2]$.

1.1. Metric Laplacians. Let $\Gamma = (V, \mathcal{E})$ be a connected graph, possibly having loops and multiple edges, where V is the set of its vertices and $\mathcal{E}$ is the set of its unoriented edges. The graphs under consideration are embedded into $\mathbb{R}^d$. An edge connecting vertices u and v from

Date: February 27, 2022.

Key words and phrases. spectral bands, flat bands, Laplace operator, periodic equilateral metric graph.

V will be denoted as the unordered pair $(u, v)_e \in \mathcal{E}$ and is said to be *incident* to the vertices. Vertices $u, v \in V$ will be called *adjacent* and denoted by $u \sim v$, if $(u, v)_e \in \mathcal{E}$. We define the degree κ_v of the vertex $v \in V$ as the number of all its incident edges from $\mathcal{E}$ (here a loop is counted twice). Below we consider locally finite $\mathbb{Z}^d$ -periodic metric equilateral graphs Γ , i.e., graphs satisfying the following conditions:

- 1) the number of vertices from V in any bounded domain $\subset \mathbb{R}^d$ is finite;
- 2) the degree of each vertex is finite;
- 3) there exists a basis $a_1, \dots, a_d$ in $\mathbb{R}^d$ such that Γ is invariant under translations through the vectors $a_1, \dots, a_d$:

$$\Gamma + a_s = \Gamma, \quad \forall s \in \mathbb{N}_d = \{1, \dots, d\}.$$

The vectors $a_1, \dots, a_d$ are called the *periods* of Γ .

- 4) All edges of the graph have the unit length.

In the space $\mathbb{R}^d$ we consider a coordinate system with the origin at some point O . The coordinate axes of this system are directed along the vectors $a_1, \dots, a_d$. Below the coordinates of all vertices of Γ will be expressed in this coordinate system. From the definition it follows that a $\mathbb{Z}^d$ -periodic graph Γ is invariant under translations through any integer vector:

$$\Gamma + m = \Gamma, \quad \forall m \in \mathbb{Z}^d.$$

Each edge $\mathbf{e}$ of Γ will be identified with the segment $[0, 1]$. This identification introduces a local coordinate $t \in [0, 1]$ along each edge. Thus, we give an orientation on the edge. Note that the spectrum of Laplacians on metric graphs does not depend on the orientation of graph edges. For each function y on Γ we define a function $y_{\mathbf{e}} = y|_{\mathbf{e}}$, $\mathbf{e} \in \mathcal{E}$. We identify each function $y_{\mathbf{e}}$ on $\mathbf{e}$ with a function on $[0, 1]$ by using the local coordinate $t \in [0, 1]$. Let $L^2(\Gamma)$ be the Hilbert space of all function $y = (y_{\mathbf{e}})_{\mathbf{e} \in \mathcal{E}}$, where each $y_{\mathbf{e}} \in L^2(0, 1)$, equipped with the norm

$$\|y\|_{L^2(\Gamma)}^2 = \sum_{\mathbf{e} \in \mathcal{E}} \|y_{\mathbf{e}}\|_{L^2(0,1)}^2 < \infty.$$

We define the metric Laplacian Δ_M on $L^2(\Gamma)$ by

$$(\Delta_M y)_{\mathbf{e}} = -y''_{\mathbf{e}}, \quad y = (y_{\mathbf{e}})_{\mathbf{e} \in \mathcal{E}},$$

where $(y''_{\mathbf{e}})_{\mathbf{e} \in \mathcal{E}} \in L^2(\Gamma)$ and y satisfies the so-called Kirchhoff conditions:

$$y \text{ is continuous on } \Gamma, \quad \sum_{\mathbf{e}=(v,u)_e \in \mathcal{E}} \delta_{\mathbf{e}}(v) y'_{\mathbf{e}}(v) = 0, \quad \forall v \in V, \quad (1.1)$$

$$\delta_{\mathbf{e}}(v) = \begin{cases} 1, & \text{if } v \text{ is a terminal vertex of the edge } \mathbf{e}, \text{ i.e. } t = 1 \text{ at } v, \\ -1, & \text{if } v \text{ is a initial vertex of the edge } \mathbf{e}, \text{ i.e. } t = 0 \text{ at } v. \end{cases}$$

1.2. Discrete Laplacians. Let $\ell^2(V)$ be the Hilbert space of all square summable functions $f : V \rightarrow \mathbb{C}$, equipped with the norm

$$\|f\|_{\ell^2(V)}^2 = \sum_{v \in V} |f(v)|^2 < \infty.$$

We define the self-adjoint normalized Laplacian (i.e., the Laplace operator) Δ on $\ell^2(V)$ by

$$(\Delta f)(v) = -\frac{1}{\sqrt{\kappa_v}} \sum_{(v,u)_e \in \mathcal{E}} \frac{1}{\sqrt{\kappa_u}} f(u), \quad v \in V, \quad f \in \ell^2(V), \quad (1.2)$$

where κ_v is the degree of the vertex $v \in V$ and all loops in the sum (1.2) are counted twice.

We recall the basic facts about the Laplacian Δ (see [Ch97], [HS04], [MW89]), which hold true for both finite and periodic graphs:

(i) the point -1 belongs to the spectrum $\sigma(\Delta)$ and $\sigma(\Delta)$ is contained in $[-1, 1]$, i.e.,

$$-1 \in \sigma(\Delta) \subset [-1, 1]. \quad (1.3)$$

(ii) On periodic graphs the points ± 1 are never eigenvalues of Δ .

We define the fundamental graph $\Gamma_f = (V_f, \mathcal{E}_f)$ of the periodic graph Γ as a graph on the surface $\mathbb{R}^d/\mathbb{Z}^d$ by

$$\Gamma_f = \Gamma/\mathbb{Z}^d \subset \mathbb{R}^d/\mathbb{Z}^d. \quad (1.4)$$

The fundamental graph Γ_f has the vertex set V_f and the set $\mathcal{E}_f$ of unoriented edges, which are finite. We identify the vertices of the fundamental graph $\Gamma_f = (V_f, \mathcal{E}_f)$ with the vertices of the graph $\Gamma = (V, \mathcal{E})$ from the set $[0, 1)^d$ by

$$V_f = [0, 1)^d \cap V = \{v_1, \dots, v_\nu\}, \quad \nu = \#V_f < \infty, \quad (1.5)$$

where ν is the number of vertices of Γ_f . Denote by $\mathcal{B}$ the set of all edges of Γ connecting the vertices from V_f with the vertices from $V \setminus V_f$. These edges will be called *bridges*.

The discrete Laplacian Δ on $\ell^2(V)$ has the standard decomposition into a constant fiber direct integral by

$$\ell^2(V) = \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d}^{\oplus} \ell^2(V_f) d\vartheta, \quad U\Delta U^{-1} = \frac{1}{(2\pi)^d} \int_{\mathbb{T}^d}^{\oplus} \Delta(\vartheta) d\vartheta, \quad (1.6)$$

$\mathbb{T}^d = \mathbb{R}^d/(2\pi\mathbb{Z})^d$, for some unitary operator U . Here $\ell^2(V_f) = \mathbb{C}^\nu$ is the fiber space and $\Delta(\vartheta)$ is the Floquet fiber $\nu \times \nu$ matrix. The precise form of the matrix $\Delta(\vartheta) = \{\Delta_{jk}(\vartheta)\}_{j,k=1}^\nu$ was determined in [KS14].

Each Floquet matrix $\Delta(\vartheta)$, $\vartheta \in \mathbb{T}^d$, has ν real eigenvalues $\lambda_n(\vartheta)$, $n \in \mathbb{N}_\nu$, labeled by

$$\lambda_1(\vartheta) \leq \dots \leq \lambda_\nu(\vartheta). \quad (1.7)$$

Each $\lambda_n(\cdot)$, $n \in \mathbb{N}_\nu$, is a continuous function on the torus $\mathbb{T}^d$ and creates the spectral band $\sigma_n(\Delta)$ given by

$$\sigma_n = \sigma_n(\Delta) = [\lambda_n^-, \lambda_n^+] = \lambda_n(\mathbb{T}^d). \quad (1.8)$$

Note that if $\lambda_n(\cdot) = C_n = \text{const}$ on some set $\mathcal{B} \subset \mathbb{T}^d$ of positive Lebesgue measure, then the operator Δ on Γ has the eigenvalue C_n with infinite multiplicity. We call C_n a *flat band*. The spectrum of the Laplace operator Δ on the periodic graph Γ has the form

$$\sigma(\Delta) = \bigcup_{n=1}^\nu \sigma_n(\Delta) = \sigma_{ac}(\Delta) \cup \sigma_{fb}(\Delta). \quad (1.9)$$

Here $\sigma_{ac}(\Delta)$ is the absolutely continuous spectrum, which is a union of non-degenerated intervals from (1.8), and $\sigma_{fb}(\Delta) = \{\mu_1, \dots, \mu_r\}$, $r < \nu$, is the set of all flat bands (eigenvalues of infinite multiplicity). An open interval between two neighboring non-degenerated bands is called a *gap*. Let $\lambda_{\nu_1}^+$, $\nu_1 \leq \nu$, be the upper point of the absolutely continuous spectrum of the operator Δ . It is convenient for us an open interval $(\lambda_{\nu_1}^+, 1)$ also to call a gap of the operator Δ .

1.3. Main results. Instead of the Laplacian $\Delta_M \geq 0$ it is convenient for us to define the momentum operator $\sqrt{\Delta_M} \geq 0$. Due to Cattaneo Theorem (see Section 2) the spectrum of the operator $\sqrt{\Delta_M} \geq 0$ on Γ has the form

$$\sigma(\sqrt{\Delta_M}) = \sigma_{ac}(\sqrt{\Delta_M}) \cup \sigma_{fb}(\sqrt{\Delta_M}). \quad (1.10)$$

Both the sets $\sigma_{ac}(\sqrt{\Delta_M})$ and $\sigma_{fb}(\sqrt{\Delta_M})$ are 2π -periodic on the half-line $(0, \infty)$ and are symmetric on the interval $(0, 2\pi)$ with respect to the point π . Thus, in order to study Δ_M it is sufficient to study its restriction Ω on the spectral interval $[0, \pi]$ given by

$$\Omega = \sqrt{\Delta_M} \chi_{[0, \pi]}(\sqrt{\Delta_M}), \quad (1.11)$$

where $\chi_A(\cdot)$ is the characteristic function of the set A . Due to Cattaneo Theorem (see Section 2) the spectrum of the operator Ω on a periodic metric graph Γ has the form

$$\begin{aligned} \sigma(\Omega) &= \bigcup_{n=1}^{\nu} \sigma_n(\Omega) = \sigma_{ac}(\Omega) \cup \sigma_{fb}(\Omega), \\ \sigma_n(\Omega) &= [z_n^-, z_n^+], \quad -\cos(z_n^\pm) = \lambda_n^\pm, \quad n \in \mathbb{N}_\nu. \end{aligned} \quad (1.12)$$

Here $\sigma_{ac}(\Omega)$ is a union of non-degenerated spectral bands $\sigma_n(\Omega)$ with $z_n^- < z_n^+$ and $\sigma_{fb}(\Omega)$ is the flat band spectrum (for more details see Section 2).

Now we formulate our main results. Let

$$\beta = \sum_{n=1}^{\nu} \frac{\beta_n}{\varkappa_n}, \quad (1.13)$$

β_n is the bridge degree (the number of bridges incident to v_n) and $\varkappa_n$ is the degree of $v_n \in \Gamma_f$.

Theorem 1.1. *i) All spectral bands $\sigma_n(\Omega)$ and $\sigma_n(\Delta)$, $n \in \mathbb{N}_\nu$, of the momentum operator Ω and the discrete Laplacian Δ , respectively, satisfy*

$$|\sigma_n(\Delta)| \leq |\sigma_n(\Omega)| \leq \frac{\pi}{\sqrt{2}} |\sigma_n(\Delta)|^{\frac{1}{2}}. \quad (1.14)$$

ii) The Lebesgue measure $|\sigma(\Omega)|$ and $|\sigma(\Delta)|$ of the spectrum of Ω and Δ , respectively, satisfies

$$|\sigma(\Delta)| \leq |\sigma(\Omega)| \leq \frac{\pi}{\sqrt{2}} |\sigma(\Delta)|^{\frac{1}{2}} \leq \pi\sqrt{\beta}, \quad (1.15)$$

where β is defined by (1.13). Moreover, if there exist spectral gaps $\gamma_1(\Omega), \dots, \gamma_s(\Omega)$, in the spectrum $\sigma(\Omega)$, then the following estimate holds true:

$$\sum_{n=1}^s |\gamma_n(\Omega)| \geq \pi(1 - \sqrt{\beta}). \quad (1.16)$$

iii) Let $\beta < 1$. Then the spectrum $\sigma(\Delta_M)$ has infinitely many gaps.

Remark. 1) The estimate $|\sigma(\Omega)| \leq \pi\sqrt{\beta}$ is not trivial iff $\beta < 1$. The condition $\beta < 1$ holds true when the number of bridges in each vertex $v \in V_f$ is sufficiently small compared to the degree of the vertex. If we change the coordinate system then the spectrum of the operator does not change, but the number of bridges does, in general. In order to get the best estimate in (1.15) we have to choose a coordinate system when the number β is minimal.

2) The Bethe-Sommerfeld conjecture states that each Schrödinger operator $-\Delta + Q$ with a periodic potential Q in $\mathbb{R}^d$, $d \geq 2$ has only finitely many gaps in the spectrum. This conjecture was proved by Skriganov [S85]. On an equilateral metric graph the spectrum of the Laplacian

Δ_M has no gaps iff $\sigma(\Delta) = [-1, 1]$. If $\sigma(\Delta) \neq [-1, 1]$, then in the spectrum of the Laplacian Δ_M on a quantum graph there exist **infinitely many gaps** $\gamma_1, \gamma_2, \dots$ and $|\gamma_n| \rightarrow \infty$ as $n \rightarrow \infty$.

Definition of Loop Graphs. *i) A periodic graph Γ is called a loop graph if each bridge $\mathbf{e}$ has the form $\mathbf{e} = (v_j, v_j + \tau(\mathbf{e}))$ for some $j \in \mathbb{N}_\nu$ and $\tau(\mathbf{e}) \in \mathbb{Z}^d$.*

ii) A loop graph Γ is called a precise loop graph if $\cos\langle\tau(\mathbf{e}), \vartheta_0\rangle = -1$ for all bridges $\mathbf{e} \in \mathcal{B}$ and some $\vartheta_0 \in \mathbb{T}^d$. This point ϑ_0 is called a precise point of the loop graph Γ .

Remark. 1) If $\langle\vartheta_0, \tau(\mathbf{e})\rangle/\pi$ is odd for all bridges $\mathbf{e} \in \mathcal{B}$ and some vector $\vartheta_0 \in \{0, \pi\}^d$, then ϑ_0 is a precise point of the loop graph Γ .

2) The class of all precise loop graphs is large enough. The simplest example of precise loop graphs is the d -dimensional lattice. More complicated examples are discussed in Proposition 2.3 in [KS13].

3) There exists a loop graph, which is not a precise loop graph. The simplest example of such graph is the triangular lattice (see Proposition 2.3 in [KS13]).

We now describe all bands for precise loop periodic graphs.

Theorem 1.2. *i) Let Γ be a loop graph. Then the spectral bands $\sigma_n(\Omega) = [z_n^-, z_n^+]$ of the operator Ω satisfy*

$$-\cos(z_n^-) = \lambda_n(0) = \lambda_n^-, \quad \forall n \in \mathbb{N}_\nu. \quad (1.17)$$

ii) Let, in addition, Γ be precise with a precise point $\vartheta_0 \in \mathbb{T}^d$. Then

$$-\cos(z_n^+) = \lambda_n(\vartheta_0) = \lambda_n^+, \quad \forall n \in \mathbb{N}_\nu, \quad (1.18)$$

$$2\beta = \sum_{n=1}^\nu |\sigma_n(\Delta)| \leq \sum_{n=1}^\nu |\sigma_n(\Omega)|, \quad (1.19)$$

where β is defined by (1.13).

We present the plan of our paper. In section 2 we estimate the Lebesgue measure of the spectrum of the operator $\sqrt{\Delta_M}$ on the finite interval $[0, \pi]$ in terms of geometric parameters of the graph and discuss some spectral properties of metric Laplacians on loop graphs and bipartite periodic graphs. In Appendix we collect spectral properties of the discrete Laplacian from [KS14], needed to prove our main results.

2. PROOFS OF THE MAIN THEOREMS

2.1. Cattaneo Correspondence. Cattaneo obtained a correspondence between the spectrum of the Laplacian Δ_M on the equilateral metric graph and the spectrum of the Laplacian Δ on the corresponding discrete graph [C97]. For the sake of completeness and the reader's convenience we recall this correspondence.

Consider the eigenvalues problem with Dirichlet boundary conditions

$$-y'' = Ey, \quad y(0) = y(1) = 0. \quad (2.1)$$

It is known that the spectrum of this problem is given by $\sigma_D = \{(\pi n)^2 : n \in \mathbb{N}\}$. Here $(\pi n)^2$ is the so-called Dirichlet eigenvalue of the problem (2.1).

We formulate Cattaneo's result [C97] in the form convenient for us. This theorem gives a basis for describing the spectrum of the operator Δ_M in terms of Δ , and conversely.

Theorem (Cattaneo) *i) The spectrum of the operator $\sqrt{\Delta_M} \geq 0$ on the periodic metric graph Γ has the form*

$$\sigma(\sqrt{\Delta_M}) = \sigma_{ac}(\sqrt{\Delta_M}) \cup \sigma_{fb}(\sqrt{\Delta_M}), \quad (2.2)$$

$$\sigma_{ac}(\sqrt{\Delta_M}) = \{z \in \mathbb{R}_+ : -\cos z \in \sigma_{ac}(\Delta)\}, \quad (2.3)$$

$$\sigma_{fb}(\sqrt{\Delta_M}) = \{z \in \mathbb{R}_+ : -\cos z \in \sigma_{fb}(\Delta)\} \cup \{\pi n : n \in \mathbb{N}\}. \quad (2.4)$$

- ii) Each flat band $2\pi n$, $n \in \mathbb{N}$, is embedded in the absolutely continuous spectrum $\sigma_{ac}(\sqrt{\Delta_M})$.
 iii) Both the sets $\sigma_{ac}(\sqrt{\Delta_M})$ and $\sigma_{fb}(\sqrt{\Delta_M})$ are 2π -periodic on the half-line $(0, \infty)$ and are symmetric on the interval $(0, 2\pi)$ with respect to the point π .
 iv) The spectrum of the operator Ω on a periodic metric graph Γ has the form*

$$\begin{aligned} \sigma(\Omega) &= \bigcup_{n=1}^{\nu} \sigma_n(\Omega) = \sigma_{ac}(\Omega) \cup \sigma_{fb}(\Omega), \\ \sigma_n(\Omega) &= [z_n^-, z_n^+], \quad -\cos(z_n^\pm) = \lambda_n^\pm, \quad n \in \mathbb{N}_\nu. \end{aligned} \quad (2.5)$$

Here $\sigma_{ac}(\Delta)$ is a union of non-degenerated spectral bands $\sigma_n(\Omega)$ with $z_n^- < z_n^+$. Moreover, the first spectral band $\sigma_1(\Omega) = [0, z_1^+]$ is open. The flat band spectrum has the form

$$\sigma_{fb}(\Omega) = \{z_1, \dots, z_r, \pi\}, \quad -\cos(z_k) = \mu_k \neq 1, \quad k \in \mathbb{N}_r. \quad (2.6)$$

v) $\sigma(\Omega) = [0, \pi]$ iff $\sigma(\Delta) = [-1, 1]$.

vi) The spectrum of the operator Ω has exactly k gaps iff $\sigma(\Delta)$ has exactly k gaps.

Remark. 1) Cattaneo considered the Laplacian Δ_M on connected locally finite graphs (including periodic). In general, some points of the Dirichlet spectrum σ_D are not flat bands of the Laplacian Δ_M .

2) Von Below [B85] considered the Laplacian Δ_M on connected finite graphs when the number of edges is greater than number of vertices. For each $n \in \mathbb{N}$ he constructed an eigenfunction with the eigenvalue $(\pi n)^2$. Without changing the proof this construction can be applied to a connected $\mathbb{Z}^d$ -periodic graph ($d \geq 2$) to obtain an eigenfunction with a compact support and the eigenvalue $(\pi n)^2$, $n \in \mathbb{N}$. Thus, $\sigma_D \subset \sigma_{fb}(\Delta_M)$ (see (2.4)). Note that for $\mathbb{Z}$ -periodic graph some (all) points of the Dirichlet spectrum σ_D may not be flat bands of the operator.

3) The relation between the spectra of Δ and $\sqrt{\Delta_M}$ is shown in Fig.1.


4) The flat bands πn , $n \in \mathbb{N}$, of the operator $\sqrt{\Delta_M}$ will be called *Dirichlet flat bands*.

5) The number of flat bands of the operator Ω is $r + 1$. Flat bands $z_1, \dots, z_r$ corresponds to r flat bands of the discrete Laplacian and the flat band π is a Dirichlet flat band.

6) Let $z_{\nu_1}^+$, $\nu_1 \leq \nu$, be the upper point of the absolutely continuous spectrum of the operator Ω . If $z_{\nu_1}^+ < \pi$, then there is the gap $(z_{\nu_1}^+, 2\pi - z_{\nu_1}^+)$ in the spectrum of $\sqrt{\Delta_M}$. For the convenience the interval $(z_{\nu_1}^+, \pi)$ is also called a gap of the operator Ω .

Proof of Theorem 1.1. i) Consider the spectral band $\sigma_n(\Delta) = [\lambda_n^-, \lambda_n^+]$ of the discrete Laplacian Δ for some $n \in \mathbb{N}_\nu$. Due to Cattaneo Theorem.iv the corresponding spectral band $\sigma_n(\Omega)$ of the momentum operator Ω has the form

$$\sigma_n(\Omega) = [z_n^-, z_n^+], \quad \text{where} \quad \cos z_n^\pm = -\lambda_n^\pm. \quad (2.7)$$

FIGURE 1. Relation between the spectra of Δ and $\sqrt{\Delta_M}$.

Applying Proposition 3.1.i to the spectral band $\sigma_n(\Delta)$ and its preimage $\sigma_n(\Omega)$ under the function $\phi(z) = -\cos z$, $z \in [0, \pi]$, we obtain (1.14).

ii) The spectrum of the discrete Laplacian, defined by (1.9), is a union of non-overlapping segments $\mathfrak{S}_1, \dots, \mathfrak{S}_{\nu_*}$, where $\nu_* \leq \nu$, i.e.,

$$\sigma(\Delta) = \bigcup_{n=1}^{\nu_*} \mathfrak{S}_n, \quad \mathfrak{S}_m \cap \mathfrak{S}_n = \emptyset, \quad m \neq n. \quad (2.8)$$

Then, due to Cattaneo Theorem.iv, the spectrum of the operator Ω has the form

$$\sigma(\Omega) = \bigcup_{n=1}^{\nu_*} \phi^{-1}(\mathfrak{S}_n), \quad \text{where} \quad \phi^{-1}(\mathfrak{S}_m) \cap \phi^{-1}(\mathfrak{S}_n) = \emptyset, \quad m \neq n. \quad (2.9)$$

The identities (2.8), (2.9) give

$$|\sigma(\Delta)| = \sum_{n=1}^{\nu_*} |\mathfrak{S}_n|, \quad |\sigma(\Omega)| = \sum_{n=1}^{\nu_*} |\phi^{-1}(\mathfrak{S}_n)|. \quad (2.10)$$

Applying Proposition 3.1.i to each segment $\mathfrak{S}_n$, $n \in \mathbb{N}_{\nu_*}$, we obtain

$$|\mathfrak{S}_n| \leq |\phi^{-1}(\mathfrak{S}_n)|, \quad \forall n \in \mathbb{N}_{\nu_*}. \quad (2.11)$$

Summing these inequalities and using (2.10), we obtain

$$|\sigma(\Delta)| = \sum_{n=1}^{\nu_*} |\mathfrak{S}_n| \leq \sum_{n=1}^{\nu_*} |\phi^{-1}(\mathfrak{S}_n)| = |\sigma(\Omega)|, \quad (2.12)$$

which gives the first inequality in (1.15).

We show the second inequality in (1.15). Let $\mathfrak{S}_* \subset [-1, 1]$ be the subset of the segment $[-1, 1]$ with the Lebesgue measure $|\mathfrak{S}_*| = |\sigma(\Delta)|$, having the form

$$\mathfrak{S}_* = [-1, -\lambda_*] \cup [\lambda_*, 1], \quad \lambda_* = 1 - \frac{|\sigma(\Delta)|}{2}, \quad 0 \leq \lambda_* < 1. \quad (2.13)$$

The preimage of the subset $\mathfrak{S}_*$ under the function $\phi(z) = -\cos z$, $z \in [0, \pi]$, is given by

$$\phi^{-1}(\mathfrak{S}_*) = [0, z_*] \cup [\pi - z_*, \pi], \quad \lambda_* = \cos z_*, \quad 0 < z_* \leq \frac{\pi}{2}. \quad (2.14)$$

Combining Cattaneo Theorem.iv, Proposition 3.1 and the identity (2.13), we obtain

$$|\sigma(\Omega)| = |\phi^{-1}(\sigma(\Delta))| \leq |\phi^{-1}(\mathfrak{S}_*)| \leq \frac{\pi}{\sqrt{2}} |\sigma(\Delta)|^{\frac{1}{2}}.$$

Thus, the second inequality in (1.15) has been proved. The last inequality in (1.15) follows from Theorem 3.2.i.

Now we will prove (1.16). Since $\sigma(\Omega) \subset [0, \pi]$, we have

$$\sum_{n=1}^s |\gamma_n(\Omega)| = \pi - |\sigma(\Omega)| \geq \pi(1 - \sqrt{\beta}). \quad (2.15)$$

Here we have used the estimate (1.15).

iii) Let $\beta < 1$. Then, due to (1.15) we have $|\sigma(\Omega)| < \pi$, which yields that in the spectrum of Ω there exists a gap. Hence, due to the periodicity of the spectrum of $\sqrt{\Delta_M}$, the spectrum of Δ_M has infinitely many gaps.

Now we give another proof of item iii). If $\beta < 1$, then Theorem 3.2.i gives that $|\sigma(\Delta)| < 2$, i.e., $\sigma(\Delta) \neq [-1, 1]$, which, by Cattaneo Theorem.iii,v, yields that the spectrum of Δ_M has infinitely many gaps. ■

Proof of Theorem 1.2. i) Let Γ be a loop graph. Then, due to Theorem 3.3.i, the spectral bands $\sigma_n(\Delta) = [\lambda_n^-, \lambda_n^+]$ of the Laplacian Δ satisfy $\lambda_n^- = \lambda_n(0)$, $\forall n \in \mathbb{N}_\nu$. This and (2.5) give that the spectral bands $\sigma_n(\Omega) = [z_n^-, z_n^+]$ of the operator Ω satisfy (1.17).

ii) Let Γ be a precise loop graph with a precise point $\vartheta_0 \in \mathbb{T}^d$. Cattaneo Theorem.iv and formula (3.17) give (1.18).

Combining (3.17), (3.18) and (1.18) with the simple estimate $\cos z_n^- - \cos z_n^+ \leq z_n^+ - z_n^-$, we obtain

$$2\beta = \sum_{n=1}^{\nu} |\sigma_n(\Delta)| = \sum_{n=1}^{\nu} (\lambda_n(\vartheta_0) - \lambda_n(0)) = \sum_{n=1}^{\nu} (\cos z_n^- - \cos z_n^+) \leq \sum_{n=1}^{\nu} (z_n^+ - z_n^-) = \sum_{n=1}^{\nu} |\sigma_n(\Omega)|.$$

Thus, (1.19) has been proved. ■

A graph is called *bipartite* if its vertex set is divided into two disjoint sets (called *parts* of the graph) such that each edge connects vertices from distinct sets. It is known ([BKS13], [HS04]) that *a periodic graph is bipartite $\Leftrightarrow$ the point $1 \in \sigma(\Delta) \Leftrightarrow$ the spectrum $\sigma(\Delta)$ is symmetric with respect to the point 0.*

Now we formulate some simple spectral properties of the Laplacian Δ_M and the momentum operator Ω on bipartite graphs.

Theorem 2.1. *The following statements hold true.*

i) *If Γ is bipartite, then the flat bands $\pi^2(2n+1)^2$, $n \in \mathbb{N}$, of the Laplacian Δ_M are embedded in $\sigma_{ac}(\Delta_M)$. If Γ is non-bipartite, then each flat band $\pi^2(n+1)^2$, $n \in \mathbb{N}$, lies in a gap.*

ii) *On the interval $(0, \pi)$ the spectrum of Ω is symmetric with respect to the point $\pi/2$ iff Γ is bipartite iff the point $\pi \in \sigma_{ac}(\Omega)$.*

iii) *Let a fundamental graph Γ_f be bipartite. If ν is odd, then $z = \pi/2$ is a flat band of Ω .*

iv) Let Γ be a loop bipartite graph (Γ_f is not bipartite, since there is a loop on Γ_f). Then each spectral band of the operator Ω on Γ has the form $\sigma_n(\Omega) = [z_n^-, z_n^+]$, $n \in \mathbb{N}_\nu$, where

$$-\cos z_n^- = \lambda_n(0), \quad \cos z_n^+ = \lambda_{\nu-n+1}(0). \quad (2.16)$$

Proof. i) If Γ is bipartite, then $1 \in \sigma(\Delta)$. Since 1 is never a flat band, the point $1 \in \sigma_{ac}(\Delta)$. Then (2.3) gives that $\pi^2(n+1)^2 \in \sigma_{ac}(\Delta_M)$ for all $n \in \mathbb{N}$. Similarly, if Γ is non-bipartite, then $1 \notin \sigma_{ac}(\Delta)$ and $\pi^2(n+1)^2 \notin \sigma_{ac}(\Delta_M)$ for all $n \in \mathbb{N}$. Thus, each flat band $\pi^2(n+1)^2$, $n \in \mathbb{N}$, lies in a gap.

ii) Firstly, we show that the graph Γ is bipartite iff the point $\pi \in \sigma_{ac}(\Omega)$. Indeed, the graph Γ is bipartite iff the point $1 \in \sigma_{ac}(\Delta)$. Due to (2.3), the condition $1 \in \sigma_{ac}(\Delta)$ is equivalent to $\pi \in \sigma_{ac}(\Omega)$.

Secondly, we prove that on the interval $(0, \pi)$ the spectrum of Ω is symmetric with respect to the point $\pi/2$ iff Γ is bipartite. Let Γ be bipartite. Then the spectrum of the discrete Laplacian Δ is symmetric with respect to 0. Assume that $z \in (0, \pi)$ and $z \in \sigma(\Omega)$. Then Cattaneo Theorem gives that $-\cos z \in \sigma(\Delta)$. Due to the symmetry of the spectrum $\sigma(\Delta)$, the point $-\cos(\pi - z) = \cos z \in \sigma(\Delta)$. This and (2.3) yield that $\pi - z \in \sigma(\Omega)$.

Conversely, let on the interval $(0, \pi)$ the spectrum of Ω be symmetric with respect to the point $\pi/2$. Since $-1 \in \sigma_{ac}(\Delta)$, due to Cattaneo Theorem $0 \in \sigma_{ac}(\Omega)$. Then the symmetry of $\sigma(\Omega)$ gives that $\pi \in \sigma_{ac}(\Omega)$ and Γ is bipartite.

iii) Let a fundamental graph Γ_f be bipartite and let ν be odd. By Theorem 3.2.ii, $\mu = 0$ is a flat band of the discrete Laplacian Δ on Γ . This and Cattaneo Theorem.iv yield that $z = \pi/2$ is a flat band of Ω .

iv) By Theorem 3.3.iii each spectral band of the discrete Laplacian on Γ has the form $\sigma_n(\Delta) = [\lambda_n^-, \lambda_n^+]$, $n \in \mathbb{N}_\nu$, where $\lambda_n^\pm$ are the eigenvalues of the matrix $\mp \Delta(0)$, i.e., $\lambda_n^- = \lambda_n(0)$, $\lambda_n^+ = -\lambda_{\nu-n+1}(0)$. Then Cattaneo Theorem.iv gives that each spectral band of the operator Ω has the form $\sigma_n(\Omega) = [z_n^-, z_n^+]$, $n \in \mathbb{N}_\nu$, where $-\cos z_n^- = \lambda_n^- = \lambda_n(0)$, $-\cos z_n^+ = \lambda_n^+ = -\lambda_{\nu-n+1}(0)$. ■

Remark. Item iii) gives a simple sufficient condition for existence of the flat band.

3. APPENDIX

We formulate some simple facts needed to prove our main results.

Proposition 3.1. *Let $\phi(z) = -\cos z$, $z \in [0, \pi]$. Then the following statements hold true.*

i) *If $\mathfrak{S} = [\lambda^-, \lambda^+] \subset [-1, 1]$, then*

$$|\mathfrak{S}| \leq |\phi^{-1}(\mathfrak{S})| \leq \frac{\pi}{\sqrt{2}} |\mathfrak{S}|^{\frac{1}{2}}. \quad (3.1)$$

ii) *If $\mathfrak{S} \subset [-1, 1]$ is any subset of the segment $[-1, 1]$ and $\mathfrak{S}_* \subset [-1, 1]$ is the subset with the same Lebesgue measure $|\mathfrak{S}_*| = |\mathfrak{S}|$, having the form*

$$\mathfrak{S}_* = [-1, -\lambda_*] \cup [\lambda_*, 1], \quad \lambda_* = 1 - \frac{1}{2} |\mathfrak{S}|, \quad (3.2)$$

then

$$|\phi^{-1}(\mathfrak{S})| \leq |\phi^{-1}(\mathfrak{S}_*)| \leq \frac{\pi}{\sqrt{2}} |\mathfrak{S}|^{\frac{1}{2}}. \quad (3.3)$$

Proof. i) The function ϕ is an increasing bijection of the segment $[0, \pi]$ onto the segment $[-1, 1]$. The preimage of the segment $\mathfrak{S} = [\lambda^-, \lambda^+] \subset [-1, 1]$ under the function ϕ has the

form

$$\phi^{-1}(\mathfrak{S}) = [z^-, z^+], \quad \text{where } z^\pm = \phi^{-1}(\lambda^\pm). \quad (3.4)$$

We have the simple inequality

$$|\mathfrak{S}| = \lambda^+ - \lambda^- = -\cos z^+ + \cos z^- = \int_{z^-}^{z^+} \sin t \, dt \leq z^+ - z^- = |\phi^{-1}(\mathfrak{S})|. \quad (3.5)$$

Thus, the first estimate in (3.1) has been proved.

We will show that

$$|\phi^{-1}(\mathfrak{S})| \leq |\phi^{-1}(\mathfrak{S}_*)|, \quad (3.6)$$

where $\mathfrak{S}_* \subset [-1, 1]$ is given by

$$\mathfrak{S}_* = [-1, -\lambda_*] \cup [\lambda_*, 1], \quad \lambda_* = 1 - \frac{1}{2}(\lambda^+ - \lambda^-) \in [0, 1]. \quad (3.7)$$

Note that from the definitions of $\mathfrak{S}$ and $\mathfrak{S}_*$ it follows that

$$|\mathfrak{S}_*| = |\mathfrak{S}|. \quad (3.8)$$

The preimage of the segment $\mathfrak{S}_*$ under the function ϕ has the form

$$\phi^{-1}(\mathfrak{S}_*) = [0, z_*] \cup [\pi - z_*, \pi], \quad \text{where } z_* = \phi^{-1}(-\lambda_*) \in [0, \frac{\pi}{2}]. \quad (3.9)$$

Using (3.4) and (3.9) we can rewrite the inequality (3.6) in the equivalent form

$$z^+ - z^- \leq 2z_*. \quad (3.10)$$

In order to prove (3.10) we use the definition of z_* in (3.9), (3.7) and consider the difference

$$\begin{aligned} \cos \frac{z^+ - z^-}{2} - \cos z_* &= \cos \frac{z^+ - z^-}{2} - 1 + \frac{1}{2}(\lambda^+ - \lambda^-) = \cos \frac{z^+ - z^-}{2} - 1 + \frac{1}{2}(-\cos z^+ + \cos z^-) \\ &= \cos \frac{z^+}{2} \cos \frac{z^-}{2} + \sin \frac{z^+}{2} \sin \frac{z^-}{2} - \cos^2 \frac{z^+}{2} - \sin^2 \frac{z^-}{2}. \end{aligned}$$

From this identity, using the simple inequalities

$$\cos \frac{z^-}{2} \geq \cos \frac{z^+}{2} \geq 0, \quad \sin \frac{z^+}{2} \geq \sin \frac{z^-}{2} \geq 0, \quad 0 \leq z^- \leq z^+ \leq \pi, \quad (3.11)$$

we obtain

$$\cos \frac{z^+ - z^-}{2} \geq \cos z_* \quad \Rightarrow \quad z_* \geq \frac{z^+ - z^-}{2}, \quad (3.12)$$

since $\frac{z^+ - z^-}{2}, z_* \in [0, \frac{\pi}{2}]$ and the function $\cos z$ decreases on this interval. Thus, (3.10) (and, consequently, (3.6)) has been proved. Using the estimate $\sin x \geq \frac{2\sqrt{2}}{\pi}x$, $x \in [0, \frac{\pi}{4}]$, we obtain that the Lebesgue measure of the set $\mathfrak{S}_*$, defined by (3.7), satisfies

$$|\mathfrak{S}_*| = 2(1 - \lambda_*) = 2(1 - \cos z_*) = 4 \sin^2 \frac{z_*}{2} \geq \frac{8}{\pi^2} z_*^2 = \frac{2}{\pi^2} |\phi^{-1}(\mathfrak{S}_*)|^2, \quad (3.13)$$

since $\frac{z_*}{2} \in [0, \frac{\pi}{4}]$.

Combining (3.6), (3.13) and (3.8), we obtain

$$|\phi^{-1}(\mathfrak{S})|^2 \leq |\phi^{-1}(\mathfrak{S}_*)|^2 = 4z_*^2 \leq \frac{\pi^2}{2} |\mathfrak{S}_*| = \frac{\pi^2}{2} |\mathfrak{S}|, \quad (3.14)$$

which yields the second estimate in (3.1).

ii) The preimage $\phi^{-1}(\mathfrak{S}_*)$ of the subset $\mathfrak{S}_*$ is given by (3.9). Since the derivative ϕ' of the function $\phi(z) = -\cos z$ increases from the point 0 to the point $\frac{\pi}{2}$ and $|\phi^{-1}([-1, -\lambda_*])| = |\phi^{-1}([\lambda_*, 1])|$, we obtain the first inequality in (3.3).

Identity (3.8) and estimate (3.13) imply the second inequality in (3.3). $\blacksquare$

We collect properties of the discrete Laplacian on periodic graphs from [KS14], which we need to prove our main results.

Theorem 3.2. *i) The Lebesgue measure $|\sigma(\Delta)|$ of the spectrum of the Laplace operator Δ satisfies*

$$|\sigma(\Delta)| \leq \sum_{n=1}^{\nu} |\sigma_n(\Delta)| \leq 2\beta, \quad (3.15)$$

where β is defined by (1.13).

ii) Let a fundamental graph Γ_f be bipartite. If ν is odd, then $\mu = 0$ is a flat band of the discrete Laplacian Δ on Γ .

Theorem 3.3. *i) Let Γ be a loop graph. Then the spectral bands $\sigma_n(\Delta) = [\lambda_n^-, \lambda_n^+]$ of the Laplace operator Δ satisfy*

$$\lambda_n^- = \lambda_n(0), \quad \forall n \in \mathbb{N}_\nu. \quad (3.16)$$

ii) Let Γ be a precise loop graph with a precise point $\vartheta_0 \in \mathbb{T}^d$. Then

$$\sigma_n(\Delta) = [\lambda_n^-, \lambda_n^+] = [\lambda_n(0), \lambda_n(\vartheta_0)], \quad \forall n \in \mathbb{N}_\nu, \quad (3.17)$$

$$\sum_{n=1}^{\nu} |\sigma_n(\Delta)| = 2\beta, \quad (3.18)$$

where β is defined by (1.13).

iii) Let in addition Γ be bipartite (Γ_f is not bipartite, since there is a loop on Γ_f). Then each spectral band of the Laplacian on Γ has the form $\sigma_n(\Delta) = [\lambda_n^-, \lambda_n^+]$, $n \in \mathbb{N}_\nu$, where $\lambda_n^\pm$ are the eigenvalues of the matrix $\mp \Delta(0)$.

Acknowledgments. Various parts of this paper were written during Evgeny Korotyaev's stay in the Mathematical Institute of Tsukuba University, Japan and Mittag-Leffler Institute, Sweden and Centre for Quantum Geometry of Moduli spaces (QGM), Aarhus University, Denmark. He is grateful to the institutes for the hospitality. His study was partly supported by The Ministry of education and science of Russian Federation, project 07.09.2012 No 8501 and the RFFI grant "Spectral and asymptotic methods for studying of the differential operators" No 11-01-00458 and the Danish National Research Foundation grant DNRF95 (Centre for Quantum Geometry of Moduli Spaces - QGM).

REFERENCES

- [BKS13] Badanin A.; Korotyaev, E.; Saburova, N. Laplacians on $\mathbb{Z}^2$ -periodic discrete graphs, preprint 2013.
- [B85] von Below, J. A characteristic equation associated to an eigenvalue problem on c^2 -networks, Linear Algebra Appl. 71 (1985), 309-325.
- [BKu12] Berkolaiko G.; Kuchment, P. Introduction to Quantum Graphs, Mathematical Surveys and Monographs, V. 186 AMS, 2012.
- [C97] Cattaneo, C. The spectrum of the continuous Laplacian on a graph, Monatsh. Math. 124 (1997), 215-235.
- [Ch97] Chung, F. Spectral graph theory, AMS, Providence, Rhode. Island, 1997.
- [HS04] Higuchi, Y.; Shirai, T. Some spectral and geometric properties for infinite graphs, AMS Contemp. Math. 347 (2004), 29-56.
- [K98] Korotyaev, E. Estimates of periodic potentials in terms of gap lengths. Comm. Math. Phys. 197 (1998), no. 3, 521-526.
- [K03] Korotyaev, E. Characterization of the spectrum of Schrödinger operators with periodic distributions. Int. Math. Res. Not. 2003, no. 37, 2019-2031.
- [KS13] Korotyaev, E.; Saburova, N. Schrödinger operators on periodic discrete graphs, preprint 2013.
- [KS14] Korotyaev, E.; Saburova, N. Spectral estimates for normalized Laplacian and its perturbations on periodic discrete graphs, preprint 2014.

- [LP08] Lledó, F.; Post, O. Eigenvalue bracketing for discrete and metric graphs, *J. Math. Anal. Appl.* 348 (2008), 806–833.
- [MW89] Mohar, B.; Woess, W. A survey on spectra of infinite graphs, *Bull. London Math. Soc.*, 21 (1989), 209–234.
- [P12] Post, O. Spectral analysis on graph-like spaces. *Lecture Notes in Mathematics*, 2039. Springer, Heidelberg, 2012.
- [S85] Skriganov, M. The spectrum band structure of three-dimensional Schrödinger operator with periodic potential, *Invent. Math.* 80 (1985), 107–121.

MATHEMATICAL PHYSICS DEPARTMENT, FACULTY OF PHYSICS, ULIANOVSKAYA 2, ST. PETERSBURG STATE UNIVERSITY, ST. PETERSBURG, 198904, KOROTYAEV@GMAIL.COM,

DEPARTMENT OF MATHEMATICAL ANALYSIS, ALGEBRA AND GEOMETRY, INSTITUTE OF MATHEMATICS, INFORMATION AND SPACE TECHNOLOGIES, URITSKOGO St. 68, NORTHERN (ARCTIC) FEDERAL UNIVERSITY, ARKHANGELSK, 163002, N.SABUROVA@GMAIL.COM